ORGANIC CHEMISTRY

Fifth Edition



JANICE GORZYNSKI SMITH

Mc Graw Hill Education

Organic Chemistry

Fifth Edition

Janice Gorzynski Smith

University of Hawai'i at Mānoa





ORGANIC CHEMISTRY, FIFTH EDITION

Published by McGraw-Hill Education, 2 Penn Plaza, New York, NY 10121. Copyright © 2017 by McGraw-Hill Education. All rights reserved. Printed in the United States of America. Previous editions © 2014, 2011, 2008, and 2006. No part of this publication may be reproduced or distributed in any form or by any means, or stored in a database or retrieval system, without the prior written consent of McGraw-Hill Education, including, but not limited to, in any network or other electronic storage or transmission, or broadcast for distance learning.

Some ancillaries, including electronic and print components, may not be available to customers outside the United States.

This book is printed on acid-free paper.

1 2 3 4 5 6 7 8 9 0 DOW/DOW 1 0 9 8 7 6

ISBN 978-0-07-802155-8
MHID 0-07-802155-3

Senior Vice President, Products & Markets: *Kurt L. Strand*
Vice President, General Manager, Products & Markets: *Marty Lange*
Vice President, Content Design & Delivery: *Kimberly Meriwether David*
Director of Development: *Rose Koos*
Managing Director: *Thomas Timp*
Director: *David Spurgeon, Ph.D.*
Brand Manager: *Andrea M. Pellerito, Ph.D.*
Product Developer: *Mary E. Hurley*
Director of Digital Content Development: *Justin Wyatt, Ph.D.*
Digital Product Analyst: *Patrick Diller*
Marketing Manager: *Matthew Garcia*
Director, Content Design & Delivery: *Linda Avenarius*
Program Manager: *Lora Neyens*
Content Project Manager: *Peggy J. Selle*
Assessment Content Project Manager: *Tammy Juran*
Buyer: *Sandy Ludovissy*
Designer: *Matthew Backhaus*
Content Licensing Specialist (Text): *DeAnna Dausener*
Content Licensing Specialist (Photo): *Carrie Burger*
Cover Image: *CDC/James Gathany*
Compositor: *Lachina Publishing*
Printer: *R.R. Donnelley*

All credits appearing on page or at the end of the book are considered to be an extension of the copyright page.

Library of Congress Cataloging-in-Publication Data

Smith, Janice G.
Organic chemistry / by Janice Gorzynski Smith. — 5th edition.
p. cm.
Includes index.
ISBN 978-0-07-802155-8 — ISBN 0-07-802155-8 (hard copy : alk. paper) 1. Chemistry, Organic—Textbooks. I. Title
QD253.2 .S63 2017
547—dc23

2015037323

The Internet addresses listed in the text were accurate at the time of publication. The inclusion of a website does not indicate an endorsement by the author or McGraw-Hill Education, and McGraw-Hill Education does not guarantee the accuracy of the information presented at these sites.

mheducation.com/highered

About the Author



Janice Gorzynski Smith was born in Schenectady, New York. She became interested in chemistry in high school and went on to major in chemistry at Cornell University, where she received an A.B. degree *summa cum laude*. Jan earned a Ph.D. in Organic Chemistry from Harvard University under the direction of Nobel Laureate E. J. Corey, and she also spent a year as a National Science Foundation National Needs Postdoctoral Fellow at Harvard. During her tenure with the Corey group, she completed the total synthesis of the plant growth hormone gibberellic acid.

Following her postdoctoral work, Jan joined the faculty of Mount Holyoke College, where she was employed for 21 years. During this time she was active in teaching organic chemistry lecture and lab courses, conducting a research program in organic synthesis, and serving as department chair. Her organic chemistry class was named one of Mount Holyoke's "Don't-miss courses" in a survey by *Boston* magazine. After spending two sabbaticals amidst the natural beauty and diversity in Hawai'i in the 1990s, Jan and her family moved there permanently in 2000. She is currently a faculty member at the University of Hawai'i at Mānoa, where she teaches the two-semester organic chemistry lecture and lab courses. In 2003, she received the Chancellor's Citation for Meritorious Teaching.

Jan resides in Hawai'i with her husband Dan, an emergency medicine physician, pictured with her hiking in New Zealand in 2015. She has four children and three grandchildren. When not teaching, writing, or enjoying her family, Jan bikes, hikes, snorkels, and scuba dives in sunny Hawai'i, and time permitting, enjoys travel and Hawaiian quilting.

For Megan Sarah

Contents in Brief


Prologue 1
1 Structure and Bonding 7
2 Acids and Bases 61
3 Introduction to Organic Molecules and Functional Groups 91
4 Alkanes 128
5 Stereochemistry 174
6 Understanding Organic Reactions 213
7 Alkyl Halides and Nucleophilic Substitution 247
8 Alkyl Halides and Elimination Reactions 297
9 Alcohols, Ethers, and Related Compounds 331
10 Alkenes 383
11 Alkynes 426
12 Oxidation and Reduction 455
13 Mass Spectrometry and Infrared Spectroscopy 495
14 Nuclear Magnetic Resonance Spectroscopy 527
15 Radical Reactions 570
16 Conjugation, Resonance, and Dienes 604
17 Benzene and Aromatic Compounds 641
18 Reactions of Aromatic Compounds 677
19 Carboxylic Acids and the Acidity of the O–H Bond 729
20 Introduction to Carbonyl Chemistry; Organometallic Reagents; Oxidation and Reduction 764
21 Aldehydes and Ketones—Nucleophilic Addition 817
22 Carboxylic Acids and Their Derivatives—Nucleophilic Acyl Substitution 868
23 Substitution Reactions of Carbonyl Compounds at the α Carbon 924
24 Carbonyl Condensation Reactions 962
25 Amines 996
26 Carbon–Carbon Bond-Forming Reactions in Organic Synthesis 1049
27 Pericyclic Reactions 1076
28 Carbohydrates 1106
29 Amino Acids and Proteins 1152
30 Synthetic Polymers 1198
31 Lipids 1231 (Available online)
Appendices A-1
Glossary G-1
Credits C-1
Index I-1

Contents

Preface xiii
Acknowledgments xxi
List of *How To's* xxiii
List of Mechanisms xxiv
List of Selected Applications xxvii


Prologue 1

What Is Organic Chemistry? 1
Some Representative Organic Molecules 2
Organic Chemistry and Malaria 4



1 Structure and Bonding 7

1.1 The Periodic Table 8
1.2 Bonding 11
1.3 Lewis Structures 13
1.4 Isomers 18
1.5 Exceptions to the Octet Rule 19
1.6 Resonance 19
1.7 Determining Molecular Shape 25
1.8 Drawing Organic Structures 30
1.9 Hybridization 36
1.10 Ethane, Ethylene, and Acetylene 40
1.11 Bond Length and Bond Strength 45
1.12 Electronegativity and Bond Polarity 47
1.13 Polarity of Molecules 49
1.14 L-Dopa—A Representative Organic Molecule 50
Key Concepts 52
Problems 53



2 Acids and Bases 61

2.1 Brønsted–Lowry Acids and Bases 62
2.2 Reactions of Brønsted–Lowry Acids and Bases 63
2.3 Acid Strength and pK_a 66
2.4 Predicting the Outcome of Acid–Base Reactions 68
2.5 Factors That Determine Acid Strength 70
2.6 Common Acids and Bases 78
2.7 Aspirin 80
2.8 Lewis Acids and Bases 81
Key Concepts 84
Problems 85




3 Introduction to Organic Molecules and Functional Groups 91

3.1 Functional Groups 92
3.2 An Overview of Functional Groups 93
3.3 Intermolecular Forces 99
3.4 Physical Properties 103
3.5 Application: Vitamins 109
3.6 Application of Solubility: Soap 111
3.7 Application: The Cell Membrane 113
3.8 Functional Groups and Reactivity 116
3.9 Biomolecules 117
Key Concepts 119
Problems 121



4 Alkanes 128

4.1 Alkanes—An Introduction 129
4.2 Cycloalkanes 132
4.3 An Introduction to Nomenclature 132
4.4 Naming Alkanes 133
4.5 Naming Cycloalkanes 138
4.6 Common Names 141
4.7 Fossil Fuels 141
4.8 Physical Properties of Alkanes 143
4.9 Conformations of Acyclic Alkanes—Ethane 144
4.10 Conformations of Butane 148
4.11 An Introduction to Cycloalkanes 151
4.12 Cyclohexane 152
4.13 Substituted Cycloalkanes 156
4.14 Oxidation of Alkanes 161
4.15 Lipids—Part 1 164
Key Concepts 166
Problems 167



5 Stereochemistry 174

5.1 Starch and Cellulose 175
5.2 The Two Major Classes of Isomers 177
5.3 Looking Glass Chemistry—Chiral and Achiral Molecules 178
5.4 Stereogenic Centers 181
5.5 Stereogenic Centers in Cyclic Compounds 183
5.6 Labeling Stereogenic Centers with *R* or *S* 185
5.7 Diastereomers 190
5.8 Meso Compounds 193
5.9 *R* and *S* Assignments in Compounds with Two or More Stereogenic Centers 194
5.10 Disubstituted Cycloalkanes 195
5.11 Isomers—A Summary 196
5.12 Physical Properties of Stereoisomers 197
5.13 Chemical Properties of Enantiomers 202
Key Concepts 204
Problems 205




6 Understanding Organic Reactions 213

6.1 Writing Equations for Organic Reactions 214
6.2 Kinds of Organic Reactions 215
6.3 Bond Breaking and Bond Making 217
6.4 Bond Dissociation Energy 221
6.5 Thermodynamics 225
6.6 Enthalpy and Entropy 227
6.7 Energy Diagrams 229
6.8 Energy Diagram for a Two-Step Reaction Mechanism 231
6.9 Kinetics 233
6.10 Catalysts 236
6.11 Enzymes 237
Key Concepts 239
Problems 240



7 Alkyl Halides and Nucleophilic Substitution 247

7.1 Introduction to Alkyl Halides 248
7.2 Nomenclature 249
7.3 Physical Properties 250
7.4 Interesting Alkyl Halides 251
7.5 The Polar Carbon–Halogen Bond 252
7.6 General Features of Nucleophilic Substitution 253
7.7 The Leaving Group 255
7.8 The Nucleophile 257
7.9 Possible Mechanisms for Nucleophilic Substitution 261
7.10 Two Mechanisms for Nucleophilic Substitution 262
7.11 The S_N2 Mechanism 263
7.12 The S_N1 Mechanism 269
7.13 Carbocation Stability 273
7.14 The Hammond Postulate 275
7.15 When Is the Mechanism S_N1 or S_N2? 278
7.16 Biological Nucleophilic Substitution 283
7.17 Vinyl Halides and Aryl Halides 286
7.18 Organic Synthesis 286
Key Concepts 288
Problems 290



8 Alkyl Halides and Elimination Reactions 297

8.1 General Features of Elimination 298
8.2 Alkenes—The Products of Elimination Reactions 299
8.3 The Mechanisms of Elimination 303
8.4 The E2 Mechanism 303
8.5 The Zaitsev Rule 308
8.6 The E1 Mechanism 310
8.7 S_N1 and E1 Reactions 314
8.8 Stereochemistry of the E2 Reaction 315
8.9 When Is the Mechanism E1 or E2? 319
8.10 E2 Reactions and Alkyne Synthesis 319
8.11 When Is the Reaction S_N1, S_N2, E1, or E2? 321
Key Concepts 325
Problems 326



9 Alcohols, Ethers, and Related Compounds 331


9.1 Introduction 332
9.2 Structure and Bonding 333
9.3 Nomenclature 334
9.4 Physical Properties 337

9.5 Interesting Alcohols, Ethers, and Epoxides 338
9.6 Preparation of Alcohols, Ethers, and Epoxides 341
9.7 General Features—Reactions of Alcohols, Ethers, and Epoxides 343
9.8 Dehydration of Alcohols to Alkenes 345
9.9 Carbocation Rearrangements 348
9.10 Dehydration Using $POCl_3$ and Pyridine 351
9.11 Conversion of Alcohols to Alkyl Halides with HX 352
9.12 Conversion of Alcohols to Alkyl Halides with $SOCl_2$ and PBr_3 356
9.13 Tosylate—Another Good Leaving Group 359
9.14 Reaction of Ethers with Strong Acid 362
9.15 Thiols and Sulfides 364
9.16 Reactions of Epoxides 367
9.17 Application: Epoxides, Leukotrienes, and Asthma 371
9.18 Application: Benzo[*a*]pyrene, Epoxides, and Cancer 373
Key Concepts 373
Problems 376



10 Alkenes 383

10.1 Introduction 384
10.2 Calculating Degrees of Unsaturation 385
10.3 Nomenclature 387
10.4 Physical Properties 391
10.5 Interesting Alkenes 391
10.6 Lipids—Part 2 393
10.7 Preparation of Alkenes 395
10.8 Introduction to Addition Reactions 396
10.9 Hydrohalogenation—Electrophilic Addition of HX 397
10.10 Markovnikov's Rule 400
10.11 Stereochemistry of Electrophilic Addition of HX 402
10.12 Hydration—Electrophilic Addition of Water 404
10.13 Halogenation—Addition of Halogen 405
10.14 Stereochemistry of Halogenation 406
10.15 Halohydrin Formation 408
10.16 Hydroboration–Oxidation 411
10.17 Keeping Track of Reactions 415
10.18 Alkenes in Organic Synthesis 417
Key Concepts 418
Problems 419



11 Alkynes 426

11.1 Introduction 427
11.2 Nomenclature 428
11.3 Physical Properties 429
11.4 Interesting Alkynes 430
11.5 Preparation of Alkynes 431
11.6 Introduction to Alkyne Reactions 432
11.7 Addition of Hydrogen Halides 434
11.8 Addition of Halogen 436
11.9 Addition of Water 437
11.10 Hydroboration–Oxidation 439
11.11 Reaction of Acetylide Anions 441
11.12 Synthesis 444
Key Concepts 447
Problems 448



12 Oxidation and Reduction 455


12.1 Introduction 456
12.2 Reducing Agents 457
12.3 Reduction of Alkenes 458
12.4 Application: Hydrogenation of Oils 461
12.5 Reduction of Alkynes 463
12.6 The Reduction of Polar C–X σ Bonds 466
12.7 Oxidizing Agents 467
12.8 Epoxidation 469
12.9 Dihydroxylation 472
12.10 Oxidative Cleavage of Alkenes 474
12.11 Oxidative Cleavage of Alkynes 476
12.12 Oxidation of Alcohols 476
12.13 Green Chemistry 479
12.14 Biological Oxidation 481
12.15 Sharpless Epoxidation 482
Key Concepts 485
Problems 487




13 Mass Spectrometry and Infrared Spectroscopy 495

13.1 Mass Spectrometry 496
13.2 Alkyl Halides and the M + 2 Peak 500
13.3 Fragmentation 501
13.4 Other Types of Mass Spectrometry 504

13.5 Electromagnetic Radiation 506
13.6 Infrared Spectroscopy 508
13.7 IR Absorptions 510
13.8 IR and Structure Determination 517
Key Concepts 519
Problems 520



14 Nuclear Magnetic Resonance Spectroscopy 527
14.1 An Introduction to NMR Spectroscopy 528
14.2 ^{1}H NMR: Number of Signals 531
14.3 ^{1}H NMR: Position of Signals 535
14.4 The Chemical Shift of Protons on sp^2 and sp Hybridized Carbons 539
14.5 ^{1}H NMR: Intensity of Signals 541
14.6 ^{1}H NMR: Spin–Spin Splitting 542
14.7 More Complex Examples of Splitting 546
14.8 Spin–Spin Splitting in Alkenes 549
14.9 Other Facts About ^{1}H NMR Spectroscopy 551
14.10 Using ^{1}H NMR to Identify an Unknown 554
14.11 ^{13}C NMR Spectroscopy 556
14.12 Magnetic Resonance Imaging (MRI) 561
Key Concepts 561
Problems 562




15 Radical Reactions 570
15.1 Introduction 571
15.2 General Features of Radical Reactions 572
15.3 Halogenation of Alkanes 574
15.4 The Mechanism of Halogenation 575
15.5 Chlorination of Other Alkanes 578
15.6 Chlorination Versus Bromination 578
15.7 Halogenation as a Tool in Organic Synthesis 581
15.8 The Stereochemistry of Halogenation Reactions 582
15.9 Application: The Ozone Layer and CFCs 584
15.10 Radical Halogenation at an Allylic Carbon 585
15.11 Application: Oxidation of Unsaturated Lipids 588
15.12 Application: Antioxidants 589
15.13 Radical Addition Reactions to Double Bonds 590
15.14 Polymers and Polymerization 593
Key Concepts 595
Problems 596



16 Conjugation, Resonance, and Dienes 604
16.1 Conjugation 605
16.2 Resonance and Allylic Carbocations 607
16.3 Common Examples of Resonance 608
16.4 The Resonance Hybrid 610
16.5 Electron Delocalization, Hybridization, and Geometry 612
16.6 Conjugated Dienes 613
16.7 Interesting Dienes and Polyenes 614
16.8 The Carbon–Carbon σ Bond Length in Buta-1,3-diene 614
16.9 Stability of Conjugated Dienes 615
16.10 Electrophilic Addition: 1,2- Versus 1,4-Addition 616
16.11 Kinetic Versus Thermodynamic Products 618
16.12 The Diels–Alder Reaction 621
16.13 Specific Rules Governing the Diels–Alder Reaction 623
16.14 Other Facts About the Diels–Alder Reaction 627
16.15 Conjugated Dienes and Ultraviolet Light 630
Key Concepts 632
Problems 634




17 Benzene and Aromatic Compounds 641
17.1 Background 642
17.2 The Structure of Benzene 643
17.3 Nomenclature of Benzene Derivatives 644
17.4 Spectroscopic Properties 647
17.5 Interesting Aromatic Compounds 648
17.6 Benzene's Unusual Stability 649
17.7 The Criteria for Aromaticity—Hückel's Rule 651
17.8 Examples of Aromatic Compounds 654
17.9 What Is the Basis of Hückel's Rule? 660
17.10 The Inscribed Polygon Method for Predicting Aromaticity 663
17.11 Buckminsterfullerene—Is It Aromatic? 666
Key Concepts 667
Problems 668




18 Reactions of Aromatic Compounds 677

18.1 Electrophilic Aromatic Substitution 678
18.2 The General Mechanism 679
18.3 Halogenation 681
18.4 Nitration and Sulfonation 682
18.5 Friedel–Crafts Alkylation and Friedel–Crafts Acylation 684
18.6 Substituted Benzenes 691
18.7 Electrophilic Aromatic Substitution of Substituted Benzenes 694
18.8 Why Substituents Activate or Deactivate a Benzene Ring 696
18.9 Orientation Effects in Substituted Benzenes 698
18.10 Limitations on Electrophilic Substitution Reactions with Substituted Benzenes 701
18.11 Disubstituted Benzenes 703
18.12 Synthesis of Benzene Derivatives 705
18.13 Nucleophilic Aromatic Substitution 706
18.14 Halogenation of Alkyl Benzenes 709
18.15 Oxidation and Reduction of Substituted Benzenes 711
18.16 Multistep Synthesis 715
Key Concepts 718
Problems 721




19 Carboxylic Acids and the Acidity of the O–H Bond 729

19.1 Structure and Bonding 730
19.2 Nomenclature 731
19.3 Physical Properties 734
19.4 Spectroscopic Properties 735
19.5 Interesting Carboxylic Acids 736
19.6 Aspirin, Arachidonic Acid, and Prostaglandins 737
19.7 Preparation of Carboxylic Acids 739
19.8 Reactions of Carboxylic Acids—General Features 740
19.9 Carboxylic Acids—Strong Organic Brønsted–Lowry Acids 741
19.10 Inductive Effects in Aliphatic Carboxylic Acids 744
19.11 Substituted Benzoic Acids 746
19.12 Extraction 749
19.13 Sulfonic Acids 751
19.14 Amino Acids 752
Key Concepts 755
Problems 756



20 Introduction to Carbonyl Chemistry; Organometallic Reagents; Oxidation and Reduction 764

20.1 Introduction 765
20.2 General Reactions of Carbonyl Compounds 766
20.3 A Preview of Oxidation and Reduction 769
20.4 Reduction of Aldehydes and Ketones 771
20.5 The Stereochemistry of Carbonyl Reduction 773
20.6 Enantioselective Carbonyl Reductions 774
20.7 Reduction of Carboxylic Acids and Their Derivatives 777
20.8 Oxidation of Aldehydes 782
20.9 Organometallic Reagents 782
20.10 Reaction of Organometallic Reagents with Aldehydes and Ketones 786
20.11 Retrosynthetic Analysis of Grignard Products 790
20.12 Protecting Groups 792
20.13 Reaction of Organometallic Reagents with Carboxylic Acid Derivatives 794
20.14 Reaction of Organometallic Reagents with Other Compounds 797
20.15 α,β-Unsaturated Carbonyl Compounds 799
20.16 Summary—The Reactions of Organometallic Reagents 802
20.17 Synthesis 802
Key Concepts 805
Problems 808



21 Aldehydes and Ketones—Nucleophilic Addition 817

21.1 Introduction 818
21.2 Nomenclature 819
21.3 Physical Properties 822
21.4 Spectroscopic Properties 823
21.5 Interesting Aldehydes and Ketones 825

21.6 Preparation of Aldehydes and Ketones 826
21.7 Reactions of Aldehydes and Ketones—General Considerations 828
21.8 Nucleophilic Addition of H^- and R^-—A Review 831
21.9 Nucleophilic Addition of ^-CN 833
21.10 The Wittig Reaction 835
21.11 Addition of 1° Amines 840
21.12 Addition of 2° Amines 844
21.13 Addition of H_2O—Hydration 845
21.14 Addition of Alcohols—Acetal Formation 849
21.15 Acetals as Protecting Groups 852
21.16 Cyclic Hemiacetals 854
21.17 An Introduction to Carbohydrates 857
Key Concepts 858
Problems 863



22 Carboxylic Acids and Their Derivatives—Nucleophilic Acyl Substitution 868

22.1 Introduction 869
22.2 Structure and Bonding 871
22.3 Nomenclature 873
22.4 Physical Properties 877
22.5 Spectroscopic Properties 878
22.6 Interesting Esters and Amides 880
22.7 Introduction to Nucleophilic Acyl Substitution 882
22.8 Reactions of Acid Chlorides 885
22.9 Reactions of Anhydrides 887
22.10 Reactions of Carboxylic Acids 889
22.11 Reactions of Esters 894
22.12 Application: Lipid Hydrolysis 896
22.13 Reactions of Amides 899
22.14 Application: The Mechanism of Action of β-Lactam Antibiotics 900
22.15 Summary of Nucleophilic Acyl Substitution Reactions 901
22.16 Natural and Synthetic Fibers 902
22.17 Biological Acylation Reactions 904
22.18 Nitriles 906
Key Concepts 911
Problems 914




23 Substitution Reactions of Carbonyl Compounds at the α Carbon 924

23.1 Introduction 925
23.2 Enols 926
23.3 Enolates 928
23.4 Enolates of Unsymmetrical Carbonyl Compounds 934
23.5 Racemization at the α Carbon 936
23.6 A Preview of Reactions at the α Carbon 937
23.7 Halogenation at the α Carbon 938
23.8 Direct Enolate Alkylation 942
23.9 Malonic Ester Synthesis 946
23.10 Acetoacetic Ester Synthesis 950
Key Concepts 953
Problems 955



24 Carbonyl Condensation Reactions 962


24.1 The Aldol Reaction 963
24.2 Crossed Aldol Reactions 967
24.3 Directed Aldol Reactions 971
24.4 Intramolecular Aldol Reactions 973
24.5 The Claisen Reaction 975
24.6 The Crossed Claisen and Related Reactions 977
24.7 The Dieckmann Reaction 979
24.8 The Michael Reaction 980
24.9 The Robinson Annulation 982
Key Concepts 986
Problems 987



25 Amines 996

25.1 Introduction 997
25.2 Structure and Bonding 997
25.3 Nomenclature 999
25.4 Physical Properties 1001
25.5 Spectroscopic Properties 1002
25.6 Interesting and Useful Amines 1004
25.7 Preparation of Amines 1007
25.8 Reactions of Amines—General Features 1014
25.9 Amines as Bases 1014

25.10 Relative Basicity of Amines and Other Compounds 1016
25.11 Amines as Nucleophiles 1022
25.12 Hofmann Elimination 1024
25.13 Reaction of Amines with Nitrous Acid 1027
25.14 Substitution Reactions of Aryl Diazonium Salts 1029
25.15 Coupling Reactions of Aryl Diazonium Salts 1034
25.16 Application: Synthetic Dyes and Sulfa Drugs 1036
Key Concepts 1038
Problems 1041




26 Carbon–Carbon Bond-Forming Reactions in Organic Synthesis 1049

26.1 Coupling Reactions of Organocuprate Reagents 1050
26.2 Suzuki Reaction 1052
26.3 Heck Reaction 1056
26.4 Carbenes and Cyclopropane Synthesis 1058
26.5 Simmons–Smith Reaction 1061
26.6 Metathesis 1062
Key Concepts 1067
Problems 1068



27 Pericyclic Reactions 1076

27.1 Types of Pericyclic Reactions 1077
27.2 Molecular Orbitals 1078
27.3 Electrocyclic Reactions 1080
27.4 Cycloaddition Reactions 1087
27.5 Sigmatropic Rearrangements 1091
27.6 Summary of Rules for Pericyclic Reactions 1097
Key Concepts 1098
Problems 1099



28 Carbohydrates 1106

28.1 Introduction 1107
28.2 Monosaccharides 1108
28.3 The Family of D-Aldoses 1113
28.4 The Family of D-Ketoses 1115
28.5 Physical Properties of Monosaccharides 1116
28.6 The Cyclic Forms of Monosaccharides 1116
28.7 Glycosides 1124
28.8 Reactions of Monosaccharides at the OH Groups 1127
28.9 Reactions at the Carbonyl Group—Oxidation and Reduction 1128
28.10 Reactions at the Carbonyl Group—Adding or Removing One Carbon Atom 1131
28.11 Disaccharides 1134
28.12 Polysaccharides 1138
28.13 Other Important Sugars and Their Derivatives 1140
Key Concepts 1144
Problems 1147



29 Amino Acids and Proteins 1152

29.1 Amino Acids 1153
29.2 Synthesis of Amino Acids 1156
29.3 Separation of Amino Acids 1159
29.4 Enantioselective Synthesis of Amino Acids 1163
29.5 Peptides 1164
29.6 Peptide Sequencing 1169
29.7 Peptide Synthesis 1172
29.8 Automated Peptide Synthesis 1177
29.9 Protein Structure 1179
29.10 Important Proteins 1186
Key Concepts 1189
Problems 1191



30 Synthetic Polymers 1198

30.1 Introduction 1199
30.2 Chain-Growth Polymers—Addition Polymers 1200
30.3 Anionic Polymerization of Epoxides 1207
30.4 Ziegler–Natta Catalysts and Polymer Stereochemistry 1208
30.5 Natural and Synthetic Rubbers 1210
30.6 Step-Growth Polymers—Condensation Polymers 1211
30.7 Polymer Structure and Properties 1216
30.8 Green Polymer Synthesis 1217
30.9 Polymer Recycling and Disposal 1220
Key Concepts 1223
Problems 1225



31 Lipids 1231 (Available online)

31.1 Introduction 1232
31.2 Waxes 1233
31.3 Triacylglycerols 1234
31.4 Phospholipids 1238
31.5 Fat-Soluble Vitamins 1241
31.6 Eicosanoids 1242
31.7 Terpenes 1245
31.8 Steroids 1250
Key Concepts 1255
Problems 1256

Appendix A pK_a Values for Selected Compounds A-1
Appendix B Nomenclature A-3
Appendix C Bond Dissociation Energies for Some Common Bonds [A–B → A• + •B] A-7
Appendix D Reactions That Form Carbon–Carbon Bonds A-8
Appendix E Characteristic IR Absorption Frequencies A-9
Appendix F Characteristic NMR Absorptions A-10
Appendix G General Types of Organic Reactions A-12
Appendix H How to Synthesize Particular Functional Groups A-14

Glossary G-1
Credits C-1
Index I-1

Preface

My goal in writing *Organic Chemistry* was to create a text that showed students the beauty and logic of organic chemistry by giving them a book that they would *use*. This text is based on lecture notes and handouts that were developed in my own organic chemistry courses over my 30-year teaching career. I have followed two guiding principles: use relevant and interesting applications to illustrate chemical phenomena, and present the material in a student-friendly fashion using bulleted lists, solved problems, and extensive illustrations and summaries. *Organic Chemistry* is my attempt to simplify and clarify a course that intimidates many students—to make organic chemistry interesting, relevant, and accessible to *all* students, both chemistry majors and those interested in pursuing careers in biology, medicine, and other disciplines, without sacrificing the rigor they need to be successful in the future.

The Basic Features

- **Style** This text is different—by design. Today's students rely more heavily on visual imagery to learn than ever before. The text uses less prose and more diagrams, equations, tables, and bulleted summaries to introduce and reinforce the major concepts and themes of organic chemistry.
- **Content** *Organic Chemistry* accents basic themes in an effort to keep memorization at a minimum. Relevant examples from everyday life are used to illustrate concepts, and this material is integrated throughout the chapter rather than confined to a boxed reading. Each topic is broken down into small chunks of information that are more manageable and easily learned. Sample problems are used as a tool to illustrate stepwise problem solving. Exceptions to the rule and older, less useful reactions are omitted to focus attention on the basic themes.
- **Organization** *Organic Chemistry* uses functional groups as the framework within which chemical reactions are discussed. Thus, the emphasis is placed on the reactions that different functional groups undergo, not on the reactions that prepare them. Moreover, similar reactions are grouped together, so that parallels can be emphasized. These include acid–base reactions (Chapter 2), oxidation and reduction (Chapters 12 and 20), radical reactions (Chapter 15), and reactions of organometallic reagents (Chapter 20).

By introducing one new concept at a time, keeping the basic themes in focus, and breaking complex problems down into small pieces, I have found that many students find organic chemistry an intense but learnable subject. Many, in fact, end the year-long course surprised that they have actually *enjoyed* their organic chemistry experience.

Organization and Presentation

For the most part, the overall order of topics in the text is consistent with the way most instructors currently teach organic chemistry. There are, however, some important differences in the way topics are presented to make the material logical and more accessible. This can especially be seen in the following areas.

- **Review material** Chapter 1 presents a healthy dose of review material covering Lewis structures, molecular geometry and hybridization, bond polarity, and types of bonding. While many of these topics are covered in general chemistry courses, they are presented here from an organic chemist's perspective. I have found that giving students a firm grasp of these fundamental concepts helps tremendously in their understanding of later material.

- **Acids and bases** Chapter 2 on acids and bases serves two purposes. It gives students experience with curved arrow notation using some familiar proton transfer reactions. It also illustrates how some fundamental concepts in organic structure affect a reaction, in this case an acid–base reaction. Since many mechanisms involve one or more acid–base reactions, I emphasize proton transfer reactions early and come back to this topic often throughout the text.
- **Functional groups** Chapter 3 uses the functional groups to introduce important properties of organic chemistry. Relevant examples—PCBs, vitamins, soap, and the cell membrane—illustrate fundamental solubility concepts. In this way, practical topics that are sometimes found in the last few chapters of an organic chemistry text (and thus often omitted because instructors run out of time) are introduced early, so that students can better grasp why they are studying the discipline.
- **Stereochemistry** Stereochemistry (the three-dimensional structure of molecules) is introduced early (Chapter 5) and reinforced often, so students have every opportunity to learn and understand a crucial concept in modern chemical research, drug design, and synthesis.
- **Modern reactions** While there is no shortage of new chemical reactions to present in an organic chemistry text, I have chosen to concentrate on new methods that introduce a particular three-dimensional arrangement in a molecule, so-called asymmetric or enantioselective reactions. Examples include Sharpless epoxidation (Chapter 12), CBS reduction (Chapter 20), and enantioselective synthesis of amino acids (Chapter 29).
- **Grouping reactions** Since certain types of reactions have their own unique characteristics and terminology that make them different from the basic organic reactions, I have grouped these reactions together in individual chapters. These include acid–base reactions (Chapter 2), oxidation and reduction (Chapters 12 and 20), radical reactions (Chapter 15), and reactions of organometallic reagents (Chapter 20). I have found that focusing on a group of reactions that share a common theme helps students to better see their similarities.
- **Synthesis** Synthesis, one of the most difficult topics for a beginning organic student to master, is introduced in small doses, beginning in Chapter 7 and augmented with a detailed discussion of retrosynthetic analysis in Chapter 11. In later chapters, special attention is given to the retrosynthetic analysis of compounds prepared by carbon–carbon bond-forming reactions (for example, Sections 20.11 and 21.10C).
- **Spectroscopy** Since spectroscopy is such a powerful tool for structure determination, four methods are discussed over two chapters (Chapters 13 and 14).
- **Key Concepts** End-of-chapter summaries succinctly summarize the main concepts and themes of the chapter, making them ideal for review prior to working the end-of-chapter problems or taking an exam.

New to this Edition


- Chemical structures were updated throughout the text for a more modern and consistent look.
- Color has also been used in many areas to help students better understand three-dimensional structure, stereochemistry, and reactions.
- All nomenclature has been updated in accord with newer IUPAC nomenclature recommendations and the 1993 nomenclature rules.
- The design of the mechanism boxes has been revised, so that students can more readily see how one intermediate is converted to another.
- In response to reviewer feedback, new material has been added to several chapters. Topics include a section on biological nucleophilic substitution with phosphorus leaving groups (Section 7.16) and a section on thiols and sulfides (Section 9.15). The section on biological oxidation was revised to include the oxidizing agent NAD^+, with new structures in the mechanism of oxidation of an alcohol, resulting in a more biological flavor to this material (Section 12.14). A new section on biological reactions with allylic diphosphates and a new mechanism on biological reactions with allylic diphosphates have been added to Section 16.2. New material on biological reduction appears in Section 20.6, and the discussion of ultraviolet spectroscopy has been expanded in Section 16.15.

- Material on classifying carbons, hydrogens, alcohols, alkyl halides, amines, and amides was moved from later chapters to earlier in the text (Section 3.2), so that it is included in the discussion of functional groups.
- Over 350 new problems have been added to the new edition, increasing the variety of problems for instructors and students alike.
- The chapter on lipids now appears online and is available in customizable versions of the text in McGraw-Hill Create.
- An online supplement covering imine derivatives is also available on the Online Learning Center's Instructor Resources, via the Library tab in Connect.
- New *How To*'s, Sample Problems, and micro-to-macro illustrations have also been added throughout the new edition to clarify topics and enhance the student learning experience.

Tools to Make Learning Organic Chemistry Easier


Illustrations

Organic Chemistry is supported by a well-developed illustration program. Besides traditional skeletal (line) structures and condensed formulas, there are numerous ball-and-stick molecular models and electrostatic potential maps to help students grasp the three-dimensional structure of molecules (including stereochemistry) and to better understand the distribution of electronic charge.




Micro-to-Macro Illustrations

Unique to *Organic Chemistry* are micro-to-macro illustrations, where line art and photos combine with chemical structures to reveal the underlying molecular structures giving rise to macroscopic properties of common phenomena. Examples include starch and cellulose (Chapter 5), adrenaline (Chapter 7), partial hydrogenation of vegetable oil (Chapter 12), and dopamine (Chapter 25).




Spectra

Over 100 spectra created specifically for *Organic Chemistry* are presented throughout the text. The spectra are color-coded by type and generously labeled. Mass spectra are green; infrared spectra are red; and proton and carbon nuclear magnetic resonance spectra are blue.



Mechanisms


Curved arrow notation is used extensively to help students follow the movement of electrons in reactions.



Problem Solving


Sample Problems

Sample Problems show students how to solve organic chemistry problems in a logical, stepwise manner. More than 800 follow-up problems are located throughout the chapters to test whether students understand concepts covered in the Sample Problems.



How To's


How To's provide students with detailed instructions on how to work through key processes.



Applications and Summaries

Key Concept Summaries

Succinct summary tables reinforcing important principles and concepts are provided at the end of each chapter.



Margin Notes

Margin notes are placed carefully throughout the chapters, providing interesting information relating to topics covered in the text. Some margin notes are illustrated with photos to make the chemistry more relevant.



All soaps are salts of fatty acids. The main difference between soaps is the addition of other ingredients that do not alter their cleaning properties: dyes for color, scents for a pleasing odor, and oils for lubrication. Soaps that float are aerated, so that they are less dense than water.

Mc Graw Hill Education

connect®


Required=Results



McGraw-Hill Connect®
Learn Without Limits

Connect is a teaching and learning platform that is proven to deliver better results for students and instructors.


Connect empowers students by continually adapting to deliver precisely what they need, when they need it, and how they need it, so your class time is more engaging and effective.



Using **Connect** improves passing rates by **10.8%** and retention by **16.4%.**

88% of instructors who use **Connect** require it; instructor satisfaction **increases** by 38% when **Connect** is required.

Analytics



Connect Insight®

Connect Insight is Connect's new one-of-a-kind visual analytics dashboard—now available for both instructors and students—that provides at-a-glance information regarding student performance, which is immediately actionable. By presenting assignment, assessment, and topical performance results together with a time metric that is easily visible for aggregate or individual results, Connect Insight gives the user the ability to take a just-in-time approach to teaching and learning, which was never before available. Connect Insight presents data that empowers students and helps instructors improve class performance in a way that is efficient and effective.


Students can view their results for any **Connect** course.

Mobile

Connect's new, intuitive mobile interface gives students and instructors flexible and convenient, anytime–anywhere access to all components of the Connect platform.



Adaptive




THE FIRST AND ONLY **ADAPTIVE READING EXPERIENCE** DESIGNED TO TRANSFORM THE WAY STUDENTS READ

More students earn **A's** and **B's** when they use McGraw-Hill Education **Adaptive** products.

SmartBook®

Proven to help students improve grades and study more efficiently, SmartBook contains the same content within the print book, but actively tailors that content to the needs of the individual. SmartBook's adaptive technology provides precise, personalized instruction on what the student should do next, guiding the student to master and remember key concepts, targeting gaps in knowledge and offering customized feedback, and driving the student toward comprehension and retention of the subject matter. Available on smartphones and tablets, SmartBook puts learning at the student's fingertips—anywhere, anytime.

Over **4 billion questions** have been answered, making McGraw-Hill Education products more intelligent, reliable, and precise.



Learning Resources for Instructors and Students

The following items may accompany this text. Please consult your McGraw-Hill representative for policies, prices, and availability as some restrictions may apply.

Presentation Tools

Within the Instructor's Presentation Tools, instructors have access to editable PowerPoint lecture outlines, which appear as ready-made presentations that combine art and lecture notes for each chapter of the text. For instructors who prefer to create their lecture notes from scratch, all illustrations, photos, tables, *How To*'s, and Sample Problems are pre-inserted by chapter into a separate set of PowerPoint slides. They are also available as individual .jpg files.

An online digital library contains photos, artwork, animations, and other media types that can be used to create customized lectures, visually enhanced tests and quizzes, compelling course websites, or attractive printed support materials. All assets are copyrighted by McGraw-Hill Higher Education, but can be used by instructors for classroom purposes. The visual resources in this collection include:

- **Art** Full-color digital files of all illustrations in the book can be readily incorporated into lecture presentations, exams, or custom-made classroom materials.
- **Photos** The photo collection contains digital files of photographs from the text, which can be reproduced for multiple classroom uses.
- **Tables** Every table that appears in the text has been saved in electronic form for use in classroom presentations and/or quizzes.
- **Animations** Numerous full-color animations illustrating important processes are also provided. Harness the visual impact of concepts in motion by importing these files into classroom presentations or online course materials.

Student Study Guide/Solutions Manual

Written by Janice Gorzynski Smith and Erin R. Smith, the Student Study Guide/Solutions Manual provides step-by-step solutions to all in-chapter and end-of-chapter problems. Each chapter begins with an overview of key concepts and includes a short-answer practice test on the fundamental principles and new reactions.

Acknowledgments

When I started working on the first edition of *Organic Chemistry* in the fall of 1999, I had no sense of the magnitude of the task, or any idea of just how many people I would rely upon to complete it. Fortunately, I have had the steadfast support of a dedicated team of publishing professionals at McGraw-Hill.

I am especially thankful for the opportunity to work with Senior Product Developer Mary Hurley, who skillfully and efficiently guided me through the process of updating this fifth edition. Mary has been my rock through the many months of re-drawing chemical structures and re-designing mechanisms and art. I am grateful to once again work with Lead Content Project Manager Peggy Selle, who managed the production of this updated and re-designed text. *Organic Chemistry* has also benefited greatly from the expertise and market-based feedback provided by Marketing Manager Matthew Garcia.

Special thanks go out to Brand Manager Andrea Pellerito, who gave me the day-to-day editorial support crucial in producing a revision of *Organic Chemistry.* Thanks also to Managing Director Thomas Timp, who efficiently directed the editorial team that produced this revision. I also appreciate the work of Matt Backhaus (Designer) and Carrie Burger (Photo Researcher) who are responsible for the visually pleasing appearance of this edition. Thanks are again due to freelance Developmental Editor John Murdzek for his meticulous editing and humorous insights on my project.

My immediate family has experienced the day-to-day demands of living with a busy author. Thanks go to my husband Dan, my children Erin, Jenna, Matthew, and Zachary, and my grandchildren Max, Koa, and Alijah, all of whom keep me grounded during the time-consuming process of writing and publishing a textbook.

Among the many others that go unnamed but who have profoundly affected this work are the thousands of students I have been lucky to teach over the last 30 years. I have learned so much from my daily interactions with them, and I hope that the wider chemistry community can benefit from this experience by the way I have presented the material in this text.

This fifth edition has evolved based on the helpful feedback of many people who reviewed the fourth edition text and digital products, class-tested the book, and attended focus groups or symposiums. These many individuals have collectively provided constructive improvements to the project.

Listed below are the reviewers of the fourth edition text:

Steven Castle, *Brigham Young University*
Ihsan Erden, *San Francisco State University*
Andrew Frazer, *University of Central Florida, Orlando*
Tiffany Gierasch, *University of Maryland, Baltimore County*
Anne Gorden, *Auburn University*
Michael Lewis, *Saint Louis University*
Eugene A. Mash, Jr., *University of Arizona*
Mark McMills, *Ohio University*
Joan Mutanyatta–Comar, *Georgia State University*
Felix Ngassa, *Grand Valley State University*
Michael Rathke, *Michigan State University*
Jacob Schroeder, *Clemson University*
Keith Schwartz, *Portland State University*
John Selegue, *University of Kentucky*
Paul J. Toscano, *University at Albany, SUNY*
Jane E. Wissinger, *University of Minnesota*

The following contributed to the editorial direction of the fifth edition text by responding to our survey on the MCAT and the organic chemistry course student population:

Chris Abelt, *College of William and Mary*
Orlando Acevedo, *Auburn University*
Kim Albizati, *University of California, San Diego*
Merritt Andrus, *Brigham Young University*
Ardeshir Azadnia, *Michigan State University*
Susan Bane, *Binghamton University*
Russell Barrows, *Metropolitan State University of Denver*
Peter Beak, *University of Illinois, Urbana Champaign*
Phil Beauchamp, *Cal Poly, Pomona*
Michael Berg, *Virginia Tech*
K. Darrell Berlin, *Oklahoma State University*
Thomas Bertolini, *University of South Carolina*
Ned Bowden, *University of Iowa*
David W. Brown, *Florida Gulf Coast University*
Rebecca Broyer, *University of Southern California*
Arthur Bull, *Oakland University*
K. Nolan Carter, *University of Central Arkansas*
Steven Castle, *Brigham Young University*
Victor Cesare, *St. John's University*
Manashi Chatterjee, *University of Nebraska, Lincoln*
Melissa Cichowicz, *West Chester University*
Jeff Corkill, *Eastern Washington University, Cheney*
Sulekha Coticone, *Florida Gulf Coast University*
Michael Crimmins, *University of North Carolina at Chapel Hill*
Eric Crumpler, *Valencia College*
David Dalton, *Temple University*
Rick Danheiser, *Massachusetts Institute of Technology*
Tammy Davidson, *University of Florida*
Brenton DeBoef, *University of Rhode Island*
Amy Deveau, *University of New England*
Kenneth M. Doxsee, *University of Oregon*

Larissa D'Souza, *Johns Hopkins University*
Philip Egan, *Texas A&M University, Corpus Christi*
Seth Elsheimer, *University of Central Florida*
John Esteb, *Butler University*
Steve Fleming, *Temple University*
Marion Franks, *North Carolina A&T State University*
Andy Frazer, *University of Central Florida*
Brian Ganley, *University of Missouri, Columbia*
Robert Giuliano, *Villanova University*
Anne Gorden, *Auburn University*
Carlos G. Gutierrez, *California State University, Los Angeles*
Scott Handy, *Middle Tennessee State University*
Rick Heldrich, *College of Charleston*
James Herndon, *New Mexico State University*
Kathleen Hess, *Brown University*
Sean Hickey, *University of New Orleans*
Carl Hoeger, *University of California, San Diego*
Javier Horta, *University of Massachusetts, Lowell*
Bob A. Howell, *Central Michigan University*
Jennifer Irvin, *Texas State University*
Phil Janowicz, *Cal State, Fullerton*
Mohamad Karim, *Tennessee State University*
Mark L. Kearley, *Florida State University*
Amy Keirstead, *University of New England*
Margaret Kerr, *Worcester State University*
James Kiddle, *Western Michigan University*
Jisook Kim, *University of Tennessee at Chattanooga*
Angela King, *Wake Forest University*
Margaret Kline, *Santa Monica College*
Dalila G. Kovacs, *Grand Valley State University*
Deborah Lieberman, *University of Cincinnati*
Carl Lovely, *University of Texas, Arlington*
Kristina Mack, *Grand Valley State University*
Daniel Macks, *Towson University*
Vivian Mativo, *Georgia Perimeter College, Clarkston*
Mark McMills, *Ohio University*
Stephen Mills, *Xavier University*
Robert Minto, *Indiana University–Purdue University, Indianapolis*
Debbie Mohler, *James Madison University*
Kathleen Morgan, *Xavier University of Louisiana*
Paul Morgan, *Butler University*
James C. Morris, *Georgia Institute of Technology*
Linda Munchausen, *Southeastern Louisiana University*
Toby Nelson, *Oklahoma State University*
Felix Ngassa, *Grand Valley State University*
George A. O'Doherty, *Northeastern University*
Anne Padias, *University of Arizona*
Dan Paschal, *Georgia Perimeter College*
Richard Pennington, *Georgia Gwinnett College*
John Pollard, *University of Arizona*
Gloria Proni, *John Jay College*
Khalilah Reddie, *University of Massachusetts, Lowell*
Joel M. Ressner, *West Chester University of Pennsylvania*
Christine Rich, *University of Louisville*
Carmelo Rizzo, *Vanderbilt University*
Harold R. Rogers, *California State University, Fullerton*
Paul B. Savage, *Brigham Young University*
Deborah Schwyter, *Santa Monica College*
Holly Sebahar, *University of Utah*
Laura Serbulea, *University of Virginia*
Abid Shaikh, *Georgia Southern University*
Kevin Shaughnessy, *The University of Alabama*
Joel Shulman, *University of Cincinnati*
Joseph M. Simard, *University of New England*
Rhett Smith, *Clemson University*
Priyantha Sugathapala, *University at Albany, SUNY*
Claudia Taenzler, *University of Texas at Dallas*
Robin Tanke, *University of Wisconsin, Stevens Point*
Richard T. Taylor, *Miami University, Oxford*
Edward Turos, *University of South Florida*
Ted Wood, *Pierce College*
Kana Yamamoto, *University of Toledo*

The following individuals helped write and review learning goal-oriented content for **LearnSmart for Organic Chemistry**: David G. Jones, Vistamar School; Adam I. Keller, Columbus State Community College; and Parul D. Root, Henry Ford Community College. Harpreet Malhotra of Florida State College at Jacskonville reviewed the Connect content for accuracy, and Ujjwal Chakraborty, also of Florida State College at Jacksonville, revised the PowerPoint Lectures and Test Bank for the fifth edition.

Although every effort has been made to make this text and its accompanying Student Study Guide/Solutions Manual as error-free as possible, some errors undoubtedly remain and, for them, I am solely responsible. Please feel free to email me about any inaccuracies, so that subsequent editions may be further improved.

With much aloha,

Janice Gorzynski Smith
jgsmith@hawaii.edu

List of *How To's*

How To boxes provide detailed instructions for key procedures that students need to master. Below is a list of each *How To* and where it is presented in the text.

Chapter 1 Structure and Bonding
How To Draw a Lewis Structure 14
How To Interpret a Skeletal Structure 33

Chapter 2 Acids and Bases
How To Determine Relative Acidity of Protons 77

Chapter 4 Alkanes
How To Name an Alkane Using the IUPAC System 135
How To Name a Cycloalkane Using the IUPAC System 139
How To Draw a Newman Projection 145
How To Draw the Chair Form of Cyclohexane 154
How To Draw the Two Conformations for a Substituted Cyclohexane 156
How To Draw Two Conformations for a Disubstituted Cyclohexane 159

Chapter 5 Stereochemistry
How To Assign *R* or *S* to a Stereogenic Center 187
How To Find and Draw All Possible Stereoisomers for a Compound with Two Stereogenic Centers 191

Chapter 7 Alkyl Halides and Nucleophilic Substitution
How To Name an Alkyl Halide Using the IUPAC System 249

Chapter 9 Alcohols, Ethers, and Related Compounds
How To Name an Alcohol Using the IUPAC System 334

Chapter 10 Alkenes
How To Name an Alkene 387
How To Assign the Prefixes *E* and *Z* to an Alkene 389

Chapter 11 Alkynes
How To Develop a Retrosynthetic Analysis 445

Chapter 13 Mass Spectrometry and Infrared Spectroscopy
How To Use MS and IR for Structure Determination 518

Chapter 14 Nuclear Magnetic Resonance Spectroscopy
How To Use ^{1}H NMR Data to Determine a Structure 554

Chapter 16 Conjugation, Resonance, and Dienes
How To Draw the Product of a Diels–Alder Reaction 622

Chapter 17 Benzene and Aromatic Compounds
How To Use the Inscribed Polygon Method to Determine the Relative Energies of MOs for Cyclic, Completely Conjugated Compounds 664

Chapter 18 Reactions of Aromatic Compounds
How To Determine the Directing Effects of a Particular Substituent 698

Chapter 21 Aldehydes and Ketones—Nucleophilic Addition
How To Determine the Starting Materials for a Wittig Reaction Using Retrosynthetic Analysis 838

Chapter 22 Carboxylic Acids and Their Derivatives—Nucleophilic Acyl Substitution
How To Name an Ester (RCO_2R') Using the IUPAC System 874
How To Name a 2° or 3° Amide 874

Chapter 24 Carbonyl Condensation Reactions
How To Synthesize a Compound Using the Aldol Reaction 967
How To Synthesize a Compound Using the Robinson Annulation 985

Chapter 25 Amines
How To Name 2° and 3° Amines with Different Alkyl Groups 999

Chapter 28 Carbohydrates
How To Draw a Haworth Projection from an Acyclic Aldohexose 1119

Chapter 29 Amino Acids and Proteins
How To Use (*R*)-α-Methylbenzylamine to Resolve a Racemic Mixture of Amino Acids 1161
How To Synthesize a Dipeptide from Two Amino Acids 1173
How To Synthesize a Peptide Using the Merrifield Solid Phase Technique 1178

List of Mechanisms

Mechanisms are the key to understanding the reactions of organic chemistry. For this reason, great care has been given to present mechanisms in a detailed, step-by-step fashion. The list below indicates when each mechanism in the text is presented for the first time.

Chapter 7 Alkyl Halides and Nucleophilic Substitution

7.1 The S_N2 Mechanism 264
7.2 The S_N1 Mechanism 269

Chapter 8 Alkyl Halides and Elimination Reactions

8.1 The E2 Mechanism 304
8.2 The E1 Mechanism 310

Chapter 9 Alcohols, Ethers, and Related Compounds

9.1 Dehydration of 2° and 3° ROH—An E1 Mechanism 346
9.2 Dehydration of a 1° ROH—An E2 Mechanism 347
9.3 A 1,2-Methyl Shift—Carbocation Rearrangement During Dehydration 349
9.4 Dehydration Using $POCl_3$ + Pyridine—An E2 Mechanism 351
9.5 Reaction of a 1° ROH with HX—An S_N2 Mechanism 353
9.6 Reaction of 2° and 3° ROH with HX—An S_N1 Mechanism 354
9.7 Reaction of ROH with $SOCl_2$ + Pyridine—An S_N2 Mechanism 356
9.8 Reaction of ROH with PBr_3—An S_N2 Mechanism 357
9.9 Mechanism of Ether Cleavage in Strong Acid—$(CH_3)_3COCH_3 + HI \rightarrow (CH_3)_3CI + CH_3I + H_2O$ 363

Chapter 10 Alkenes

10.1 Electrophilic Addition of HX to an Alkene 399
10.2 Electrophilic Addition of H_2O to an Alkene—Hydration 404
10.3 Addition of X_2 to an Alkene—Halogenation 406
10.4 Addition of X and OH—Halohydrin Formation 408
10.5 Addition of H and BH_2—Hydroboration 411

Chapter 11 Alkynes

11.1 Electrophilic Addition of HX to an Alkyne 435
11.2 Addition of X_2 to an Alkyne—Halogenation 436
11.3 Tautomerization in Acid 438
11.4 Hydration of an Alkyne 438

Chapter 12 Oxidation and Reduction

12.1 Addition of H_2 to an Alkene—Hydrogenation 459
12.2 Dissolving Metal Reduction of an Alkyne to a Trans Alkene 465
12.3 Reduction of RX with $LiAlH_4$ 467
12.4 Epoxidation of an Alkene with a Peroxyacid 469
12.5 Oxidation of an Alcohol with CrO_3 478
12.6 Oxidation of a 1° Alcohol to a Carboxylic Acid 478

Chapter 15 Radical Reactions

15.1 Radical Halogenation of Alkanes 576
15.2 Allylic Bromination with NBS 586
15.3 Radical Addition of HBr to an Alkene 591
15.4 Radical Polymerization of CH_2=CHZ 595

Chapter 16 Conjugation, Resonance, and Dienes

16.1 Biological Formation of Geranyl Diphosphate 608
16.2 Electrophilic Addition of HBr to a 1,3-Diene—1,2- and 1,4-Addition 617

Chapter 18 Reactions of Aromatic Compounds

18.1 General Mechanism—Electrophilic Aromatic Substitution 679
18.2 Bromination of Benzene 681
18.3 Formation of the Nitronium Ion ($^+NO_2$) for Nitration 682
18.4 Formation of the Electrophile $^+SO_3H$ for Sulfonation 683

18.5 Formation of the Electrophile in Friedel–Crafts Alkylation—Two Possibilities 685
18.6 Friedel–Crafts Alkylation Using a 3° Carbocation 685
18.7 Formation of the Electrophile in Friedel–Crafts Acylation 686
18.8 Friedel–Crafts Alkylation Involving Carbocation Rearrangement 687
18.9 A Rearrangement Reaction Beginning with a 1° Alkyl Chloride 688
18.10 Nucleophilic Aromatic Substitution by Addition–Elimination 707
18.11 Nucleophilic Aromatic Substitution by Elimination–Addition: Benzyne 708
18.12 Benzylic Bromination 710

Chapter 20 Introduction to Carbonyl Chemistry; Organometallic Reagents; Oxidation and Reduction

20.1 Nucleophilic Addition—A Two-Step Process 767
20.2 Nucleophilic Substitution—A Two-Step Process 768
20.3 $LiAlH_4$ Reduction of RCHO and $R_2C=O$ 772
20.4 Reduction of RCOCl and RCOOR' with a Metal Hydride Reagent 778
20.5 Reduction of an Amide to an Amine with $LiAlH_4$ 780
20.6 Nucleophilic Addition of R"MgX to RCHO and $R_2C=O$ 786
20.7 Reaction of R"MgX or R"Li with RCOCl and RCOOR' 795
20.8 Carboxylation—Reaction of RMgX with CO_2 798
20.9 1,2-Addition to an α,β-Unsaturated Carbonyl Compound 800
20.10 1,4-Addition to an α,β-Unsaturated Carbonyl Compound 800

Chapter 21 Aldehydes and Ketones—Nucleophilic Addition

21.1 General Mechanism—Nucleophilic Addition 829
21.2 General Mechanism—Acid-Catalyzed Nucleophilic Addition 829
21.3 Nucleophilic Addition of ^-CN—Cyanohydrin Formation 833
21.4 The Wittig Reaction 837
21.5 Imine Formation from an Aldehyde or Ketone 841
21.6 Enamine Formation from an Aldehyde or Ketone 843
21.7 Base-Catalyzed Addition of H_2O to a Carbonyl Group 846
21.8 Acid-Catalyzed Addition of H_2O to a Carbonyl Group 846
21.9 Acetal Formation 849
21.10 Acid-Catalyzed Cyclic Hemiacetal Formation 853
21.11 A Cyclic Acetal from a Cyclic Hemiacetal 854

Chapter 22 Carboxylic Acids and Their Derivatives—Nucleophilic Acyl Substitution

22.1 General Mechanism—Nucleophilic Acyl Substitution 882
22.2 Conversion of Acid Chlorides to Anhydrides 886
22.3 Conversion of Acid Chlorides to Carboxylic Acids 887
22.4 Conversion of an Anhydride to an Amide 888
22.5 Conversion of Carboxylic Acids to Acid Chlorides 890
22.6 Fischer Esterification—Acid-Catalyzed Conversion of Carboxylic Acids to Esters 891
22.7 Conversion of Carboxylic Acids to Amides with DCC 893
22.8 Acid-Catalyzed Hydrolysis of an Ester to a Carboxylic Acid 895
22.9 Base-Promoted Hydrolysis of an Ester to a Carboxylic Acid 895
22.10 Amide Hydrolysis in Base 899
22.11 Hydrolysis of a Nitrile in Base 908
22.12 Reduction of a Nitrile with $LiAlH_4$ 909
22.13 Reduction of a Nitrile with DIBAL-H 910
22.14 Addition of Grignard and Organolithium Reagents (R–M) to Nitriles 910

Chapter 23 Substitution Reactions of Carbonyl Compounds at the α Carbon

23.1 Tautomerization in Acid 927
23.2 Tautomerization in Base 927
23.3 Acid-Catalyzed Halogenation at the α Carbon 938
23.4 Halogenation at the α Carbon in Base 939
23.5 The Haloform Reaction 940

Chapter 24 Carbonyl Condensation Reactions

24.1 The Aldol Reaction 964
24.2 Dehydration of β-Hydroxy Carbonyl Compounds with Base 966
24.3 The Intramolecular Aldol Reaction 974
24.4 The Claisen Reaction 976

24.5 The Dieckmann Reaction 980
24.6 The Michael Reaction 981
24.7 The Robinson Annulation 985

Chapter 25 Amines
25.1 The E2 Mechanism for the Hofmann Elimination 1025
25.2 Formation of a Diazonium Salt from a 1° Amine 1028
25.3 Formation of an *N*-Nitrosamine from a 2° Amine 1029
25.4 Azo Coupling 1034

Chapter 26 Carbon–Carbon Bond-Forming Reactions in Organic Synthesis
26.1 Suzuki Reaction 1055
26.2 Heck Reaction 1058
26.3 Formation of Dichlorocarbene 1059
26.4 Addition of Dichlorocarbene to an Alkene 1060
26.5 Simmons–Smith Reaction 1062
26.6 Olefin Metathesis: 2 $RCH{=}CH_2 \rightarrow RCH{=}CHR + CH_2{=}CH_2$ 1064

Chapter 28 Carbohydrates
28.1 Glycoside Formation 1125
28.2 Glycoside Hydrolysis 1126

Chapter 29 Amino Acids and Proteins
29.1 Formation of an α-Amino Nitrile 1159
29.2 Edman Degradation 1170

Chapter 30 Synthetic Polymers
30.1 Radical Polymerization of $CH_2{=}CHPh$ 1201
30.2 Forming Branched Polyethylene During Radical Polymerization 1203
30.3 Cationic Polymerization of $CH_2{=}CHZ$ 1204
30.4 Anionic Polymerization of $CH_2{=}CHZ$ 1206
30.5 Ziegler–Natta Polymerization of $CH_2{=}CH_2$ 1209

Chapter 31 Lipids (Available online)
31.1 Biological Formation of Farnesyl Diphosphate 1248
31.2 Isomerization of Geranyl Diphosphate to Neryl Diphosphate 1249

List of Selected Applications

Applications make any subject seem more relevant and interesting—for nonmajors and majors alike. The following is a list of the biological, medicinal, and environmental applications that have been integrated throughout *Organic Chemistry.* Each chapter opener showcases an interesting and current application relating to the chapter's topic. (Code: G = general; M = medicinal; B = biological; E = environmental)

Prologue

- G Methane, the main component of natural gas
- G Ethanol, the alcohol in beverages
- E Trichlorofluoromethane, a CFC responsible for destroying the stratospheric ozone layer
- M Amoxicillin, a widely used antibiotic
- M Fluoxetine, the antidepressant Prozac
- M AZT, a drug used to treat HIV
- M Capsaicin, a compound found in topical pain relief creams
- E DDT, a nonspecific pesticide that persists in the environment
- M The antimalarial drugs quinine, chloroquine, and artemisinin

Chapter 1 Structure and Bonding

- M L-Dopa, a drug used to treat Parkinson's disease (Chapter opener and Section 1.14)
- M Alendronic acid (Fosamax), a drug used to prevent osteoporosis (Section 1.5)
- B Enanthotoxin, a poisonous compound isolated from hemlock water dropwort (Section 1.7)
- G Vanillin, the principal component in the extract of the vanilla bean (Section 1.8B)
- M Structures of active ingredients in common sunscreens (Section 1.8B)
- G Ethane, a component of natural gas (Section 1.10A)
- G Ethylene, a hydrocarbon used to make the plastic polyethylene (Section 1.10B)
- G Acetylene, a gas used in welding torches (Section 1.10C)
- G Cucumber aldehyde, the compound responsible for the odor of freshly cut cucumbers (Section 1.10C)
- M Sinemet, a drug used to treat Parkinson's disease that combines L-dopa and carbidopa (Section 1.14)
- B Vitamin B_6 (Section 1.14)

Chapter 2 Acids and Bases

- M Aspirin, a common analgesic and antipyretic (Chapter opener and Section 2.7)
- M The acid–base chemistry of morphine (Section 2.1)
- M The nasal decongestant pseudoephedrine (Section 2.5, Problem 2.17)
- M Glycolic acid, an α-hydroxy acid used in skin care products (Section 2.5, Problem 2.20)
- E Sulfuric acid, a major contributor to acid rain (Section 2.6)
- M Salicin, an analgesic found in willow bark

Chapter 3 Introduction to Organic Molecules and Functional Groups

- B Vitamin C, a water-soluble vitamin that is important in the formation of collagen (Chapter opener and Section 3.5B)
- M The local anesthetic chloroethane (Section 3.2B)
- E Hemibrevetoxin B, a neurotoxin produced by algal blooms ("red tides") (Section 3.2B)
- M Diethyl ether, the first common general anesthetic (Section 3.2B)
- B Bilobalide, a compound isolated from the *Ginkgo biloba* extracts used in Chinese medicine (Section 3.2B, Problem 3.3)
- M Dexamethasone, a synthetic steroid (Section 3.2B, Problem 3.5)
- B Spermine, isolated from semen, and meperidine, the narcotic Demerol (Section 3.2B, Problem 3.6)
- M Atenolol, a β blocker used to treat high blood pressure, and donepezil, used to treat Alzheimer's disease (Section 3.2C)
- M Dolastatin, an anticancer compound isolated from the seahare *Dolabella auricularia* (Section 3.2C, Problem 3.8)
- M Tamiflu, an antiviral drug used to treat influenza (Section 3.2C, Problem 3.9)
- G How geckos use van der Waals forces to stick to walls (Section 3.3B)
- G MTBE, a high-octane additive in unleaded gasoline, and 4,4'-dichlorobiphenyl, a PCB (Section 3.4C)
- B Norethindrone, an oral contraceptive, and arachidonic acid, a fatty acid (Section 3.4C, Problem 3.18)
- B Vitamin A (retinol), a fat-soluble vitamin found in the vision receptors of the eyes (Section 3.5A)
- B β-Carotene, a precursor to vitamin A (Section 3.5A)
- B Vitamin B_3 and vitamin K_1 (Section 3.5B, Problem 3.19)
- B Avocados as a source of pantothenic acid, vitamin B_5 (Section 3.5B, Problem 3.20)

M Morphine and heroin (Section 3.7A, Problem 3.23)
M The antibiotics nonactin and valinomycin (Section 3.7B)
B Biomolecules, such as glucose, oleic acid, alanine, and dAMP (Section 3.9)
B The artificial sweetener aspartame (Section 3.9, Problem 3.28)

Chapter 4 Alkanes

E Oil slicks that result from crude petroleum being spilled into the ocean from oil tankers or oil wells (Chapter opener)
B The cockroach pheromone undecane (Section 4.1)
B Cyclohexane, one component of mangoes (Section 4.1)
B Allicin, a compound responsible for the odor of garlic (Section 4.3)
M Systematic names, generic names, and trade names in over-the-counter drugs like Motrin (Section 4.3)
G Fossil fuels such as natural gas and petroleum (Section 4.7)
E The combustion of alkanes and how it contributes to global warming (Section 4.14B)
B Lipids such as fat-soluble vitamins, phospholipids, waxes, prostaglandins, and steroids (Section 4.15)
B Pristane, a high molecular weight alkane found in shark liver oil (Section 4.15, Problem 4.33)
B End-of-chapter problems: 4.66 and 4.69

Chapter 5 Stereochemistry

M, B Paclitaxel (Taxol), a drug used to treat ovarian, breast, and other cancers (Chapter opener)
B How differences in the three-dimensional structure of starch and cellulose affect their shape and function (Section 5.1)
M, B Identifying stereogenic centers in Darvon (an analgesic), ephedrine (a decongestant), and fructose (a simple sugar) (Section 5.4A)
M The three-dimensional structure of thalidomide, an anti-nausea drug that caused catastrophic birth defects (Section 5.5)
M, B Identifying stereogenic centers in paclitaxel (anticancer agent) and sucrose (Section 5.5)
M Identifying stereogenic centers in gabapentin (a drug used to treat seizures and chronic pain), gabapentin enacarbil, cholesterol, and Zocor (cholesterol-lowering drug) (Section 5.5, Problems 5.9 and 5.10)
M Assigning *R* and *S* configurations in the drugs Plavix and Zestril (Section 5.6, Problems 5.14 and 5.15)
B The sweetener sorbitol (Section 5.9, Problem 5.24)
B The specific rotation of MSG, a common flavor enhancer (Section 5.12D, Problem 5.32)
M Chiral drugs and how mirror image isomers can have drastically different properties—the analgesic ibuprofen, the antidepressant fluoxetine, and the anti-inflammatory agent naproxen (Section 5.13A)
B The sense of smell and how mirror image isomers (e.g., carvone and celery ketone) can smell differently (Section 5.13B and Problem 5.35)
M, B End-of-chapter problems: 5.36, 5.43, 5.49, 5.50, 5.53, 5.55, 5.60, and 5.65–5.71

Chapter 6 Understanding Organic Reactions

B Entropy changes in the metabolism of glucose (Chapter opener and Section 6.4)
B The synthesis of capsaicin by a substitution reaction (Section 6.2)
B Precursors to the female sex hormone estrone (Section 6.2C, Problem 6.2)
G The reaction of gasoline with O_2 (Section 6.9A)
G Refrigeration and spoilage (Section 6.9A)
B Enzymes, biological catalysts (Section 6.11)
B End-of-chapter problems: 6.33, 6.55, and 6.59

Chapter 7 Alkyl Halides and Nucleophilic Substitution

M Flonase, a synthetic steroid used to treat seasonal allergies (Chapter opener)
B, M Telfairine (insecticide) and halomon (antitumor agent), halogenated compounds isolated from red algae (Section 7.1, Problem 7.1)
B, M Simple alkyl halides—chloromethane (found in emissions from volcanoes), dichloromethane (once used to decaffeinate coffee), and halothane (a general anesthetic) (Section 7.4)
E CFCs and DDT, two polyhalogenated compounds once widely used, now discontinued because of adverse environmental effects (Section 7.4)
B, M Ma‘ilione and plocoralide B, halogenated compounds isolated from red algae (Section 7.4)
B Chondrocole A, a marine natural product isolated from red seaweed (Section 7.4, Problem 7.5)
M The antiseptic CPC (Section 7.6)
M Nucleophilic substitutions in the syntheses of Myambutol (used to treat tuberculosis) and Prozac (an antidepressant) (Section 7.11)
M The synthesis of imatinib, an anticancer drug, by a nucleophilic substitution reaction (Section 7.11, Problem 7.22)
B, M Biological nucleophilic substitution reactions: phosphate leaving groups and *S*-adenosylmethionine (SAM) (Section 7.16)

B The biological synthesis of adrenaline using SAM (Section 7.16)
B The synthesis of nicotine using SAM (Section 7.16, Problem 7.36)
M The importance of organic synthesis in preparing useful drugs such as aspirin (Section 7.18)
B, M End-of-chapter problems: 7.64–7.66, 7.70, 7.76

Chapter 8 Alkyl Halides and Elimination Reactions

E DDE, a degradation product of the pesticide DDT (Chapter opener and Section 8.1)
B Ethylene, a hormone that regulates plant growth and fruit ripening (Section 8.2)
B Classifying alkenes using vitamins A and D (Section 8.2, Problem 8.2)
B Identifying stereoisomerism in alkenes using (*E*)-ocimene, found in lilacs (Section 8.2, Problem 8.4)
B, M Elimination reactions in the syntheses of a prostaglandin, quinine, and estradiol (Section 8.4)
B, M End-of-chapter problems: 8.29 and 8.66

Chapter 9 Alcohols, Ethers, and Related Compounds

B Linalool, an alcohol used in scented soaps and lotions and as an insecticide for controlling fleas and cockroaches (Chapter opener)
B Classifying alcohols using cortisol (Section 9.1)
B Classifying ethers and alcohols using brevenal, a marine natural product formed in red tides (Section 9.1, Problem 9.1)
G, E Ethanol, a gasoline additive and renewable fuel source that can be produced from the fermentation of carbohydrates in grains (Section 9.5A)
G Useful simple alcohols: methanol (wood alcohol), isopropanol (rubbing alcohol), and ethylene glycol (antifreeze) (Section 9.5A)
M Diethyl ether, a general anesthetic (Section 9.5B)
M Sevoflurane, a halogenated ether currently used as a general anesthetic (Section 9.5B)
M Medicinal epoxides: eplerenone (a drug that reduces cardiovascular risk in patients who have already had a heart attack) and tiotropium bromide (a bronchodilator) (Section 9.5C)
M A Williamson ether synthesis in the preparation of paroxetine (antidepressant) (Section 9.6, Problem 9.9)
G The syntheses of vitamin A and patchouli alcohol (used in perfumery) using a dehydration reaction (Section 9.10)
G, B The unpleasant odors related to skunks, onions, and human sweat (Section 9.15A)
B The oxidation of a thiol to a disulfide using grapefruit mercaptan (Section 9.15A, Problem 9.31)
B The synthesis of SAM from methionine and ATP by an S_N2 reaction (Section 9.15B)
M The syntheses of salmeterol and albuterol (two bronchodilators) by the opening of an epoxide ring (Section 9.16)
M The design of asthma drugs that block the synthesis of leukotrienes, highly potent molecules that contribute to the asthmatic response (Section 9.17)
B The metabolism of polycyclic aromatic hydrocarbons (PAHs) to carcinogens that disrupt normal cell function, resulting in cancer or cell death (Section 9.18)
M End-of-chapter problems: 9.49, 9.73, and 9.81

Chapter 10 Alkenes

B The unsaturated fatty acids found in kukui nuts (Chapter opener)
M Degrees of unsaturation in the drugs Ambien and mefloquine (Section 10.2, Problem 10.3)
B 11-*cis*-Retinal, the light-sensitive aldehyde involved in the vision of all vertebrates, arthropods, and mollusks (Section 10.3B, Problem 10.7)
B The sex pheromone of the codling moth (Section 10.3B, Problem 10.9)
G Ethylene, the starting material for preparing polyethylene and a variety of other polymers (Section 10.5)
B The naturally occurring alkenes β-carotene, zingiberene, (*R*)-limonene, and α-farnesene (Section 10.5)
B Triacylglycerols, fatty acids, fats, and oils (Section 10.6)
B Omega-3 fatty acids (Section 10.6, Problem 10.11)
B The synthesis of the female sex hormone estrone (Section 10.15B)
M The synthesis of artemisinin, an antimalarial drug, by a hydroboration–oxidation step (Section 10.16B)
B, M End-of-chapter problems: 10.37, 10.43, 10.44, 10.45, 10.69, and 10.71

Chapter 11 Alkynes

M Oral contraceptives (Chapter opener and Section 11.4)
B Nepheliosyne B, a novel acetylenic fatty acid (Section 11.1, Problem 11.1)
M Synthetic hormones mifepristone and Plan B, drugs that prevent pregnancy (Section 11.4)
B Histrionicotoxin, a diyne isolated from the skin of a frog, used as a poison on arrow tips by the Choco tribe of South America (Section 11.4)
B Acetylide anion reactions in the synthesis of two marine natural products (Section 11.11)
M, B End-of-chapter problems: 11.25 and 11.43

Chapter 12 Oxidation and Reduction

B The metabolism of ethanol, the alcohol in alcoholic beverages (Chapter opener and Section 12.14)
B The partial hydrogenation of vegetable oils and the formation of "trans fats" (Section 12.4)
B The reduction of an alkyne to form *cis*-jasmone, a component of perfume (Section 12.5B, Problem 12.10)
B The use of disparlure, a sex hormone, in controlling the spread of gypsy moths (Section 12.8B)
G The production of ozone from O_2 during electrical storms (Section 12.10)
G Blood alcohol screening (Section 12.12)
E Green chemistry—environmentally benign oxidation reactions (Section 12.13)
B Biological oxidations (Section 12.14)
B The synthesis of insect pheromones using asymmetric epoxidation (Section 12.15)
B, M End-of-chapter problems: 12.37, 12.41, 12.51, 12.53, 12.55, 12.56, 12.60, and 12.61

Chapter 13 Mass Spectrometry and Infrared Spectroscopy

M Infrared spectroscopy and the structure determination of penicillin (Chapter opener and Section 13.8)
M Applying the nitrogen rule to 3-methylfentanyl and MPPP, two drugs that mimic the effects of heroin (Section 13.1)
B Determining the molecular formula of nootkatone (found in grapefruit) (Section 13.1, Problem 13.3)
M Using instrumental analysis to detect THC, the active compound in marijuana, and other drugs (Section 13.4B)
B Mass spectrometry and high molecular weight biomolecules (Section 13.4C)
B End-of-chapter problems: 13.29, 13.30, 13.44, and 13.62

Chapter 14 Nuclear Magnetic Resonance Spectroscopy

B Modern spectroscopic methods and the structure of palau'amine, a complex natural product isolated from a sea sponge (Chapter opener and Problem 14.23)
E The high-octane gasoline additive MTBE, which has contaminated water supplies (Section 14.1B)
B Esters of chrysanthemic acid (from chrysanthemum flowers) as insecticides (Section 14.11, Problem 14.29)
M Magnetic resonance imaging (Section 14.12)
B End-of-chapter problem: 14.37

Chapter 15 Radical Reactions

G Polystyrene, a common synthetic polymer used in packaging materials and beverage cups (Chapter opener)
E Ozone destruction and CFCs (Section 15.9)
B The oxidation of unsaturated lipids by radical reactions (Section 15.11)
M, B Two antioxidants—naturally occurring vitamin E and synthetic BHT (Section 15.12)
B The antioxidant rosmarinic acid (Section 15.12)
G The formation of useful polymers from monomers by radical reactions (Section 15.14)
B, G, M End-of-chapter problems: 15.63, 15.66–15.70, and 15.79

Chapter 16 Conjugation, Resonance, and Dienes

M The laboratory synthesis of morphine by a Diels–Alder reaction (Chapter opener)
B Allylic carbocations in biological reactions, such as the formation of geranyl diphosphate (Section 16.2B)
B Isoprene, a conjugated compound that helps plants tolerate heat stress (Section 16.7)
M The antioxidant lycopene (Sections 16.7 and 16.15A)
M Simvastatin (Zocor) and calcitriol (Rocaltrol), two drugs with conjugated double bonds (Section 16.7)
B The synthesis of tetrodotoxin (found in Japanese puffer fish) by a Diels–Alder reaction (Section 16.12)
B The trienes zingiberene and β-sesquiphellandrene found in ginger root (Section 16.13A, Problem 16.21)
B The Diels–Alder reaction in the synthesis of steroids (Section 16.14C)
G Why lycopene and other highly conjugated compounds are colored (Section 16.15A)
G How sunscreens work (Section 16.15B)
B, M End-of-chapter problems: 16.54, 16.61, 16.69, 16.73, and 16.75

Chapter 17 Benzene and Aromatic Compounds

B, M Capsaicin, the spicy component of hot peppers and the active ingredient in topical creams for the treatment of chronic pain (Chapter opener)
G Polycyclic aromatic hydrocarbons (PAHs), constituents of cigarette smoke and diesel exhaust (Section 17.5)
M Examples of common drugs that contain an aromatic ring—Zoloft, Valium, Novocain, Viracept, Viagra, and Claritin (Section 17.5)
B Histamine and scombroid fish poisoning (Section 17.8)
M Quinine, an antimalarial drug (Section 17.8, Problem 17.13)
M Januvia, a drug used to treat type 2 diabetes (Section 17.8, Problem 17.14)
G Diamond, graphite, and buckminsterfullerene (Section 17.11)
M, B End-of-chapter problems: 17.37, 17.57, 17.60–17.63, and 17.67

Chapter 18 Reactions of Aromatic Compounds

B Vitamin K_1, a fat-soluble vitamin that regulates the synthesis of proteins needed for blood to clot (Chapter opener and Section 18.5E)
M, E Biologically active aryl chlorides: the drugs bupropion and chlorpheniramine, and 2,4-D and 2,4,5-T, herbicide components of the defoliant Agent Orange (Section 18.3)
M Intramolecular Friedel–Crafts acylation in the synthesis of LSD (Section 18.5D)
M The synthesis of sertraline (Zoloft), an SSRI antidepressant (Section 18.5D, Problem 18.10)
B A biological Friedel–Crafts reaction (Section 18.5E)
M Nucleophilic aromatic substitution by addition–elimination in the synthesis of Prozac (Section 18.13A, Problem 18.25)
M Benzocaine, the active ingredient in the over-the-counter topical anesthetic Orajel (Section 18.15C)
M, G, B End-of-chapter problems: 18.42–18.44, 18.61, 18.63, 18.67, 18.68, 18.70, 18.73, and 18.77

Chapter 19 Carboxylic Acids and the Acidity of the O–H Bond

B The essential amino acid lysine (Chapter opener)
B Hexanoic acid, the foul-smelling carboxylic acid in ginkgo seeds (Section 19.2B)
B Biologically significant diacids: oxalic acid, malonic acid, and succinic acid (Section 19.2C)
M Depakote (used to treat seizures) (Section 19.2C, Problem 19.5)
B Biologically significant carboxylic acids: formic acid (ant stings), acetic acid (vinegar), butanoic acid (body odor), oxalic acid (spinach), and lactic acid (sour milk) (Section 19.5)
B GHB (4-hydroxybutanoic acid), an illegal recreational intoxicant used as a "date rape" drug (Section 19.5)
M Isotretinoin, a fatty acid used to treat severe acne (Section 19.5, Problem 19.8)
M, B How NSAIDs block the synthesis of prostaglandins to prevent inflammation (Section 19.6)
B Mandelic acid, a naturally occurring carboxylic acid in plums and peaches (Section 19.9, Problem 19.15)
M The irritant urushiol in poison ivy (Section 19.11, Problem 19.19)
B An introduction to amino acids, the building blocks of proteins; why vegetarians must have a balanced diet (Section 19.14)
B, M End-of-chapter problems: 19.31, 19.41, 19.52, 19.62–19.68, 19.71, and 19.72

Chapter 20 Introduction to Carbonyl Chemistry; Organometallic Reagents; Oxidation and Reduction

B The use of a reduction reaction to synthesize the marine neurotoxin ciguatoxin CTX3C (Chapter opener and Section 20.7A)
B The aldehyde α-sinensal, a component of mandarin oil (Section 20.1, Problem 20.1)
M The anticancer drug Taxol and nucleophilic substitution (Section 20.2, Problem 20.2)
B, M Reduction reactions in the synthesis of the analgesic ibuprofen and the perfume component muscone (Section 20.4)
M The synthesis of the long-acting bronchodilator salmeterol (Section 20.6A)
M The use of CBS reagents in the synthesis of cholesterol-lowering drugs (Section 20.6A, Problem 20.9)
B Biological oxidation–reduction reactions with the coenzymes NADH and NAD^+ (Section 20.6B)
B The synthesis of NAD^+ from the vitamin niacin (Section 20.6B)
M The use of organometallic reagents to synthesize the oral contraceptive ethynylestradiol (Section 20.10C)
B The use of Grignard reagents in the synthesis of C_{18} juvenile hormones and the use of juvenile hormone mimics to regulate the life cycle of insects (Section 21.10C)
B The use of organolithium reagents in the synthesis of two components of lavender oil (Section 20.11, Problem 20.24)
M The use of protecting groups in the conversion of estrone to ethynylestradiol (Section 20.12, Problem 20.26)
M, B End-of-chapter problems: 20.50, 20.56, 20.61, 20.68, 20.75, and 20.78

Chapter 21 Aldehydes and Ketones—Nucleophilic Addition

M Digitoxin, a naturally occurring drug isolated from the woolly foxglove plant and used to treat congestive heart failure (Chapter opener and Problem 21.37)
B Determining the IUPAC names of neral (from lemon grass) and cucumber aldehyde (Section 21.2E, Problem 21.7)
G Formaldehyde and acetone, an industrially useful aldehyde and ketone (Section 21.5)
B Examples of naturally occurring compounds that contain aldehydes or ketones—vanillin, citronellal, cinnamaldehyde, and geranial (Section 21.5)
M Cortisone and prednisone, steroids that contain ketones (Section 21.5)
B Naturally occurring cyanohydrin derivatives: linamarin, from cassava root; and amygdalin, from apricot, peach, and wild cherry pits (Section 21.9B)
B The use of the Wittig reaction in the synthesis of β-carotene, the orange pigment in carrots (Section 21.10B)
B The role of rhodopsin in the chemistry of vision (Section 21.11B)

B The acid-catalyzed hydrolysis of safrole, a carcinogen once used in root beer (Section 21.14B, Problem 21.33)
B, M The acid-catalyzed hydrolysis of the acetal in oleandrin (Section 21.14B, Problem 21.34)
B The carbohydrates glucose and lactose (Section 21.17)
M The role of carbohydrates in diabetes (Section 21.17)
B The carbohydrate galactose (Section 21.17, Problem 21.39)
M, B End-of-chapter problems: 21.52, 21.65, 21.69–21.71, 21.79, 21.80, 21.82, and 21.84–21.86

Chapter 22 Carboxylic Acids and Their Derivatives—Nucleophilic Acyl Substitution

B, M Ginkgolide B, a major constituent of the extracts of the ginkgo tree, *Ginkgo biloba* (Chapter opener and Problem 22.21)
B The esters responsible for the odors of banana, mango, and pineapple (Section 22.6A)
M, B Compounds that contain an ester: vitamin C, cocaine, and the immunosuppressant FK506 (Section 22.6A)
M, B Useful amides: proteins, met-enkephalin, the anticancer drug Gleevec, the penicillin antibiotics, and the cephalosporin antibiotics (Section 22.6B)
G The synthesis of the insect repellant DEET (Section 22.8)
B Mechanism for the synthesis of blattellaquinone, the sex pheromone of the female German cockroach (Section 22.8, Problem 22.13)
M Acylation in the syntheses of aspirin, acetaminophen, and heroin (Section 22.9)
M The cholesterol-lowering drug fenofibrate (Section 22.11B, Problem 22.20)
B The hydrolysis of triacylglycerols in the metabolism of lipids (Section 22.12A)
G Olestra, a fake fat (Section 22.12A)
G The synthesis of soap (Section 22.12B)
M The mechanism of action of β-lactam antibiotics like penicillin (Section 22.14)
G Natural and synthetic fibers: nylon and polyesters (Section 22.16)
B Biological acylation reactions (Section 22.17)
M Cholesteryl esters in plaque, the deposits that form on the inside walls of arteries (Section 22.17)
B The acylation of glucosamine to form NAG, the monomer in chitin (Section 22.17, Problem 22.30)
B, M End-of-chapter problems: 22.48, 22.52, 22.53, 22.56–22.61, 22.67, 22.68, 22.72, 22.77, and 22.83–22.85

Chapter 23 Substitution Reactions of Carbonyl Compounds at the α Carbon

M The synthesis of the anticancer drug tamoxifen (Chapter opener and Section 23.8C)
B Keto–enol tautomerizations in glycolysis (Section 23.2A, Problem 23.2)
M The synthesis of the antimalarial drug quinine by an intramolecular substitution reaction (Section 23.7C)
M The heterocyclic ring system in some antitumor agents (Section 23.8C, Problem 23.19)
M The use of the acetoacetic ester synthesis in the synthesis of illudin-S, an antitumor agent (Section 23.10, Problem 23.27)
M Retrosynthesis of the pain reliever nabumetone (Section 23.10, Problem 23.28)
B, M End-of-chapter problems: 23.38, 23.40, 23.45, 23.53, 23.54, 23.61, 23.64, 23.68, 23.72, and 23.74

Chapter 24 Carbonyl Condensation Reactions

M The synthesis of ibuprofen (Chapter opener and Problem 24.20)
B The perfume component flosal, an α,β-unsaturated aldehyde (Section 24.2B, Problem 24.6)
B The synthesis of periplanone B, sex pheromone of the female American cockroach (Section 24.3)
B The synthesis of *ar*-turmerone, a component of turmeric, a principal ingredient in curry powder (Section 24.3)
B The conversion of zingerone to gingerol, components of ginger, using a directed aldol reaction (Section 24.3, Problem 24.11)
M A directed aldol reaction in the synthesis of the drug donepezil (for treating dementia) (Section 24.3, Problem 24.12)
B The synthesis of the steroid progesterone by an intramolecular aldol reaction (Section 24.4)
M Avobenzone, a common ingredient in commercial sunscreens (Section 24.6A, Problem 24.18)
B The synthesis of the female sex hormone estrone by a Michael reaction (Section 24.8)
M, B End-of-chapter problems: 24.34, 24.44, 24.50, 24.53–24.56, 24.58, 24.66, 24.72, and 24.73

Chapter 25 Amines

M Scopolamine, an alkaloid used to treat the nausea and vomiting associated with motion sickness (Chapter opener)
M The stereogenic centers in dobutamine, an amine used in stress tests (Section 25.2, Problem 25.1)
B Poisonous diamines with putrid odors: putrescine and cadaverine (Section 25.6A)
B Naturally occurring alkaloids: atropine, nicotine, and coniine (Section 25.6A)
M Histamine, antihistamines, and antiulcer drugs like Tagamet (cimetidine) (Section 25.6B)
B, M Biologically active derivatives of 2-phenylethanamine: adrenaline, noradrenaline, methamphetamine, mescaline, and dopamine (Section 25.6C)
B, M The neurotransmitter serotonin and SSRI antidepressants (Section 25.6C)
B Bufotenin and psilocin (hallucinogens) (Section 25.6C)

M The synthesis of methamphetamine (Section 25.7C)
M The synthesis of enalapril, an antihypertensive, by reductive amination (Section 25.7C, Problem 25.14)
M The synthesis of the drugs rimantadine and pseudoephedrine by reductive amination (Section 25.7C, Problem 25.15)
M The systematic name of a component of the diet drug fen–phen (Section 25.7C, Problem 25.16)
M Drugs, such as the antihistamine diphenhydramine, sold as water-soluble ammonium salts (Section 25.9)
M Hybridization effects on the basicity of nicotine (Section 25.10E, Problem 25.22)
M Acid–base properties of the drugs chloroquine, matrine, tacrine, and quinine (Section 25.10F and Problem 25.23)
G Azo dyes (Section 25.15)
G Perkin's mauveine and synthetic dyes (Section 25.16A)
M Sulfa drugs (Section 25.16B)
M End-of-chapter problems: 25.37, 25.42, 25.44, 25.54, 25.57, 25.58, 25.68, 25.70, 25.77, and 25.78

Chapter 26 Carbon–Carbon Bond-Forming Reactions in Organic Synthesis

M Ingenol mebutate, used to treat the skin condition actinic keratosis (Chapter opener and Section 26.6, Problem 26.16)
B The synthesis of C_{18} juvenile hormone (Section 26.1A, Problem 26.2)
B, E Use of the Suzuki reaction to prepare bombykol, the sex pheromone of the female silkworm moth, and humulene, a lipid isolated from hops (Section 26.2B)
E Pyrethin I, a biodegradable insecticide isolated from chrysanthemums, and decamethrin, a synthetic analogue (Section 26.4)
M Ring-closing metathesis and the synthesis of epothilone A, an anticancer drug, and Sch38516, an antiviral agent (Section 26.6)
M, B, G End-of-chapter problems: 26.25, 26.26, 26.33, 26.37, 26.38, 26.50

Chapter 27 Pericyclic Reactions

B One synthesis of periplanone B (sex pheromone of the female American cockroach) using pericyclic reactions (Chapter opener and Section 27.5B, Problem 27.22)
B The role of photochemical electrocyclic ring opening and sigmatropic rearrangements in the formation of vitamin D_3 from 7-dehydrocholesterol (Section 27.3C, Problem 27.9)
M The synthesis of the alkaloid reserpine by a [4 + 2] cycloaddition reaction (Section 27.4B, Problem 27.15)
M Garsubellin A and the synthesis of the neurotransmitter acetylcholine (Section 27.5B, Problem 27.25)
B End-of-chapter problems: 27.43, 27.48, and 27.62

Chapter 28 Carbohydrates

B Solanine, the defensive toxin found in the leaves, stems, and green spots of potatoes (Chapter opener and Section 28.7C)
B The use of fructose in "lite" foods (Section 28.2)
B Dihydroxyacetone, the active ingredient in many artificial tanning agents (Section 28.2)
B Glucose, the most common simple sugar (Section 28.6)
G Honey, a mixture of D-fructose and D-glucose (Section 28.6D)
B, M The naturally occurring glycosides salicin and solanine (Section 28.7C)
G Rebaudioside A, a sweet glycoside from the stevia plant (Section 28.7C, Problem 28.19)
B Glucitol (sorbitol), a sucrose substitute (Section 28.9A)
B The common disaccharides maltose, lactose, and sucrose (Section 28.11)
M Lactose intolerance (Section 28.11B)
G Artificial sweeteners (Section 28.11C)
B The common polysaccharides cellulose, starch, and glycogen (Section 28.12)
B, M Glucosamine, an over-the-counter remedy for osteoarthritis, and chitin, the carbohydrate that gives rigidity to crab shells (Section 28.13A)
B *N*-Glycosides and the structure of DNA (Section 28.14B)
B, M End-of-chapter problems: 28.66 and 28.69

Chapter 29 Amino Acids and Proteins

B Myoglobin, the protein that stores oxygen in tissues (Chapter opener and Section 29.10C)
B The naturally occurring amino acids (Section 29.1)
M L-Thyroxine, used to treat thyroid hormone deficiency (Section 29.1B, Problem 29.4)
B The structures of the hormones bradykinin, oxytocin, and vasopressin (Section 29.5C)
B The artificial sweetener aspartame (Section 29.5C)
B The amino acid sequence of leu-enkephalin, an analgesic and opiate (Section 29.5C, Problem 29.17)
B The structure of glutathione, a powerful antioxidant in cells (Section 29.5C, Problem 29.18)
B The Merrifield method of automated protein synthesis (Section 29.8)
B The structures of lysozyme and spider silk (Section 29.9B)

M The structure of insulin (Section 29.9C)
B α-Keratin, the protein in hair, hooves, nails, skin, and wool (Section 29.10A)
B Collagen, the protein in connective tissue (Section 29.10B)
B, M Hemoglobin and the structure of sickle cell hemoglobin (Section 29.10C)
M, B End-of-chapter problems: 29.32, 29.46, 29.48, 29.50, 29.54, 29.56, 29.67, and 29.70

Chapter 30 Synthetic Polymers

G Polyethylene terephthalate, an easily recycled synthetic polymer used in transparent soft drink containers (Chapter opener and Sections 30.6B and 30.9A)
G Consumer products made from Lexan, nylon 6,6, rubber, and polyethylene (Section 30.1)
G Polyethylene, the plastic in milk jugs and plastic bags, and other chain-growth polymers (Section 30.2)
G ABS, a copolymer used in crash helmets, small appliances, and toys (Section 30.3, Problem 30.11)
G Using Ziegler–Natta catalysts to make high-density polyethylene (Section 30.4)
G Dyneema, a strong fiber made of ultra high-density polyethylene (Section 30.4)
B Natural and synthetic rubber (Section 30.5)
G The synthesis of the step-growth polymers nylon, Kevlar, Dacron, spandex, and Lexan (Section 30.6)
M Dissolving sutures (Section 30.6B)
E Polyethylene furanoate, a polymer synthesized from renewable resources (Section 30.6B, Problem 30.16)
G Spandex for active wear (Section 30.6C)
G Lexan for bike helmets, goggles, catcher's masks, and bulletproof glass (Section 30.6D)
G Epoxy resins (Section 30.6E)
G Bakelite for bowling balls (Section 30.7)
E Green polymer synthesis: environmentally benign methods for preparing polymers (Section 30.8)
E Polymer recycling (Section 30.9A)
E Biodegradable polymers (Section 30.9B)
G, E, M End-of-chapter problems: 30.34, 30.35, 30.50, 30.52, and 30.56–30.58

Chapter 31 Lipids (Available online)

B Cholesterol, the most prominent steroid (Chapter opener and Section 31.8B)
B Structure of spermaceti wax (Section 31.2)
B Waxes obtained from jojoba seeds that are used in cosmetics and personal care products (Section 31.2, Problem 31.1)
B Triacylglycerols, the components of fats and oils (Section 31.3)
B Essential fatty acids (Section 31.3)
B The saturated versus unsaturated fatty acid content of fats and oils (Section 31.3)
B Energy storage and the metabolism of fats (Section 31.3)
B The phospholipids in cell membranes (Section 31.4)
B Fat-soluble vitamins: A, D, E, and K (Section 31.5)
B The eicosanoids, a group of biologically active lipids that includes the prostaglandins and leukotrienes (Section 31.6)
M Misoprostol, an analogue of PGE_1 used to prevent gastric ulcers, and unoprostone isopropyl, a prostaglandin analogue used to treat glaucoma (Section 31.6)
M NSAIDs like aspirin and ibuprofen and the COX-1 and COX-2 enzymes (Section 31.6)
M The anti-inflammatory drugs Vioxx, Bextra, and Celebrex (Section 31.6)
B Essential oils that are terpenes and terpenoids (Section 31.7)
B Locating isoprene units in geraniol, vitamin A, grandisol (pheromone), and camphor (Section 31.7, Problem 31.10)
B Biformene, a terpenoid from amber (Section 31.7, Problem 31.11)
B, M The structures of steroids: cholesterol, sex hormones (female and male), adrenal cortical steroids, anabolic steroids, and oral contraceptives (Section 31.8)
M Cholesterol and the cholesterol-lowering drugs Lipitor and Zocor (Section 31.8B)
B, M Anabolic steroids (Section 31.8C)
B, M End-of-chapter problems: 31.20, 31.26–31.28, 31.30, 31.31, 31.35, 31.36, and 31.39

Prologue

What is organic chemistry?
Some representative organic molecules
Organic chemistry and malaria

Organic chemistry. You might wonder how a discipline that conjures up images of eccentric old scientists working in basement laboratories is relevant to you, a student in the twenty-first century.

Consider for a moment the activities that occupied your past 24 hours. You likely showered with soap, drank a caffeinated beverage, ate at least one form of starch, took some medication, listened to a CD, and traveled in a vehicle that had rubber tires and was powered by fossil fuels. If you did any *one* of these, your life was touched by organic chemistry.

What Is Organic Chemistry?

- **Organic chemistry is the chemistry of compounds that contain the element carbon.**

It is one branch in the entire field of chemistry, which encompasses many classical subdisciplines including inorganic, physical, and analytical chemistry, and newer fields such as bioinorganic chemistry, physical biochemistry, polymer chemistry, and materials science.

Some compounds that contain the element carbon are *not* organic compounds. Examples include carbon dioxide (CO_2), sodium carbonate (Na_2CO_3), and sodium bicarbonate ($NaHCO_3$).

Organic chemistry was singled out as a separate discipline for historical reasons. Originally, it was thought that compounds in living things, termed ***organic compounds,*** were fundamentally different from those in nonliving things, called ***inorganic compounds.*** Although we have known for more than 150 years that this distinction is artificial, the name *organic* persists. Today the term refers to the study of the compounds that contain carbon, many of which, incidentally, are found in living organisms.

It may seem odd that a whole discipline is devoted to the study of a single element in the periodic table, when more than 100 elements exist. It turns out, though, that there are far more organic compounds than any other type. **Organic chemicals affect virtually every facet of our lives, and for this reason, it is important and useful to know something about them.**

Clothes, foods, medicines, gasoline, refrigerants, and soaps are composed almost solely of organic compounds. Some, like cotton, wool, or silk are naturally occurring; that is, they can be isolated directly from natural sources. Others, such as nylon and polyester, are synthetic, meaning they are produced by chemists in the laboratory. By studying the principles and concepts of organic chemistry, you can learn more about compounds such as these and how they affect the world around you.

Realize, too, what organic chemistry has done for us. Organic chemistry has made available both comforts and necessities that were previously nonexistent, or reserved for only the wealthy. We have seen an enormous increase in life span, from 47 years in 1900 to over 70 years currently. To a large extent this is due to the isolation and synthesis of new drugs to fight infections and the availability of vaccines for childhood diseases. Chemistry has also given us the tools to control

insect populations that spread disease, and there is more food for all because of fertilizers, pesticides, and herbicides. Our lives would be vastly different today without the many products that result from organic chemistry (Figure 1).

Figure 1
Products of organic chemistry used in medicine


a. Oral contraceptives



b. Plastic syringes



c. Antibiotics



d. Synthetic heart valves



- Organic chemistry has given us contraceptives, plastics, antibiotics, and the knitted material used in synthetic heart valves.


Some Representative Organic Molecules

Perhaps the best way to appreciate the variety of organic molecules is to look at a few. Three simple organic compounds are **methane, ethanol,** and **trichlorofluoromethane.**




methane

- **Methane,** the simplest of all organic compounds, contains one carbon atom. Methane—the main component of natural gas—occurs widely in nature. Like other **hydrocarbons**—organic compounds that contain only carbon and hydrogen—methane is combustible; that is, it burns in the presence of oxygen. Methane is the product of the anaerobic (without air) decomposition of organic matter by bacteria. The natural gas we use today was formed by the decomposition of organic material millions of years ago. Hydrocarbons such as methane are discussed in Chapter 4.



ethanol




trichlorofluoromethane

- **Ethanol,** the alcohol present in beer, wine, and other alcoholic beverages, is formed by the fermentation of sugar, quite possibly the oldest example of organic synthesis. Ethanol can also be made in the lab by a totally different process, but **the ethanol produced in the lab is *identical* to the ethanol produced by fermentation.** Alcohols including ethanol are discussed in Chapter 9.
- **Trichlorofluoromethane** is a member of a class of molecules called **c**hloro**f**luoro**c**arbons or **CFCs,** which contain one or two carbon atoms and several halogens. Trichlorofluoromethane is an unusual organic molecule in that **it contains no hydrogen atoms.** Because it has a low molecular weight and is easily vaporized, trichlorofluoromethane has been used as an aerosol propellant and refrigerant. It and other CFCs have been implicated in the destruction of the stratospheric ozone layer, a topic discussed in Chapter 15.

Three complex organic molecules that are important medications are **amoxicillin, fluoxetine,** and **AZT.**


- **Amoxicillin** is one of the most widely used antibiotics in the penicillin family. The discovery and synthesis of such antibiotics in the twentieth century have made routine the treatment of infections that were formerly fatal. You were likely given some amoxicillin to treat an ear infection when you were a child. The penicillin antibiotics are discussed in Chapter 22.

Complex organic structures are drawn with shorthand conventions described in Chapter 1.




amoxicillin

- **Fluoxetine** is the generic name for the antidepressant **Prozac.** Prozac was designed and synthesized by chemists in the laboratory, and is now produced on a large scale in chemical factories. Because it is safe and highly effective in treating depression, Prozac is widely prescribed. Over 40 million individuals worldwide have used Prozac since 1986.



fluoxetine


- **AZT, az**idodeoxy**t**hymidine, is a drug that treats human immunodeficiency virus (HIV), the virus that causes acquired immune deficiency syndrome (AIDS). Also known by its generic name **zidovudine,** AZT represents a chemical success to a different challenge: synthesizing agents that combat viral infections.



AZT


Other complex organic compounds having interesting properties are **capsaicin** and **DDT.**

- **Capsaicin,** one member of a group of compounds called *vanilloids,* is responsible for the characteristic spiciness of hot peppers. It is the active ingredient in pepper sprays used for personal defense and topical creams used for pain relief.



capsaicin

- **DDT, d**ichloro**d**iphenyl**t**richloroethane, is a pesticide once called "miraculous" by Winston Churchill because of the many lives it saved by killing disease-carrying mosquitoes. DDT use is now banned in the United States and many developed countries because it is a non-specific insecticide that persists in the environment.



DDT

What are the common features of these organic compounds?


- All organic compounds contain carbon atoms and most contain hydrogen atoms.
- All the carbon atoms have four bonds. A stable carbon atom is said to be *tetravalent*.
- Other elements may also be present. Any atom that is not carbon or hydrogen is called a *heteroatom*. Common heteroatoms include N, O, S, P, and the halogens.
- Some compounds have chains of atoms and some compounds have rings.

These features explain why there are so many organic compounds: **Carbon forms four strong bonds with itself and other elements. Carbon atoms combine together to form rings and chains.**

Organic Chemistry and Malaria


A vast array of organic compounds is now available to fight malaria, a mosquito-borne infectious disease that affects an estimated 200 million people worldwide. Antimalarial medications include organic compounds isolated from natural sources or those synthesized by chemists in the laboratory. Two common antimalarial drugs shown in Figure 2 are **quinine,** a centuries-old remedy obtained from the bark of the cinchona tree native to the Andes Mountains, and **chloroquine,** a synthetic drug introduced in the late 1940s.

Figure 2
Antimalarial drugs



quinine

buds and leaves of *Cinchona pubescens*




chloroquine



Because malaria is caused by a variety of closely related parasitic microorganisms and drug-resistant strains have developed, currently recommended therapy consists of a combination of drugs that includes **artemisinin** or a related compound. Artemisinin is a complex compound isolated from sweet wormwood, *Artemisia annua,* a plant used for hundreds of years in traditional Chinese medicine. Although artemisinin can be obtained by extracting the active drug from the dried leaves of *Artemisia annua,* this process does not meet the worldwide demand. As a result, artemisinin can now be obtained using genetic engineering and fermentation processes.

The 2015 Nobel Prize in Physiology or Medicine was awarded to Youyou Tu for her discovery of artemisinin as an antimalarial drug.



artemisinin



Artemisia annua, sweet wormwood


Malaria continues to present a major public health challenge for chemists, health professionals, and biologists. Despite extensive efforts to prevent and control the disease in the equatorial regions of Asia, Africa, and Latin America, it is estimated that malaria was responisble for over 450,000 deaths in 2012.

In this introduction, we have seen a variety of molecules that have diverse structures. They represent a miniscule fraction of the organic compounds currently known and the many thousands that are newly discovered or synthesized each year. The principles you learn in organic chemistry will apply to all of these molecules, from simple ones like methane and ethanol, to complex ones like capsaicin and artemisinin. It is these beautiful molecules, their properties, and their reactions that we will study in organic chemistry.

WELCOME TO THE WORLD OF ORGANIC CHEMISTRY!

Structure and Bonding

1



1.1 The periodic table
1.2 Bonding
1.3 Lewis structures
1.4 Isomers
1.5 Exceptions to the octet rule
1.6 Resonance
1.7 Determining molecular shape
1.8 Drawing organic structures
1.9 Hybridization
1.10 Ethane, ethylene, and acetylene
1.11 Bond length and bond strength
1.12 Electronegativity and bond polarity
1.13 Polarity of molecules
1.14 L-Dopa—A representative organic molecule

L-Dopa, also called levodopa, was first isolated from seeds of the broad bean plant *Vicia faba* in 1913. Since 1967 it has been the drug of choice for the treatment of Parkinson's disease, a debilitating illness that results from the degeneration of neurons that produce the neurotransmitter dopamine in the brain. L-Dopa is an oral medication that is transported to the brain by the bloodstream, where it is converted to dopamine. Since L-dopa must be taken in large doses with some serious side effects, today it is often given with other drugs that lessen its negative impact. In Chapter 1, we learn about the structure, bonding, and properties of organic molecules like L-dopa.


Before examining organic molecules in detail, we must review topics about structure and bonding learned in previous chemistry courses. We will discuss these concepts primarily from an organic chemist's perspective, and spend time on only the particulars needed to understand organic compounds.

Important topics in Chapter 1 include drawing Lewis structures, predicting the shape of molecules, determining what orbitals are used to form bonds, and how electronegativity affects bond polarity. Equally important is Section 1.8 on drawing organic molecules, both shorthand methods routinely used for simple and complex compounds, as well as three-dimensional representations that allow us to more clearly visualize them.

1.1 The Periodic Table

All matter is composed of the same building blocks called **atoms.** There are two main components of an atom.

- The **nucleus** contains positively charged **protons** and uncharged **neutrons.** Most of the mass of the atom is contained in the nucleus.
- The **electron cloud** is composed of negatively charged **electrons.** The electron cloud comprises most of the volume of the atom.



The charge on a proton is equal in magnitude but opposite in sign to the charge on an electron. In a neutral atom, the **number of protons in the nucleus equals the number of electrons.** This quantity, called the **atomic number,** is unique to a particular element. For example, every neutral carbon atom has an atomic number of six, meaning it has six protons in its nucleus and six electrons surrounding the nucleus.

In addition to neutral atoms, we will also encounter **charged ions.**

- A *cation* is positively charged and has fewer electrons than protons.
- An *anion* is negatively charged and has more electrons than protons.

The number of neutrons in the nucleus of a particular element can vary. **Isotopes** are two atoms of the same element having a different number of neutrons. The **mass number** of an atom is the total number of protons and neutrons in the nucleus. Isotopes have different mass numbers. The **atomic weight** of a particular element is the weighted average of the mass of all its isotopes, reported in atomic mass units (amu).

Isotopes of carbon and hydrogen are sometimes used in organic chemistry. The most common isotope of hydrogen has one proton and no neutrons in the nucleus, but 0.02% of hydrogen atoms have one proton and one neutron. This isotope of hydrogen is called **deuterium,** and is sometimes symbolized by the letter **D.**

Each atom is identified by a one- or two-letter abbreviation that is the characteristic symbol for that element. Carbon is identified by the single letter **C.** Sometimes the atomic number is indicated as a subscript to the left of the element symbol, and the mass number is indicated as a superscript. Using this convention, the most common isotope of carbon, which contains six protons and six neutrons, is designated as $^{12}_{6}C$.

A **row** in the periodic table is also called a ***period,*** and a **column** is also called a ***group.*** A periodic table is located on the inside front cover for your reference.


Long ago it was realized that groups of elements have similar properties, and that these atoms could be arranged in a schematic way called the **periodic table.** There are more than 100 known elements, arranged in the periodic table in order of increasing atomic number. The periodic table is composed of rows and columns.

- **Elements in the same row are similar in *size.***
- **Elements in the same column have similar *electronic and chemical properties.***

Each column in the periodic table is identified by a **group number,** an Arabic (1 to 8) or Roman (I to VIII) numeral followed by the letter A or B. Carbon is located in group **4A** in the periodic table in this text.


Although more than 100 elements exist, most are not common in organic compounds. Figure 1.1 contains a truncated periodic table, indicating the handful of elements that are routinely seen in this text. **Most of these elements are located in the first and second rows of the periodic table.**

Figure 1.1
A periodic table of the common elements seen in organic chemistry



- Carbon is located in the second row, group 4A.

Carbon's entry in the periodic table:




Across each row of the periodic table, electrons are added to a particular shell of orbitals around the nucleus. The shells are numbered 1, 2, 3, and so on. Adding electrons to the first shell forms the first row. Adding electrons to the second shell forms the second row. **Electrons are first added to the shells closest to the nucleus.**

Each shell contains a certain number of **orbitals.** An orbital is a region of space that is high in electron density. There are four different kinds of orbitals, called ***s, p, d,*** and ***f.*** The first shell has only one orbital, called an ***s*** orbital. The second shell has two kinds of orbitals, ***s*** and ***p,*** and so on. Each type of orbital has a particular shape.

For the first- and second-row elements, we must consider only ***s*** **orbitals** and ***p*** **orbitals.**

- **An *s* orbital has a sphere of electron density. It is *lower in energy* than other orbitals of the same shell, because electrons are kept closer to the positively charged nucleus.**
- **A *p* orbital has a dumbbell shape. It contains a node of electron density at the nucleus. A node means there is *no* electron density in this region. A *p* orbital is *higher in energy* than an *s* orbital (in the same shell) because its electron density is farther away from the nucleus.**



An *s* orbital is filled with electrons before a *p* orbital in the same shell.

1.1A The First Row

The first row of the periodic table is formed by adding electrons to the first shell of orbitals around the nucleus. There is only one orbital in the first shell, called the **1*s* orbital.**

- **Each orbital can have a maximum of two electrons.**


As a result, there are **two elements in the first row,** one having one electron added to the 1*s* orbital, and one having two. The element hydrogen (H) has what is called a $1s^1$ configuration with one electron in the 1*s* orbital, and helium (He) has a $1s^2$ configuration with two electrons in the 1*s* orbital.




1.1B The Second Row

Every element in the second row has a filled first shell of electrons. Thus, all second-row elements have a $1s^2$ configuration. Each element in the second row of the periodic table also has four orbitals available to accept additional electrons:

- **one 2*s* orbital,** the *s* orbital in the second shell
- **three 2*p* orbitals,** all dumbbell-shaped and perpendicular to each other along the *x*, *y*, and *z* axes



Because each of the four orbitals in the second shell can hold two electrons, there is a **maximum capacity of *eight* electrons** for elements in the second row. The second row of the periodic table consists of eight elements, obtained by adding electrons to the 2*s* and three 2*p* orbitals.



The outermost electrons are called **valence electrons.** The valence electrons are more loosely held than the electrons closer to the nucleus, and as such, they participate in chemical reactions. **The group number of a second-row element reveals its number of valence electrons.** For example, carbon in group 4A has four valence electrons, and oxygen in group 6A has six.

Problem 1.1 While the most common isotope of nitrogen has a mass number of 14 (nitrogen-14), a radioactive isotope of nitrogen has a mass number of 13 (nitrogen-13). Nitrogen-13 is used in PET (positron emission tomography) scans by physicians to monitor brain activity and diagnose dementia. For each isotope, give the following information: (a) the number of protons; (b) the number of neutrons; (c) the number of electrons in the neutral atom; (d) the group number; and (e) the number of valence electrons.

1.2 Bonding

Until now our discussion has centered on individual atoms, but it is more common in nature to find two or more atoms joined together.

- *Bonding* is the joining of two atoms in a stable arrangement.

Bonding is a favorable process because it always leads to ***lowered energy and increased stability.*** Joining two or more elements forms **compounds.** Although only about 100 elements exist, more than 50 million compounds are known. Examples of compounds include hydrogen gas (H_2), formed by joining two hydrogen atoms, and methane (CH_4), the simplest organic compound, formed by joining a carbon atom with four hydrogen atoms.

One general rule governs the bonding process.

- Through bonding, atoms attain a complete outer shell of valence electrons.

Alternatively, because the noble gases in group 8A of the periodic table are especially stable as atoms having a filled shell of valence electrons, the general rule can be restated.

- Through bonding, atoms gain, lose, or share electrons to attain the electronic configuration of the noble gas closest to them in the periodic table.

What does this mean for first- and second-row elements? **A first-row element like hydrogen can accommodate *two electrons* around it.** This would make it like the noble gas helium at the end of the same row. **A second-row element is generally most stable with *eight valence electrons* around it** like neon. Elements that behave in this manner are said to follow the **octet rule.**

There are two different kinds of bonding: **ionic bonding** and **covalent bonding.**


- *Ionic bonds* result from the *transfer* of electrons from one element to another.
- *Covalent bonds* result from the *sharing* of electrons between two nuclei.



Atoms readily form ionic bonds when they can attain a noble gas configuration by gaining or losing just one or two electrons. NaCl and KI are ionic compounds.

The type of bonding is determined by the location of an element in the periodic table. An ionic bond generally occurs when elements on the **far left** side of the periodic table combine with elements on the **far right** side, ignoring the noble gases, which form bonds only rarely. **The resulting ions are held together by extremely strong electrostatic interactions.** A positively charged cation formed from the element on the left side attracts a negatively charged anion formed from the element on the right side. Examples of ionic inorganic compounds include sodium chloride (NaCl), common table salt, and potassium iodide (KI), an essential nutrient added to make iodized salt.


Ionic compounds form extended crystal lattices that maximize the positive and negative electrostatic interactions. In NaCl, each positively charged Na^+ ion is surrounded by six negatively charged Cl^- ions, and each Cl^- ion is surrounded by six Na^+ ions.



NaCl

Lithium fluoride, LiF, is an example of an ionic compound.

- The element **lithium,** located in group 1A of the periodic table, has one valence electron in its second shell. Loss of this electron forms the cation Li^+ having no electrons in the second shell and two electrons in the first shell like helium.
- The element **fluorine,** located in group 7A of the periodic table, has seven valence electrons. By gaining one it forms the anion F^-, which has a filled valence shell (an octet of electrons), like neon.
- Thus, lithium fluoride is a stable ionic compound composed of Li^+ cations and F^- anions.



- **The transfer of electrons forms stable salts composed of cations and anions.**


A **compound** may have either ionic or covalent bonds. A **molecule** has only covalent bonds.

The second type of bonding, **covalent bonding,** occurs with elements like carbon in the middle of the periodic table, which would otherwise have to gain or lose several electrons to form an ion with a complete valence shell. **A covalent bond is a two-electron bond,** and a compound with covalent bonds is called a **molecule.** Covalent bonds also form between two elements from the same side of the table, such as two hydrogen atoms or two chlorine atoms. H_2, Cl_2, and CH_4 are all examples of covalent molecules.

Problem 1.2 Label each bond in the following compounds as ionic or covalent.

a. F_2 b. LiBr c. CH_3CH_3 d. $NaNH_2$ e. $NaOCH_3$

How many covalent bonds will a particular atom typically form? As you might expect, it depends on the location of the atom in the periodic table. In the first row, **hydrogen forms one covalent bond** using its one valence electron. When two hydrogen atoms are joined in a bond, each has a filled valence shell of two electrons.



Second-row elements can have no more than eight valence electrons around them. For neutral molecules, two consequences result.

- **Atoms with one, two, three, or four valence electrons form one, two, three, or four bonds,** respectively, in neutral molecules.
- **Atoms with five or more valence electrons** form enough bonds to give an octet. In this case, **the predicted number of bonds = 8 – the number of valence electrons.**

For example, B has three valence electrons, so it forms three bonds, as in BF_3. N has five valence electrons, so it also forms three bonds (8 – 5 = 3 bonds), as in NH_3.


Nonbonded pair of electrons = unshared pair of electrons = lone pair

These guidelines are used in Figure 1.2 to summarize the usual number of bonds formed by the common atoms in organic compounds. When second-row elements form fewer than four bonds their octets consist of both **bonding (shared) electrons** and **nonbonding (unshared) electrons.** Unshared electrons are also called **lone pairs.**

Problem 1.3 How many covalent bonds are predicted for each atom?

a. O b. Al c. Br d. Si

Figure 1.2
The usual number of bonds of common neutral atoms




1.3 Lewis Structures

Lewis structures **are electron dot representations for molecules.** Three rules are used for drawing Lewis structures.

1. Draw only the valence electrons.
2. Give every second-row element no more than eight electrons.
3. Give each hydrogen two electrons.

The letter **X** is often used to represent one of the halogens in group 7A: F, Cl, Br, or I.

To draw a Lewis structure for a diatomic molecule like **HF,** recall that hydrogen has one valence electron and fluorine has seven. H and F each donate one electron to form a two-electron bond. The resulting molecule gives both H and F a filled valence shell. In a Lewis structure, **a *solid line* indicates a two-electron covalent bond.**




1.3A A Procedure for Drawing Lewis Structures

Drawing a Lewis structure for larger molecules is easier if you follow a stepwise procedure.


How To Draw a Lewis Structure

Step [1] Arrange atoms next to each other that you think are bonded together.

- Always place hydrogen atoms and halogen atoms on the periphery because H and X (X = F, Cl, Br, and I) form only one bond each.



- As a first approximation, use the common bonding patterns in Figure 1.2 to arrange the atoms.



- In truth, the proper arrangement of atoms may not be obvious, or more than one arrangement may be possible (Section 1.4). Even in many simple molecules, the connectivity between atoms must be determined experimentally.

Step [2] Count the electrons.

- Count the number of valence electrons from all atoms.
- Add one electron for each negative charge.
- Subtract one electron for each positive charge.
- This sum gives the total number of electrons that must be used in drawing the Lewis structure.

Step [3] Arrange the electrons around the atoms.

- Place a bond between every two atoms, giving two electrons to each H and no more than eight to any second-row atom.
- Use all remaining electrons to fill octets with lone pairs.
- If all valence electrons are used and an atom does not have an octet, form multiple bonds, as shown in Sample Problem 1.2.

Step [4] Assign formal charges to all atoms.


- Formal charges are discussed in Section 1.3C.

Sample Problem 1.1 illustrates how to draw a Lewis structure of a simple organic molecule.

Sample Problem 1.1 Draw a Lewis structure for methanol, a compound with molecular formula CH_4O.

Solution

Step [1] Arrange the atoms.




- Place the second-row elements, C and O, in the middle.
- Place three H's around C to surround C by four atoms.
- Place one H next to O to surround O by two atoms.

Step [2] Count the electrons.

$$\begin{aligned} 1\text{ C} \times 4\text{ e}^- &= 4\text{ e}^- \\ 1\text{ O} \times 6\text{ e}^- &= 6\text{ e}^- \\ 4\text{ H} \times 1\text{ e}^- &= 4\text{ e}^- \\ &\textbf{14 e}^- \textbf{ total} \end{aligned}$$

Step [3] Add the bonds and lone pairs.

- Add five two-electron bonds to form the C–H, C–O, and O–H bonds, using 10 of the 14 electrons.
- Place two lone pairs on the O atom to use the remaining four electrons and give the O atom an octet.



This Lewis structure is valid because it uses all 14 electrons, each H is surrounded by two electrons, and each second-row element is surrounded by no more than eight electrons.

Problem 1.4 Draw a valid Lewis structure for each species.

a. CH_3CH_3 b. CH_5N c. CH_3Cl

1.3B Multiple Bonds

Sample Problem 1.2 illustrates an example of a Lewis structure with a double bond.

Sample Problem 1.2 Draw a Lewis structure for ethylene, a compound of molecular formula C_2H_4, in which each carbon is bonded to two hydrogens.

Solution

Follow Steps [1] to [3] to draw a Lewis structure.

Step [1] Arrange the atoms.




- Each C gets 2 H's.

Step [2] Count the electrons.


$$\begin{aligned} 2\text{ C} \times 4\text{ e}^- &= 8\text{ e}^- \\ 4\text{ H} \times 1\text{ e}^- &= 4\text{ e}^- \\ &\textbf{12 e}^- \textbf{ total} \end{aligned}$$

Step [3] Add the bonds and lone pairs.



After placing five bonds between the atoms and adding the two remaining electrons as a lone pair, one C still has no octet.

To give both C's an octet, **change *one* lone pair into *one* bonding pair of electrons between the two C's, forming a double bond.**




ethylene
a valid Lewis structure

This uses all 12 electrons, each C has an octet, and each H has two electrons. The Lewis structure is valid. **Ethylene contains a carbon–carbon double bond.**


- After placing all electrons in bonds and lone pairs, use a lone pair to form a multiple bond if an atom does not have an octet.

Carbon always forms four bonds in stable organic molecules. Carbon forms single, double, and triple bonds to itself and other elements.

You must change *one* lone pair into *one* new bond for each *two* electrons needed to complete an octet. In acetylene, a compound with molecular formula C_2H_2, placing the 10 valence electrons gives a Lewis structure in which one or both of the C's lack an octet.



In this case, **change *two* lone pairs into *two* bonding pairs of electrons, forming a triple bond.**



acetylene
a valid Lewis structure

Problem 1.5 Draw an acceptable Lewis structure for each compound, assuming the atoms are connected as arranged. Hydrogen cyanide (HCN) is a poison, formaldehyde (H_2CO) is a preservative, and glycolic acid ($HOCH_2CO_2H$) is used to make dissolving sutures.

a. HCN H C N b. H_2CO H C O (H) c. $HOCH_2CO_2H$ H O C (H) C O H (H O)

1.3C Formal Charge

To manage electron bookkeeping in a Lewis structure, chemists use **formal charge.**

- *Formal charge* is the charge assigned to individual atoms in a Lewis structure.

By calculating formal charge, we determine how the number of electrons around a particular atom compares to its number of valence electrons. Formal charge is calculated as follows:

formal charge = **number of valence electrons** − **number of electrons an atom "owns"**

The number of electrons "owned" by an atom is determined by its number of bonds and lone pairs.

- An atom "owns" *all* of its unshared electrons and *half* of its shared electrons.

$$\text{number of electrons owned} = \text{number of unshared electrons} + \frac{1}{2}\left[\text{number of shared electrons}\right]$$

The number of electrons "owned" by different carbon atoms is indicated in the following examples:



- C shares eight electrons.
- C "owns" **four** electrons.




- Each C shares eight electrons.
- Each C "owns" **four** electrons.



- C shares six electrons.
- C has two unshared electrons.
- C "owns" **five** electrons.

Sample Problem 1.3 illustrates how formal charge is calculated on the atoms of a polyatomic ion. **The sum of the formal charges on the individual atoms equals the net charge on the molecule or ion.**


Sample Problem 1.3 Determine the formal charge on each atom in the ion H_3O^+.



Solution

To calculate the formal charge on each atom:

- Determine the number of valence electrons from the group number.
- Determine the number of electrons an atom "owns" from the number of bonding and nonbonding electrons it has.
- Subtract the second quantity from the first to give the formal charge.



For the O atom (group 6A):

- number of valence electrons = 6
- number of bonding electrons = 6
- number of nonbonding electrons = 2

$$\text{formal charge} = 6 - \left[2 + \frac{1}{2}(6)\right] = +1$$


For each H atom (group 1A):

- number of valence electrons = 1
- number of bonding electrons = 2
- number of nonbonding electrons = 0


$$\text{formal charge} = 1 - \left[0 + \frac{1}{2}(2)\right] = 0$$

The formal charge on the O atom is +1 and the formal charge on each H is 0. The overall charge on the ion H_3O^+ is the sum of all of the formal charges on the atoms: 1 + 0 + 0 + 0 = +1.

Problem 1.6 Calculate the formal charge on each second-row atom.

a. 

b. $CH_3{-}N{\equiv}C:$

c. 

Problem 1.7 Draw a Lewis structure for each ion.


a. CH_3O^- b. HC_2^- c. $(CH_3NH_3)^+$ d. $(CH_3NH)^-$

When you first add formal charges to Lewis structures, use the procedure in Sample Problem 1.3. With practice, you will notice that certain bonding patterns always result in the same formal charge. For example, any N atom with four bonds (and, thus no lone pairs) has a +1 formal charge. Table 1.1 lists the bonding patterns and resulting formal charges for carbon, nitrogen, and oxygen.

Table 1.1 Formal Charge Observed with Common Bonding Patterns for C, N, and O


Atom	Number of valence electrons	Formal charge		
		+1	0	–1
C	4			
N	5			
O	6			

Problem 1.8 What is the formal charge on the O atom in each of the following species that contains a multiple bond to O?

a. b.  c.

1.4 Isomers

In drawing a Lewis structure for a molecule with several atoms, sometimes more than one arrangement of atoms is possible for a given molecular formula. For example, there are two acceptable arrangements of atoms for the molecular formula C_2H_6O.



ethanol dimethyl ether

same molecular formula
C_2H_6O
isomers

Both are valid Lewis structures, and both molecules exist. One is called ethanol, and the other, dimethyl ether. These two compounds are called **isomers.**


- *Isomers* are different molecules having the same molecular formula.

Ethanol and dimethyl ether are **constitutional isomers** because they have the same molecular formula, but the ***connectivity of their atoms is different.*** Ethanol has one C−C bond and one O−H bond, whereas dimethyl ether has two C−O bonds. A second class of isomers, called **stereoisomers,** is introduced in Section 4.13B.

Problem 1.9 Draw Lewis structures for each molecular formula.

a. $C_2H_4Cl_2$ (two isomers) b. C_3H_8O (three isomers) c. C_3H_6 (two isomers)

1.5 Exceptions to the Octet Rule




Alendronic acid, sold as a sodium salt under the trade name of **Fosamax,** is used to prevent osteoporosis in women. Osteoporosis decreases bone density, as shown by comparing normal bone (top) with brittle bone (bottom).

Most of the common elements in organic compounds—**C, N, O, and the halogens**—follow the octet rule. Hydrogen is a notable exception, because it accommodates only two electrons in bonding. Additional exceptions include boron and beryllium (second-row elements in groups 3A and 2A, respectively), and elements in the third row (particularly phosphorus and sulfur).

Elements in groups 2A and 3A of the periodic table, such as beryllium and boron, do not have enough valence electrons to form an octet in a neutral molecule. Lewis structures for BeH_2 and BF_3 show that these atoms have only four and six electrons, respectively, around the central atom. There simply aren't enough electrons to form an octet. Because the Be and B atoms each have less than an octet of electrons, these molecules are highly reactive.




A second exception to the octet rule occurs with some elements located in the third row and later in the periodic table. These elements have empty *d* orbitals available to accept electrons, and thus they may have ***more than eight*** **electrons** around them. For organic chemists, the two most common elements in this category are **phosphorus** and **sulfur,** which can have 10 or even 12 electrons around them, as shown in dimethyl sulfoxide, sulfuric acid, and alendronic acid.



1.6 Resonance

Some molecules can't be adequately represented by a single Lewis structure. For example, two valid Lewis structures can be drawn for the anion $(HCONH)^-$. One structure has a negatively charged N atom and a C−O double bond; the other has a negatively charged O atom and a C−N double bond. These structures are called **resonance structures** or **resonance forms.** A **double-headed arrow** is used to separate two resonance structures.



- *Resonance structures* are two Lewis structures having the *same* placement of atoms but a *different* arrangement of electrons.

Which resonance structure is an accurate representation for $(HCONH)^-$? **The answer is *neither* of them.** The true structure is a composite of both resonance forms, and is called a **resonance hybrid.** The hybrid shows characteristics of ***both*** resonance structures.


Each resonance structure implies that electron pairs are localized in bonds or on atoms. In actuality, resonance allows certain electron pairs to be ***delocalized*** over two or more atoms, and this delocalization of electron density adds stability. **A molecule with two or more resonance structures is said to be *resonance stabilized.***

1.6A An Introduction to Resonance Theory


Keep in mind the following basic principles of resonance theory.

- Resonance structures are not real. An individual resonance structure does not accurately represent the structure of a molecule or ion.
- Resonance structures are *not* in equilibrium with each other. There is no movement of electrons from one form to another.
- Resonance structures are *not* isomers. Two isomers differ in the arrangement of *both* atoms and electrons, whereas resonance structures differ *only* in the *arrangement of electrons.*

For example, ions **A** and **B** are resonance structures because the atom position is the same in both compounds, but the location of an electron pair is different. In contrast, compounds **C** and **D** are isomers because the atom placement is different; **C** has an O−H bond, and **D** has an additional C−H bond.




A




B

- **A** and **B** are **resonance** structures.
- The position of one electron pair (in red) is different.





C





D


- **C** and **D** are **isomers.**
- The position of a H atom (in red) is different.

Problem 1.10 Classify each pair of compounds as isomers or resonance structures.


a.  and 

b.  and 


Problem 1.11 Considering structures **A–D,** classify each pair of compounds as isomers, resonance structures, or neither: (a) **A** and **B;** (b) **A** and **C;** (c) **A** and **D;** (d) **B** and **D.**




A



B



C




D

1.6B Drawing Resonance Structures

To draw resonance structures, use three rules.


Rule [1] **Two resonance structures differ in the position of multiple bonds and nonbonded electrons. The placement of atoms and single bonds always stays the same.**



A **B**

- The position of a double bond (in blue) is different.
- The position of a lone pair (in red) is different.

Rule [2] **Two resonance structures must have the same number of unpaired electrons.**




A **C**

no unpaired electrons — two unpaired electrons

not a resonance structure of **A** (or **B**)

Rule [3] **Resonance structures must be valid Lewis structures. Hydrogen must have two electrons and no second-row element can have more than eight electrons.**



A


10 electrons around C

not a valid resonance structure of **A**

Curved arrow notation is a convention that shows how electron position differs between the two resonance forms.

- ***Curved arrow notation shows the movement of an electron pair.*** **The tail of the arrow always begins at an electron pair, either in a bond or lone pair. The head points to where the electron pair "moves."**


A curved arrow always begins at an electron pair. It ends at an atom or a bond.



A **B**


Resonance structures **A** and **B** differ in the location of two electron pairs, so two curved arrows are needed. To convert **A** to **B,** take the lone pair on N and form a double bond between C and N. Then, move an electron pair in the C−O double bond to form a lone pair on O. Curved arrows thus show how to reposition the electrons in converting one resonance form to another. **The electrons themselves do not actually move.** Sample Problem 1.4 illustrates the use of curved arrows to convert one resonance structure to another.

Sample Problem 1.4 Follow the curved arrows to draw a second resonance structure for each ion.




Solution

a. The curved arrow tells us to move **one** electron pair in the double bond to the adjacent C – C bond. Then determine the formal charge on any atom whose bonding is different.




Positively charged carbon atoms are called **carbocations.** Carbocations are unstable intermediates because they contain a carbon atom that is lacking an octet of electrons.

b. **Two** curved arrows tell us to move **two** electron pairs. The second resonance structure has a formal charge of (–1) on O.




This type of resonance-stabilized anion is called an **enolate anion.** Enolates are important intermediates in many organic reactions, and all of Chapters 23 and 24 is devoted to their preparation and reactions.

Problem 1.12 Follow the curved arrows to draw a second resonance structure for each species.




Problem 1.13 Use curved arrow notation to show how the first resonance structure can be converted to the second.




Two resonance structures can have exactly the same kinds of bonds, as they do in the carbocation in Sample Problem 1.4a, or they may have different types of bonds, as they do in the enolate in Sample Problem 1.4b. Either possibility is fine as long as the individual resonance structures are valid Lewis structures.

A resonance structure can have an atom with *fewer* than eight electrons around it. **B** is a resonance structure of **A** even though the carbon atom is surrounded by only six electrons.



valid resonance structure

In contrast, a resonance structure can *never* have a second-row element with more than eight electrons. **C** is *not* a resonance structure of **A** because the carbon atom is now surrounded by 10 electrons.




not a valid resonance structure

We will learn much more about resonance in Chapter 16.

The ability to draw and manipulate resonance structures is a necessary skill that will be used throughout your study of organic chemistry. With practice, you will begin to recognize certain common bonding patterns for which more than one Lewis structure can be drawn. For instance, both the carbocation in Sample Problem 1.4a and the enolate anion in Sample Problem 1.4b are specific examples of one general type of resonance observed in certain three-atom systems.

- **In a group of three atoms having a multiple bond X=Y joined to an atom Z having a *p* orbital with zero, one, or two electrons, two resonance structures can be drawn.**




The * corresponds to a charge, a lone pair, or a single electron.

* = +, –, ·, or :

Recall from the Prologue that a ***heteroatom* is an atom other than carbon or hydrogen.**

X, Y, and Z may all be carbon atoms or they may be **heteroatoms** such as nitrogen or oxygen. The atom Z can be charged (positive or negative) or neutral (with a lone pair or a single electron), corresponding to the [*] in the general structure X=Y–Z*. The two resonance structures differ in the location of the multiple bond and the [*].


In the enolate anion in Sample Problem 1.4b, X corresponds to oxygen and [*] is a lone pair, which gives carbon a net negative charge. Moving the double bond and the lone pair and readjusting charges gives the second resonance structure.



- The position of the double bond changes.
- The location of a lone pair changes.

In Chapter 16, we learn more about the orbitals involved in this type of resonance.


Sample Problem 1.5 Draw a second resonance structure for acetamide.



acetamide

Solution

Always look for a three-atom system that contains a multiple bond joined to an atom Z with zero, one, or two electrons in a *p* orbital. Move the double bond (from X = Y to Y = Z) and move the [*] from Z to X. Recalculate formal charges on X and Z.




acetamide
- C=O
- lone pair on N

second resonance structure
- C=N
- additional lone pair on O

In this example, the three-atom system for resonance (X = Y − Z*) is O = C − N with a lone pair on N. After moving the double bond and the lone pair, the formal charges on O and N are –1 and +1, respectively, calculated using the procedure for determining formal charges.

Problem 1.14 Draw a second resonance structure for each species in parts (a) and (b). Draw two additional resonance structures for the ion in part (c).



1.6C The Resonance Hybrid

The **resonance hybrid** is the composite of all possible resonance structures. In the resonance hybrid, the electron pairs drawn in different locations in individual resonance structures are ***delocalized.***

- **The resonance hybrid is more stable than any resonance structure because it delocalizes electron density over a larger volume.**

What does the hybrid look like? When all resonance forms are identical, as they were in the carbocation in Sample Problem 1.4a, each resonance form contributes **equally** to the hybrid.

When two resonance structures are different, the hybrid looks more like the "better" resonance structure. The "better" resonance structure is called the **major contributor** to the hybrid, and all others are **minor contributors.** The hybrid is the weighted average of the contributing resonance structures. What makes one resonance structure "better" than another? There are many factors, but for now, we will learn one fact.

- **A "better" resonance structure is one that has *more bonds* and *fewer charges.***



X
more bonds
fewer charges


major contributor

Y

minor contributor

Comparing resonance structures **X** and **Y, X** is the major contributor because it has more bonds and fewer charges. Thus, the hybrid looks more like **X** than **Y.**

How can we draw a hybrid, which has delocalized electron density? First, we must determine what is different in the resonance structures. Two differences commonly seen are the **position of a multiple bond** and the **site of a charge.** The anion $(HCONH)^-$ illustrates two conventions for drawing resonance hybrids.



A **B**

individual resonance structures

resonance hybrid

- The (–) charge is delocalized on N and O.
- The double bond is delocalized between O, C, and N.


Common symbols and conventions used in organic chemistry are listed on the inside back cover.

- **Double bond position.** Use a dashed line for a bond that is single in one resonance structure and double in another.
- **Location of charge.** Use a δ– (partial negative charge) or δ+ (partial positive charge) for an atom that is neutral in one resonance structure and charged in another.

The hybrid for $(HCONH)^-$ shows two dashed bonds, indicating that both the C–O and C–N bonds have partial double bond character. Both the O and N atoms bear a partial negative charge (δ–) because these atoms are neutral in one resonance structure and negatively charged in the other.

This discussion of resonance is meant to serve as an introduction only. You will learn many more facets of resonance theory in later chapters. In Chapter 2, for example, the enormous effect of resonance on acidity is discussed.

Problem 1.15 Label the resonance structures in each pair as major, minor, or equal contributors to the hybrid. Then draw the hybrid.



Problem 1.16 (a) Draw a second resonance structure for **A.** (b) Why can't a second resonance structure be drawn for **B?**



A **B**

1.7 Determining Molecular Shape


Consider the H_2O molecule. The Lewis structure tells us which atoms are connected to each other, but it implies nothing about the geometry. What does the overall molecule look like? Is H_2O a bent or linear molecule? Two variables define a molecule's structure: **bond length** and **bond angle.**

1.7A Bond Length

Although the SI unit for bond length is the picometer (pm), the angstrom (Å) is still widely used in the chemical literature; 1 Å = 10^{-10} m. As a result, 1 pm = 10^{-2} Å, and 95.8 pm = 0.958 Å.

***Bond length* is the average distance between the centers of two bonded nuclei.** Bond lengths are typically reported in picometers (pm), where 1 pm = 10^{-12} m. For example, the O−H bond length in H_2O is 95.8 pm. Average bond lengths for common bonds are listed in Table 1.2.

- Bond length *decreases* across a row of the periodic table as the size of the atom *decreases.*



- Bond length *increases* down a column of the periodic table as the size of an atom *increases.*




Table 1.2 Average Bond Lengths

Bond	Length (pm)	Bond	Length (pm)	Bond	Length (pm)
H−H	74	H−F	92	C−F	133
C−H	109	H−Cl	127	C−Cl	177
N−H	101	H−Br	141	C−Br	194
O−H	96	H−I	161	C−I	213

1.7B Bond Angle

Bond angle determines the shape around any atom bonded to two other atoms. To determine the bond angle and shape around a given atom, first count how many groups surround the atom. **A group is either an atom or a lone pair of electrons.** Then use the **valence shell electron pair repulsion (VSEPR) theory** to determine the shape. VSEPR is based on the fact that electron pairs repel each other; thus:

- The most stable arrangement keeps these groups as far away from each other as possible.

A second-row element has only three possible arrangements, defined by the number of groups surrounding it.


To determine geometry: [1] Draw a valid Lewis structure; [2] count groups around a given atom.

Number of groups	Geometry	Bond angle
• two groups	linear	180°
• three groups	trigonal planar	120°
• four groups	tetrahedral	109.5°

Let's examine several molecules to illustrate this phenomenon. We first need a valid Lewis structure, and then we count groups around a given atom to predict its geometry.

Two Groups Around an Atom

Any atom surrounded by only two groups is linear and has a bond angle of 180°. Two examples illustrating this geometry are **BeH_2** (beryllium hydride) and **HC≡CH** (acetylene). We consider each carbon atom in acetylene *separately.* Because each C is surrounded by two atoms and no lone pairs, each H−C−C bond angle in acetylene is 180°, and therefore all four atoms are linear.



two groups **linear** molecule

two groups **linear** carbons

Most students in organic chemistry find that building models helps them visualize the shape of molecules. Invest in a set of models *now*.

Common element colors are also shown on the inside back cover.


Acetylene illustrates another important feature: ***ignore multiple bonds in predicting geometry.* Count only atoms and lone pairs.**

We will represent molecules with models having balls for atoms and sticks for bonds, as in the ball-and-stick model of acetylene just shown. These representations are analogous to a set of molecular models. Balls are color-coded using accepted conventions: carbon (black), hydrogen (white or gray), oxygen (red), and so forth, as shown.



Three Groups Around an Atom

Any atom surrounded by three groups is trigonal planar and has bond angles of 120°. Two examples illustrating this geometry are **BF_3** (boron trifluoride) and **$CH_2=CH_2$** (ethylene). All three B−F bonds lie in one plane. *Each* carbon atom of ethylene is surrounded by three atoms and no lone pairs, making *each* H−C−C bond angle 120°. All six atoms of $CH_2=CH_2$ lie in one plane.




three groups **trigonal planar** molecule

three groups **trigonal planar** carbons

Four Groups Around an Atom


Any atom surrounded by four groups is tetrahedral and has bond angles of approximately 109.5°. The simple organic compound methane, **CH_4,** has a central carbon atom with bonds to four hydrogen atoms, each pointing to a corner of a tetrahedron. This arrangement keeps four groups farther apart than a square planar arrangement in which all bond angles would be only 90°.



four groups **tetrahedral** molecule

How can we represent the three-dimensional geometry of a tetrahedron on a two-dimensional piece of paper? **Place two of the bonds in the plane of the paper, one bond in front and one bond behind,** using the following conventions:

- A *solid line* is used for a bond *in* the plane.
- A *wedge* is used for a bond in *front* of the plane.
- A *dashed wedge* is used for a bond *behind* the plane.




This is just one way to draw a tetrahedron for CH_4. We can turn the molecule in many different ways, generating many equivalent representations. All of the following are acceptable drawings for CH_4, because each drawing has two solid lines, one wedge, and one dashed wedge.




Finally, **wedges and dashed wedges are used for groups that are really *aligned one behind another.*** It does not matter in the following two drawings whether the wedge or dashed wedge is skewed to the left or right, because the two H atoms are really aligned as shown in the three-dimensional model.

All carbons in stable molecules are ***tetravalent,*** but the geometry varies with the number of groups around the particular carbon.




- These representations are equivalent.
- The wedge can be skewed to the left or the right of the dashed wedge.

Ammonia (NH_3) and water (H_2O) both have atoms surrounded by four groups, some of which are lone pairs. In **NH_3,** the three H atoms and one lone pair around N point to the corners of a tetrahedron. The H−N−H bond angle of 107° is close to the theoretical tetrahedral bond angle of 109.5°. This molecular shape is referred to as **trigonal pyramidal,** because one of the groups around the N is a nonbonded electron pair, not another atom.



In **H_2O,** the two H atoms and two lone pairs around O point to the corners of a tetrahedron. The H−O−H bond angle of 105° is close to the theoretical tetrahedral bond angle of 109.5°. Water has a **bent** molecular shape, because two of the groups around oxygen are lone pairs of electrons.



In both NH_3 and H_2O the bond angle is somewhat smaller than the theoretical tetrahedral bond angle because of repulsion of the lone pairs of electrons. The bonded atoms are compressed into a smaller space with a smaller bond angle.

Predicting geometry based on counting groups is summarized in Table 1.3.

Table 1.3 Summary: Determining Geometry Based on the Number of Groups


Number of groups around an atom	Geometry	Bond angle	Examples
2	linear	180°	BeH_2, HC≡CH
3	trigonal planar	120°	BF_3, $CH_2=CH_2$
4	tetrahedral	109.5°	CH_4, NH_3, H_2O

Sample Problem 1.6 Determine the geometry around the highlighted atom in each species.

a. :Ö=C=Ö: b. H−N⁺−H (H above and below N)

Solution

a.




two atoms around C
no lone pairs

two groups

linear

b.



four atoms around N
no lone pairs

four groups ⟶ **tetrahedral**

Problem 1.17 Determine the geometry around all second-row elements in each compound drawn as a Lewis structure with no implied geometry.

a. $CH_3-\overset{\overset{:O:}{\|}}{C}-CH_3$ b. $CH_3-\ddot{O}-CH_3$ c. $\ddot{N}H_2^-$ d. $CH_3-C\equiv N:$


Problem 1.18 Predict the indicated bond angles in each compound drawn as a Lewis structure with no implied geometry.

a. $CH_3-C\equiv C-\ddot{C}l:$ b. $CH_2=C(H)-\ddot{C}l:$ c. $CH_3-C(H)-\ddot{C}l:$

Problem 1.19 Using the principles of VSEPR theory, you can predict the geometry around any atom in any molecule, no matter how complex. Enanthotoxin is a poisonous compound isolated from a common variety of hemlock grown in England. Predict the geometry around the highlighted atoms in enanthotoxin.



The hemlock water dropwort is the source of the poison enanthotoxin.



enanthotoxin

1.8 Drawing Organic Structures


Drawing organic molecules presents a special challenge. Because they often contain many atoms, we need shorthand methods to simplify their structures. The two main types of shorthand representations used for organic compounds are **condensed structures** and **skeletal structures.**


1.8A Condensed Structures

Condensed structures can be used for compounds having a chain of atoms bonded together. The following conventions are used:

- All of the atoms are drawn in, but the two-electron bond lines are generally omitted.
- Atoms are usually drawn next to the atoms to which they are bonded.
- Parentheses are used around similar groups bonded to the same atom.
- Lone pairs are omitted.


To interpret a condensed formula, it is usually best to start at the *left side* of the molecule and remember that the ***carbon atoms must be tetravalent.*** A carbon bonded to three H atoms becomes **CH_3;** a carbon bonded to two H atoms becomes **CH_2;** and a carbon bonded to one H atom becomes **CH.**





Other examples of condensed structures with heteroatoms and carbon–carbon multiple bonds are given in Figure 1.3.


Figure 1.3
Examples of condensed structures



- Entry [1]: Draw the H atom next to the C to which it is bonded, and use parentheses around CH_3 to show it is bonded to the carbon chain.
- Entry [2]: Keep the carbon–carbon double bond and draw the H atoms after each C to which they are bonded.
- Entry [3]: Omit the lone pairs on the O atom in the condensed structure.
- Entry [4]: Omit the lone pairs on Cl and O and draw the two CH_2 groups as $(CH_2)_2$.

Translating some condensed formulas is not obvious, and it will come only with practice. This is especially true for compounds containing a carbon–oxygen double bond. Some noteworthy examples in this category are given in Figure 1.4. While carbon–carbon double bonds are generally drawn in condensed structures, carbon–oxygen double bonds are usually omitted.

Figure 1.4
Condensed structures containing a C–O double bond



- In **A,** the H atom is bonded to C, *not* O.
- In **B,** each CH_3 group is bonded to C, *not* O.
- In **C** and **D,** the C atom is doubly bonded to one O and singly bonded to the other O.


Sample Problem 1.7 Convert each condensed formula to a Lewis structure.

a. $(CH_3)_2CHOCH_2CH_2CH_2OH$ b. $CH_3(CH_2)_2CO_2C(CH_3)_3$


Solution

Start at the left and proceed to the right, making sure that each carbon has four bonds. Give each O atom two lone pairs to have an octet.

a. $(CH_3)_2CHOCH_2CH_2CH_2OH$ b. $CH_3(CH_2)_2CO_2C(CH_3)_3$



One C atom (labeled in blue) is bonded to 2 CH_3's, 1 H, and 1 O.



One C atom (labeled in blue) is bonded to both O's.

Problem 1.20 Convert each condensed formula to a Lewis structure.

a. $CH_3(CH_2)_4CH(CH_3)_2$
b. $(CH_3)_3CCH(OH)CH_2CH_3$
c. $(CH_3)_2CHCHO$
d. $(HOCH_2)_2CH(CH_2)_3C(CH_3)_2CH_2CH_3$


Problem 1.21 During periods of strenuous exercise, the buildup of lactic acid [$CH_3CH(OH)CO_2H$] causes the aching feeling in sore muscles. Convert this condensed structure to a Lewis structure of lactic acid.

1.8B Skeletal Structures


Skeletal structures are used for organic compounds containing both rings and chains of atoms. Three rules are used to draw them.

- **Assume a carbon atom is located at the junction of any two lines or at the end of any line.**
- **Assume each carbon has enough hydrogens to make it tetravalent.**
- **Draw in all heteroatoms and the hydrogens directly bonded to them.**

Carbon chains are drawn in a zigzag fashion, and rings are drawn as polygons, as shown for hexane and cyclohexane.




hexane



Lewis structure
C_6H_{12}


Each C is bonded to 2 H's.



skeletal structure
cyclohexane

How To Interpret a Skeletal Structure

Example Draw in all C atoms, H atoms, and lone pairs in the following molecule:




Step [1] **Place a C atom at the intersection of any two lines and at the end of any line.**




- This molecule has six carbons, including the C labeled in red at the left end of the chain.
- There are two C's (labeled in green) between the C = C and the OH group.

Step [2] **Add enough H's to make each C tetravalent.**



- The end C labeled in red needs three H's to be tetravalent.
- Each C on the C = C has three bonds already, so only one H must be drawn.
- There are two CH_2 groups between the C = C and the OH group.


Step [3] **Add lone pairs to give each heteroatom an octet.**



Each O needs **two** lone pairs for an octet.

Figure 1.5 shows other examples of skeletal structures, and Sample Problem 1.8 illustrates how to interpret the skeletal structure for a more complex cyclic compound.

Figure 1.5
Interpreting skeletal structures




- C labeled in red needs 3 H's.
- C's labeled in blue need 1 H.
- All other C's need 2 H's.
- N needs one lone pair.




- C's labeled in red need 3 H's.
- C's labeled in green have four bonds to C.

Sample Problem 1.8 Draw a complete structure for vanillin showing all C atoms, H atoms, and lone pairs, and give the molecular formula. Vanillin is the principal component of the extract of the vanilla bean.




vanillin




vanilla bean

Solution

Skeletal structures have a C atom at the junction of any two lines and at the end of any line. Each C must have enough H's to make it tetravalent. Each O atom needs two lone pairs to have a complete octet.



=




- C in blue has three H's.
- C in green has four bonds to other C's.
- C in red is doubly bonded to O.

$C_8H_8O_3$


Problem 1.22 How many hydrogen atoms are present around each highlighted carbon atom in the following molecules? What is the molecular formula for each molecule? Both compounds are active ingredients in some common sunscreens.

a.



octinoxate
(2-ethylhexyl 4-methoxycinnamate)

b.




avobenzone




Problem 1.23 Convert each skeletal structure to a complete structure with all C's, H's, and lone pairs drawn in.

a. 


b. 

c. 


d. 

Problem 1.24 What is the molecular formula of quinine, the antimalarial medication whose skeletal structure appears in Figure 2 of the Prologue?

When heteroatoms are bonded to a carbon skeleton, the heteroatom is joined *directly* to the carbon to which it is bonded, with no H atoms in between. Thus, an OH group is drawn as OH or HO depending on where the OH is located. In contrast, when carbon appendages are bonded to a carbon skeleton, the H atoms will be drawn to the *right* of the carbon to which they are bonded regardless of the location.



Place the O and N atoms directly joined to the ring.



Two C atoms in red are bonded to the middle C.





Two C atoms in red are bonded to the ring.

1.8C Skeletal Structures with Charged Carbon Atoms


Take care in interpreting skeletal structures for positively and negatively charged carbon atoms, because *both* the hydrogen atoms *and* the lone pairs are omitted. Keep in mind the following:

- **A charge on a carbon atom takes the place of one hydrogen atom.**
- **The charge determines the number of lone pairs. Negatively charged carbon atoms have one lone pair and positively charged carbon atoms have none.**






Skeletal structures often leave out lone pairs on heteroatoms, but *don't forget about them.* Use the formal charge on an atom to determine the number of lone pairs. For example, a neutral O atom with two bonds needs two additional lone pairs, and a positively charged O atom with three bonds needs only one lone pair.





neutral O atom
two lone pairs





positively charged O atom
one lone pair

Problem 1.25 Draw in all hydrogens and lone pairs on the charged carbons in each ion.

a. 


b. 


c. 


d. 

Problem 1.26 Draw a skeletal structure for the molecules in parts (a) and (b), and a condensed structure for the molecules in parts (c) and (d).

a. $(CH_3)_2C{=}CH(CH_2)_4CH_3$

b. 

c. 


d. 

1.9 Hybridization

What orbitals do the first- and second-row atoms use to form bonds?

1.9A Hydrogen

Recall from Section 1.2 that two hydrogen atoms share each of their electrons to form H_2. Thus, the 1*s* orbital on one H overlaps with the 1*s* orbital on the other H to form a bond that concentrates electron density between the two nuclei. This type of bond, called a **σ** (sigma) **bond,** is cylindrically symmetrical because the electrons forming the bond are distributed symmetrically about an imaginary line connecting the two nuclei.




- A σ bond concentrates electron density on the axis that joins two nuclei. All single bonds are σ bonds.

1.9B Bonding in Methane

To account for the bonding patterns observed in more complex molecules, we must take a closer look at the 2*s* and 2*p* orbitals of atoms of the second row. Let's illustrate this with methane, CH_4.

Carbon has **four valence electrons.** To fill atomic orbitals in the most stable arrangement, electrons are placed in the orbitals of lowest energy. For carbon, this places two electrons in the 2*s* orbital and one each in two 2*p* orbitals.




ground state for
carbon's four valence electrons

- This lowest energy arrangement of electrons for an atom is called its **ground state.**

In this description, **carbon should form *only two bonds*** because it has only two unpaired valence electrons, and CH_2 should be a stable molecule. In reality, however, CH_2 is a highly reactive species because carbon does not have an octet of electrons.

Alternatively, promotion of an electron from a 2*s* to a vacant 2*p* orbital would form **four** unpaired electrons for bonding. This process requires energy because it moves an electron to a higher energy orbital. This higher energy electron configuration is called an electronically **excited state.**




In this description carbon would form two different types of bonds: three with 2*p* orbitals and one with a 2*s* orbital. **But experimental evidence points to carbon forming *four identical bonds* in methane.**


To solve this dilemma, chemists have proposed that atoms like carbon do not use pure *s* and pure *p* orbitals in forming bonds. Instead, atoms use a set of new orbitals called **hybrid orbitals.** The mathematical process by which these orbitals are formed is called **hybridization.**

- *Hybridization* is the combination of two or more atomic orbitals to form the same number of hybrid orbitals, each having the same shape and energy.


Hybridization of one 2*s* orbital and three 2*p* orbitals for carbon forms four hybrid orbitals, each with one electron. These new hybrid orbitals are intermediate in energy between the 2*s* and 2*p* orbitals.



- These hybrid orbitals are called sp^3 *hybrids* because they are formed from *one s* orbital and *three p* orbitals.



What do these new hybrid orbitals look like? Mixing a spherical 2*s* orbital and three dumbbell-shaped 2*p* orbitals together produces four orbitals having one large lobe and one small lobe, oriented toward the corners of a tetrahedron. Each large lobe concentrates electron density in the bonding direction between two nuclei. This makes bonds formed from hybrid orbitals ***stronger*** than bonds formed from pure *p* orbitals.




The four hybrid orbitals form four equivalent bonds. We can now explain the observed bonding in CH_4.

- Each bond in CH_4 is formed by overlap of an sp^3 hybrid orbital of carbon with a 1*s* orbital of hydrogen. These four bonds point to the corners of a tetrahedron.

All four C – H bonds in methane are **σ bonds,** because the electron density is concentrated on the axis joining C and H. An orbital picture of the bonding in CH_4 is given in Figure 1.6.

Figure 1.6
Bonding in CH_4 using sp^3 hybrid orbitals



- All four C – H bonds are σ bonds.

Problem 1.27 What orbitals are used to form each of the C – C and C – H bonds in $CH_3CH_2CH_3$ (propane)? How many σ bonds are present in this molecule?


1.9C Other Hybridization Patterns—*sp* and *sp*^{2} Hybrid Orbitals

Forming sp^3 hybrid orbitals is just one way that 2*s* and 2*p* orbitals can hybridize. Three common modes of hybridization are seen in organic molecules. The number of orbitals is always conserved in hybridization; that is, a **given number of atomic orbitals hybridizes to form an equivalent number of hybrid orbitals.**

- *One* 2*s* orbital and *three* 2*p* orbitals form *four* sp^3 hybrid orbitals.
- *One* 2*s* orbital and *two* 2*p* orbitals form *three* sp^2 hybrid orbitals.
- *One* 2*s* orbital and *one* 2*p* orbital form *two sp* hybrid orbitals.


We have already seen pictorially how four sp^3 hybrid orbitals are formed from one 2*s* and three 2*p* orbitals. Figures 1.7 and 1.8 illustrate the same process for *sp* and sp^2 hybrids. Each *sp* and sp^2 hybrid orbital has one large and one small lobe, much like an sp^3 hybrid orbital. Note, however, that both sp^2 and *sp* hybridization **leave one and two 2*p* orbitals *unhybridized,*** respectively, on each atom.

Figure 1.7
Forming two *sp* hybrid orbitals



- Forming **two *sp* hybrid orbitals** uses **one 2*s*** and **one 2*p* orbital,** leaving **two 2*p* orbitals *unhybridized.***

Figure 1.8
Forming three sp^2 hybrid orbitals



- Forming **three sp^2 hybrid orbitals** uses **one 2*s*** and **two 2*p* orbitals,** leaving **one 2*p* orbital *unhybridized.***

The **superscripts** for hybrid orbitals correspond to the **number of atomic orbitals** used to form them. The number "1" is understood.

For example: $sp^3 = s^1p^3$

one 2*s* + **three** 2*p* orbitals used to make each hybrid orbital

To determine the hybridization of an atom in a molecule, we count groups (atoms and lone pairs) around the atom, just as we did in determining geometry.

- **The number of groups around an atom equals the number of atomic orbitals that are hybridized to form hybrid orbitals (Table 1.4).**


Table 1.4 Three Types of Hybrid Orbitals

Number of groups	Number of orbitals used	Type of hybrid orbital
2	2	two *sp* hybrid orbitals
3	3	three sp^2 hybrid orbitals
4	4	four sp^3 hybrid orbitals


Let's illustrate this phenomenon with BeH_2, BF_3, NH_3, and H_2O, whose geometry was determined by counting groups in Section 1.7.

In **BeH_2,** the **Be atom is *sp* hybridized** because it is surrounded by two groups (two H atoms). Each Be−H bond is formed by overlap of an *sp* hybrid orbital from Be and a 1*s* orbital from H. The *sp* hybrid orbitals are oriented 180° away from each other.

In **BF_3,** the **B atom is sp^2 hybridized** because it is surrounded by three groups (three F atoms). Each B−F bond is formed by overlap of an sp^2 hybrid orbital from B and a 2*p* orbital from F. The sp^2 hybrid orbitals all lie in a plane, and are oriented 120° apart. The B atom also has a vacant unhybridized 2*p* orbital. This orbital is located ***above and below the plane*** of the BF_3 molecule.




two Be−H bonds



BF_3 from above

The three B−F bonds all lie in a plane, 120° apart.




BF_3 from the side

The **unhybridized *p* orbital** extends above and below the plane.

The N atom in **NH_3** and the O atom in **H_2O** are both surrounded by four groups, making them sp^3 hybridized. Each N−H and O−H bond in these molecules is formed by overlap of an sp^3 hybrid orbital with a 1*s* orbital from H. The lone pairs of electrons on N and O also occupy sp^3 hybrid orbitals, as shown in Figure 1.9.


Figure 1.9
Hybrid orbitals of NH_3 and H_2O



Sample Problem 1.9 What orbitals are used to form each bond in methanol, CH_3OH?

Solution

To solve this problem, draw a valid Lewis structure and count groups around each atom. Then, use the rule to determine hybridization: two groups = *sp,* three groups = sp^2, and four groups = sp^3.




- All C–H bonds are formed from C_{sp^3}–H_{1s}.
- The C–O bond is formed from C_{sp^3}–O_{sp^3}.
- The O–H bond is formed from O_{sp^3}–H_{1s}.

Problem 1.28 What orbitals are used to form each bond in the following molecules?

a. CH_3BeH b. $(CH_3)_3B$ c. CH_3OCH_3

1.10 Ethane, Ethylene, and Acetylene

The principles of hybridization determine the type of bonds in **ethane, ethylene,** and **acetylene.**




1.10A Ethane—CH_3CH_3

According to the Lewis structure for **ethane, CH_3CH_3,** each carbon atom is singly bonded to four other atoms. As a result:


- **Each carbon is tetrahedral.**
- **Each carbon is sp^3 hybridized.**




Ethane is a constituent of natural gas.



All of the bonds in ethane are σ bonds. The C–H bonds are formed from the overlap of one of the three sp^3 hybrid orbitals on each carbon atom with the 1*s* orbital on hydrogen. The C–C bond is formed from the overlap of an sp^3 hybrid orbital on each carbon atom.



A model of ethane shows that **rotation can occur around the central C−C σ bond.** The relative position of the H atoms on the adjacent CH_3 groups changes with bond rotation, as seen in the location of the labeled red H atom before and after rotation. This process is discussed in greater detail in Chapter 4.




1.10B Ethylene—C_2H_4




Ethylene is an important starting material in the preparation of the plastic polyethylene.

Based on the Lewis structure of **ethylene, $CH_2{=}CH_2$,** each carbon atom is singly bonded to two H atoms and doubly bonded to the other C atom, so each C is surrounded by three groups. As a result:


- **Each carbon is trigonal planar (Section 1.7B).**
- **Each carbon is sp^2 hybridized.**



What orbitals are used to form the two bonds of the C−C double bond? Recall from Section 1.9 that ***sp*² hybrid orbitals** are formed from **one 2*s* and two 2*p* orbitals,** leaving one **2*p* orbital unhybridized.** Because carbon has four valence electrons, **each of these orbitals has one electron** that can be used to form a bond.



Each C−H bond results from the end-on overlap of an sp^2 hybrid orbital on carbon and the 1*s* orbital on hydrogen. Similarly, one of the C−C bonds results from the end-on overlap of an sp^2 hybrid orbital on each carbon atom. Each of these bonds is a **σ bond.** All five σ bonds lie in the same plane, viewed from above in the following representation, and from the side in Figure 1.10a.




- Each C has three sp^2 hybrid orbitals.
- The C−H bonds and the C−C bond are σ bonds.

The second C−C bond results from the side-by-side overlap of the 2*p* orbitals on each carbon. Side-by-side overlap creates an area of electron density above and below the plane containing the

sp^2 hybrid orbitals (that is, the plane containing the six atoms in the σ bonding system), as shown in Figure 1.10b.


Figure 1.10
The σ and π bonds in ethylene

a. 2*p* orbitals



- Overlap of the two sp^2 hybrid orbitals forms the C−C σ bond.

b. π bond



- Overlap of the two 2*p* orbitals forms the C−C π bond.
- The π bond extends **above and below the plane** of the molecule.

In this second bond, the electron density is ***not*** concentrated on the axis joining the two nuclei. This new type of bond is called a **π bond.** Because the electron density in a π bond is farther from the two nuclei, **π bonds are usually weaker and therefore more easily broken than σ bonds.**

Thus, a carbon–carbon double bond has two components:

- a σ bond, formed by end-on overlap of two sp^2 hybrid orbitals;
- a π bond, formed by side-by-side overlap of two 2*p* orbitals.

Unlike the C−C single bond in ethane, rotation about the C−C double bond in ethylene is **restricted.** It can occur only if the π bond first breaks and then re-forms, a process that requires considerable energy.

All double bonds are composed of one σ and one π bond.

Rotation around a C=C bond does *not* occur.




1.10C Acetylene—C_2H_2



Because acetylene produces a very hot flame on burning, it is often used in welding torches. The fire is very bright, too, so it was once used in the lamps worn by spelunkers—people who study and explore caves.

Based on the Lewis structure of **acetylene, HC≡CH,** each carbon atom is singly bonded to one hydrogen atom and triply bonded to the other carbon atom, so each carbon atom is surrounded by two groups. As a result:


- Each carbon is linear (Section 1.7B).
- Each carbon is *sp* hybridized.




acetylene

What orbitals are used to form the bonds of the C−C triple bond? Recall from Section 1.9 that ***sp* hybrid orbitals** are formed from **one 2*s* and one 2*p* orbital,** leaving **two 2*p* orbitals**

unhybridized. Because carbon has four valence electrons, **each of these orbitals has one electron** that can be used to form a bond.



Each C−H bond results from the end-on overlap of an *sp* hybrid orbital on carbon and the 1*s* orbital on hydrogen. Similarly, one of the C−C bonds results from the end-on overlap of an *sp* hybrid orbital on each carbon atom. Each of these bonds is a **σ bond.**




- Each C has two *sp* hybrid orbitals.
- The C−H bonds and C−C bond are σ bonds.

Each carbon atom also has two **unhybridized 2*p* orbitals** that are perpendicular to each other and to the *sp* hybrid orbitals (Figure 1.11a). Side-by-side overlap between the two 2*p* orbitals on one carbon with the two 2*p* orbitals on the other carbon creates the second and third bonds of the C−C triple bond (Figure 1.11b). The electron density from one of these two bonds is above and below the axis joining the two nuclei, and the electron density from the second of these two bonds is in front of and behind the axis, so both of these bonds are **π bonds.**


Figure 1.11
The σ and π bonds in acetylene

a. 2*p* orbitals



- Overlap of the two *sp* hybrid orbitals forms the C−C σ bond.

b. one π bond



- Overlap of two sets of two 2*p* orbitals forms two C−C π bonds.
- Two π bonds extend out from the axis of the linear molecule.

The side-by-side overlap of two *p* orbitals always forms a π bond.




All triple bonds are composed of one σ and two π bonds.

Thus, a carbon–carbon triple bond has three components:


- a σ bond, formed by end-on overlap of two *sp* hybrid orbitals;
- two π bonds, formed by side-by-side overlap of two sets of 2*p* orbitals.

Table 1.5 summarizes the three possible types of bonding in carbon compounds.

Table 1.5 A Summary of Covalent Bonding Seen in Carbon Compounds

Number of groups bonded to C	Hybridization	Bond angle	Example	Observed bonding
4	sp^3	109.5°	**CH_3CH_3** ethane	 **one σ bond** C_{sp^3}–C_{sp^3}
3	sp^2	120°	**$CH_2{=}CH_2$** ethylene	 **one σ bond** + **one π bond** C_{sp^2}–C_{sp^2} C_{2p}–C_{2p}
2	sp	180°	**HC≡CH** acetylene	 **one σ bond** + **two π bonds** C_{sp}–C_{sp} C_{2p}–C_{2p} C_{2p}–C_{2p}


Sample Problem 1.10 Answer each question for the molecule acetone.



acetone

a. Determine the hybridization of the highlighted atoms.
b. What orbitals are used to form the C–O double bond?
c. In what type of orbital does each lone pair reside?


Solution




b. • The σ bond is formed from the end-on overlap of C_{sp^2}–O_{sp^2}.
• The π bond is formed from the side-by-side overlap of C_{2p}–O_{2p}.

c. The O atom has three sp^2 hybrid orbitals.
• One is used for the σ bond of the double bond.
• The remaining two sp^2 hybrids are occupied by the lone pairs.

Problem 1.29 Determine the hybridization around the highlighted atoms in each molecule.

a. $CH_3-C\equiv CH$ b.  c. $CH_2=C=CH_2$

Problem 1.30 The unmistakable odor of a freshly cut cucumber is largely due to cucumber aldehyde. (a) How many sp^2 hybridized carbon atoms does cucumber aldehyde contain? (b) What is the hybridization of the O atom? (c) What orbitals are used to form the carbon–oxygen double bond? (d) How many σ bonds does cucumber aldehyde contain? (e) How many π bonds does it contain?




cucumber aldehyde

1.11 Bond Length and Bond Strength

Let's now examine the relative bond length and bond strength of the C−C and C−H bonds in ethane, ethylene, and acetylene.

1.11A A Comparison of Carbon–Carbon Bonds

While the SI unit of energy is the **joule** (J), organic chemists often report energy values in **calories** (cal). For this reason, energy values in the tables in this text are reported in joules, followed by the number of calories in parentheses.
1 cal = 4.18 J



- As the number of electrons between two nuclei *increases,* bonds become shorter and stronger.
- Triple bonds are shorter and stronger than double bonds, which are shorter and stronger than single bonds.

An inverse relationship exists between bond length and bond strength. The shorter the bond, the closer the electron density is kept to the nucleus, and the harder the bond is to break. ***Shorter* bonds are *stronger* bonds.**

Values for bond lengths and bond strengths for CH_3CH_3, $CH_2{=}CH_2$, and $HC{\equiv}CH$ are listed in Table 1.6. Be careful not to confuse two related but different principles regarding multiple bonds such as C−C double bonds. **Double bonds, consisting of both a σ and a π bond, are *strong*. The π component** of the double bond, however, is usually much ***weaker* than the σ component.** This is a particularly important consideration when studying alkenes in Chapter 10.

Table 1.6 Bond Lengths and Bond Strengths for Ethane, Ethylene, and Acetylene

Compound	C−C bond length (pm)	Bond strength kJ/mol (kcal/mol)
$CH_3{-}CH_3$	153	368 (88)
$CH_2{=}CH_2$	134	635 (152)
$HC{\equiv}CH$	121	837 (200)


Increasing bond length ↑ Increasing bond strength ↓

Compound	C−H bond length (pm)	Bond strength kJ/mol (kcal/mol)
$CH_3CH_2{-}H$	111	410 (98)
$CH_2{=}C(H)H$	110	435 (104)
$HC{\equiv}C{-}H$	109	523 (125)

Increasing bond length ↑ Increasing bond strength ↓

1.11B A Comparison of Carbon–Hydrogen Bonds

The length and strength of a C−H bond vary slightly depending on the hybridization of the carbon atom.



To understand why this is so, we must look at the atomic orbitals used to form each type of hybrid orbital. A single 2*s* orbital is always used, but the number of 2*p* orbitals varies with the type of hybridization. The **percent *s*-character** indicates the fraction of a hybrid orbital due to the 2*s* orbital used to form it.


***sp* hybrid**	$\frac{\text{one 2}s\text{ orbital}}{\text{two hybrid orbitals}}$	**= 50% *s*-character**
***sp*^{2} hybrid**	$\frac{\text{one 2}s\text{ orbital}}{\text{three hybrid orbitals}}$	**= 33% *s*-character**
***sp*^{3} hybrid**	$\frac{\text{one 2}s\text{ orbital}}{\text{four hybrid orbitals}}$	**= 25% *s*-character**


Why should the percent *s*-character of a hybrid orbital affect the length of a C−H bond? A 2*s* orbital keeps electron density closer to a nucleus compared to a 2*p* orbital. As the **percent *s*-character increases,** a hybrid orbital holds its electrons closer to the nucleus, and the **bond becomes shorter and stronger.**


- **Increased percent *s*-character ⟶ Increased bond strength ⟶ Decreased bond length**

Problem 1.31 Which of the bonds shown in red in each compound or pair of compounds is shorter?

a. 

b. 

c. 

d. 


1.12 Electronegativity and Bond Polarity

***Electronegativity* is a measure of an atom's attraction for electrons in a bond.** Electronegativity indicates how much a particular atom "*wants*" electrons.

- Electronegativity *increases* across a row of the periodic table as the nuclear charge increases (excluding the noble gases).
- Electronegativity *decreases* down a column of the periodic table as the atomic radius increases, pushing the valence electrons farther from the nucleus.

As a result, the *most* electronegative elements are located at the **upper right-hand corner** of the periodic table, and the *least* electronegative elements in the **lower left-hand corner.** A scale has been established to represent electronegativity values arbitrarily, from 0 to 4, as shown in Figure 1.12.

Figure 1.12
Electronegativity values for some common elements




Electronegativity values are relative, so they can be used for comparison purposes only. When comparing two different elements, one is **more electronegative** than the other if it attracts electron density toward itself. One is less electronegative—**more electro*positive***—if it gives up electron density to the other element.


Problem 1.32 Rank the following atoms in order of increasing electronegativity. Label the most electronegative and most electropositive atom in each group.

a. Se, O, S b. P, Na, Cl c. Cl, S, F d. O, P, N

Electronegativity values are used as a guideline to indicate whether the electrons in a bond are **equally shared** or **unequally shared** between two atoms. Whenever two identical atoms are bonded together, each atom attracts the electrons in the bond to the same extent. The electrons are equally shared, and the **bond is *nonpolar.*** Thus, a **carbon–carbon bond is nonpolar.** Whenever two different atoms having similar electronegativities are bonded together, the bond is also **nonpolar. C − H bonds are considered to be nonpolar,** because the electronegativity difference between C (2.5) and H (2.2) is small.



nonpolar bond




nonpolar bond

The small electronegativity difference between C and H is ignored.

Bonding between atoms of different electronegativity values results in the **unequal sharing** of electrons. In a C − O bond, the electrons are pulled away from C (2.5) toward O (3.4), the element of higher electronegativity. **The bond is *polar,*** or ***polar covalent.*** The bond is said to have a **dipole;** that is, **a partial separation of charge.**

A C − O bond is a **polar** bond.





a bond dipole

The direction of polarity in a bond is often indicated by an arrow, with the head of the arrow pointing toward the more electronegative element. The tail of the arrow, with a perpendicular line drawn through it, is positioned at the less electronegative element. Alternatively, the symbols δ+ and δ– indicate this unequal sharing of electron density.

- **δ+ means an atom is electron deficient (has a partial positive charge).**
- **δ– means the atom is electron rich (has a partial negative charge).**

Problem 1.33 Show the direction of the dipole in each bond. Label the atoms with δ+ and δ–.


a. H–F b.  c.  d. 

Students often wonder how large an electronegativity difference must be to consider a bond polar. That's hard to say. We will set an arbitrary value for this difference and use it as an *approximation.* **Usually, a polar bond will be one in which the electronegativity difference between two atoms is ≥ 0.5 units.**

The distribution of electron density in a molecule can be shown using an **electrostatic potential map.** These maps are color coded to illustrate areas of high and low electron density. Electron-rich regions are indicated in red, and electron-deficient sites are indicated in blue. Regions of intermediate electron density are shown in orange, yellow, and green.

An electrostatic potential map of CH_3Cl indicates the polar nature of the C–Cl bond (Figure 1.13). The more electronegative Cl atom pulls electron density toward it, making it electron rich. This is indicated by the red around the Cl in the plot. The carbon is electron deficient, and this is shown with blue. When comparing two maps, the comparison is useful only if they are plotted *using the same scale* of color gradation. For this reason, whenever we compare two plots in this text, they will be drawn side by side using the same scale.

Figure 1.13
Electrostatic potential plot of CH_3Cl



1.13 Polarity of Molecules

A **polar molecule** has either one polar bond, or two or more bond dipoles that reinforce. A **nonpolar molecule** has either no polar bonds, or two or more bond dipoles that cancel.

Whenever C or H is bonded to N, O, and all halogens, the bond is ***polar.*** Thus, the C–I bond is considered polar even though the electronegativity difference between C and I is small. Remember, electronegativity is just an approximation.

Thus far, we have been concerned with the polarity of one bond. To determine whether a molecule has a net dipole, use the following two-step procedure:

[1] Use electronegativity differences to **identify all of the polar bonds and the directions of the bond dipoles.**

[2] **Determine the geometry** around individual atoms by counting groups, and decide if individual dipoles **cancel** or **reinforce each other in space.**

The two molecules **H_2O** and **CO_2** illustrate different outcomes of this process. In H_2O, each O–H bond is polar because the electronegativity difference between O (3.4) and H (2.2) is large. Since H_2O is a **bent** molecule, the two dipoles reinforce (both point *up*). Thus, **H_2O has a net dipole, making it a polar molecule.** CO_2 also has polar C–O bonds because the electronegativity difference between O (3.4) and C (2.5) is large. However, CO_2 is a **linear** molecule, so the two dipoles, which are equal and opposite in direction, **cancel.** Thus, CO_2 is a **nonpolar molecule** with **no net dipole.**



Electrostatic potential plots for H_2O and CO_2 appear in Figure 1.14. Additional examples of polar and nonpolar molecules are given in Figure 1.15.

Problem 1.34 Indicate which of the following molecules is polar because it possesses a net dipole. Show the direction of the net dipole if one exists.





a. CH_3Br b. CH_2Br_2 c. CF_4 d.  e. 

Figure 1.14
Electrostatic potential plots for H_2O and CO_2




- The electron-rich (red) region is concentrated on the more electronegative O atom. Both H atoms are electron deficient (blue-green).



- Both electronegative O atoms are electron rich (red) and the central C atom is electron deficient (blue).

Figure 1.15
Examples of polar and nonpolar molecules



one polar bond — a **polar** molecule

three polar bonds
All dipoles cancel.
no net dipole — a **nonpolar** molecule

three polar bonds
All dipoles reinforce. — a **polar** molecule


two polar bonds
Both dipoles reinforce. — a **polar** molecule

four polar bonds
All dipoles cancel.
no net dipole — a **nonpolar** molecule

1.14 L-Dopa—A Representative Organic Molecule

The principles learned in this chapter apply to all organic molecules regardless of size or complexity. We now know a great deal about the structure of **L-dopa,** a drug used to treat Parkinson's disease described in the chapter opener.

Sample Problem 1.11 Answer each question about L-dopa.



L-dopa


a. Convert the skeletal structure to a Lewis structure.
b. What is the hybridization and geometry around each labeled atom?
c. Label three polar bonds.



Sinemet, the trade name of a drug used to treat Parkinson's disease, contains a combination of L-dopa (Sample Problem 1.11) and carbidopa (Problem 1.36). Carbidopa increases the effectiveness of L-dopa by inhibiting its metabolism prior to crossing the blood–brain barrier and entering the brain.

Solution

a, b. Convert the skeletal structure to a Lewis structure as in Sample Problem 1.8. Each O atom needs two lone pairs and each N atom needs one for an octet. Count groups around an atom to determine hybridization and shape.



Lewis structure of L-dopa


c. All C–O, O–H , N–H, and C–N bonds are polar because the electronegativity difference between the atoms is large.

Problem 1.35 Provide the following information about L-dopa.

a. Label all sp^3 hybridized C atoms.
b. Label all H atoms that bear a partial positive charge (δ+).
c. Draw another resonance structure.

Sample Problem 1.12 illustrates how to derive structural information from a ball-and-stick model.

Sample Problem 1.12 Use the ball-and-stick model of vitamin B_6 to answer each question.




vitamin B_6

a. Draw a skeletal structure of vitamin B_6.
b. How many sp^2 hybridized carbons are present?
c. What is the hybridization of the N atom in the ring?

Solution


Use the element colors shown in Section 1.7B to convert the 3-D model to a skeletal structure [black (C), gray (H), red (O), blue (N)]. H atoms on carbon are omitted, but H atoms on heteroatoms are drawn. Count groups to determine hybridization. Each O atom needs two lone pairs and each N needs one to give an octet of electrons.

a.




skeletal structure of vitamin B_6

b, c.



- Each C labeled in blue is sp^2 hybridized.
- The N atom is surrounded by three groups (two atoms and one lone pair), making it sp^2 hybridized.

Problem 1.36 Use the ball-and-stick model to answer each question about carbidopa, a drug used in combination with L-dopa to treat Parkinson's disease



carbidopa

a. Draw a skeletal structure of carbidopa.
b. Determine the hybridization around each carbon atom.
c. What is the hybridization and geometry around each N atom?
d. How many polar bonds are present?

KEY CONCEPTS

Structure and Bonding

Important Facts

- **The general rule of bonding:** Atoms "strive" to attain a complete outer shell of valence electrons (1.2). H "wants" two electrons. Second-row elements "want" eight electrons.
- **Formal charge** is the difference between the number of valence electrons of an atom and the number of electrons it "owns" (1.3C). See Sample Problem 1.3 for a stepwise example.
- **Curved arrow notation** shows the movement of an electron pair. The tail of the arrow always begins at an electron pair, either in a bond or a lone pair. The head points to where the electron pair "moves" (1.6).
- **Electrostatic potential plots** are color-coded maps of electron density, indicating electron-rich and electron-deficient regions (1.12).

The Importance of Lewis Structures (1.3–1.5)

A properly drawn Lewis structure shows the number of bonds and lone pairs present around each atom in a molecule. In a valid Lewis structure, each H has two electrons, and each second-row element has no more than eight. This is the first step needed to determine many properties of a molecule.

Lewis structure ----→
- **Geometry:** linear, trigonal planar, or tetrahedral (1.7)
- **Hybridization:** *sp*, sp^2, or sp^3 (1.9)
- **Types of bonds:** single, double, or triple (1.3, 1.10)


Resonance (1.6)

The basic principles:

- Resonance exists when a compound cannot be represented by a single Lewis structure.
- Resonance structures differ in the position of only nonbonded electrons and π bonds, not atoms.
- The resonance hybrid is the only accurate representation for a resonance-stabilized compound. A hybrid represents the compound better than any single resonance structure because electron density is delocalized.

The difference between resonance structures and isomers:

- Two **isomers** differ in the arrangement of *both* atoms and electrons.
- **Resonance structures** differ *only* in the arrangement of electrons.




Geometry and Hybridization

The number of groups around an atom determines both its geometry (1.7) and hybridization (1.9).

Number of groups	Geometry	Bond angle (°)	Hybridization
2	linear	180	*sp*
3	trigonal planar	120	sp^2
4	tetrahedral	109.5	sp^3

Drawing Organic Molecules (1.8)

- Shorthand methods are used to abbreviate the structure of organic molecules.



skeletal structure isooctane condensed structure

- A carbon bonded to four atoms is tetrahedral. The best way to represent a tetrahedron is to draw two bonds in the plane, one bond in front, and one bond behind.

Bond Length

- Bond length decreases across a row and increases down a column of the periodic table (1.7A).
- Bond length decreases as the number of electrons between two nuclei increases (1.11A).
- Bond length decreases as the percent *s*-character increases (1.11B).
- Bond length and bond strength are inversely related. In general, shorter bonds are stronger bonds (1.11).
- Sigma (σ) bonds are generally stronger than π bonds (1.10).


Electronegativity and Polarity (1.12, 1.13)

- Electronegativity increases across a row and decreases down a column of the periodic table.
- A polar bond results when two atoms of different electronegativity values are bonded together. Whenever C or H is bonded to N, O, or any halogen, the bond is polar.
- A polar molecule has either one polar bond, or two or more bond dipoles that reinforce.

PROBLEMS

Problems Using Three-Dimensional Models


1.37 Citric acid is responsible for the tartness of citrus fruits, especially lemons and limes.



citric acid

a. What is the molecular formula for citric acid?
b. How many lone pairs are present?
c. Draw a skeletal structure.
d. How many sp^2 hybridized carbons are present?
e. What orbitals are used to form each indicated bond ([1]–[4])?

1.38 Zingerone gives ginger its pungent taste.



zingerone





a. What is the molecular formula for zingerone?
b. How many lone pairs are present?
c. Draw a skeletal structure.
d. How many sp^2 hybridized carbons are present?
e. What orbitals are used to form each indicated bond ([1]–[4])?

Lewis Structures and Formal Charge

1.39 Assign formal charges to each carbon atom in the given species. All lone pairs have been drawn in.

a. $CH_2{=}\ddot{C}H$ b. $H{-}\ddot{C}{-}H$ c. $H{-}\dot{C}{-}H$ (H) d. $H{-}C(H,H){-}C(H,H)$


1.40 Assign formal charges to each N and O atom in the given molecules. All lone pairs have been drawn in.


a.  b.  c.  d. 

1.41 Draw one valid Lewis structure for each compound. Assume the atoms are arranged as drawn.

a. CH_2N_2 

b. CH_3NO_2 

c. CH_3CNO 


d. $(CH_2CN)^-$ 

1.42 Draw an acceptable Lewis structure from each condensed structure, such that all atoms have zero formal charge.

a. diethyl ether, $(CH_3CH_2)_2O$, the first general anesthetic used in medical procedures
b. acrylonitrile, CH_2CHCN, starting material used to manufacture synthetic Orlon fibers
c. dihydroxyacetone, $(HOCH_2)_2CO$, an ingredient in sunless tanning products
d. acetic anhydride, $(CH_3CO)_2O$, a reagent used to synthesize aspirin

Isomers and Resonance Structures

1.43 Creatine is a dietary supplement used by some athletes to boost their athletic performance. (a) Draw in all lone pairs in creatine. (b) Draw two additional resonance structures showing all lone pairs and formal charges.




creatine

1.44 Draw all possible isomers for each molecular formula.

a. C_3H_7Cl (two isomers) b. C_2H_4O (three isomers) c. C_3H_9N (four isomers)


1.45 Draw Lewis structures for the nine isomers having molecular formula C_3H_6O, with all atoms having a zero formal charge.

1.46 With reference to compound **A** drawn below, label each compound as an isomer, a resonance structure, or neither.




A


1.47 With reference to cation **B,** label each species as an isomer, a resonance structure, or neither.




B

1.48 How are the molecules or ions in each pair related? Classify them as resonance structures, isomers, or neither.


a. 

b. 


c. 

d. $CH_3CH_2CH_3$ and $CH_3CH_2\ddot{\underset{\cdot\cdot}{C}}H_2$


1.49 Add curved arrows to show how the first resonance structure can be converted into the second.




1.50 Follow the curved arrows to draw a second resonance structure for each species.




1.51 Draw a second resonance structure for each ion. Then, draw the resonance hybrid.



1.52 Draw all reasonable resonance structures for each species.


a. O_3 b. NO_3^- (a central N atom) c. N_3^- d.  e. 

1.53 Consider compounds **A–D,** which contain both a heteroatom and a double bond. (a) For which compounds are no additional Lewis structures possible? (b) When two or more Lewis structures can be drawn, draw all additional resonance structures.




A **B** **C** **D**

1.54 Draw four additional resonance structures for the following cation. Then draw the resonance hybrid.




1.55 Which of the given resonance structures (**A, B,** or **C**) contributes most to the resonance hybrid? Which contributes least?




A **B** **C**

1.56 Consider the compounds and ions with curved arrows drawn below. When the curved arrows give a second valid resonance structure, draw the resonance structure. When the curved arrows generate an invalid Lewis structure, explain why the structure is unacceptable.


a.  b.  c. $CH_3CH_2-C\equiv N:$ d. 

Geometry

1.57 Predict all bond angles in each compound.

a. CH_3Cl b. NH_2OH c. $CH_2{=}NCH_3$ d. $HC{\equiv}CCH_2OH$ e. 


1.58 Predict the geometry around each highlighted atom.

a.  b. $(CH_3)_2N^-$ c.  d. 


e. $(CH_3)_3N$

Drawing Organic Molecules

1.59 How many hydrogens are present around each carbon atom in the following molecules?


a. 

capsaicin
(spicy component of hot peppers)


b. 

fexofenadine


1.60 Draw in all the carbon and hydrogen atoms in each molecule.


a. 

menthol
(isolated from
peppermint oil)

c. 

ethambutol
(drug used to treat tuberculosis)

b. 
myrcene
(isolated from bayberry)

d. 


estradiol
(a female sex hormone)

1.61 Convert each molecule into a skeletal structure.

a. $(CH_3)_2CHCH_2CH_2CH(CH_3)_2$

b. $CH_3CH(Cl)CH(OH)CH_3$

c. $CH_3(CH_2)_2C(CH_3)_2CH(CH_3)CH(CH_3)CH(Br)CH_3$

d. 


limonene
(oil of lemon)

1.62 Convert the following condensed formulas into skeletal structures.

a. $CH_3CONHCH_3$ c. $(CH_3)_3COH$ e. $CH_3COCH_2CO_2H$


b. CH_3COCH_2Br d. CH_3COCl f. $HO_2CCH(OH)CO_2H$

1.63 Draw in all the hydrogen atoms and nonbonded electron pairs in each ion.




Hybridization

1.64 Predict the hybridization and geometry around each highlighted atom.




1.65 What orbitals are used to form each highlighted bond? For multiple bonds, indicate the orbitals used in individual bonds.




1.66 Ketene, $CH_2=C=O$, is an unusual organic molecule that has a single carbon atom doubly bonded to two different atoms. Determine the hybridization of both C atoms and the O in ketene. Then, draw a diagram showing what orbitals are used to form each bond (similar to Figures 1.10 and 1.11).

Bond Length and Strength

1.67 Rank the following bonds in order of *increasing* bond length.



1.68 Answer the following questions about compound **A.**




A

a. Label the shortest C – C single bond.
b. Label the longest C – C single bond.
c. Considering all the bonds, label the shortest C – C bond.
d. Label the weakest C – C bond.
e. Label the strongest C – H bond.
f. Explain why bond [1] and bond [2] are different in length, even though they are both C – C single bonds.

1.69 Two useful organic compounds that contain Cl atoms are vinyl chloride ($CH_2=CHCl$) and chloroethane (CH_3CH_2Cl). Vinyl chloride is the starting material used to prepare poly(vinyl chloride), a plastic used in insulation, pipes, and bottles. Chloroethane (ethyl chloride) is a local anesthetic. Why is the C – Cl bond of vinyl chloride stronger than the C – Cl bond in chloroethane?


Bond Polarity


1.70 Use the symbols $\delta+$ and $\delta-$ to indicate the polarity of the highlighted bonds.



1.71 Label the polar bonds in each molecule. Indicate the direction of the net dipole (if there is one).


a. $CHBr_3$
b. $CH_3CH_2OCH_2CH_3$

c. 


d. 

General Questions

1.72 Anacin is an over-the-counter pain reliever that contains aspirin and caffeine. Answer the following questions about each compound.



aspirin
(acetylsalicylic acid)



caffeine

a. What is the molecular formula?
b. How many lone pairs are present on heteroatoms?
c. Label the hybridization state of each carbon.
d. Draw three additional resonance structures.

1.73 Answer the following questions about acetonitrile ($CH_3C{\equiv}N$:).


a. Determine the hybridization of both C atoms and the N atom.
b. Label all bonds as σ or π.
c. In what type of orbital does the lone pair on N reside?
d. Label all bonds as polar or nonpolar.

1.74 Benzene is the simplest member of a whole class of compounds called aromatic hydrocarbons.




benzene

a. How is each carbon atom hybridized?
b. What is the geometry around each carbon atom? What is the overall geometry of the benzene ring?
c. Follow the indicated curved arrow notation to draw a second resonance structure.



d. Benzene and other aromatic hydrocarbons are shown in Chapter 17 to be very stable. Offer an explanation.


1.75 The principles of this chapter can be applied to organic molecules of any size. Answer the following questions about amoxicillin, an antibiotic from the penicillin family.



amoxicillin

a. Predict the hybridization and geometry around each highlighted atom.
b. Label five polar bonds using the symbols δ+ and δ–.
c. How many π bonds does amoxicillin have? Label them.
d. Find a C–H bond containing a carbon atom having a hybrid orbital with 33% *s*-character.


1.76



nicotine

a. What is the hybridization of each N atom in nicotine?
b. What is the geometry around each N atom?
c. In what type of orbital does the lone pair on each N atom reside?
d. Draw a constitutional isomer of nicotine.
e. Draw a resonance structure of nicotine.

1.77 Stalevo is the trade name for a medication used for Parkinson's disease, which contains L-dopa, carbidopa, and entacapone.




entacapone


a. Draw a Lewis structure for entacapone.
b. Which C–C bond in entacapone is the longest?
c. Which C–C single bond is the shortest?
d. Which C–N bond is the longest?
e. Which C–N bond is the shortest?
f. Use curved arrows to draw a resonance structure that is an equal contributor to the resonance hybrid.
g. Use curved arrows to draw a resonance structure that is a minor contributor to the resonance hybrid.

1.78 CH_3^+ and CH_3^- are two highly reactive carbon species.

a. What is the predicted hybridization and geometry around each carbon atom?
b. Two electrostatic potential plots are drawn for these species. Which ion corresponds to which diagram and why?



A





B


Challenge Problems

1.79 The N atom in CH_3CONH_2 (acetamide) is sp^2 hybridized, even though it is surrounded by four groups. Using this information, draw a diagram that shows the orbitals used by the atoms in the $-CONH_2$ portion of acetamide, and offer an explanation as to the observed hybridization.


1.80 Use the observed bond lengths to answer each question. (a) Why is bond [1] longer than bond [2] (143 pm versus 136 pm)? (b) Why are bonds [3] and [4] equal in length (127 pm), and shorter than bond [2]?







1.81 Draw at least 10 more resonance structures for acetaminophen, the active pain reliever in Tylenol.



acetaminophen

1.82 When two carbons having different hybridization are bonded together, the C – C bond contains a slight dipole. In a $C_{sp^2} - C_{sp^3}$ bond, what is the direction of the dipole? Which carbon is considered more electronegative?


1.83 Draw all possible isomers having molecular formula C_4H_8 that contain one π bond.

1.84 Use the principles of resonance theory to explain why carbocation **A** is more stable than carbocation **B.**



A **B**

1.85 The curved arrow notation introduced in Section 1.6 is a powerful method used by organic chemists to show the movement of electrons not only in resonance structures, but also in chemical reactions. Since each curved arrow shows the movement of two electrons, following the curved arrows illustrates what bonds are broken and formed in a reaction. Consider the following three-step process. (a) Add curved arrows in Step [1] to show the movement of electrons. (b) Use the curved arrows drawn in Step [2] to identify the structure of **X. X** is converted in Step [3] to phenol and HCl.



Acids and Bases

2



Aspirin is one of the most widely used over-the-counter drugs. Whether you purchase Anacin, Bufferin, Bayer, or a generic, the active ingredient is the same—**acetylsalicylic acid.** Aspirin, a synthetic compound that does not occur in nature, was first marketed to the general public in 1899, and is now used regularly by over 100 million people throughout the world. Like many drugs, aspirin undergoes a proton transfer reaction after ingestion. In Chapter 2, we learn about acidity and the role of acid–base reactions in aspirin's chemistry.

2.1 Brønsted–Lowry acids and bases
2.2 Reactions of Brønsted–Lowry acids and bases
2.3 Acid strength and pK_a
2.4 Predicting the outcome of acid–base reactions
2.5 Factors that determine acid strength
2.6 Common acids and bases
2.7 Aspirin
2.8 Lewis acids and bases

Chemical terms such as *anion* and *cation* may be unfamiliar to most nonscientists, but *acid* has found a place in everyday language. Commercials advertise the latest remedy for the heartburn caused by excess stomach *acid.* The nightly news may report the latest environmental impact of *acid* rain. Wine lovers know that wine sours because its alcohol has turned to *acid. Acid* comes from the Latin word *acidus,* meaning sour, because when tasting compounds was a routine method of identification, these compounds were sour.

In Chapter 2, we concentrate on two definitions of acids and bases: the **Brønsted–Lowry** definition, which describes acids as **proton donors** and bases as **proton acceptors,** and the **Lewis** definition, which describes acids as **electron pair acceptors** and bases as **electron pair donors.**

2.1 Brønsted–Lowry Acids and Bases

The general words "acid" and "base" usually mean a *Brønsted–Lowry* acid and *Brønsted–Lowry* base.

H^+ = proton.

HA = Brønsted–Lowry acid.
B: = Brønsted–Lowry base.

The Brønsted–Lowry definition describes acidity in terms of protons: positively charged **hydrogen ions, H^+.**

- A Brønsted–Lowry acid is a *proton donor.*
- A Brønsted–Lowry base is a *proton acceptor.*

A Brønsted–Lowry acid must contain a *hydrogen* atom. This definition of an acid is often familiar to students, because many inorganic acids in general chemistry are Brønsted–Lowry acids. The symbol **HA** is used for a general Brønsted–Lowry acid.

A Brønsted–Lowry base must be able to form a bond to a proton. Because a proton has no electrons, **a base must contain an "available" electron pair** that can be easily donated to form a new bond. These include **lone pairs** or electron pairs in **π bonds.** The symbol **B:** is used for a general Brønsted–Lowry base. Examples of Brønsted–Lowry acids and bases are given in Figure 2.1.

Charged species such as ^-OH and $^-NH_2$ are used as **salts,** with cations such as Li^+, Na^+, or K^+ to balance the negative charge. These cations are called **counterions** or **spectator ions,** and their **identity is usually inconsequential.** For this reason, the counterion is often omitted.


NaOH	=	Na^+	^-OH
KOH	=	K^+	^-OH
salt		**counterion**	**base**

Compounds like H_2O and CH_3OH that contain both hydrogen atoms and lone pairs may be either an acid or a base, depending on the particular reaction. These fundamental principles are true no matter how complex the compound. For example, the addictive pain reliever **morphine** is a

Figure 2.1
Examples of Brønsted–Lowry acids and bases

a. **Brønsted–Lowry acids (HA)**


HCl
H_2SO_4
HSO_4^-
H_2O
H_3O^+



acetic acid

- All Brønsted–Lowry acids contain a proton.
- The net charge may be zero, (+), or (–).

b. **Brønsted–Lowry bases (B:)**




- All Brønsted–Lowry bases contain a lone pair of electrons or a π bond.
- The net charge may be zero or (–).



Morphine is obtained from the opium poppy.

Brønsted–Lowry acid because it contains many hydrogen atoms. It is also a Brønsted–Lowry base because it has lone pairs on O and N, and four π bonds.



morphine

- H atoms on O make morphine an acid.
- Lone pairs and π bonds (in blue) make morphine a base.

Problem 2.1

a. Which compounds are Brønsted–Lowry acids: HBr, NH_3, CCl_4?

b. Which compounds are Brønsted–Lowry bases: CH_3CH_3, $(CH_3)_3CO^-$, HC≡CH?

c. Classify each compound as an acid, a base, or both: CH_3CH_2OH, $CH_3CH_2CH_2CH_3$, $CH_3CO_2CH_3$.


2.2 Reactions of Brønsted–Lowry Acids and Bases

A Brønsted–Lowry acid–base reaction results in transfer of a proton from an acid to a base. These acid–base reactions, also called ***proton transfer reactions,*** are fundamental to the study of organic chemistry.

Consider, for example, the reaction of the acid HA with the base :B. **In an acid–base reaction, one bond is broken and one is formed.**

- **The electron pair of the base B: forms a new bond to the proton of the acid.**
- **The acid HA loses a proton, leaving the electron pair in the HA bond on A.**

Recall from Section 1.6 that a curved arrow shows the movement of an **electron pair.** The tail of the arrow always begins at an electron pair and the head points to where that electron pair "moves."



This "movement" of electrons in reactions can be illustrated using curved arrow notation. Because **two electron pairs** are involved in this reaction, **two curved arrows** are needed. Two products are formed.

- Loss of a proton from an acid forms its *conjugate base.*
- Gain of a proton by a base forms its *conjugate acid.*


The **net charge must be the same** on both sides of any equation. In this example, the net charge on each side is zero. Individual charges can be calculated using formal charges. **A double reaction arrow** is used between starting materials and products to indicate that the reaction can proceed in the forward and reverse directions. These are **equilibrium arrows.**

A double reaction arrow indicates equilibrium.

equilibrium arrows

Remove H^+ from an acid to form its conjugate base. Add H^+ to a base to form its conjugate acid.

Two examples of proton transfer reactions are drawn here with curved arrow notation.



The acid loses H^+. **The base gains H^+.**


- **Brønsted–Lowry acid–base reactions always result in the transfer of a proton from an acid to a base.**

The ability to identify and draw a conjugate acid or base from a given starting material is illustrated in Sample Problems 2.1 and 2.2.


Sample Problem 2.1 a. What is the conjugate acid of CH_3O^-?
b. What is the conjugate base of NH_3?

Solution

a. **Add H^+ to CH_3O^- to form its conjugate acid.**




b. **Remove H^+ from NH_3 to form its conjugate base.**



Problem 2.2 a. Draw the conjugate acid of each base: NH_3, Cl^-, $(CH_3)_2C{=}O$.
b. Draw the conjugate base of each acid: HBr, HSO_4^-, CH_3OH.

Problem 2.3 Label each statement as True or False.
a. $CH_3CH_2^+$ is the conjugate acid of $CH_2{=}CH_2$.
b. $CH_3CH_2^-$ is the conjugate base of $CH_3CH_2^+$.
c. $CH_2{=}CH_2$ is the conjugate base of $CH_3CH_2^-$.
d. $CH_2{=}CH^-$ is the conjugate base of $CH_2{=}CH_2$.
e. CH_3CH_3 is the conjugate acid of $CH_3CH_2^-$.


Sample Problem 2.2 Label the acid and base, and the conjugate acid and base in the following reaction. Use curved arrow notation to show the movement of electron pairs.



A **B** **C** **D**

Solution

A is the base because it accepts a proton, forming its conjugate acid, **C. B** is the acid because it donates a proton, forming its conjugate base, **D.** Two curved arrows are needed. One shows that the lone pair on **A** bonds to a proton of **B,** and the second shows that the electron pair in the O – H bond remains on O.




A **B** **C** **D**

base acid conjugate acid conjugate base

The base gains H^+. **The acid loses H^+.**

Problem 2.4 Label the acid and base, and the conjugate acid and base in the following reactions. Use curved arrows to show the movement of electron pairs.



In all proton transfer reactions, the **electron-rich base** donates an electron pair to the acid, which usually has a polar HA bond. The H of the acid bears a partial positive charge, making it **electron deficient.** This is the first example of a general pattern of reactivity.

- **Electron-rich species react with electron-deficient ones.**


Given two starting materials, how do you know which is the acid and which is the base in a proton transfer reaction? Use the following generalizations:

[1] Common acids and bases introduced in general chemistry are used in the same way in organic reactions. HCl and H_2SO_4 are strong acids, and ^-OH is a strong base.


[2] When only one starting material contains a hydrogen, it must be the acid. If only one starting material has a lone pair or a π bond, it must be the base.

[3] A starting material with a net positive charge is usually the acid. A starting material with a negative charge is usually the base.

Problem 2.5 Decide which compound is the acid and which is the base, and draw the products of each proton transfer reaction.



Problem 2.6 Draw the products formed from the acid–base reaction of HCl with each compound.




2.3 Acid Strength and pK_a

Acid strength is the tendency of an acid to donate a proton.

- The more readily a compound donates a proton, the stronger the acid.

Acidity is measured by an equilibrium constant. When a Brønsted–Lowry acid HA is dissolved in water, an acid–base reaction occurs, and an equilibrium constant K_{eq} can be written for the reaction.



$$K_{eq} = \frac{[\text{products}]}{[\text{starting materials}]} = \frac{[H_3O^+][A:^-]}{[HA][H_2O]}$$

Because the concentration of the solvent H_2O is essentially constant, the equation can be re-arranged and a new equilibrium constant, called the **acidity constant, K_a,** can be defined.

$$K_a = [H_2O]K_{eq} = \frac{[H_3O^+][A:^-]}{[HA]}$$

How is the magnitude of K_a related to acid strength?

- The *stronger the acid,* the further the equilibrium lies to the right, and the *larger the* K_a.

For most organic compounds, K_a is small, typically 10^{-5} to 10^{-50}. This contrasts with the K_a values for many inorganic acids, which range from 10^0 to 10^{10}. Because using exponents can be cumbersome, it is often more convenient to use pK_a values instead of K_a values.

$$pK_a = -\log K_a$$

How does pK_a relate to acid strength?



K_a values of typical organic acids		**pK_a values** of typical organic acids	
10^{-5} →	10^{-50}	+5 →	+50
larger number stronger acid	smaller number weaker acid	smaller number stronger acid	larger number weaker acid

Recall that a **log** is an **exponent;** for example, log $10^{-5} = -5$.

- The *smaller* the pK_a, the *stronger* the acid.

Problem 2.7 Which compound in each pair is the stronger acid?

a. $CH_3CH_2CH_3$ (pK_a = 50) or CH_3CH_2OH (pK_a = 16)

b.  ($K_a = 10^{-10}$) or  ($K_a = 10^{-41}$)

Problem 2.8 Use a calculator when necessary to answer the following questions.

a. What is the pK_a for each K_a: 10^{-10}, 10^{-21}, and 5.2×10^{-5}?

b. What is the K_a for each pK_a: 7, 11, and 3.2?

An inverse relationship exists between acidity and basicity.

- A *strong acid* readily donates a proton, forming a *weak conjugate base.*
- A *strong base* readily accepts a proton, forming a *weak conjugate acid.*


Table 2.1 is a brief list of pK_a values for some common compounds, ranked in order of ***increasing* pK_a** and therefore ***decreasing* acidity.** Because strong acids form weak conjugate bases, this list also ranks their conjugate bases, in order of ***increasing* basicity.** CH_4 is the weakest acid in the list, because it has the highest pK_a (50). Its conjugate base, CH_3^-, is therefore the strongest conjugate base. An extensive pK_a table is located in Appendix A.

Table 2.1 Selected pK_a Values

	Acid	pK_a	Conjugate base	
Increasing acidity ↑	H–Cl	–7	Cl^-	Increasing basicity ↓
	CH_3CO_2–H	4.8	$CH_3CO_2^-$	
	HO–H	15.7	HO^-	
	CH_3CH_2O–H	16	$CH_3CH_2O^-$	
	HC≡CH	25	$HC{\equiv}C^-$	
	H–H	35	H^-	
	H_2N–H	38	H_2N^-	
	$CH_2{=}CH_2$	44	$CH_2{=}\bar{C}H$	
	CH_3–H	50	CH_3^-	

Comparing pK_a values tells us the **relative acidity of two acids,** and the **relative basicity of their conjugate bases,** as shown in Sample Problem 2.3.

Sample Problem 2.3 Rank the following compounds in order of increasing acidity, and then rank their conjugate bases in order of increasing basicity.




H–Cl



Solution

Use the pK_a values in Table 2.1 and the rule: **the lower the pK_a, the stronger the acid.**



pK_a = 44




pK_a = 4.8

H–Cl

pK_a = –7

Increasing acidity

Because strong acids form weak conjugate bases, the **basicity of conjugate bases increases with increasing pK_a** of their acids.







Increasing basicity

Problem 2.9 Rank the conjugate bases of each group of acids in order of increasing basicity.


a. NH_3, H_2O, CH_4 b. $CH_2{=}CH_2$, $HC{\equiv}CH$, CH_4


Problem 2.10 Consider two acids: HCO_2H (formic acid, $pK_a = 3.8$) and pivalic acid [$(CH_3)_3CCO_2H$, $pK_a = 5.0$]. (a) Which acid has the larger K_a? (b) Which acid is the stronger acid? (c) Which acid forms the stronger conjugate base? (d) When each acid is dissolved in water, for which acid does the equilibrium lie further to the right?


The pK_a values in Table 2.1 span a large range (–7 to 50). The pK_a scale is logarithmic, so a small difference in pK_a translates into a large numerical difference. The difference between the pK_a values of NH_3 (38) and $CH_2{=}CH_2$ (44) is six pK_a units, so NH_3 is 10^6 or *one million times more acidic* than $CH_2{=}CH_2$.

Although Table 2.1 is abbreviated, it is a useful tool for *estimating* the pK_a of a compound similar though not identical to one in the table. Suppose you are asked to estimate the pK_a of the N–H bond of CH_3NH_2. Although CH_3NH_2 is not listed in the table, we have enough information to *approximate* its pK_a. Because the pK_a of the N–H bond of NH_3 is 38, we can estimate the pK_a of the N–H bond of CH_3NH_2 to be 38. Its actual pK_a is 40, so this is a good first approximation.

Problem 2.11 Estimate the pK_a of each of the indicated bonds.

a. 

b. 

c. 

2.4 Predicting the Outcome of Acid–Base Reactions

In a proton transfer reaction, the **stronger acid reacts with the stronger base** to form the weaker acid and the weaker base.

A proton transfer reaction represents an equilibrium. Because an acid donates a proton to a base, forming a conjugate acid and conjugate base, there are always two acids and two bases in the reaction mixture. Which pair of acids and bases is favored at equilibrium? **The position of the equilibrium depends on the relative strengths of the acids and bases.**

- **Equilibrium always favors formation of the weaker acid and base.**

Because a strong acid readily donates a proton and a strong base readily accepts one, these two species react to form a weaker conjugate acid and base that do not donate or accept a proton as readily. Comparing pK_a values allows us to determine the position of equilibrium, as illustrated in Sample Problem 2.4.

Sample Problem 2.4 Determine the direction of equilibrium when acetylene ($HC{\equiv}CH$) reacts with $^-NH_2$ in a proton transfer reaction.


Solution

Follow three steps to determine the position of equilibrium:

Step [1] Identify the acid and base in the starting materials.


- $^-NH_2$ is the base because it bears a net negative charge, so $HC{\equiv}CH$ is the acid.

Step [2] Draw the products of proton transfer and identify the conjugate acid and base in the products.



Step [3] **Compare the pK_a values of the acid and the conjugate acid. Equilibrium favors formation of the weaker acid with the higher pK_a.**


Use unequal equilibrium arrows (⇀↽ or ↽⇀) for a reversible reaction in which products or reactants are favored at equilibrium.



Equilibrium favors the products, forming the weaker acid.

- Because the pK_a of the starting acid (25) is ***lower*** than the pK_a of the conjugate acid (38), HC≡CH is a ***stronger*** acid and equilibrium favors the products.

Problem 2.12 Draw the products of each reaction and determine the direction of equilibrium.




How can we know if a particular base is strong enough to deprotonate a given acid, so that the equilibrium lies to the right? The pK_a table readily gives us this information, as shown in Sample Problem 2.5.

Sample Problem 2.5 Which of the following bases is strong enough to deprotonate *N,N*-dimethylacetamide [$CH_3CON(CH_3)_2$, pK_a = 30], so that equilibrium favors the products: (a) $NaNH_2$; (b) NaOH?

Solution


- Draw the structure of the conjugate acid of each base, and determine its pK_a from Table 2.1 or Appendix A.
- Compare the pK_a values of the starting acid and the conjugate acid. If the conjugate acid has a *higher* pK_a than the starting acid, the conjugate acid is the *weaker* acid and equilibrium favors the *products.* **The base is strong enough** to deprotonate the acid.
- If the conjugate acid has a *lower* pK_a than the starting acid, the conjugate acid is the *stronger* acid and equilibrium favors the *starting materials.* **The base is not strong enough** to deprotonate the acid.

a. Na^+ is a counterion and $^-NH_2$ is the base in $NaNH_2$.



The conjugate acid (NH_3) of the base is a weaker acid than $CH_3CON(CH_3)_2$ (pK_a = 30), so the base is strong enough to deprotonate the acid, and equilibrium favors the products.

b. Na^+ is a counterion and ^-OH is the base in NaOH.



The conjugate acid (H_2O) of the base is a stronger acid than $CH_3CON(CH_3)_2$ (pK_a = 30), so the base is *not* strong enough to deprotonate the acid, and equilibrium favors the starting materials.

Because Table 2.1 is arranged from low to high pK_a, **an acid can be deprotonated by the conjugate base of any acid below it in the table.**

Sample Problem 2.5 illustrates a fundamental principle in acid–base reactions.

- **An acid can be deprotonated by the conjugate base of any acid having a *higher* pK_a.**

Problem 2.13 Using the data in Appendix A, determine which of the following bases is strong enough to deprotonate acetonitrile (CH_3CN), so that equilibrium favors the products: (a) NaH; (b) Na_2CO_3; (c) NaOH; (d) $NaNH_2$; (e) $NaHCO_3$.

2.5 Factors That Determine Acid Strength

The wide range of pK_a values in Table 2.1 illustrates that a tremendous difference in acidity exists among compounds. HCl (pK_a < 0) is an extremely strong acid, water (pK_a = 15.7) is moderate in acidity, and CH_4 (pK_a = 50) is an extremely weak acid. How are these differences explained? One general rule governs acid strength.

- Anything that stabilizes a conjugate base $A:^-$ makes the starting acid HA more acidic.

Four factors affect the acidity of HA:

[1] Element effects
[2] Inductive effects
[3] Resonance effects
[4] Hybridization effects

No matter which factor is discussed, follow the same procedure. To compare the acidity of any two acids:

- **Draw the conjugate bases.**
- **Determine which conjugate base is more stable.**
- **The *more stable* the conjugate base, the *more acidic* the acid.**


2.5A Element Effects—Trends in the Periodic Table

The most important factor determining the acidity of HA is the location of A in the periodic table.

Comparing Elements in the Same Row of the Periodic Table


To examine acidity trends **across a row** of the periodic table, we compare CH_4 and H_2O, two compounds having H atoms bonded to a second-row element. We know from Table 2.1 that **H_2O has a much *lower* pK_a and therefore is much *more acidic* than CH_4,** but why is this the case?

To answer this question, first draw both conjugate bases and then determine which is more stable. Each conjugate base has a net negative charge, but the negative charge in ^-OH is on oxygen and in CH_3^- it is on carbon.



Because the oxygen atom is much **more electronegative** than carbon, oxygen more readily accepts a negative charge, making ^-OH much more stable than CH_3^-. **H_2O is a stronger acid than CH_4 because ^-OH is a more stable conjugate base than CH_3^-.** This is a specific example of a general trend.

- Across a row of the periodic table, the acidity of HA increases as the electronegativity of A increases.



The enormity of this effect is evident by comparing the pK_a values for these bonds. **A C–H bond is approximately 10^{47} times *less acidic* than H–F.**

Comparing Elements Down a Column of the Periodic Table

To examine acidity trends down a column of the periodic table, we compare H–F and H–Br. Draw both conjugate bases and then determine which is more stable. In this case, removal of a proton forms F^- and Br^-.



conjugate base conjugate base

There are two important differences between F^- and Br^-—electronegativity and size. In this case, **size is more important than electronegativity.** The size of an atom or ion increases down a column of the periodic table, so Br^- is much larger than F^-, and this stabilizes the negative charge.

- Positive or negative charge is stabilized when it is spread over a larger volume.

Because Br^- is larger than F^-, Br^- is more stable than F^-, and H–Br is a stronger acid than H–F.




smaller anion **larger anion**

less stable conjugate base **more stable conjugate base**

This again is a specific example of a general trend.

- Down a column of the periodic table, the acidity of HA increases as the size of A increases.



This is ***opposite*** to what would be expected on the basis of electronegativity differences between F and Br, because F is more electronegative than Br. **Size and *not* electronegativity determines acidity down a column.** Combining both trends together:

- The acidity of HA increases both left-to-right across a row and down a column of the periodic table.

Because of carbon's position in the periodic table (in the second row and to the left of O, N, and the halogens), **C–H bonds are usually the *least acidic* bonds in a molecule.**

Sample Problem 2.6

Without reference to a pK_a table, decide which compound in each pair is the stronger acid:


a. H_2O or HF b. H_2S or H_2O

Solution


a. H_2O and HF both have H atoms bonded to a second-row element. Because the acidity of HA increases across a row of the periodic table, the H–F bond is more acidic than the H–O bond. **HF is a stronger acid than H_2O.**

b. H_2O and H_2S both have H atoms bonded to elements in the same column. Because the acidity of HA increases down a column of the periodic table, the H–S bond is more acidic than the H–O bond. **H_2S is a stronger acid than H_2O.**

Problem 2.14 Without reference to a pK_a table, decide which compound in each pair is the stronger acid.

a.  or H_2O b.  or H_2S

Problem 2.15 Rank the labeled H atoms in the following compound in order of increasing acidity.






Because the pseudoephedrine (Problem 2.17) in Sudafed can be readily converted to the illegal, addictive drug methamphetamine, products that contain pseudoephedrine are now stocked behind the pharmacy counter, so their sale can be more closely monitored. Sudafed PE is a related product that contains a decongestant less easily converted to methamphetamine.




When discussing acidity, the most acidic proton in a compound is the one removed first by a base. Although four factors determine the overall acidity of a particular hydrogen atom, **the element effect—the identity of A—is the single most important factor in determining the acidity of the HA bond.**

To decide which hydrogen is most acidic, **first determine what element each hydrogen is bonded to and then decide its acidity based on periodic trends.** For example, $CH_3NHCH_2CH_2CH_2CH_3$ contains only C−H and N−H bonds. Because the acidity of HA increases across a row of the periodic table, the single H on N is the most acidic H in this compound.




most acidic H shown in red

Problem 2.16 Which hydrogen in each molecule is most acidic?




a.  b.  c. 

Problem 2.17 Which hydrogen in pseudoephedrine, the nasal decongestant in the commercial medication Sudafed, is most acidic?



pseudoephedrine


Problem 2.18 Which compound in each pair of isomers is the stronger acid?

a.  or  b.  or 


2.5B Inductive Effects

A second factor affecting the acidity of HA is the presence of atoms more electronegative than carbon. To illustrate this phenomenon, compare ethanol (CH_3CH_2OH) and 2,2,2-trifluoroethanol (CF_3CH_2OH), two compounds containing O−H bonds. The pK_a table in Appendix A

indicates that CF_3CH_2OH is a stronger acid than CH_3CH_2OH. We are comparing the acidity of the O−H bond in both compounds, so what causes the difference?



ethanol
$pK_a = 16$




2,2,2-trifluoroethanol
$pK_a = 12.4$

stronger acid

Draw both conjugate bases and then determine which is more stable. Both bases have a negative charge on an electronegative oxygen, but the second anion has three very electronegative fluorine atoms. These fluorine atoms withdraw electron density from the carbon to which they are bonded, making it electron deficient. Furthermore, this electron-deficient carbon pulls electron density through σ bonds from the negatively charged oxygen atom, stabilizing the negative charge. This is called an **inductive effect.**



No additional electronegative atoms stabilize the conjugate base.



CF_3 withdraws electron density, stabilizing the conjugate base.

- An *inductive effect* is the pull of electron density through σ bonds caused by electronegativity differences of atoms.


In this case, the electron density is pulled away from the negative charge through σ bonds by the very electronegative fluorine atoms, so it is called an **electron-*withdrawing* inductive effect. Thus, the three very electronegative fluorine atoms stabilize the negatively charged conjugate base $CF_3CH_2O^-$, making CF_3CH_2OH a stronger acid than CH_3CH_2OH.** We have learned two important principles from this discussion:

- More electronegative atoms stabilize regions of high electron density by an electron-withdrawing inductive effect.
- The acidity of HA increases with the presence of electron-withdrawing groups in A.


Inductive effects result because an electronegative atom stabilizes the negative charge of the conjugate base. **The *more electronegative* the atom and the *closer* it is to the site of the negative charge, the greater the effect.** This effect is discussed in greater detail in Chapter 19.

Electrostatic potential plots in Figure 2.2 compare the electron density around the oxygen atoms in $CH_3CH_2O^-$ and $CF_3CH_2O^-$. The darker red region around the O atom of $CH_3CH_2O^-$ indicates a higher concentration of electron density compared to the O atom of $CF_3CH_2O^-$.

Figure 2.2
Electrostatic potential plots of $CH_3CH_2O^-$ and $CF_3CH_2O^-$







$CH_3CH_2O^-$
The dark red of the O atom indicates a region of high electron density.






$CF_3CH_2O^-$
The O atom is yellow, indicating it is less electron rich.

Problem 2.19 Which compound in each pair is the stronger acid?

a.  or 

b.  or 

c.  or 




α-Hydroxy acids (Problem 2.20) are used in skin care products that purportedly smooth fine lines and improve skin texture by reacting with the outer layer of skin cells, causing them to loosen and flake off.

Problem 2.20 Glycolic acid, $HOCH_2CO_2H$, is the simplest member of a group of compounds called α-hydroxy acids, ingredients in skin care products that have an OH group on the carbon adjacent to a CO_2H group. Would you expect $HOCH_2CO_2H$ to be a stronger or weaker acid than acetic acid, CH_3CO_2H?


Problem 2.21 Explain the apparent paradox. HBr is a stronger acid than HCl, but HOCl is a stronger acid than HOBr.

2.5C Resonance Effects

A third factor that determines acidity is resonance. Recall from Section 1.6 that resonance occurs whenever two or more different Lewis structures can be drawn for the same arrangement of atoms. To illustrate this phenomenon, compare ethanol (CH_3CH_2OH) and acetic acid (CH_3CO_2H), two compounds containing O−H bonds. Based on Table 2.1, CH_3CO_2H is a stronger acid than CH_3CH_2OH.



ethanol
$pK_a = 16$



acetic acid
$pK_a = 4.8$
stronger acid

Resonance structures are two Lewis structures having the same placement of atoms but a different arrangement of electrons.


Draw the conjugate bases of these acids to illustrate the importance of resonance. For ethoxide ($CH_3CH_2O^-$), the conjugate base of ethanol, only one Lewis structure can be drawn. The negative charge of this conjugate base is *localized* on the O atom.



ethanol
acid

ethoxide
conjugate base
only **one** Lewis structure

With acetate ($CH_3CO_2^-$), however, two resonance structures can be drawn.



acetic acid

acetate
conjugate base
two resonance structures

hybrid
resonance-stabilized
conjugate base

These two resonance structures differ in the **position of a π bond** and a **lone pair.** Although each resonance structure of acetate implies that the negative charge is localized on an O atom, in actuality, charge is ***delocalized*** over both O atoms. **Delocalization of electron density stabilizes acetate, making it a weaker base.**

Resonance delocalization often produces a larger effect on pK_a than the inductive effects discussed in Section 2.5B. Resonance makes CH_3CO_2H (pK_a = 4.8) a much stronger acid than CH_3CH_2OH (pK_a = 16), while the inductive effects due to three electronegative F atoms make CF_3CH_2OH (pK_a = 12.4) a somewhat stronger acid than CH_3CH_2OH.


Remember that neither resonance form adequately represents acetate. The true structure is a **hybrid** of both structures. In the hybrid, the electron pairs drawn in different locations in individual resonance structures are ***delocalized.*** With acetate, a dashed line is used to show that each C–O bond has partial double bond character. The symbol δ– (partial negative) indicates that the charge is delocalized on both O atoms in the hybrid.

Thus, **resonance delocalization makes $CH_3CO_2^-$ more stable than $CH_3CH_2O^-$, so CH_3CO_2H is a stronger acid than CH_3CH_2OH.** This is another example of a general rule.


- **The acidity of HA increases when the conjugate base $A:^-$ is resonance stabilized.**

Electrostatic potential plots of $CH_3CH_2O^-$ and $CH_3CO_2^-$ in Figure 2.3 indicate that the negative charge is concentrated on a single O in $CH_3CH_2O^-$, but delocalized over the O atoms in $CH_3CO_2^-$.

Figure 2.3
Electrostatic potential plots of $CH_3CH_2O^-$ and $CH_3CO_2^-$



$CH_3CH_2O^-$
The negative charge is concentrated on the single oxygen atom, making this anion *less stable*.




$CH_3CO_2^-$
The negative charge is delocalized over both oxygen atoms, making this anion *more stable*.

Problem 2.22 The C–H bond in acetone, $(CH_3)_2C{=}O$, has a pK_a of 19.2. Draw two resonance structures for its conjugate base. Then, explain why acetone is much more acidic than propane, $CH_3CH_2CH_3$ (pK_a = 50).


Problem 2.23 Acetonitrile (CH_3CN) has a pK_a of 25, making it more acidic than many other compounds having only C–H bonds. Draw Lewis structures for acetonitrile and its conjugate base. Use resonance structures to account for the acidity of acetonitrile.

2.5D Hybridization Effects

The final factor affecting the acidity of HA is the hybridization of A. To illustrate this phenomenon, compare ethane (CH_3CH_3), ethylene ($CH_2{=}CH_2$), and acetylene ($HC{\equiv}CH$). Appendix A indicates that there is a considerable difference in the pK_a values of these compounds.



The conjugate bases formed by removing a proton from ethane, ethylene, and acetylene are **carbanions—species with a negative charge on carbon.**



The hybridization of the carbon bearing the negative charge is different in each anion, so the lone pair of electrons occupies an orbital with a different percent *s*-character in each case. A higher percent *s*-character means a hybrid orbital has a larger fraction of the lower energy *s* orbital.


- The *higher* the percent *s*-character of the hybrid orbital, the **more stable** the conjugate base.

Thus, **acidity increases from CH_3CH_3 to $CH_2{=}CH_2$ to $HC{\equiv}CH$ as the negative charge of the conjugate base is stabilized by increasing percent *s*-character.** Once again this is a specific example of a general trend.

- The acidity of HA increases as the percent *s*-character of $A:^-$ increases.


Electrostatic potential plots of these carbanions appear in Figure 2.4.

Figure 2.4
Electrostatic potential plots of three carbanions



- As the lone pair of electrons is pulled closer to the nucleus, the negatively charged carbon appears less intensely red.

Problem 2.24 For each pair of compounds: [1] Which indicated H is more acidic? [2] Draw the conjugate base of each acid. [3] Which conjugate base is stronger?



2.5E Summary of Factors Determining Acid Strength

The ability to recognize the most acidic site in a molecule will be important throughout the study of organic chemistry. All the factors that determine acidity are therefore summarized in Figure 2.5. The following two-step procedure shows how these four factors can be used to determine the relative acidity of protons.

Figure 2.5
Summary of the factors that determine acidity

Factor	Example		
1. **Element effects:** The acidity of HA increases both left-to-right across a row and down a column of the periodic table.	CH_4	and	H_2O **more acidic**
2. **Inductive effects:** The acidity of HA increases with the presence of electron-withdrawing groups in A.	CH_3CH_2O-H	and	CF_3CH_2O-H **more acidic**
3. **Resonance effects:** The acidity of HA increases when the conjugate base $A:^-$ is resonance stabilized.	CH_3CH_2O-H	and	CH_3CO_2-H **more acidic**
4. **Hybridization effects:** The acidity of HA increases as the percent *s*-character of $A:^-$ increases.	$CH_2=CH_2$	and	$H-C\equiv C-H$ **more acidic**

How To Determine the Relative Acidity of Protons

Step [1] **Identify the atoms bonded to hydrogen, and use periodic trends to assign relative acidity.**


- The most common HA bonds in organic compounds are C – H, N – H, and O – H. Because acidity increases left-to-right across a row, the relative acidity of these bonds is C – H < N – H < O – H. Therefore, H atoms bonded to C atoms are usually *less acidic* than H atoms bonded to any heteroatom.

Step [2] **If the two H atoms in question are bonded to the same element, draw the conjugate bases and look for other points of difference. Ask three questions:**


- Do electron-withdrawing groups stabilize the conjugate base?
- Is the conjugate base resonance stabilized?
- How is the conjugate base hybridized?

Sample Problem 2.7 shows how to apply this procedure to actual compounds.


Sample Problem 2.7 Rank the following compounds in order of increasing acidity of their most acidic hydrogen atom.



A



B




C


Solution

[1] Compounds **A, B,** and **C** contain C – H, N – H, and O – H bonds. Because acidity increases left-to-right across a row of the periodic table, the O – H bonds are most acidic. Compound **C** is thus the least acidic because it has *no* O – H bonds.


[2] The only difference between compounds **A** and **B** is the presence of an electronegative Cl in **A.** The Cl atom stabilizes the conjugate base of **A,** making it more acidic than **B.** Thus,



Problem 2.25 Rank the compounds in each group in order of increasing acidity.



Problem 2.26 Which proton in each of the following drugs is most acidic? THC is the active component in marijuana, and ketoprofen is an anti-inflammatory agent.



Problem 2.27 Which anion (**A** or **B**) is the stronger base?



2.6 Common Acids and Bases

Many strong or moderately strong acids and bases are used as reagents in organic reactions.

2.6A Common Acids

Several organic reactions are carried out in the presence of strong inorganic acids, most commonly **HCl** and **H_2SO_4.** These strong acids, with **pK_a values $\leq$ 0,** should be familiar from previous chemistry courses.



Sulfuric acid is the most widely produced industrial chemical. It is also formed when sulfur oxides, emitted into the atmosphere by burning fossil fuels high in sulfur content, dissolve in water. This makes rainwater acidic, forming acid rain, which has destroyed acres of forests worldwide.

Two organic acids are also commonly used, namely **acetic acid** and ***p*-toluenesulfonic acid** (usually abbreviated as **TsOH**). Although acetic acid has a higher pK_a than the inorganic acids, making it a weaker acid, it is more acidic than most organic compounds. *p*-Toluenesulfonic acid is similar in acidity to the strong inorganic acids. Because it is a solid, small quantities can be easily weighed on a balance and then added to a reaction mixture.



acetic acid
pK_a = 4.8

p-toluenesulfonic acid
pK_a = –7

2.6B Common Bases

Three common kinds of strong bases include:

[1] Negatively charged oxygen bases: $^-$**OH** (hydroxide) and its organic derivatives
[2] Negatively charged nitrogen bases: $^-$**NH$_2$** (amide) and its organic derivatives
[3] Hydride (**H$^-$**)

Figure 2.6 gives examples of these strong bases. Each negatively charged base is used as a salt with a spectator ion (usually Li^+, Na^+, or K^+) that serves to balance charge.

- **Strong bases have weak conjugate acids with high pK_a values, usually > 12.**

Figure 2.6
Some common negatively charged bases

Oxygen bases			Nitrogen bases		
Na^+	$^-$:OH	sodium hydroxide	Na^+	$^-$:NH$_2$	sodium amide
Na^+	$^-$:OCH$_3$	sodium methoxide	Li^+	$^-$:N[CH(CH$_3$)$_2$]$_2$	lithium diisopropylamide
Na^+	$^-$:OCH$_2$CH$_3$	sodium ethoxide			
K^+	$^-$:OC(CH$_3$)$_3$	potassium *tert*-butoxide	Hydride		
			Na^+	H:$^-$	sodium hydride


Strong bases have a net negative charge, but not all negatively charged species are strong bases. For example, none of the halides, F^-, Cl^-, Br^-, or I^-, is a strong base. These anions have very strong conjugate acids and have little affinity for donating their electron pairs to a proton.

Carbanions, negatively charged carbon atoms discussed in Section 2.5D, are especially strong bases. Perhaps the most common example is **butyllithium.** Butyllithium and related compounds are discussed in greater detail in Chapter 20.


$CH_3CH_2CH_2CH_2^-$ Li^+

butyllithium

Two other weaker organic bases are **triethylamine** and **pyridine.** These compounds have a lone pair on nitrogen, making them basic, but they are considerably weaker than the amide bases because they are neutral, not negatively charged.



triethylamine




pyridine

Problem 2.28 Draw the products formed when propan-2-ol [$(CH_3)_2CHOH$], the main ingredient in rubbing alcohol, is treated with each acid or base: (a) NaH; (b) H_2SO_4; (c) $Li^+{}^-N[CH(CH_3)_2]_2$; (d) CH_3CO_2H.

2.7 Aspirin

Aspirin, or acetylsalicylic acid, is the most well known member of a group of compounds called **salicylates.** Although aspirin was first used in medicine for its analgesic (pain-relieving), antipyretic (fever-reducing), and anti-inflammatory properties, today it is commonly used as an antiplatelet agent in the treatment and prevention of heart attacks and strokes. **Aspirin is a synthetic compound;** it does not occur in nature, although some related salicylates are found in willow bark and meadowsweet blossoms (Figure 2.7).

Figure 2.7
Salicin, an analgesic in willow bark




salicin



willow tree

- The modern history of aspirin dates back to 1763 when Reverend Edmund Stone reported on the analgesic effect of chewing on the bark of the willow tree. Willow bark is now known to contain *salicin,* which is structurally related to aspirin.

Like many drugs, aspirin undergoes a proton transfer reaction. Its most acidic proton is the H bonded to O, and in the presence of base, this H is readily removed.



aspirin
acetylsalicylic acid
neutral form

This form exists in the **stomach.**

conjugate base
ionic form


This form exists in the **intestines.**

Why is this acid–base reaction important? After ingestion, aspirin first travels into the stomach and then the intestines. In the acidic environment of the stomach, aspirin remains in its neutral form, but in the basic environment of the small intestine, aspirin is deprotonated to form its conjugate base, an ion. Likewise, in the slightly basic environment of the blood, aspirin exists primarily as its ionic conjugate base.

We will learn more about solubility and the cell membrane in Section 3.7.

Whether aspirin is a neutral acid or an ionic conjugate base affects its transport throughout the body and its ability to pass through a cell membrane. In its ionic form, aspirin is readily soluble in the aqueous environment of the blood, so it is transported in the bloodstream to tissues. Once aspirin has reached its target location, however, its conjugate base must be re-protonated to form the neutral acid that can pass through the nonpolar interior of a cell membrane where it inhibits prostaglandin synthesis, as we will learn in Chapter 19. Thus, in the body, aspirin undergoes acid–base reactions and these reactions are crucial in determining its properties and action.

Problem 2.29



amphetamine

Compounds like amphetamine that contain nitrogen atoms are protonated by the HCl in the gastric juices of the stomach, and the resulting salt is then deprotonated in the basic environment of the intestines to regenerate the neutral form. Write proton transfer reactions for both of these processes. In which form will amphetamine pass through a cell membrane?

2.8 Lewis Acids and Bases

The Lewis definition of acids and bases is more general than the Brønsted–Lowry definition.

- A Lewis acid is an *electron pair acceptor.*
- A Lewis base is an *electron pair donor.*


All Brønsted–Lowry bases are Lewis bases.

Lewis bases are structurally the same as Brønsted–Lowry bases. Both have an **available electron pair**—a lone pair or an electron pair in a π bond. A Brønsted–Lowry base always donates this electron pair to a proton, but a Lewis base donates this electron pair to anything that is electron deficient. Simple Lewis bases are shown in Figure 2.8.

A Lewis acid must be able to accept an electron pair, but there are many ways for this to occur. **All Brønsted–Lowry acids are also Lewis acids, but the reverse is not necessarily true.** Any species that is electron deficient and capable of accepting an electron pair is also a Lewis acid, as shown in Figure 2.8.

Common examples of Lewis acids (which are not Brønsted–Lowry acids) include BF_3 and $AlCl_3$. These compounds contain elements in group 3A of the periodic table that can accept an electron pair because they do not have filled valence shells of electrons.

Figure 2.8
Simple Lewis acids and bases



Problem 2.30 Which species are Lewis bases?

a. NH_3 b. $CH_3CH_2CH_3$ c. H^- d. $H-C\equiv C-H$

Problem 2.31 Which species are Lewis acids?


a. BBr_3 b. CH_3CH_2OH c. $(CH_3)_3C^+$ d. Br^-

Any reaction in which one species donates an electron pair to another species is a Lewis acid–base reaction.

In a Lewis acid–base reaction, a Lewis base donates an electron pair to a Lewis acid. Most reactions in organic chemistry involving movement of electron pairs can be classified as Lewis acid–base reactions. Lewis acid–base reactions illustrate a general pattern of reactivity.

- **Electron-rich species react with electron-poor species.**

In the simplest Lewis acid–base reaction one bond is formed and no bonds are broken. This is illustrated with the reaction of BF_3 with H_2O. BF_3 has only six electrons around B, so it is the electron-deficient Lewis acid. H_2O has two lone pairs on O, so it is the electron-rich Lewis base.




H_2O donates an electron pair to BF_3 to form one new bond. The electron pair in the new B–O bond comes from the oxygen atom, and a single product, a Lewis acid–base complex, is formed. Both B and O bear formal charges in the product, but the overall product is neutral.

Nucleophile = nucleus loving.
Electrophile = electron loving.


- **A Lewis acid is called an *electrophile.***
- **When a Lewis base reacts with an electrophile other than a proton, the Lewis base is called a *nucleophile.***

In this Lewis acid–base reaction, **BF_3 is the electrophile** and **H_2O is the nucleophile.**

Two other examples are drawn. In each reaction the **electron pair is not removed from the Lewis base;** instead, the electron pair is donated to an atom of the Lewis acid, and one new covalent bond is formed.




Problem 2.32 For each reaction, label the Lewis acid and base. Use curved arrow notation to show the movement of electron pairs.

a. 

b. 

Problem 2.33 Draw the products of each reaction, and label the nucleophile and electrophile.

a. 

Problem 2.34 Draw the product formed when $(CH_3CH_2)_3N$:, a Lewis base, reacts with each Lewis acid: (a) $B(CH_3)_3$; (b) $(CH_3)_3C^+$; (c) $AlCl_3$.

In some Lewis acid–base reactions, one bond is formed and one bond is broken. To draw the products of these reactions, keep the following steps in mind.

[1] Always identify the Lewis acid and base first.
[2] Draw a curved arrow from the electron pair of the base to the electron-deficient atom of the acid.
[3] Count electron pairs and break a bond when needed to keep the correct number of valence electrons.

For example, draw the Lewis acid–base reaction between cyclohexene and H−Cl. The Brønsted–Lowry acid HCl is also a Lewis acid, and cyclohexene, having a π bond, is the Lewis base.




To draw the product of this reaction, the electron pair in the π bond of the Lewis base forms a new bond to the proton of the Lewis acid, forming a carbocation. The H−Cl bond must break, giving its two electrons to Cl, forming Cl^-. Because two electron pairs are involved, two curved arrows are needed.



The Lewis acid–base reaction of cyclohexene with HCl is a specific example of a fundamental reaction of compounds containing C−C double bonds, as discussed in Chapter 10.

Recall from Chapter 1 that a positively charged carbon atom is called a **carbocation.**



In the reaction of cyclohexene with HCl, the new bond to H could form at **either carbon of the double bond,** because the same carbocation results.

Problem 2.35 Label the Lewis acid and base. Use curved arrow notation to show the movement of electron pairs.



KEY CONCEPTS


Acids and Bases

A Comparison of Brønsted–Lowry and Lewis Acids and Bases


Type	Definition	Structural feature	Examples
Brønsted–Lowry acid (2.1)	proton donor	a proton	HCl, H_2SO_4, H_2O, CH_3CO_2H, TsOH
Brønsted–Lowry base (2.1)	proton acceptor	a lone pair *or* a π bond	^-OH, $^-OCH_3$, H^-, $^-NH_2$, NH_3, $CH_2=CH_2$
Lewis acid (2.8)	electron pair acceptor	a proton, *or* an unfilled valence shell, *or* a partial (+) charge	BF_3, $AlCl_3$, HCl, CH_3CO_2H, H_2O
Lewis base (2.8)	electron pair donor	a lone pair *or* a π bond	^-OH, $^-OCH_3$, H^-, $^-NH_2$, NH_3, $CH_2=CH_2$

Acid–Base Reactions

[1] A Brønsted–Lowry acid donates a proton to a Brønsted–Lowry base (2.2).



[2] A Lewis base donates an electron pair to a Lewis acid (2.8).




- Electron-rich species react with electron-poor ones.
- Nucleophiles react with electrophiles.

Important Facts

- Definition: **$pK_a = -\log K_a$. The lower the pK_a, the stronger the acid** (2.3).
- The stronger the acid, the weaker the conjugate base (2.3).
- In proton transfer reactions, equilibrium favors the weaker acid and weaker base (2.4).
- An acid can be deprotonated by the conjugate base of any acid having a higher pK_a (2.4).

Periodic Trends in Acidity and Basicity (2.5A)




Factors That Determine Acidity (2.5)

[1] **Element effects** (2.5A)	The acidity of HA increases both left-to-right across a row and down a column of the periodic table.
[2] **Inductive effects** (2.5B)	The acidity of HA increases with the presence of electron-withdrawing groups in A.
[3] **Resonance effects** (2.5C)	The acidity of HA increases when the conjugate base $A:^-$ is resonance stabilized.
[4] **Hybridization effects** (2.5D)	The acidity of HA increases as the percent *s*-character of $A:^-$ increases.

PROBLEMS


Problems Using Three-Dimensional Models

2.36 Propranolol is an antihypertensive agent—that is, it lowers blood pressure. (a) Which proton in propranolol is most acidic? (b) What products are formed when propranolol is treated with NaH? (c) Which atom is most basic? (d) What products are formed when propranolol is treated with HCl?



propranolol


2.37 Amphetamine is a powerful stimulant of the central nervous system. (a) Which proton in amphetamine is most acidic? (b) What products are formed when amphetamine is treated with NaH? (c) What products are formed when amphetamine is treated with HCl?






amphetamine

Brønsted–Lowry Acids and Bases

2.38 What is the conjugate acid of each base?





a. HCO_3^- b.  c.  d. 

2.39 What is the conjugate base of each acid?





a. HCO_3^- b.  c.  d. 

Reactions of Brønsted–Lowry Acids and Bases

2.40 Draw the products formed from the acid–base reaction of H_2SO_4 with each compound.


a.  b.  c.  d. 

2.41 Draw the products formed from the acid–base reaction of KOH with each compound.


a.  b.  c.  d. 

2.42 Draw the products of each proton transfer reaction. Label the acid and base in the starting materials, and the conjugate acid and base in the products.


a. 

b. 

c. 


d. 

2.43 Draw the products of each acid–base reaction.

a. 

+ NaOH ⇌


naproxen
anti-inflammatory agent

b. 


+ HCl ⇌

fluoxetine
antidepressant

2.44 Fenfluramine and phentermine are two components of fen–phen, an appetite suppressant withdrawn from the market in 1997 after it was shown to damage the heart valves in some patients. What products are formed when fenfluramine and phentermine are each treated with acetic acid (CH_3CO_2H)?



fenfluramine



phentermine

pK_a, K_a, and the Direction of Equilibrium


2.45 What is K_a for each compound? Use a calculator when necessary.

a. H_2S pK_a = 7.0


b. $ClCH_2COOH$ pK_a = 2.8

c. HCN pK_a = 9.1


2.46 What is the pK_a for each compound?

a. 

$K_a = 4.7 \times 10^{-10}$

b. 


$K_a = 2.3 \times 10^{-5}$

c. 

$K_a = 5.9 \times 10^{-1}$

2.47 Which of the following bases are strong enough to deprotonate $CH_3CH_2CH_2C{\equiv}CH$ (pK_a = 25), so that equilibrium favors the products: (a) H_2O; (b) NaOH; (c) $NaNH_2$; (d) NH_3; (e) NaH; (f) CH_3Li?

2.48 Which compounds can be deprotonated by ^-OH, so that equilibrium favors the products? Refer to the pK_a table in Appendix A.

a. HCOOH b. H_2S c. 


d. CH_3NH_2

2.49 Draw the products of each reaction. Use the pK_a table in Appendix A to decide if the equilibrium favors the starting materials or products.

a. CH_3NH_2 + H_2SO_4 ⇌

b. 

+ NaCl ⇌


c. 


+ $NaHCO_3$ ⇌

d. H–C≡C–H + $CH_3CH_2^-Li^+$ ⇌


Relative Acid Strength


2.50 Rank the following compounds in order of increasing acidity.


a. 









b. 





c. 








d. 



2.51 Rank the following ions in order of increasing basicity.



a. $CH_3\bar{C}H_2$, CH_3O^-, $CH_3\bar{N}H$

b. CH_3^-, HO^-, Br^-


c.   

d.   

2.52 The pK_a values of the two ammonium cations drawn below are 8.33 and 11.1. Which pK_a corresponds to which cation? Explain your choice.




 

2.53 Explain why the C – H_a bond is much more acidic than the C – H_b bond in pentan-2-one.



pentan-2-one

2.54 The pK_a of three C – H bonds is given below.


pK_a = 50 pK_a = 43 pK_a = 19.2

a. For each compound, draw the conjugate base, including all possible resonance structures.

b. Explain the observed trend in pK_a.

2.55 a. What is the conjugate acid of **A?**

b. What is the conjugate base of **A?**




A

2.56 Draw the structure of a constitutional isomer of compound **B** that fits each description.

a. an isomer that is at least 10^5 times more acidic than **B**

b. an isomer that is at least 10^5 times less acidic than **B**

c. an isomer that is comparable in acidity to **B**




B


2.57 Many drugs are Brønsted–Lowry acids or bases.

a. What is the most acidic proton in the analgesic ibuprofen? Draw the conjugate base.

b. What is the most basic electron pair in cocaine? Draw the conjugate acid.




ibuprofen



cocaine

2.58 Dimethyl ether (CH_3OCH_3) and ethanol (CH_3CH_2OH) are isomers, but CH_3OCH_3 has a pK_a of 40 and CH_3CH_2OH has a pK_a of 16. Why are these pK_a values so different?


2.59 Atenolol is a β (beta) blocker, a drug used to treat high blood pressure. Which of the indicated N – H bonds is more acidic? Explain your reasoning.



atenolol


2.60 Use the principles in Section 2.5 to label the most acidic hydrogen in each drug. Explain your choice.

a.




valproic acid
(used to treat epilepsy)

b.



paroxetine
trade name Paxil
(used to treat depression)


c.



metoprolol
(used to treat high blood pressure)

2.61 Label the three most acidic hydrogen atoms in lactic acid, $CH_3CH(OH)CO_2H$, and rank them in order of decreasing acidity. Explain your reasoning.

2.62 Bupivacaine (trade name Marcaine) is a quick-acting anesthetic often used during labor and delivery. Which nitrogen atom in bupivacaine is more basic? Explain your reasoning.



bupivacaine

Lewis Acids and Bases

2.63 Classify each compound as a Lewis base, a Brønsted–Lowry base, both, or neither.


a.  b.  c.  d. 

2.64 Classify each species as a Lewis acid, a Brønsted–Lowry acid, both, or neither.

a. H_3O^+ b. Cl_3C^+ c. BCl_3 d. BF_4^-


Lewis Acid–Base Reactions

2.65 Label the Lewis acid and Lewis base in each reaction. Use curved arrows to show the movement of electron pairs.

a. $Cl^- + BCl_3 \longrightarrow$ 

b.  $+ {}^-OH \longrightarrow$ 

2.66 Draw the products of each Lewis acid–base reaction. Label the electrophile and nucleophile.

a.  $+ AlCl_3 \longrightarrow$


b.  $+ BF_3 \longrightarrow$

c.  $+ H_2O \longrightarrow$

2.67 Draw the product formed when the Lewis acid $(CH_3CH_2)_3C^+$ reacts with each Lewis base: (a) CH_3OH; (b) $(CH_3)_2O$; (c) $(CH_3)_2NH$.

General Problems


2.68 Answer the following questions about the four species **A–D.**



A **B** **C** **D**


a. Which two species represent a conjugate acid–base pair?
b. Which two species represent resonance structures?
c. Which two species represent constitutional isomers?

2.69 Classify each reaction as either a proton transfer reaction, or a reaction of a nucleophile with an electrophile. Use curved arrows to show how the electron pairs move.



2.70 Hydroxide ($^-$OH) can react as a Brønsted–Lowry base (and remove a proton), or a Lewis base (and attack a carbon atom). (a) What organic product is formed when $^-$OH reacts with the carbocation $(CH_3)_3C^+$ as a Brønsted–Lowry base? (b) What organic product is formed when $^-$OH reacts with $(CH_3)_3C^+$ as a Lewis base?

2.71 Answer the following questions about esmolol, a drug used to treat high blood pressure sold under the trade name Brevibloc.




esmolol

a. Label the most acidic hydrogen atom in esmolol.
b. What products are formed when esmolol is treated with NaH?
c. What products are formed when esmolol is treated with HCl?
d. Label all sp^2 hybridized C atoms.
e. Label the only trigonal pyramidal atom.
f. Label all C's that bear a $\delta+$ charge.

Challenge Problems


2.72 DBU, 1,8-diazabicyclo[5.4.0]undec-7-ene, is a base we will encounter in elimination reactions in Chapter 8. Which N atom is more basic in DBU? Explain your choice.



DBU

2.73 Molecules like acetamide (CH_3CONH_2) can be protonated on either their O or N atoms when treated with a strong acid like HCl. Which site is more readily protonated and why?


2.74 Two pK_a values are reported for malonic acid, a compound with two COOH groups. Explain why one pK_a is lower and one pK_a is higher than the pK_a of acetic acid (CH_3COOH, pK_a = 4.8).



malonic acid

pK_a = 2.86 pK_a = 5.70


2.75 Amino acids such as glycine are the building blocks of large molecules called proteins that give structure to muscle, tendon, hair, and nails.




glycine zwitterion form

a. Explain why glycine does not actually exist in the form with all atoms uncharged, but actually exists as a salt called a zwitterion.
b. What product is formed when glycine is treated with concentrated HCl?
c. What product is formed when glycine is treated with NaOH?

2.76 Write a stepwise reaction sequence using proton transfer reactions to show how the following reaction occurs. (Hint: As a first step, use ^-OH to remove a proton from the CH_2 group between the C=O and C=C.)




2.77 Which H atom in vitamin C (ascorbic acid) is most acidic?



vitamin C
ascorbic acid


2.78 Which compound, **M** or **N,** is the stronger acid? Explain your choice.



M **N**

Introduction to Organic Molecules and Functional Groups

3



Vitamin C, or **ascorbic acid,** is important in the formation of collagen, a protein that holds together the connective tissues of skin, muscle, and blood vessels. Citrus fruits—oranges, grapefruit, and lemons—are well known sources of vitamin C. A deficiency of vitamin C causes scurvy, a common disease of sailors in the 1600s when they had no access to fresh fruits on long voyages. Early Arctic explorers, however, were able to obtain vitamin C by eating scurvy grass, a low-growing plant found along the coasts of the Arctic region of the northern hemisphere. In Chapter 3, we learn why some vitamins like vitamin A can be stored in the fat cells in the body, whereas others like vitamin C are excreted in urine.

3.1 Functional groups
3.2 An overview of functional groups
3.3 Intermolecular forces
3.4 Physical properties
3.5 Application: Vitamins
3.6 Application of solubility: Soap
3.7 Application: The cell membrane
3.8 Functional groups and reactivity
3.9 Biomolecules

Having learned some basic concepts about structure, bonding, and acid–base chemistry in Chapters 1 and 2, we will now concentrate on organic molecules.

- What are the characteristic features of an organic compound?
- What determines the properties of an organic compound?

After these questions are answered, we can understand some common phenomena. For example, why do we store some vitamins in the body and readily excrete others? How does soap clean away dirt? We will also use the properties of organic molecules to explain some basic biological phenomena, such as the structure of cell membranes and the transport of species across these membranes.

3.1 Functional Groups

What are the characteristic features of an organic compound? Most organic molecules have C−C and C−H σ bonds. These bonds are strong, nonpolar, and not readily broken. Organic molecules may have the following structural features as well:


- **Heteroatoms—atoms other than carbon or hydrogen.** Common heteroatoms are nitrogen, oxygen, sulfur, phosphorus, and the halogens.
- **π Bonds.** The most common π bonds occur in C−C and C−O double bonds.

These structural features distinguish one organic molecule from another. They determine a molecule's geometry, physical properties, and reactivity, and comprise what is called a **functional group.**

- A *functional group* is an atom or a group of atoms with characteristic chemical and physical properties. It is the *reactive part* of the molecule.

Why do heteroatoms and π bonds confer reactivity on a particular molecule?

- Heteroatoms have lone pairs and create electron-deficient sites on carbon.
- π Bonds are easily broken in chemical reactions. A π bond makes a molecule a base and a nucleophile.




- Lone pairs make O a base and a nucleophile.
- The C atom is electron deficient, making it an electrophile.




- The π bond is easily broken.
- The π bond makes a compound a base and a nucleophile.

Don't think, though, that the C−C and C−H σ bonds are unimportant. They form the **carbon backbone** or **skeleton** to which the functional groups are bonded. A functional group usually behaves the same whether it is bonded to a carbon skeleton having as few as two or as many as 20 carbons. For this reason, we often abbreviate the carbon and hydrogen portion of the molecule by a capital letter **R,** and draw the **R** bonded to a particular functional group.



Ethane, for example, has only C−C and C−H σ bonds, so it has *no* functional group. Ethane has no polar bonds, no lone pairs, and no π bonds, so it has **no reactive sites.** Because of this, ethane and molecules like it are very unreactive.

Ethanol, on the other hand, has two carbons and five hydrogens in its carbon backbone, as well as an OH group, a functional group called a **hydroxy** group. Ethanol has lone pairs and polar bonds that make it reactive with a variety of reagents, including the acids and bases discussed in Chapter 2. The hydroxy group makes the properties of ethanol very different from the properties of ethane. Moreover, any organic molecule containing a hydroxy group has properties similar to ethanol.



ethane

- all C–C and C–H σ bonds
- no functional group

ethanol

- polar C–O and O–H bonds
- two lone pairs

Most organic compounds can be grouped into a relatively small number of categories, based on the structure of their functional group. Ethane, for example, is an **alkane,** whereas ethanol is a simple **alcohol.**

Problem 3.1 What reaction occurs when CH_3CH_2OH is treated with (a) H_2SO_4? (b) NaH? What happens when CH_3CH_3 is treated with these same reagents?

3.2 An Overview of Functional Groups

We can subdivide the most common functional groups into three types. A list of functional groups is presented on the inside front cover.

- **Hydrocarbons**
- **Compounds containing a C – Z σ bond** where Z = an electronegative element
- **Compounds containing a C = O group**

3.2A Hydrocarbons

To review the structure and bonding of the simple aliphatic hydrocarbons, return to Section 1.10.

The word *aliphatic* is derived from the Greek word *aleiphas* meaning *fat.* Aliphatic compounds have physical properties similar to fats.


Hydrocarbons **are compounds made up of only the elements carbon and hydrogen. They may be aliphatic or aromatic.**

[1] Aliphatic hydrocarbons. Aliphatic hydrocarbons can be divided into three subgroups.

- **Alkanes** have only C – C σ bonds and no functional group. Ethane, CH_3CH_3, is a simple alkane.
- **Alkenes** have a C – C double bond as a functional group. Ethylene, $CH_2=CH_2$, is a simple alkene.
- **Alkynes** have a C – C triple bond as a functional group. Acetylene, $HC\equiv CH$, is a simple alkyne.

[2] Aromatic hydrocarbons. This class of hydrocarbons was so named because many of the earliest known aromatic compounds had strong, characteristic odors.

The simplest aromatic hydrocarbon is **benzene.** The six-membered ring and three π bonds of benzene comprise a *single* functional group. Benzene is a component of the **BTX** mixture (**B** for **b**enzene) added to gasoline to boost octane ratings.



benzene
molecular formula C_6H_6

phenyl group
C_6H_5–
phenylcyclohexane


When a benzene ring is bonded to another group, it is called a **phenyl group.** In phenylcyclohexane, for example, a phenyl group is bonded to the six-membered cyclohexane ring. Table 3.1 summarizes the four different types of hydrocarbons.

Table 3.1 Hydrocarbons

Type of compound	General structure	Example	Functional group
Alkane	R—H	CH_3CH_3	—
Alkene			double bond
Alkyne	—C≡C—	H—C≡C—H	triple bond
Aromatic compound			phenyl group

Polyethylene is a synthetic plastic first produced in the 1930s, and initially used as insulating material for radar during World War II. It is now a plastic used in milk containers, sandwich bags, and plastic wrapping. Over 100 billion pounds of polyethylene are manufactured each year.

Alkanes, which have no functional groups, are notoriously unreactive except under very drastic conditions. For example, **polyethylene** is a synthetic plastic and high molecular weight alkane, consisting of chains of $-CH_2-$ groups bonded together, hundreds or even thousands of atoms long. Because it is an alkane with no reactive sites, it is a very stable compound that does not readily degrade and thus persists for years in landfills.



Carbon atoms in alkanes and other organic compounds are classified by the number of other carbons directly bonded to them.


- A *primary carbon* (1° carbon) is bonded to *one* other C atom.
- A *secondary carbon* (2° carbon) is bonded to *two* other C atoms.
- A *tertiary carbon* (3° carbon) is bonded to *three* other C atoms.
- A *quaternary carbon* (4° carbon) is bonded to *four* other C atoms.



One example of each type of C is labeled.


Hydrogen atoms are classified as **primary (1°), secondary (2°),** or **tertiary (3°)** depending on the **type of carbon atom** to which they are bonded.

- A *primary hydrogen* (1° H) is on a C bonded to one other C atom.
- A *secondary hydrogen* (2° H) is on a C bonded to two other C atoms.
- A *tertiary hydrogen* (3° H) is on a C bonded to three other C atoms.




One example of each type of H is labeled.

Sample Problem 3.1 Classify the designated carbon atoms in **A** as 1°, 2°, 3°, or 4°. Classify the designated hydrogen atoms in **B** as 1°, 2°, or 3°.




A




B

Solution




A




B

Problem 3.2 (a) Classify the carbon atoms in each compound as 1°, 2°, 3°, or 4°. (b) Classify the hydrogen atoms in each compound as 1°, 2°, or 3°.



Problem 3.3 Classifying a carbon atom by the number of carbons to which it is bonded can also be done in more complex molecules that contain heteroatoms. Classify each sp^3 hybridized carbon atom in bilobalide, a compound isolated from *Ginkgo biloba* extracts, as 1°, 2°, 3°, or 4°.



bilobalide



Bilobalide is obtained from *Ginkgo biloba,* the oldest seed-producing plant that currently lives on earth. Extracts from the leaves, roots, bark, and seeds of the ginkgo tree have been used in traditional Chinese medicine and currently comprise the most widely taken herbal supplements.

3.2B Compounds Containing C–Z σ Bonds

Functional groups that contain C–Z σ bonds include alkyl halides, alcohols, ethers, amines, thiols, and sulfides (Table 3.2). The electronegative heteroatom Z creates a polar bond, making carbon electron deficient. The lone pairs on Z are available for reaction with protons and other electrophiles, especially when Z = N or O.








Chloroethane, CH_3CH_2Cl, is a local anesthetic.

Several simple compounds in this category are widely used. As an example, chloroethane (CH_3CH_2Cl, commonly called ethyl chloride) is an alkyl halide used as a local anesthetic. Chloroethane quickly evaporates when sprayed on a wound, causing a cooling sensation that numbs the site of an injury.


Table 3.2 Compounds Containing C–Z σ Bonds

Type of compound	General structure	Example	3-D structure	Functional group
Alkyl halide	R–X: (X = F, Cl, Br, I)	CH_3–Br:		**–X** halo group
Alcohol	R–OH	CH_3–O–H		**–OH** hydroxy group
Ether	R–O–R	CH_3–O–CH_3		**–OR** alkoxy group
Amine	R–NH_2 or R_2NH or R_3N	CH_3–N(H)–H		**–NH_2** amino group
Thiol	R–SH	CH_3–S–H		**–SH** mercapto group
Sulfide	R–S–R	CH_3–S–CH_3		**–SR** alkylthio group

Molecules containing these functional groups may be simple or very complex. Diethyl ether, the first common general anesthetic, is a simple ether because it contains a single O atom, depicted in red, bonded to two C atoms. Hemibrevetoxin B, on the other hand, contains four ether groups, in addition to other functional groups.




diethyl ether




hemibrevetoxin B

Hemibrevetoxin B is a neurotoxin produced by algal blooms referred to as "red tides," because of the color often seen in shallow ocean waters when these algae proliferate.

Alkyl halides and alcohols are classified as **primary (1°), secondary (2°),** or **tertiary (3°)** based on the number of carbon atoms bonded to the carbon bearing the halogen or OH group.




Z = X 1° alkyl halide 2° alkyl halide 3° alkyl halide
Z = OH 1° alcohol 2° alcohol 3° alcohol



2° chloride
3° alcohol
1° alcohol


Problem 3.4 Classify each alkyl halide and alcohol as 1°, 2°, or 3°.



Problem 3.5 Classify each OH group and halogen in dexamethasone, a synthetic steroid, as 1°, 2°, or 3°.




Dexamethasone relieves inflammation and is used to treat some forms of arthritis, skin conditions, and asthma.




dexamethasone

Amines are classified as **primary (1°), secondary (2°),** or **tertiary (3°)** based on the number of carbon atoms bonded to the nitrogen atom.



Classifying amines is different from classifying alcohols and alkyl halides as primary (1°), secondary (2°), or tertiary (3°). Amines are classified by the number of carbon–*nitrogen* bonds, whereas alkyl halides and alcohols are classified by the type of *carbon* bonded to the halogen or hydroxy group.

Problem 3.6 Classify each amine in the following compounds as 1°, 2°, or 3°.




spermine
(isolated from semen)

meperidine
(a narcotic)
trade name Demerol

Problem 3.7 Draw the structure of a compound of molecular formula $C_4H_{11}NO$ that fits each description: (a) a compound that contains a 1° amine and a 3° alcohol; (b) a compound that contains a 3° amine and a 1° alcohol.

3.2C Compounds Containing a C=O Group

Many different types of functional groups possess a C−O double bond (a **carbonyl group**), including aldehydes, ketones, carboxylic acids, esters, amides, and acid chlorides (Table 3.3). The polar C−O bond makes the carbonyl carbon an electrophile, while the lone pairs on O allow it to react as a nucleophile and base. The carbonyl group also contains a π bond that is more easily broken than a C−O σ bond.



carbonyl group

Amides, compounds that contain a nitrogen atom bonded directly to the carbonyl carbon, are classified as **primary (1°), secondary (2°),** or **tertiary (3°)** based on the number of carbon atoms bonded to the nitrogen atom.


















Table 3.3 Compounds Containing a C=O Group

Type of compound	General structure	Example	Condensed structure	3-D structure	Functional group
Aldehyde			CH_3CHO		
Ketone			$(CH_3)_2CO$		 carbonyl group
Carboxylic acid			CH_3CO_2H		 carboxy group
Ester			$CH_3CO_2CH_3$		
Amide			CH_3CONH_2		
Acid chloride			CH_3COCl		

Problem 3.8 Classify the amides in dolastatin, an anticancer compound isolated from the Indian seahare *Dolabella auricularia*, as 1°, 2°, or 3°.




dolastatin

The importance of a functional group cannot be overstated. A functional group determines a molecule's bonding and shape, type and strength of intermolecular forces, physical properties, nomenclature, and chemical reactivity.

Problem 3.9 Oseltamivir can be prepared in 10 steps from shikimic acid. Identify the functional groups in oseltamivir and shikimic acid.




Tamiflu is the trade name for oseltamivir, an antiviral drug used to treat influenza.



shikimic acid — oseltamivir

Atenolol and donepezil are examples of useful drugs that contain a variety of functional groups. Atenolol is a β (beta) blocker, a group of drugs used to treat hypertension (high blood pressure). Donepezil, sold under the trade name Aricept, is used to treat mild to moderate dementia associated with Alzheimer's disease.



atenolol (used to treat high blood pressure) — donepezil (used to treat Alzheimer's disease)

Problem 3.10 Draw the structure of a compound fitting each description:

a. an aldehyde with molecular formula C_4H_8O
b. a ketone with molecular formula C_4H_8O
c. a carboxylic acid with molecular formula $C_4H_8O_2$
d. an ester with molecular formula $C_4H_8O_2$


Problem 3.11 Draw structures that fit each description and name the functional group in each molecule: (a) two constitutional isomers with molecular formula $C_5H_{10}O$ that contain different functional groups; (b) two constitutional isomers with molecular formula $C_6H_{10}O$ that contain the same functional group.

3.3 Intermolecular Forces

Intermolecular forces are also referred to as **noncovalent interactions** or **nonbonded interactions.**

Intermolecular forces are the interactions that exist *between* molecules. A functional group determines the type and strength of these interactions.

3.3A Ionic Compounds



strong electrostatic interaction between Na^+ and Cl^-

Ionic compounds, such as NaCl, contain oppositely charged particles held together by **extremely strong electrostatic interactions.** These ionic interactions are much *stronger* than the intermolecular forces present between covalent molecules, so it takes a great deal of energy to separate oppositely charged ions from each other.

3.3B Covalent Compounds

Covalent compounds are composed of discrete molecules. The nature of the forces between the molecules depends on the functional group present. There are three different types of interactions, presented here in order of *increasing strength:*

- **van der Waals forces**
- **dipole–dipole interactions**
- **hydrogen bonding**

Van der Waals Forces






Although any single van der Waals interaction is weak, a large number of van der Waals interactions creates a strong force. For example, geckos stick to walls and ceilings by van der Waals interactions of the surfaces with the 500,000 tiny hairs on each foot.

Van der Waals forces, also called **London forces,** are very weak interactions caused by the **momentary changes in electron density in a molecule.** Van der Waals forces are the only attractive forces present in nonpolar compounds.

For example, although a nonpolar CH_4 molecule has no net dipole, at any one instant its electron density may not be completely symmetrical, creating a *temporary* dipole. This can induce a temporary dipole in another CH_4 molecule, with the partial positive and negative charges arranged close to each other. **The weak interaction of these temporary dipoles constitutes van der Waals forces.** All compounds exhibit van der Waals forces.



The surface area of a molecule determines the strength of the van der Waals interactions. **The larger the surface area, the larger the attractive force between two molecules, and the stronger the intermolecular forces.** Long, sausage-shaped molecules such as $CH_3CH_2CH_2CH_2CH_3$ (pentane) have stronger van der Waals interactions than compact spherical ones like $C(CH_3)_4$ (2,2-dimethylpropane), as shown in Figure 3.1.


Another factor affecting the strength of van der Waals forces is **polarizability.**

- *Polarizability* is a measure of how the electron cloud around an atom responds to changes in its electronic environment.

Larger atoms like iodine, which have more loosely held valence electrons, are more polarizable than smaller atoms like fluorine, which have more tightly held electrons. Because larger atoms have more easily induced dipoles, compounds containing them possess stronger intermolecular interactions.


Figure 3.1
Surface area and van der Waals forces

a. $CH_3CH_2CH_2CH_2CH_3$ (pentane)




- A long, cylindrical molecule like pentane has a larger surface area, causing stronger van der Waals interactions.

b. $C(CH_3)_4$ (2,2-dimethylpropane)




- A compact, spherical molecule like 2,2-dimethylpropane has a smaller surface area, causing weaker van der Waals interactions.

Two F_2 molecules have little force of attraction between them, because the electrons are held very tightly and temporary dipoles are difficult to induce. On the other hand, two I_2 molecules exhibit a much stronger force of attraction, because the electrons are held much more loosely and temporary dipoles are easily induced.



Dipole–Dipole Interactions


Dipole–dipole interactions **are the attractive forces between the permanent dipoles of two polar molecules.** In acetone, $(CH_3)_2C{=}O$, for example, the dipoles in adjacent molecules align so that the partial positive and partial negative charges are in close proximity. These attractive forces caused by permanent dipoles are much stronger than weak van der Waals forces.



Hydrogen Bonding

Hydrogen bonding helps determine the three-dimensional shape of large biomolecules such as carbohydrates and proteins. See Chapters 28 and 29 for details.

Hydrogen bonding **typically occurs when a hydrogen atom bonded to O, N, or F is electrostatically attracted to a lone pair of electrons on an O, N, or F atom in another molecule.** Thus, H_2O molecules can hydrogen bond to each other. When they do, a H atom covalently bonded to O in one water molecule is attracted to a lone pair of electrons on the O in another water molecule. Hydrogen bonds are the strongest of the three types of intermolecular forces, though they are still much weaker than any covalent bond.




Sample Problem 3.2 illustrates how to determine the relative strength of intermolecular forces for a group of compounds. Table 3.4 summarizes the four types of interactions that affect the properties of all compounds.

Table 3.4 Summary of Types of Intermolecular Forces

Type of force	Relative strength	Exhibited by	Example
van der Waals	weak	all molecules	$CH_3CH_2CH_2CH_2CH_3$ $CH_3CH_2CH_2CHO$ $CH_3CH_2CH_2CH_2OH$
dipole–dipole	moderate	molecules with a net dipole	$CH_3CH_2CH_2CHO$ $CH_3CH_2CH_2CH_2OH$
hydrogen bonding	strong	molecules with an O – H, N – H, or H – F bond	$CH_3CH_2CH_2CH_2OH$
ion–ion	very strong	ionic compounds	NaCl, LiF

Sample Problem 3.2 Rank the following compounds in order of increasing strength of intermolecular forces: $CH_3CH_2CH_2CH_2CH_3$ (pentane), $CH_3CH_2CH_2CH_2OH$ (butan-1-ol), and $CH_3CH_2CH_2CHO$ (butanal).

Solution




pentane
- nonpolar molecule


butan-1-ol
- bent molecule around O
- polar C–O and O–H bonds
- O–H bond for hydrogen bonding

butanal
- trigonal planar C
- polar C=O bond

- Pentane has only nonpolar C – C and C – H bonds, so its molecules are held together by only **van der Waals** forces.
- Butan-1-ol is a polar bent molecule, so it can have **dipole–dipole** interactions in addition to **van der Waals** forces. Because it has an O – H bond, butan-1-ol molecules are held together by intermolecular **hydrogen bonds** as well.
- Butanal has a trigonal planar carbon with a polar C = O bond, so it exhibits **dipole–dipole** interactions in addition to **van der Waals** forces. There is *no* H atom bonded to O, so two butanal molecules *cannot* hydrogen bond to each other.



Problem 3.12 What types of intermolecular forces are present in each compound?



3.4 Physical Properties


The strength of a compound's intermolecular forces determines many of its physical properties, including its boiling point, melting point, and solubility.

3.4A Boiling Point (bp)


The *boiling point* of a compound is the temperature at which a liquid is converted to a gas. In boiling, energy is needed to overcome the attractive forces in the more ordered liquid state.

- The stronger the intermolecular forces, the higher the boiling point.

Because **ionic compounds** are held together by extremely strong interactions, they have **very high boiling points.** The boiling point of NaCl, for example, is 1413 °C. **With covalent molecules, the boiling point depends on the identity of the functional group.** For compounds of approximately the same molecular weight:



Recall from Sample Problem 3.2, for example, that the relative strength of the intermolecular forces increases from pentane to butanal to butan-1-ol. The boiling points of these compounds increase in the same order.




Because surface area and polarizability affect the strength of intermolecular forces, they also affect the boiling point. For two compounds with similar functional groups:


- The larger the surface area, the higher the boiling point.
- The more polarizable the atoms, the higher the boiling point.

Examples of each phenomenon are illustrated in Figure 3.2. In comparing two ketones that differ in size, pentan-3-one has a higher boiling point than acetone because it has a greater molecular weight and larger surface area. In comparing two alkyl halides having the same number of carbon atoms, CH_3I has a higher boiling point than CH_3F because I is more polarizable than F.

Figure 3.2
Effect of surface area and polarizability on boiling point




Sample Problem 3.3 Which compound in each pair has the higher boiling point?



Solution

a. Isomers **A** and **B** have only nonpolar C – C and C – H bonds, so they exhibit only van der Waals forces. Because **B** is more compact, it has less surface area and a lower boiling point.
b. Compounds **C** and **D** have approximately the same molecular weight but different functional groups. **C** is a nonpolar alkane, exhibiting only van der Waals forces. **D** is an alcohol with an O – H group available for hydrogen bonding, so it has stronger intermolecular forces and a higher boiling point.


Problem 3.13 Which compound in each pair has the higher boiling point?



Problem 3.14 Explain why the boiling point of propanamide, $CH_3CH_2CONH_2$, is considerably higher than the boiling point of *N,N*-dimethylformamide, $HCON(CH_3)_2$ (213 °C vs. 153 °C), even though both compounds are isomeric amides.

Liquids having different boiling points can be separated in the laboratory using a *distillation* apparatus (Figure 3.3). When a mixture of two liquids is heated in the distilling flask, the lower boiling compound, the **more volatile component,** distills first, followed by the **less volatile, higher boiling component.** By collecting the distillate in a series of receiver flasks, the two liquids can usually be separated from each other. The best separations are generally achieved when the liquids in the mixture have widely different boiling points.

Figure 3.3
Schematic of a distillation apparatus




By periodically changing the receiver flask, one can collect compounds having different boiling points in separate flasks.

3.4B Melting Point (mp)


The *melting point* is the temperature at which a solid is converted to its liquid phase. In melting, energy is needed to overcome the attractive forces in the more ordered crystalline solid. Two factors determine the melting point of a compound.

- **The stronger the intermolecular forces, the higher the melting point.**
- **Given the same functional group, the more symmetrical the compound, the higher the melting point.**


Because **ionic compounds** are held together by extremely strong interactions, they have **very high melting points.** For example, the melting point of NaCl is 801 °C. With covalent molecules, the melting point once again depends on the identity of the functional group. For compounds of approximately the same molecular weight:




The trend in the melting points of pentane, butanal, and butan-1-ol parallels the trend observed in their boiling points.



Symmetry also plays a role in determining the melting points of compounds having the same functional group and similar molecular weights, but very different shapes. A compact symmetrical molecule like 2,2-dimethylpropane packs well into a crystalline lattice whereas 2-methylbutane, which has a CH_3 group dangling from a four-carbon chain, does not. Thus, 2,2-dimethylpropane has a much higher melting point.



Problem 3.15 Predict which compound in each pair has the higher melting point.




Problem 3.16 Consider acetic acid (CH_3CO_2H) and its conjugate base, sodium acetate (CH_3CO_2Na). (a) What intermolecular forces are present in each compound? (b) Explain why the melting point of sodium acetate (324 °C) is considerably higher than the melting point of acetic acid (17 °C).

3.4C Solubility

Quantitatively, a compound may be considered soluble when 3 g of solute dissolves in 100 mL of solvent.

***Solubility* is the extent to which a compound, called the *solute*, dissolves in a liquid, called the *solvent*.** In dissolving a compound, the energy needed to break up the interactions between the molecules or ions of the solute comes from new interactions between the solute and the solvent.




Compounds dissolve in solvents having similar kinds of intermolecular forces.

- **"Like dissolves like."**
- **Polar compounds dissolve in polar solvents. Nonpolar or weakly polar compounds dissolve in nonpolar or weakly polar solvents.**

Water and **organic liquids** are two different kinds of solvents. Water is very polar because it is capable of hydrogen bonding with a solute. Many organic solvents are either nonpolar, like carbon tetrachloride (CCl_4) and hexane [$CH_3(CH_2)_4CH_3$], or weakly polar like diethyl ether ($CH_3CH_2OCH_2CH_3$).

Ionic compounds are held together by strong electrostatic forces, so they need very polar solvents to dissolve. **Most ionic compounds are soluble in water, but are insoluble in organic solvents.** To dissolve an ionic compound, the strong ion–ion interactions must be replaced by many weaker **ion–dipole interactions,** as illustrated in Figure 3.4.


Figure 3.4
Dissolving an ionic compound in H_2O



- When an ionic solid is dissolved in H_2O, the ion–ion interactions are replaced by ion–dipole interactions. Though these forces are weaker, there are so many of them that they compensate for the stronger ionic bonds.


Most organic compounds are soluble in organic solvents (remember, *like dissolves like*). **An organic compound is water soluble only if it contains one polar functional group capable of hydrogen bonding with the solvent for every five C atoms it contains.** In other words, a water-soluble organic compound has an O- or N-containing functional group that solubilizes its nonpolar carbon backbone.

Compare, for example, the solubility of butane and acetone in H_2O and CCl_4.




Because butane and acetone are both organic compounds having a C–C and C–H backbone, they are soluble in the organic solvent CCl_4. Butane, a nonpolar molecule, is insoluble in the polar solvent H_2O. Acetone, however, is H_2O soluble because it contains only three C atoms and its O atom can hydrogen bond with one H atom of H_2O. In fact, acetone is so soluble in water that acetone and water are **miscible**—they form solutions in all proportions with each other.

$(CH_3)_2C{=}O$ molecules cannot hydrogen bond to each other because they have no OH group. However, $(CH_3)_2C{=}O$ can hydrogen bond to H_2O because its O atom can hydrogen bond to one of the H atoms of H_2O.




For an organic compound with one functional group, **a compound is water soluble only if it has ≤ five C atoms and contains an O or N atom.**

The size of an organic molecule with a polar functional group determines its water solubility. A low molecular weight alcohol like **ethanol is water soluble** because it has a small carbon skeleton (≤ five C atoms) compared to the size of its polar OH group. Cholesterol, on the other hand, has 27 carbon atoms and only one OH group. Its carbon skeleton is too large for the OH group to solubilize by hydrogen bonding, so **cholesterol is insoluble in water.**



ethanol

H_2O soluble



cholesterol


H_2O insoluble

Hydrophobic = afraid of H_2O.
Hydrophilic = H_2O loving.


- The nonpolar part of a molecule that is not attracted to H_2O is said to be *hydrophobic.*
- The polar part of a molecule that can hydrogen bond to H_2O is said to be *hydrophilic.*

In cholesterol, for example, the **hydroxy group is hydrophilic,** whereas the **carbon skeleton is hydrophobic.**

MTBE (*tert*-butyl methyl ether) and 4,4'-dichlorobiphenyl (a polychlorinated biphenyl, abbreviated as PCB) demonstrate that solubility properties can help determine the fate of organic compounds in the environment.



MTBE
tert-butyl methyl ether



4,4'-dichlorobiphenyl
(a polychlorinated biphenyl, PCB)

Using **MTBE** as a high-octane additive in unleaded gasoline has had a negative environmental impact. Although MTBE is not toxic or carcinogenic, it has a distinctive, nauseating odor, and **it is water soluble.** Small amounts of MTBE have contaminated the drinking water in several communities, making it unfit for consumption. For this reason, the use of MTBE as a gasoline additive has steadily declined in the United States since 1999.


4,4'-Dichlorobiphenyl is a polychlorinated biphenyl **(PCB),** a compound that contains two benzene rings joined by a C−C bond, and substituted by one or more chlorine atoms on each ring. PCBs have been used as plasticizers in polystyrene coffee cups and coolants in transformers. They have been released into the environment during production, use, storage, and disposal, making them one of the most widespread organic pollutants. **PCBs are insoluble in H_2O, but very soluble in organic media,** so they are soluble in fatty tissue, including that found in all types of fish and birds around the world. Although PCBs are not acutely toxic, frequently ingesting large quantities of fish contaminated with PCBs has been shown to retard growth and memory retention in children.

Solubility properties of some representative compounds are summarized in Table 3.5.


Table 3.5 Summary of Solubility

Type of compound	Solubility in H_2O	Solubility in organic solvents (such as CCl_4)
Ionic		
NaCl	**soluble**	**insoluble**
Covalent		
$CH_3CH_2CH_2CH_3$	**insoluble** (no N or O atom to hydrogen bond to H_2O)	**soluble**
$CH_3CH_2CH_2OH$	**soluble** (≤ 5 C's and an O atom for hydrogen bonding to H_2O)	**soluble**
$CH_3(CH_2)_{10}OH$	**insoluble** (> 5 C's; too large to be soluble even though it has an O atom for hydrogen bonding to H_2O)	**soluble**

Sample Problem 3.4 Which compounds are water soluble?



chlorocyclopentane
A





cyclopentanol
B


Solution

A has five C atoms and a polar C – Cl bond, but it is incapable of hydrogen bonding with H_2O, making it H_2O insoluble. **B** has five C atoms and a polar OH group able to hydrogen bond to H_2O, making it H_2O soluble.


Problem 3.17 Which compounds are water soluble?

a. 


b. 

c. 

Sample Problem 3.5
a. Which of the following molecules can hydrogen bond to another molecule like itself?
b. Which of the following molecules can hydrogen bond with water?




diethyl ether
A




diethylamine
B

Solution

a. Compounds **A** and **B** have polar C – O and C – N bonds but **A** does not have an O – H bond, so it cannot hydrogen bond to another **A** molecule. **B,** on the other hand, has an N – H bond, so two **B** molecules can hydrogen bond with each other.
b. Because **A** has an electronegative O atom and **B** has an electronegative N atom, both **A** and **B** can hydrogen bond with a hydrogen atom in water.




A




B

Problem 3.18 (a) Label the hydrophobic and hydrophilic portions of each molecule. (b) At which sites can **C** hydrogen bond to another molecule like itself? (c) At which sites can **D** hydrogen bond to water?



C
norethindrone
(oral contraceptive component)



D
arachidonic acid
(fatty acid)

3.5 Application: Vitamins


***Vitamins* are organic compounds needed in small amounts for normal cell function.** Our bodies cannot synthesize these compounds, so they must be obtained in the diet. Most vitamins are identified by a letter, such as A, C, D, E, and K. There are several different B vitamins, though, so a subscript is added to distinguish them: for example, B_1, B_2, and B_{12}.

Whether a vitamin is **fat soluble** (it dissolves in organic media) or **water soluble** can be determined by applying the solubility principles discussed in Section 3.4C. Vitamins A and C illustrate the differences between fat-soluble and water-soluble vitamins.

3.5A Vitamin A

The name **vitamin** was first used in 1912 by the Polish chemist Casimir Funk, who called them *vitamines,* because he thought that they all contained an *amine* functional group. Later the word was shortened to vitamin, because some are amines but others, like vitamins A and C, are not.

Vitamin A, or **retinol,** is an essential component of the vision receptors in the eyes. It also helps to maintain the health of mucous membranes and the skin, so many anti-aging creams contain vitamin A. A deficiency of this vitamin leads to a loss of night vision.




vitamin A

Vitamin A contains 20 carbons and a single OH group, making it **water insoluble.** Because it is organic, it is **soluble in any organic medium.** To understand the consequences of these solubility characteristics, we must learn about the chemical environment of the body.


Vitamin A is synthesized from β-carotene, the orange pigment in carrots.

About 70% of the body is composed of water. Fluids such as blood, gastric juices in the stomach, and urine are largely water with dissolved ions such as Na^+ and K^+. Vitamin A is insoluble in these fluids. There are also fat cells composed of organic compounds having C−C and C−H bonds. Vitamin A is soluble in this organic environment, and thus it is readily stored in these fat cells, particularly in the liver.

Vitamin A may be obtained directly from the diet. In addition, β-carotene, the orange pigment found in many plants including carrots, is readily converted to vitamin A in our bodies.



vitamin A


Eating too many carrots does not result in an excess of stored vitamin A. If you consume more β-carotene than you need, your body stores this precursor until it needs more vitamin A. Some β-carotene reaches the surface tissues of the skin and eyes, giving them an orange color. This phenomenon may look odd, but it is harmless and reversible. When stored β-carotene is converted to vitamin A and is no longer in excess, these tissues will return to their normal hue.



Vitamin C is obtained by eating citrus fruits and a wide variety of other fruits and vegetables. Individuals can also obtain the recommended daily dose of vitamin C by taking tablets that contain vitamin C prepared in the laboratory. Both the "natural" vitamin C in oranges and the "synthetic" vitamin C in vitamin supplements are identical.

3.5B Vitamin C

Although most animal species can synthesize vitamin C, humans, guinea pigs, the Indian fruit bat, and the bulbul bird must obtain this vitamin from dietary sources. Citrus fruits, strawberries, tomatoes, and sweet potatoes are all excellent sources of vitamin C.




vitamin C
(ascorbic acid)

Vitamin C has six carbon atoms, each bonded to an oxygen atom that is capable of hydrogen bonding, making it **water soluble.** Vitamin C thus dissolves in urine. Although it has been acclaimed as a deterrent for all kinds of diseases, from the common cold to cancer, the consequences of taking large amounts of vitamin C are not really known, because any excess of the minimum daily requirement is excreted in the urine.


Problem 3.19 Predict the water solubility of each vitamin.

a.




vitamin B_3
(niacin)

b.



vitamin K_1
(phylloquinone)

Problem 3.20 (a) Identify the functional groups in the ball-and-stick model of pantothenic acid, vitamin B_5. (b) At which sites can pantothenic acid hydrogen bond to water? (c) Predict the water solubility of pantothenic acid.



pantothenic acid
vitamin B_5




Avocados are an excellent dietary source of pantothenic acid, vitamin B_5.

3.6 Application of Solubility: Soap

Soap has been used by humankind for some 2000 years. Historical records describe its manufacture in the first century and document the presence of a soap factory in Pompeii. Before this time clothes were cleaned by rubbing them on rocks in water, or by forming soapy lathers from the roots, bark, and leaves of certain plants. These plants produced natural materials called *saponins,* which act in much the same way as modern soaps.

On a molecular level, soap has two distinct parts:

- **a hydrophilic portion composed of ions called the *polar head***
- **a hydrophobic carbon chain of nonpolar C−C and C−H bonds, called the *nonpolar tail***




ionic end — long, hydrocarbon chain

polar head **nonpolar tail** **soap**

Dissolving soap in water forms ***micelles*, spherical droplets having the ionic heads on the surface and the nonpolar tails packed together in the interior,** as shown in Figure 3.5. In this arrangement, the ionic heads are solvated by the polar solvent water, thus solubilizing the nonpolar, "greasy" hydrocarbon portion of the soap.


Figure 3.5
Dissolving soap in water




- When soap is dissolved in H_2O, it forms micelles with the nonpolar tails in the interior and the polar heads on the surface. The polar heads are solvated by ion–dipole interactions with H_2O molecules.

How does soap dissolve grease and oil? Water alone cannot dissolve dirt, which is composed largely of nonpolar hydrocarbons. When soap is mixed with water, however, the nonpolar hydrocarbon tails dissolve the dirt in the interior of the micelle. The polar head of the soap remains on the surface of the micelle to interact with water. The nonpolar tails of the soap are so well sealed off from the water by the polar head groups that the micelles are water soluble, allowing them to separate from the fibers of our clothes and be washed down the drain with water. In this way, soaps do a seemingly impossible task: they remove nonpolar hydrocarbon material from skin and clothes, by solubilizing it in the polar solvent water.


Cross-section of a soap micelle with a grease particle dissolved in the interior



Problem 3.21 Which of the following structures represent soaps? Explain your answers.



Problem 3.22 Today, synthetic detergents like the compound drawn here, not soaps, are used to clean clothes. Explain how this detergent cleans away dirt.



a detergent


3.7 Application: The Cell Membrane

The cell membrane is a beautifully complex example of how the principles of organic chemistry come into play in a biological system.

3.7A Structure of the Cell Membrane

The basic unit of living organisms is the **cell.** The cytoplasm is the aqueous medium inside the cell, separated from water outside the cell by the **cell membrane.** The cell membrane serves two apparently contradictory functions. It acts as a barrier to the passage of ions, water, and other molecules into and out of the cell, and it is also selectively permeable, letting nutrients in and waste out.

A major component of the cell membrane is a group of organic compounds called **phospholipids.** Like soap, they contain a hydrophilic ionic portion and a hydrophobic hydrocarbon portion, in this case two long carbon chains composed of C−C and C−H bonds. **Phospholipids thus contain a polar head and *two* nonpolar tails.**




phospholipid

When phospholipids are mixed with water, they assemble in an arrangement called a **lipid bilayer,** with the ionic heads oriented on the outside and the nonpolar tails on the inside. The polar heads electrostatically interact with the polar solvent H_2O, while the nonpolar tails are held in close proximity by numerous van der Waals interactions. This is schematically illustrated in Figure 3.6.

Cell membranes are composed of these lipid bilayers. The charged heads of the phospholipids are oriented toward the aqueous interior and exterior of the cell. The nonpolar tails form the hydrophobic interior of the membrane, thus serving as an insoluble barrier that protects the cell from the outside.


The nonpolar interior of the cell membrane is especially important in protecting the human brain from fluctuation in the concentration of compounds in the blood, as well as the passage of unwanted substances into the brain. The blood–brain barrier consists of a tight layer of cells in the blood capillaries of the brain, and all substances must pass through the cell membrane of these capillaries to enter the brain. Because ions are not soluble in the nonpolar interior of the cell membrane, the blood–brain barrier is only slightly permeable to ions. On the other hand,


Figure 3.6
The cell membrane





- Phospholipids contain an ionic or polar head, and two long nonpolar hydrocarbon tails. In an aqueous environment, phospholipids form a lipid bilayer, with the polar heads oriented toward the aqueous exterior and the nonpolar tails forming a hydrophobic interior. Cell membranes are composed largely of this lipid bilayer.

uncharged organic molecules like nicotine, caffeine, and heroin are very soluble in the interior of the cell membrane, so they readily pass into the brain.


nicotine



caffeine



heroin


sevoflurane

General anesthetics such as sevoflurane are also weakly polar compounds that can penetrate the blood–brain barrier because they are soluble in the lipid bilayer of the blood capillaries.

Problem 3.23 (a) What types of intermolecular forces do morphine and heroin each possess? (b) Which compound can cross the blood–brain barrier more readily, and therefore serve as the more potent pain reliever?


morphine



heroin

Problem 3.24 Explain why the noble gas xenon is a general anesthetic.


3.7B Transport Across a Cell Membrane

How does a polar molecule or ion in the water outside a cell pass through the nonpolar interior of the cell membrane and enter the cell? Some nonpolar molecules like O_2 are small enough to enter and exit the cell by diffusion. Polar molecules and ions, on the other hand, may be too large or too polar to diffuse efficiently. Some ions are transported across the membrane with the help of molecules called **ionophores.**

Ionophores **are organic molecules that complex cations.** They have a hydrophobic exterior that makes them soluble in the nonpolar interior of the cell membrane, and a central cavity with several oxygen atoms whose lone pairs complex with a given ion. The size of the cavity determines the identity of the cation with which the ionophore complexes. Two naturally occurring antibiotics that act as ionophores are **nonactin** and **valinomycin.**




nonactin




valinomycin

Several synthetic ionophores have also been prepared, including one group called **crown ethers.** ***Crown ethers*** **are cyclic ethers containing several oxygen atoms that bind specific cations depending on the size of their cavity.** Crown ethers are named according to the general format ***x*-crown-*y,*** where ***x*** is the total number of atoms in the ring and ***y*** is the number of oxygen atoms. For example, 18-crown-6 contains 18 atoms in the ring, including 6 O atoms. This crown ether binds potassium ions. Sodium ions are too small to form a tight complex with the O atoms, and larger cations do not fit in the cavity.




18-crown-6



complex with K^+

How does an ionophore transfer an ion across a membrane? The ionophore binds the ion on one side of the membrane in its polar interior. It can then move across the membrane because its

Figure 3.7
Transport of ions across a cell membrane



- By binding an ion on one side of a lipid bilayer (where the concentration of the ion is high) and releasing it on the other side of the bilayer (where the concentration of the ion is low), an ionophore transports an ion across a cell membrane.

hydrophobic exterior interacts with the hydrophobic tails of the phospholipid. The ionophore then releases the ion on the other side of the membrane. This ion-transfer role is essential for normal cell function. This process is illustrated in Figure 3.7.

In this manner, antibiotic ionophores like nonactin transport ions across a cell membrane of bacteria. This disrupts the normal ionic balance in the cell, thus interfering with cell function and causing the bacteria to die.

Problem 3.25 Now that you have learned about solubility, explain why aspirin (Section 2.7) crosses a cell membrane as a neutral carboxylic acid rather than an ionic conjugate base.


3.8 Functional Groups and Reactivity


Much of Chapter 3 has been devoted to how a functional group determines the strength of intermolecular forces and, consequently, the physical properties of molecules. A functional group also determines reactivity. What type of reaction does a particular kind of organic compound undergo? Begin by recalling two fundamental concepts.

- Functional groups create reactive sites in molecules.
- Electron-rich sites react with electron-poor sites.

All functional groups contain a heteroatom, a π bond, or both, and these features make electron-deficient (or electrophilic) sites and electron-rich (or nucleophilic) sites in a molecule. To predict reactivity, first locate the functional group and then determine the resulting electron-rich or electron-deficient sites it creates. Keep three guidelines in mind.

- An electronegative heteroatom like N, O, or X makes a carbon atom *electrophilic.*






- A lone pair on a heteroatom makes it basic and nucleophilic.



base
nucleophile




base
nucleophile

- **π Bonds create *nucleophilic* sites and are more easily broken than σ bonds.**



Problem 3.26 Label the electrophilic and nucleophilic sites in each molecule.




By identifying the nucleophilic and electrophilic sites in a compound you can begin to understand how it will react. In general, electron-rich sites react with electron-deficient sites:

:Nu⁻ = a nucleophile; E⁺ = an electrophile.

- **An electron-deficient carbon atom reacts with a nucleophile, symbolized as :Nu⁻.**
- **An electron-rich carbon reacts with an electrophile, symbolized as E⁺.**


At this point we don't know enough organic chemistry to draw the products of many reactions with confidence. We do know enough, however, to begin to predict if two compounds might react together based solely on electron density arguments, and at what atoms that reaction is most likely to occur.

For example, alkenes contain an electron-rich C–C double bond, so they react with electrophiles, E^+. On the other hand, alkyl halides possess an electrophilic carbon atom, so they react with electron-rich nucleophiles.




You don't need to worry about the products of these reactions. At this point you should only be able to find reactive sites in molecules and begin to understand why a reaction might occur at these sites. After you learn more about the structure of organic molecules in Chapters 4 and 5, we will begin a detailed discussion of organic reactions in Chapter 6.

Problem 3.27 Considering only electron density, state whether the following reactions will occur.




3.9 Biomolecules


***Biomolecules* are organic compounds found in biological systems.** Many are relatively small, with molecular weights of less than 1000 g/mol. There are four main families of these small molecules—simple sugars, amino acids, lipids, and nucleotides. Many simple biomolecules are used to synthesize larger compounds that have important cellular functions.




glucose
a simple sugar



oleic acid
a fatty acid



alanine
an amino acid




deoxyadenosine 5'-monophosphate
a nucleotide

Simple sugars such as glucose combine to form the complex carbohydrates starch and cellulose, as described in Chapter 28. Alanine is an amino acid used to synthesize proteins, the subject of Chapter 29. Fatty acids such as oleic acid react with alcohols to form triacylglycerols, the most prevalent lipids, first mentioned in Chapter 10, and discussed in more detail in Chapters 22 and 31. While these biomolecules all contain more than one functional group, their properties and reactions are explained by the principles of basic organic chemistry.


Finally, deoxyadenosine 5'-monophosphate is a nucleotide that combines with thousands of other nucleotides to form DNA, deoxyribonucleic acid, the high molecular weight polynucleotide that stores the genetic information of an organism. DNA consists of two polynucleotide chains that wind together in a double helix. Figure 3.8 illustrates the importance of hydrogen bonding in the structure of DNA. The two polynucleotide chains are held together by an extensive network of hydrogen bonds in which the N−H groups on one chain intermolecularly hydrogen bond to an oxygen or nitrogen atom on the adjacent chain.

Figure 3.8
The double helix of DNA



- DNA, which is contained in the chromosomes of the nucleus of a cell, stores all of the genetic information in an organism. DNA consists of two long strands of polynucleotides held together by hydrogen bonding.

Problem 3.28 The fact that sweet-tasting carbohydrates like table sugar are also high in calories has prompted the development of sweet, low-calorie alternatives. (a) Identify the functional groups in aspartame, the artificial sweetener in Equal. (b) Label all of the sites that can hydrogen bond to the oxygen atom of water. (c) Label all of the sites that can hydrogen bond with a hydrogen atom of water.



aspartame

KEY CONCEPTS

Introduction to Organic Molecules and Functional Groups

Classifying Carbon Atoms, Hydrogen Atoms, Alcohols, Alkyl Halides, Amines, and Amides (3.2)

- Carbon atoms are classified by the number of carbons bonded to them; a 1° carbon is bonded to one other carbon, and so forth.
- Hydrogen atoms are classified by the type of carbon atom to which they are bonded; a 1° hydrogen is bonded to a 1° carbon, and so forth.
- Alkyl halides and alcohols are classified by the type of carbon to which the OH or X group is bonded; a 1° alcohol has an OH group bonded to a 1° carbon, and so forth.
- Amines and amides are classified by the number of carbons bonded to the nitrogen atom; a 1° amine has one carbon–nitrogen bond, and so forth.


Types of Intermolecular Forces (3.3)

Increasing strength

Type of force	Cause
van der Waals	Caused by the interaction of temporary dipoles • Larger surface area, stronger forces • Larger, more polarizable atoms, stronger forces
dipole–dipole	Caused by the interaction of permanent dipoles
hydrogen bonding	Caused by the electrostatic interaction of a H atom in an O – H, N – H, or H – F bond with the lone pair of another N, O, or F atom
ion–ion	Caused by the charge attraction of two ions

Physical Properties

Property	Observation
Boiling point (3.4A)	• For compounds of comparable molecular weight, the stronger the intermolecular forces the higher the bp.




VDW
bp = 36 °C

VDW, DD
bp = 76 °C

VDW, DD, HB
bp = 118 °C

Increasing strength of intermolecular forces
Increasing boiling point

- For compounds with similar functional groups, the larger the surface area, the higher the bp.




- For compounds with similar functional groups, the more polarizable the atoms, the higher the bp.




Melting point (3.4B)

- For compounds of comparable molecular weight, the stronger the intermolecular forces the higher the mp.



- For compounds with similar functional groups, the more symmetrical the compound, the higher the mp.



Solubility (3.4C)

Types of water-soluble compounds:

- Ionic compounds
- Organic compounds having ≤ 5 C's, and an O or N atom for hydrogen bonding (for a compound with one functional group)

Types of compounds soluble in organic solvents:

- Organic compounds regardless of size or functional group

Key: VDW = van der Waals, DD = dipole–dipole, HB = hydrogen bonding


Reactivity (3.8)

- **Nucleophiles react with electrophiles.**
- Electronegative heteroatoms create electrophilic carbon atoms that react with nucleophiles.
- Lone pairs and π bonds are nucleophilic sites that react with electrophiles.

PROBLEMS

Problems with Three-Dimensional Models


3.29



elemicin

a. Identify the functional groups in the ball-and-stick model of elemicin, a compound partly responsible for the flavor and fragrance of nutmeg.
b. Draw a skeletal structure of a constitutional isomer of elemicin that should have a higher boiling point and melting point.
c. Label all electrophilic carbon atoms.

3.30




neral


a. Identify the functional groups in the ball-and-stick model of neral, a compound with a lemony odor isolated from lemon grass.
b. Draw a skeletal structure of a constitutional isomer of neral that should be more water soluble.
c. Label the most electrophilic carbon atom.

Functional Groups

3.31 For each alkane: (a) classify each carbon atom as 1°, 2°, 3°, or 4°; (b) classify each hydrogen atom as 1°, 2°, or 3°.




A



B


3.32 Identify the functional groups in each molecule. Classify each alcohol, alkyl halide, amide, and amine as 1°, 2°, or 3°.

a.




Darvon
(analgesic)

c.




ibuprofen
(analgesic)

e.




penicillin G
(an antibiotic)

b.




pregabalin
trade name Lyrica
(used in treating chronic pain)

d.



histrionicotoxin
(poison secreted by a
South American frog)


f.



pyrethrin I
(potent insecticide
from chrysanthemums)

3.33 Identify each functional group located in the following rings. Which structure represents a lactone—a cyclic ester—and which represents a lactam—a cyclic amide?


a.




b.




c.



d.



3.34 (a) Identify the functional groups in salinosporamide A, an anticancer agent isolated from marine sediment. (b) Classify each alcohol, alkyl halide, amide, and amine as 1°, 2°, or 3°.




salinosporamide A


3.35 Draw seven constitutional isomers with molecular formula $C_3H_6O_2$ that contain a carbonyl group. Identify the functional group(s) in each isomer.

Intermolecular Forces


3.36 What types of intermolecular forces are exhibited by each compound?



3.37 Rank the compounds in each group in order of increasing strength of intermolecular forces.




3.38 Indinavir (trade name Crixivan) is a drug used to treat HIV. (a) At which sites can indinavir hydrogen bond to another molecule like itself? (b) At which sites can indinavir hydrogen bond to water?



indinavir

3.39 Intramolecular forces of attraction are often important in holding large molecules together. For example, some proteins fold into compact shapes, held together by attractive forces between nearby functional groups. A schematic of a folded protein is drawn here, with the protein backbone indicated by a blue-green ribbon, and various appendages drawn dangling from the chain. What types of intramolecular forces occur at each labeled site (**A–F**)?



Physical Properties

3.40 (a) Draw four compounds with molecular formula $C_6H_{12}O$, each containing at least one different functional group. (b) Predict which compound has the highest boiling point, and explain your reasoning.

3.41 Rank the compounds in each group in order of increasing boiling point.

a.

b.

3.42 Explain why $CH_3CH_2NHCH_3$ has a higher boiling point than $(CH_3)_3N$, even though they have the same molecular weight.

3.43 Menthone and menthol are both isolated from mint. Explain why menthol is a solid at room temperature but menthone is a liquid.

menthone menthol

3.44 Rank the following compounds in order of increasing melting point.

3.45 Explain why benzene has a lower boiling point but much higher melting point than toluene.

benzene
bp = 80 °C
mp = 5 °C


and


toluene
bp = 111 °C
mp = –93 °C


3.46 Rank the following compounds in order of increasing water solubility.

3.47 Which of the following molecules can hydrogen bond to another molecule of itself? Which can hydrogen bond with water?

a. 


b. 

c. 


d. 

3.48 Explain why diethyl ether ($CH_3CH_2OCH_2CH_3$) and butan-1-ol ($CH_3CH_2CH_2CH_2OH$) have similar solubility properties in water, but butan-1-ol has a much higher boiling point.


3.49 Predict the water solubility of each of the following organic molecules.

a. 


caffeine
(stimulant in coffee, tea, and many soft drinks)

c. 

sucrose
(table sugar)

b. 


mestranol
(component in oral contraceptives)

d. 


carotatoxin
(neurotoxin isolated from carrots)

Applications

3.50 Predict the solubility of each of the following vitamins in water and in organic solvents.


a. 

vitamin E


b. 

pyridoxine
vitamin B_6

3.51 Avobenzone and dioxybenzone are two commercial sunscreens. Using the principles of solubility, predict which sunscreen is more readily washed off when an individual goes swimming. Explain your choice.




avobenzone




dioxybenzone

3.52 Poly(ethylene glycol) (PEG) and poly(vinyl chloride) (PVC) are examples of polymers, large organic molecules composed of repeating smaller units covalently bonded together. Polymers have very different properties depending (in part) on their functional groups. Discuss the water solubility of each polymer and suggest why PEG is used in shampoos, whereas PVC is used to make garden hoses and pipes. Synthetic polymers are discussed in detail in Chapters 15 and 30.




poly(ethylene glycol)
PEG



poly(vinyl chloride)
PVC

3.53 THC is the active component in marijuana, and ethanol is the alcohol in alcoholic beverages. Explain why drug screenings are able to detect the presence of THC but not ethanol weeks after these substances have been introduced into the body.




tetrahydrocannabinol
THC




ethanol

3.54 Cocaine is a widely abused, addicting drug. Cocaine is usually obtained as its hydrochloride salt (cocaine hydrochloride) but can be converted to crack (the neutral molecule) by treatment with base. Which of the two compounds here has a higher boiling point? Which is more soluble in water? How does the relative solubility explain why crack is usually smoked but cocaine hydrochloride is injected directly into the bloodstream?




cocaine (crack)
neutral organic molecule



cocaine hydrochloride
a salt

3.55 Many drugs are sold as their hydrochloride salts ($R_2NH_2^+$ Cl^-), formed by reaction of an amine (R_2NH) with HCl.





acebutolol


a. Draw the product (a hydrochloride salt) formed by reaction of acebutolol with HCl. Acebutolol is a β blocker used to treat high blood pressure.
b. Discuss the solubility of acebutolol and its hydrochloride salt in water.
c. Offer a reason as to why the drug is marketed as a hydrochloride salt rather than a neutral amine.

Reactivity of Organic Molecules

3.56 Label the electrophilic and nucleophilic sites in each molecule.


a. 


b. 


c. 


d. 

3.57 By using only electron density arguments, determine whether the following reactions will occur.

a. 


c. 

b. 

d. 


Cell Membrane

3.58 The composition of a cell membrane is not uniform for all types of cells. Some cell membranes are more rigid than others. Rigidity is determined by a variety of factors, one of which is the structure of the carbon chains in the phospholipids that comprise the membrane. One example of a phospholipid was drawn in Section 3.7A, and another, having C – C double bonds in its carbon chains, is drawn here. Which phospholipid would be present in the more rigid cell membrane and why?




phospholipid

3.59 Which compound is more likely to be a general anesthetic? Explain your choice.




A



B

General Questions


3.60 Quinapril (trade name Accupril) is a drug used to treat hypertension and congestive heart failure.



quinapril

a. Identify the functional groups in quinapril.
b. Classify any alcohol, amide, or amine as 1°, 2°, or 3°.
c. At which sites can quinapril hydrogen bond to water?
d. At which sites can quinapril hydrogen bond to acetone [$(CH_3)_2CO$]?
e. Label the most acidic hydrogen atom.
f. Which site is most basic?

3.61 Answer each question about oxycodone, a narcotic analgesic used for severe pain.




oxycodone

a. Identify the functional groups in oxycodone.
b. Classify any alcohol, amide, or amine as 1°, 2°, or 3°.
c. Which proton is most acidic?
d. Which site is most basic?
e. What is the hybridization of the N atom?
f. How many sp^2 hybridized C atoms does oxycodone contain?

Challenge Problems

3.62 Although diethyl ether and tetrahydrofuran are both four-carbon ethers, one compound is much more water soluble than the other. Predict which compound has higher water solubility and offer an explanation.




diethyl ether



tetrahydrofuran


3.63 Answer the following questions by referring to the ball-and-stick model of fentanyl, a potent narcotic analgesic used in surgical procedures.




fentanyl

a. Identify the functional groups.
b. Label the most acidic proton.
c. Label the most basic atom.
d. What types of intermolecular forces are present between two molecules of fentanyl?
e. Draw an isomer predicted to have a higher boiling point.
f. Which sites in the molecule can hydrogen bond to water?
g. Label all electrophilic carbons.

3.64 Explain why **A** is less water soluble than **B,** even though both compounds have the same functional groups.




A




B

3.65 Recall from Section 1.10B that there is restricted rotation around carbon–carbon double bonds. Maleic acid and fumaric acid are two isomers with vastly different physical properties and pK_a values for loss of both protons. Explain why each of these differences occurs.



maleic acid



fumaric acid

	maleic acid	fumaric acid
mp (°C)	130	286
solubility (g/L) in H_2O at 25 °C	788	7
pK_{a1}	1.9	3.0
pK_{a2}	6.5	4.5

4 Alkanes

4.1 Alkanes—An introduction
4.2 Cycloalkanes
4.3 An introduction to nomenclature
4.4 Naming alkanes
4.5 Naming cycloalkanes
4.6 Common names
4.7 Fossil fuels
4.8 Physical properties of alkanes
4.9 Conformations of acyclic alkanes—Ethane
4.10 Conformations of butane
4.11 An introduction to cycloalkanes
4.12 Cyclohexane
4.13 Substituted cycloalkanes
4.14 Oxidation of alkanes
4.15 Lipids—Part 1



Alkanes, the simplest hydrocarbons, are found in all shapes and sizes and occur widely in nature. They are the major constituents of petroleum, a complex mixture of compounds that includes hydrocarbons such as **hexane** and **decane.** Crude petroleum spilled into the sea from a ruptured oil tanker or offshore oil well creates an insoluble oil slick on the surface. Petroleum is refined to produce gasoline, diesel fuel, home heating oil, and a myriad of other useful compounds. In Chapter 4, we learn about the properties of alkanes, how to name them (nomenclature), and oxidation—one of their important reactions.

In Chapter 4, we apply the principles of bonding, shape, and reactivity discussed in Chapters 1–3 to our first family of organic compounds, the **alkanes.** Because alkanes have no functional group, they are much less reactive than other organic compounds, and for this reason, much of Chapter 4 is devoted to learning how to name and draw them, as well as to understanding what happens when rotation occurs about their carbon–carbon single bonds.

Studying alkanes also provides an opportunity to learn about **lipids,** a group of biomolecules similar to alkanes, in that they are composed mainly of nonpolar carbon–carbon and carbon–hydrogen σ bonds. Section 4.15 serves as a brief introduction only, so we will return to lipids in Chapters 10 and 31 (online).

4.1 Alkanes—An Introduction



Secretion of **undecane** by a cockroach causes other members of the species to aggregate. Undecane is a ***pheromone,* a chemical substance used for communication** in an animal species, most commonly an insect population.




Cyclohexane is one component of the mango, the most widely consumed fruit in the world.

Recall from Section 3.2 that **alkanes are aliphatic hydrocarbons having only C−C and C−H σ bonds.** Because their carbon atoms can be joined together in chains or rings, they can be categorized as acyclic or cyclic.

- **Acyclic alkanes** have the molecular formula C_nH_{2n+2} (where n = an integer) and contain only linear and branched chains of carbon atoms. Acyclic alkanes are also called **saturated hydrocarbons** because they have the maximum number of hydrogen atoms per carbon.
- **Cycloalkanes** contain carbons joined in one or more rings. Because their general formula is C_nH_{2n}, they have two fewer H atoms than an acyclic alkane with the same number of carbons.

Undecane, an acyclic alkane, and cyclohexane, a cycloalkane, are two naturally occurring alkanes.




undecane
$C_{11}H_{24}$

cyclohexane
C_6H_{12}


4.1A Acyclic Alkanes Having One to Five C Atoms

Structures for the two simplest acyclic alkanes were given in Chapter 1. **Methane, CH_4,** has a single carbon atom, and **ethane, CH_3CH_3,** has two. All C atoms in an alkane are surrounded by four groups, making them ***sp*^{3} hybridized** and **tetrahedral,** and all bond angles are 109.5°.



To draw the structure of an alkane, join the carbon atoms together with single bonds, and add enough H atoms to make each C tetravalent.

The three-carbon alkane **$CH_3CH_2CH_3$, propane,** has molecular formula C_3H_8. Each carbon in the three-dimensional drawing has two bonds in the plane (solid lines), one bond in front (on a wedge), and one bond behind the plane (on a dashed wedge).



Problem 4.1 Three components of the sex pheromone of the female sand bee (*Ophrys sphegodes*) are saturated hydrocarbons containing 23, 25, and 27 carbon atoms. How many H atoms does each of these alkanes contain? Interestingly, the early spider orchid emits a similar hydrocarbon mixture to attract male sand bees to pollinate its flowers.

The molecular formulas for methane, ethane, and propane fit into the general molecular formula for an alkane, $\mathbf{C_nH_{2n+2}}$.

- Methane = CH_4 = $C_1H_{2(1)+2}$
- Ethane = C_2H_6 = $C_2H_{2(2)+2}$
- Propane = C_3H_8 = $C_3H_{2(3)+2}$

There are two different ways to arrange four carbons, giving two compounds with molecular formula C_4H_{10}, named **butane** and **2-methylpropane** (or isobutane).



butane
4 C's in a row
straight-chain alkane


2-methylpropane
(or isobutane)
3 C's in a row with a one-carbon branch
branched-chain alkane

Butane and 2-methylpropane are ***isomers,* two different compounds with the same molecular formula** (Section 1.4). They belong to one of the two major classes of isomers called **constitutional** or **structural isomers.** We will learn about the second major class of isomers, called **stereoisomers,** in Section 4.13B.

- *Constitutional isomers* differ in the way the atoms are connected to each other.

Butane, which has four carbons in a row, is a **straight-chain** or **normal alkane** (an ***n*-alkane**). 2-Methylpropane, on the other hand, is a **branched-chain alkane.**

With alkanes having more than four carbons, the names of the straight-chain isomers are systematic and derive from Greek roots: ***pent***ane for five C atoms, ***hex***ane for six, and so on. There are three constitutional isomers for the five-carbon alkane, each having molecular formula C_5H_{12}: **pentane, 2-methylbutane** (or isopentane), and **2,2-dimethylpropane** (or neopentane).




pentane


2-methylbutane
(isopentane)

2,2-dimethylpropane
(neopentane)

Take care in interpreting skeletal structures. Although pentane is typically drawn using a zigzag structure, the carbon skeleton can be drawn in a variety of ways, and still represent the same compound. Each of the following representations has five carbon atoms in a row, so each represents pentane, not an isomer of pentane.



Problem 4.2 Which of the following is *not* another representation for 2-methylbutane?



4.1B Acyclic Alkanes Having More Than Five C Atoms

The maximum number of possible constitutional isomers increases dramatically as the number of carbon atoms in the alkane increases, as shown in Table 4.1. For example, there are 75 possible isomers for an alkane having 10 carbon atoms, and 366,319 possible isomers for one having 20 carbons.

Each entry in Table 4.1 is formed from the preceding entry by adding a CH_2 group. **A CH_2 group is called a *methylene group*. A group of compounds that differ by only a CH_2 group is called a *homologous series*.** The names of all alkanes end in the suffix ***-ane,*** and the syllables preceding the suffix identify the number of carbon atoms in the chain.

Table 4.1 Summary: Straight-Chain Alkanes

Number of C atoms	Molecular formula	Name (*n*-alkane)	Number of constitutional isomers	Number of C atoms	Molecular formula	Name (*n*-alkane)	Number of constitutional isomers
1	CH_4	methane	—	9	C_9H_{20}	nonane	35
2	C_2H_6	ethane	—	10	$C_{10}H_{22}$	decane	75
3	C_3H_8	propane	—	11	$C_{11}H_{24}$	undecane	159
4	C_4H_{10}	butane	2	12	$C_{12}H_{26}$	dodecane	355
5	C_5H_{12}	pentane	3	13	$C_{13}H_{28}$	tridecane	802
6	C_6H_{14}	hexane	5	14	$C_{14}H_{30}$	tetradecane	1858
7	C_7H_{16}	heptane	9	15	$C_{15}H_{32}$	pentadecane	4347
8	C_8H_{18}	octane	18	20	$C_{20}H_{42}$	icosane	366,319

Problem 4.3 Draw the five constitutional isomers having molecular formula C_6H_{14}.

Problem 4.4 Review classifying carbons and hydrogens in Section 3.2, and draw the structure of an alkane with molecular formula C_7H_{16} that contains (a) one 4° carbon; (b) only 1° and 2° carbons; (c) 1°, 2°, and 3° hydrogens.


Problem 4.5 (a) Which compounds (**B–F**) are identical to **A?** (b) Which compounds (**B–F**) represent an isomer of **A?**



4.2 Cycloalkanes

Cycloalkanes have molecular formula C_nH_{2n} and contain carbon atoms arranged in a ring. Think of a cycloalkane as being formed by removing two H atoms from the end carbons of a chain, and then bonding the two carbons together. Simple cycloalkanes are named by adding the prefix **cyclo-** to the name of the acyclic alkane having the same number of carbons.

Cycloalkanes with three to six carbon atoms are shown.




Problem 4.6 Draw the five constitutional isomers that have molecular formula C_5H_{10} and contain one ring.



Garlic has been a valued commodity throughout history. It has been used in Chinese herbal medicine for more than 4000 years, as a form of currency in Siberia, and as a repellent for witches by the Saxons. Today it is used as a dietary supplement because of its reported health benefits. **Allicin,** the molecule largely responsible for garlic's odor, is not stored in the garlic bulb, but instead is produced by the action of enzymes when the bulb is crushed or bruised.

4.3 An Introduction to Nomenclature

How are organic compounds named? Long ago, the name of a compound was often based on the plant or animal source from which it was obtained. For example, the name for **formic acid,** a caustic compound isolated from certain ants, comes from the Latin word *formica,* meaning *ant;* and **allicin,** the pungent principle of garlic, is derived from the botanical name for garlic, *Allium sativum.* Other compounds were named by their discoverer for more personal reasons. Adolf von Baeyer supposedly named barbituric acid after a woman named Barbara, although speculation continues on Barbara's identity—a lover, a Munich waitress, or even St. Barbara.




With the isolation and preparation of thousands of new organic compounds it became clear that each organic compound must have an unambiguous name, derived from a set of easily remembered rules. A systematic method of naming compounds was developed by the ***I****nternational* ***U****nion of* ***P****ure and* ***A****pplied* ***C****hemistry.* It is referred to as the **IUPAC system of nomenclature;** how it can be used to name alkanes is explained in Sections 4.4 and 4.5.

The IUPAC system of nomenclature has been regularly revised since it was first adopted in 1892. Revisions in 1979 and 1993 and recent extensive recommendations in 2004 have given chemists a variety of acceptable names for compounds. Many changes are minor. For example, the 1979 nomenclature rules assign the name 1-butene to $CH_2{=}CHCH_2CH_3$, while the 1993 rules assign the name but-1-ene; that is, only the position of the number differs. In this text, more recent IUPAC conventions will be used, and often a margin note will be added to mention the differences between past and recent recommendations.

Naming organic compounds has become big business for drug companies. The IUPAC name of an organic compound can be long and complex, and may be comprehensible only to a chemist. As a result, most drugs have three names:

- **Systematic:** The systematic name follows the accepted rules of nomenclature and indicates the compound's chemical structure; this is the IUPAC name.
- **Generic:** The generic name is the official, internationally approved name for the drug.
- **Trade:** The trade name for a drug is assigned by the company that manufactures it. Trade names are often "catchy" and easy to remember. Companies hope that the public will continue to purchase a drug with an easily recalled trade name long after a cheaper generic version becomes available.

In the world of over-the-counter anti-inflammatory agents, the compound a chemist calls 2-[4-(2-methylpropyl)phenyl]propanoic acid has the generic name ibuprofen. It is marketed under a variety of trade names including Motrin and Advil.



Systematic name: 2-[4-(2-methylpropyl)phenyl]propanoic acid
Generic name: ibuprofen
Trade name: Motrin or Advil

4.4 Naming Alkanes

The name of every organic molecule has three parts.

- The **parent name** indicates the number of carbons in the longest continuous carbon chain in the molecule.
- The **suffix** indicates what functional group is present.
- The **prefix** reveals the identity, location, and number of substituents attached to the carbon chain.

prefix + **parent** + **suffix**

What and where are the substituents? | What is the longest carbon chain? | What is the functional group?

The names listed in Table 4.1 of Section 4.1B for the simple *n*-alkanes consist of the parent name, which indicates the number of carbon atoms in the longest carbon chain, and the suffix ***-ane,*** which indicates that the compounds are alkanes. The parent name for **one carbon is *meth-*,** for **two carbons is *eth-*,** and so on. Thus, we are already familiar with two parts of the name of an organic compound.

To determine the third part of a name, the prefix, we must learn how to name the carbon groups or *substituents* that are bonded to the longest carbon chain.


4.4A Naming Substituents

Carbon substituents bonded to a long carbon chain are called **alkyl groups.**

- An *alkyl group* is formed by removing one hydrogen from an alkane.

An alkyl group is a part of a molecule that is now able to bond to another atom or a functional group. **To name an alkyl group, change the *-ane* ending of the parent alkane to *-yl.*** Thus, methane (CH_4) becomes methyl (CH_3-) and ethane (CH_3CH_3) becomes ethyl (CH_3CH_2-). As we learned in Section 3.1, **R** denotes a general carbon group bonded to a functional group. **R** thus denotes any alkyl group.


Naming three- and four-carbon alkyl groups is more complicated because the parent hydrocarbons have more than one type of hydrogen atom. Propane has both 1° and 2° H atoms, and removal of each of these H atoms forms a different alkyl group, **propyl** or **isopropyl.**




The prefix **iso-** is part of the words *propyl* and *butyl,* forming a single word: **isopropyl** and **isobutyl.** The prefixes ***sec-*** and ***tert-*** are separated from the word *butyl* by a hyphen: ***sec*-butyl** and ***tert*-butyl.**

Because there are two different butane isomers to begin with, each with two different kinds of H atoms, there are four possible alkyl groups containing four carbon atoms: **butyl, *sec*-butyl, isobutyl,** and ***tert*-butyl.**

The prefix *sec-* is short for *secondary.* A *sec*-butyl group is formed by removal of a **2° H.** The prefix *tert-* is short for *tertiary.* A *tert*-butyl group is formed by removal of a **3° H.**





Abbreviations are sometimes used for certain common alkyl groups.

- methyl (Me)
- ethyl (Et)
- butyl (Bu)
- *tert*-butyl (*t*-Bu)

The names isopropyl, *sec*-butyl, isobutyl, and *tert*-butyl are recognized as acceptable substituent names in both the 1979 and 1993 revisions of IUPAC nomenclature. A general method to name these substituents, as well as alkyl groups that contain five or more carbon atoms, is described in Appendix B.

4.4B Naming an Acyclic Alkane


Four steps are needed to name an alkane.

How To Name an Alkane Using the IUPAC System

Step [1] Find the parent carbon chain and add the suffix.


- Find the longest continuous carbon chain, and name the molecule by using the parent name for that number of carbons, given in Table 4.1. To the name of the parent, add the suffix ***-ane*** for an alkane. Each functional group has its own characteristic suffix.

Correct




8 C's in the longest chain
8 C's ⇢ **octane**

Incorrect




7 C's in the longest chain

- Finding the longest chain is a matter of trial and error. Place your pencil on one end of the chain, go to the other end without picking it up, and count carbons. Repeat this procedure until you have found the chain with the largest number of carbons.
- It does not matter if the chain is *straight* or has *bends.* All of the following representations are equivalent, and each longest chain has eight carbons.



- If there are two chains of equal length, pick the chain with more substituents. In the following example, two different chains in the same alkane contain 7 C's, but the compound on the left has two alkyl groups attached to its long chain, whereas the compound to the right has only one.


Correct



7 atoms in the longest chain
2 substituents

more substituents

Incorrect




7 atoms in the longest chain
only 1 substituent

fewer substituents

Step [2] Number the atoms in the carbon chain.


- Number the longest chain to give the *first* substituent the lower number.

Correct



first substituent at C2

Incorrect




first substituent at C3

—*Continued*

How To, continued . . .


- If the first substituent is the same distance from both ends, number the chain to give the *second* substituent the lower number. Always look for the first point of difference in numbering from each end of the longest chain.

Correct



CH_3 groups at C2, C3, and C5
The second CH_3 group has the lower number (C3).


Incorrect



CH_3 groups at C2, C4, and C5
The second CH_3 group has the higher number (C4).


- When numbering a carbon chain results in the same numbers from either end of the chain, assign the lower number *alphabetically* to the first substituent.

Correct



- **e**thyl at **C3**
- **m**ethyl at **C5**


Incorrect



- **m**ethyl at **C3**
- **e**thyl at **C5**

Earlier letter ⟶ lower number

Step [3] Name and number the substituents.



8 C's in the longest chain


- Name the substituents as alkyl groups, and use the numbers from Step 2 to designate their location.
- Every carbon belongs to *either* the longest chain or a substituent, but *not both.*
- Each substituent needs its *own* number.
- If two or more identical substituents are bonded to the longest chain, use prefixes to indicate how many: **di-** for two groups, **tri-** for three groups, **tetra-** for four groups, and so forth. This molecule has two methyl substituents, so its name contains the prefix di- before the word methyl → *di*methyl.

Step [4] Combine substituent names and numbers + parent + suffix.

- Precede the name of the parent by the names of the substituents.
- Alphabetize the names of the substituents, ignoring all prefixes except *iso,* as in isopropyl and isobutyl.
- Precede the name of each substituent by the number that indicates its location. There must be **one number for each substituent.**
- Separate numbers by commas and separate numbers from letters by hyphens. The name of an alkane is a single word, with no spaces after hyphens or commas.

[1] Identify all the pieces of a compound, using Steps 1–3.

2,6-dimethyl



8 C's in the longest chain

octane

5-ethyl


[2] Then, put the pieces of the name together.

substituent names and numbers	+	parent	+	suffix
5-ethyl-2,6-dimethyl	+	**oct**	+	**ane**
Alphabetize: **e** for **ethyl,** then **m** for **methyl**		**8** C's		an **alkane**

Answer: 5-ethyl-2,6-dimethyloctane


Several additional examples of alkane nomenclature are given in Figure 4.1.

Figure 4.1
Examples of alkane nomenclature




2,3-dimethylpentane

Number to give the 1st methyl group the lower number.




4-ethyl-5-methyloctane

Assign the lower number to the 1st substituent alphabetically: the **e** of **e**thyl before the **m** of **m**ethyl.



4-ethyl-3,4-dimethyloctane

Alphabetize the **e** of **e**thyl before the **m** of **m**ethyl.




2,3,5-trimethyl-4-propylheptane

Pick the long chain with more substituents.

- The carbon atoms of each long chain are drawn in **red.**


Sample Problem 4.1 Give the IUPAC name for the following compound.



Solution


To help identify which carbons belong to the longest chain and which are substituents, box in or highlight the atoms of the long chain. Every other carbon atom then becomes a substituent that needs its own name as an alkyl group.

Step 1: Name the parent.




9 C's in the longest chain
nonane

Step 3: Name and number the substituents.



Step 2: Number the chain.




first substituent at C3


Step 4: Combine the parts.


- Alphabetize: the **b** of **butyl** before the **m** of **methyl**


Answer: 5-*tert*-butyl-3-methylnonane

Problem 4.7 Give the IUPAC name for each compound.

a. 

b. 


c. 


d. 

Problem 4.8 Give the IUPAC name for each compound.

a. $(CH_3)_3CCH_2CH(CH_2CH_3)_2$

c. $CH_3(CH_2)_3CH(CH_2CH_2CH_3)CH(CH_3)_2$

b. 

d. 

You must also know how to derive a structure from a given name. Sample Problem 4.2 illustrates a stepwise method.

Sample Problem 4.2 Give the structure corresponding to the following IUPAC name: 6-isopropyl-3,3,7-trimethyldecane.


Solution

Follow three steps to derive a structure from a name.

Step [1] Identify the parent name and functional group found at the *end* of the name.

decane ----→ **10** C's ----→ 

Step [2] Number the carbon skeleton in *either* direction.



Step [3] Add the substituents at the appropriate carbons.



Problem 4.9 Give the structure corresponding to each IUPAC name.

a. 3-methylhexane

b. 3,3-dimethylpentane

c. 3,5,5-trimethyloctane


d. 3-ethyl-4-methylhexane

e. 3-ethyl-5-isobutylnonane

Problem 4.10 Give the IUPAC name for each of the five constitutional isomers of molecular formula C_6H_{14} in Problem 4.3.

4.5 Naming Cycloalkanes


Cycloalkanes are named by using similar rules, but the prefix **cyclo-** immediately precedes the name of the parent.



How To Name a Cycloalkane Using the IUPAC System

Step [1] Find the parent cycloalkane.


- Count the number of carbon atoms in the ring and use the parent name for that number of carbons. Add the prefix **cyclo-** and the suffix ***-ane*** to the parent name.




6 C's in the ring
cyclohexane

Step [2] Name and number the substituents.

- No number is needed to indicate the location of a single substituent.




methylcyclohexane




tert-butylcyclopentane

- For rings with more than one substituent, begin numbering at one substituent and proceed around the ring clockwise or counterclockwise to give the second substituent the lower number.



CH_3 groups at C1 and C3
The 2nd substituent has a lower number.


Correct: 1,3-dimethylcyclohexane



CH_3 groups at C1 and C5


Incorrect: 1,5-dimethylcyclohexane

- With two different substituents, number the ring to assign the lower number to the substituents *alphabetically.*



earlier letter ⟶ lower number

Correct: 1-ethyl-3-methylcyclohexane




Incorrect: 3-ethyl-1-methylcyclohexane

When an alkane is composed of both a ring and a long chain, what determines whether a compound is named as an acyclic alkane or a cycloalkane? If the number of carbons in the ring is greater than or equal to the number of carbons in the longest chain, the compound is named as

a **cycloalkane,** as shown in Figure 4.2. Several examples of cycloalkane nomenclature are given in Figure 4.3.

Figure 4.2
Naming compounds containing both a ring and a chain of carbon atoms




6 C's in the ring
cyclohexane

Name as a **cyclohexane** with a substituent.

butylcyclohexane

- Name the molecule as a substituted cycloalkane when it has more C's in the ring than any single alkyl substituent.




4 C's in the ring
cyclobutyl group

Name as a **hexane** with a substituent.

1-cyclobutylhexane


- Name the molecule as a substituted alkane when it has a carbon chain with more C's than the ring.

Figure 4.3
Examples of cycloalkane nomenclature




ethylcyclobutane

No number is needed with only one substituent.




1-*sec*-butyl-3-methylcyclohexane

Assign the lower number to the 1st substituent alphabetically: the **b** of **b**utyl before the **m** of **m**ethyl.



1,2-dimethylcyclohexane


Number to give the 2nd CH_3 group the lower number: 1,2- not 1,6-.





1,2,4-triethylcyclopentane


Number to give the 2nd CH_3CH_2 group the lower number: 1,2,4- not 1,3,4- or 1,3,5-.


Problem 4.11 Give the IUPAC name for each compound.


a. 

c. 

e. 

b. 

d. 


f. 

Problem 4.12 Give the structure corresponding to each IUPAC name.

a. 1,2-dimethylcyclobutane
b. 1,1,2-trimethylcyclopropane
c. 4-ethyl-1,2-dimethylcyclohexane
d. 1-*sec*-butyl-3-isopropylcyclopentane
e. 1,1,2,3,4-pentamethylcycloheptane

4.6 Common Names

Some organic compounds are identified using **common names** that do not follow the IUPAC system of nomenclature. Many of these names were given to molecules long ago, before the IUPAC system was adopted. These names are still widely used. For example, isopentane, an older name for 2-methylbutane, is still allowed by IUPAC rules. We will follow the IUPAC system except in cases in which a common name is widely accepted.



isopentane or 2-methylbutane




dodecahedrane

In the past several years organic chemists have attempted to synthesize some unusual cycloalkanes not found in nature. **Dodecahedrane,** a beautifully symmetrical compound composed of 12 five-membered rings, is one such molecule. It was first prepared at The Ohio State University in 1982. The IUPAC name for dodecahedrane is undecacyclo[9.9.0.$0^{2,9}$.$0^{3,7}$.$0^{4,20}$.$0^{5,18}$.$0^{6,16}$.$0^{8,15}$.$0^{10,14}$.$0^{12,19}$.$0^{13,17}$]-icosane, a name so complex that few trained organic chemists would be able to identify its structure.

Because these systematic names are so unwieldy, organic chemists often assign a name to a polycyclic compound that is more descriptive of its shape and structure. Dodecahedrane is named because its 12 five-membered rings resemble a dodecahedron. Figure 4.4 shows the names and structures of several other cycloalkanes whose names were inspired by the shape of their carbon skeletons. All the names end in the suffix ***-ane,*** indicating that they refer to alkanes.

Figure 4.4
Common names for some polycyclic alkanes



For a more comprehensive list of unusual polycyclic alkanes (including windowpane, davidane, catenane, propellane, and many others), see *Organic Chemistry: The Name Game* by Alex Nickon and Ernest Silversmith, Pergamon Press, 1987.

4.7 Fossil Fuels

Natural gas is odorless. The smell observed in a gas leak is due to minute amounts of a sulfur additive such as methanethiol, CH_3SH, which provides an odor for easy detection.

Many alkanes occur in nature, primarily in natural gas and petroleum. Both of these fossil fuels serve as energy sources, formed from the degradation of organic material long ago.


Natural gas is composed largely of **methane** (60% to 80% depending on its source), with lesser amounts of ethane, propane, and butane. These organic compounds burn in the presence of oxygen, releasing energy for cooking and heating.

Petroleum is a complex mixture of compounds, most of which are hydrocarbons containing 1–40 carbon atoms. Distilling crude petroleum, a process called **refining,** separates it into usable

Figure 4.5
Refining crude petroleum into usable fuel and other petroleum products



a. **An oil refinery.** At an oil refinery, crude petroleum is separated into fractions of similar boiling point by the process of **distillation.**




b. **Schematic of a refinery tower.** As crude petroleum is heated, the lower boiling, more volatile components distill first, followed by fractions of progressively higher boiling point.

Methane is formed and used in a variety of ways. The CH_4 released from decaying vegetable matter in New York City's main landfill is used for heating homes. CH_4 generators in China convert cow manure into energy in rural farming towns.

fractions that differ in boiling point (Figure 4.5). Most products of petroleum refining provide fuel for home heating, automobiles, diesel engines, and airplanes. Each fuel type has a different composition of hydrocarbons: gasoline ($C_5H_{12}-C_{12}H_{26}$), kerosene ($C_{12}H_{26}-C_{16}H_{34}$), and diesel fuel ($C_{15}H_{32}-C_{18}H_{38}$).

Petroleum provides more than fuel. About 3% of crude oil is used to make plastics and other synthetic compounds including drugs, fabrics, dyes, and pesticides. These products are responsible for many of the comforts we now take for granted in industrialized countries. Imagine what life would be like without air conditioning, refrigeration, anesthetics, and pain relievers, all products of the petroleum industry.









products made from petroleum

Energy from petroleum is ***nonrenewable,*** and the remaining known oil reserves are limited. Given our dependence on petroleum, not only for fuel, but also for the many necessities of modern society, it becomes clear that we must both conserve what we have and find alternate energy sources.

4.8 Physical Properties of Alkanes

Alkanes contain only nonpolar C−C and C−H bonds, and as a result they exhibit only **weak van der Waals forces.** Table 4.2 summarizes how these intermolecular forces affect the physical properties of alkanes.

Table 4.2 Physical Properties of Alkanes

Property	Observation
Boiling point and melting point	• Alkanes have low bp's and mp's compared to more polar compounds of comparable size. 
	• Bp and mp increase as the number of carbons increases because of increased surface area. 
	• The bp of isomers decreases with branching because of decreased surface area. 
	• Mp increases with increased symmetry. 
Solubility	• Alkanes are soluble in organic solvents. • Alkanes are insoluble in water.


Key: bp = boiling point; mp = melting point; VDW = van der Waals; DD = dipole–dipole; HB = hydrogen bonding

The gasoline industry exploits the dependence of boiling point and melting point on alkane size by seasonally changing the composition of gasoline in locations where it gets very hot in the summer and very cold in the winter. Gasoline is refined to contain a larger fraction of higher boiling hydrocarbons in warmer weather, so it evaporates less readily. In colder weather, it is refined to contain more lower boiling hydrocarbons, so it freezes less readily.

The mutual insolubility of nonpolar oil and very polar water leads to the common expression, "Oil and water don't mix."

Because nonpolar alkanes are not water soluble, crude petroleum that leaks into the sea from an oil tanker or offshore oil well creates an insoluble oil slick on the surface. The insoluble hydrocarbon oil poses a special threat to birds whose feathers are coated with natural nonpolar oils for insulation. Because these hydrophobic oils dissolve in the crude petroleum, birds lose their layer of natural protection and many die.


Problem 4.13 Arrange the following compounds in order of increasing boiling point.



4.9 Conformations of Acyclic Alkanes—Ethane

Let's now take a closer look at the three-dimensional structure of alkanes. The three-dimensional structure of molecules is called **stereochemistry.** In Chapter 4 we examine the effect of rotation around single bonds. In Chapter 5, we will learn about other aspects of stereochemistry.

Recall from Section 1.10A that **rotation occurs around carbon–carbon σ bonds.** Thus, the two CH_3 groups of ethane rotate, allowing the hydrogens on one carbon to adopt different orientations relative to the hydrogens on the other carbon. These arrangements are called **conformations.**




Rotation occurs here.

two different conformations

- *Conformations* are different arrangements of atoms that are interconverted by rotation about single bonds.

Two different arrangements are the **eclipsed conformation** and the **staggered conformation.**

- In the *eclipsed conformation,* the C – H bonds on one carbon are directly aligned with the C – H bonds on the adjacent carbon.
- In the *staggered conformation,* the C – H bonds on one carbon bisect the H – C – H bond angle on the adjacent carbon.




eclipsed conformation

The C–H bonds are all **aligned.**

staggered conformation

The C–H bonds in front **bisect** the H–C–H bond angles in back.


Rotating the atoms on one carbon by 60° converts an eclipsed conformation into a staggered conformation, and vice versa. These conformations are often viewed end-on—that is, looking directly down the carbon–carbon bond. The angle that separates a bond on one atom from a bond on an adjacent atom is called a **dihedral angle.** For ethane in the staggered conformation, the dihedral angle for the C−H bonds is **60°.** For eclipsed ethane, it is **0°.**



End-on representations for conformations are commonly drawn using a convention called a **Newman projection.** A Newman projection is a graphic that shows the three groups bonded to each carbon atom in a particular C−C bond, as well as the dihedral angle that separates them.


How To Draw a Newman Projection

Step [1] **Look directly down the C – C bond (end-on), and draw a circle with a dot in the center to represent the carbons of the C – C bond.**




- The circle represents the back carbon and the dot represents the front carbon.

Step [2] **Draw in the bonds.**



- Draw the bonds on the **front** C as three lines **meeting at the center** of the circle.
- Draw the bonds on the **back** C as three lines coming **out of the edge** of the circle.


Step [3] **Add the atoms on each bond.**



- Each C has 3 H's in ethane.

Figure 4.6 illustrates the Newman projections for both the staggered and eclipsed **conformations** for ethane.

Figure 4.6
Newman projections for the staggered and eclipsed conformations of ethane




staggered conformation **eclipsed conformation**

Follow this procedure for any C−C bond. With a Newman projection, **always consider *one* C−C bond only and draw the atoms bonded to the carbon atoms, *not* the carbon atoms in the bond itself.** Newman projections for the staggered and eclipsed conformations of propane are drawn in Figure 4.7.


Figure 4.7
Newman projections for the staggered and eclipsed conformations of propane

Consider one C–C bond only.



propane


- 3 H's on one C
- 2 H's and 1 CH_3 on the other C




staggered conformation **eclipsed conformation**

Problem 4.14 Draw the staggered and eclipsed conformations that result from rotation around the C−C bond in CH_3-CH_2Br.

Problem 4.15 Which of the following is (are) possible Newman projections for 2-methylpentane?




2-methylpentane



The staggered and eclipsed conformations of ethane interconvert at room temperature, but **each conformation is *not* equally stable.**

- **The staggered conformations are more stable (lower in energy) than the eclipsed conformations.**

The cause of this stability difference is the subject of some debate in the chemical literature. A contributing factor may be increased electron–electron repulsion between the bonds in the eclipsed conformation compared to the staggered conformation, where the bonding electrons are farther apart.



Each H,H eclipsing increases energy by 4.0 kJ/mol.

The difference in energy between the staggered and eclipsed conformations is 12 kJ/mol (2.9 kcal/mol), a small enough difference that the rotation is still very rapid at room temperature, and the conformations cannot be separated. Because three eclipsed C−H bonds increase the energy of a conformation by 12 kJ/mol, **each eclipsed C−H bond results in an increase in energy of 4.0 kJ/mol (1.0 kcal/mol).** The energy difference between the staggered and eclipsed conformations is called **torsional energy.** Thus, eclipsing introduces **torsional strain** into a molecule.

- *Torsional strain* is an increase in energy caused by eclipsing interactions.


The graph in Figure 4.8 shows how the potential energy of ethane changes with dihedral angle as one CH_3 group rotates relative to the other. **The staggered conformation is the most stable arrangement, so it is at an *energy minimum*.** As the C−H bonds on one carbon are rotated relative to the C−H bonds on the other carbon, the energy increases as the C−H bonds get closer until a **maximum is reached after 60° rotation to the eclipsed conformation.** As rotation continues, the energy decreases until after 60° rotation, when the staggered conformation is reached once again. At any given moment, all ethane molecules do not exist in the more stable staggered conformation; rather, a higher percentage of molecules is present in the more stable staggered conformation than any other possible arrangement.

Strain results in an **increase in energy.** Torsional strain is the first of three types of strain discussed in this text. The other two are steric strain (Section 4.10) and angle strain (Section 4.11).

- An energy minimum and maximum occur every 60° as the conformation changes from staggered to eclipsed. Conformations that are neither staggered nor eclipsed are intermediate in energy.

Problem 4.16 The torsional energy in propane is 14 kJ/mol (3.4 kcal/mol). Because each H,H eclipsing interaction is worth 4.0 kJ/mol (1.0 kcal/mol) of destabilization, how much is one H,CH_3 eclipsing interaction worth in destabilization? (See Section 4.10 for an alternate way to arrive at this value.)


Figure 4.8
Graph: Energy versus dihedral angle for ethane



- Note the position of the labeled H atom after each 60° rotation. All three staggered conformations are identical (except for the position of the label), and the same is true for all three eclipsed conformations.

4.10 Conformations of Butane

Butane and higher molecular weight alkanes have several carbon–carbon bonds, all capable of rotation.



butane
Consider rotation at C2–C3.
Each C is bonded to 2 H's and 1 CH_3 group.


To analyze the different conformations that result from rotation about the C2−C3 bond, begin arbitrarily with one—for example, the staggered conformation that places two CH_3 groups 180° from each other—then,

It takes six 60° rotations to return to the original conformation.

- **Rotate one carbon atom in 60° increments either clockwise or counterclockwise, while keeping the other carbon fixed. Continue until you return to the original conformation.**

Figure 4.9 illustrates the six possible conformations that result from this process.

Figure 4.9
Six different conformations of butane



Although each 60° bond rotation converts a staggered conformation into an eclipsed conformation (or vice versa), neither all the staggered conformations nor all the eclipsed conformations are the same. For example, the dihedral angle between the methyl groups in staggered conformations **3** and **5** are both 60°, whereas it is 180° in staggered conformation **1.**


- **A staggered conformation with two larger groups 180° from each other is called *anti.***
- **A staggered conformation with two larger groups 60° from each other is called *gauche.***

Similarly, the methyl groups in conformations **2** and **6** both eclipse hydrogen atoms, whereas they eclipse each other in conformation **4.**

The staggered conformations (**1, 3,** and **5**) are lower in energy than the eclipsed conformations (**2, 4,** and **6**), but how do the energies of the individual staggered and eclipsed conformations compare to each other? The relative energies of the individual staggered conformations (or the individual eclipsed conformations) depend on their **steric strain.**


- ***Steric*** **strain is an increase in energy resulting when atoms are forced too close to one another.**

The methyl groups are farther apart in the anti conformation (**1**) than in the gauche conformations (**3** and **5**), so amongst the staggered conformations, **1** is lower in energy (more stable) than **3** and **5.** In fact, the anti conformation is 3.8 kJ/mol (0.9 kcal/mol) lower in energy than either gauche conformation because of the steric strain that results from the proximity of the methyl groups in **3** and **5.**



- **Gauche conformations are generally higher in energy than anti conformations because of steric strain.**

Steric strain also affects the relative energies of eclipsed conformations. Conformation **4** is higher in energy than **2** or **6,** because the two larger CH_3 groups are forced close to each other, introducing considerable steric strain.




To graph energy versus dihedral angle, keep in mind two considerations:


- **Staggered conformations are at energy minima and eclipsed conformations are at energy maxima.**
- **Unfavorable steric interactions increase energy.**

For butane, this means that anti conformation **1** is lowest in energy, and conformation **4** with two eclipsed CH_3 groups is the highest in energy. The relative energy of other conformations is depicted in the energy versus rotation diagram for butane in Figure 4.10.


Figure 4.10
Graph: Energy versus dihedral angle for butane



- Staggered conformations **1, 3,** and **5** are at energy minima.
- Anti conformation **1** is lower in energy than gauche conformations **3** and **5,** which possess steric strain.
- Eclipsed conformations **2, 4,** and **6** are at energy maxima.
- Eclipsed conformation **4,** which has additional steric strain due to two eclipsed CH_3 groups, is highest in energy.



CH_3,CH_3 eclipsing
11 kJ/mol destabilization



H,CH_3 eclipsing
6.0 kJ/mol destabilization

We can now use the values in Figure 4.10 to estimate the destabilization caused by other eclipsed groups. For example, conformation **4** is 19 kJ/mol less stable than the anti conformation **1.** Conformation **4** possesses two H,H eclipsing interactions, worth 4.0 kJ/mol each in destabilization (Section 4.9), and one CH_3,CH_3 eclipsing interaction. Thus, the **CH_3,CH_3 interaction** is worth 19 – 2(4.0) = **11 kJ/mol** of destabilization.


Similarly, conformation **2** is 16 kJ/mol less stable than the anti conformation **1,** and possesses one H,H eclipsing interaction (worth 4.0 kJ/mol of destabilization), and two H,CH_3 interactions. Thus **each H,CH_3 interaction** is worth 1/2(16 – 4.0) = **6.0 kJ/mol** of destabilization. These values are summarized in Table 4.3.

- **The energy difference between the lowest and highest energy conformations is called the *barrier to rotation.***

Table 4.3 Summary: Torsional and Steric Strain Energies in Acyclic Alkanes

Type of interaction	Energy increase	
	kJ/mol	kcal/mol
H,H eclipsing	4.0	1.0
H,CH_3 eclipsing	6.0	1.4
CH_3,CH_3 eclipsing	11	2.6
gauche CH_3 groups	3.8	0.9

We can use these same principles to determine conformations and relative energies for any acyclic alkane. Because the lowest energy conformation has all bonds staggered and all large groups anti, alkanes are often drawn in zigzag skeletal structures to indicate this.




Problem 4.17




a. Draw the three staggered and three eclipsed conformations that result from rotation around the bond labeled in red using Newman projections.

b. Label the most stable and least stable conformation.


Problem 4.18 Rank the following conformations in order of increasing energy.




A



B



C



D


Problem 4.19 Consider rotation around the carbon–carbon bond in 1,2-dichloroethane ($ClCH_2CH_2Cl$).

a. Using Newman projections, draw all of the staggered and eclipsed conformations that result from rotation around this bond.


b. Graph energy versus dihedral angle for rotation around this bond.

Problem 4.20 Calculate the destabilization present in each eclipsed conformation.

a.



b.




4.11 An Introduction to Cycloalkanes

Besides torsional strain and steric strain, the conformations of cycloalkanes are also affected by **angle strain.**

- ***Angle*** **strain is an increase in energy when tetrahedral bond angles deviate from the optimum angle of 109.5°.**


Originally cycloalkanes were thought to be flat rings, with the bond angles between carbon atoms determined by the size of the ring. For example, a flat cyclopropane ring would have 60° internal bond angles, a flat cyclobutane ring would have 90° angles, and large flat rings would have very large angles. It was assumed that rings with bond angles so different from the tetrahedral bond angle would be very strained and highly reactive. This is called the Baeyer strain theory.





small internal angles
angle strain


large internal angles
angle strain

Figure 4.11
Three-dimensional structure of some cycloalkanes




tetrahedrane


cubane


dodecahedrane

It turns out, though, that **cycloalkanes with more than three C atoms in the ring are not flat molecules.** They are puckered to **reduce strain,** both angle strain and torsional strain. The three-dimensional structures of some simple cycloalkanes are shown in Figure 4.11. Three- and four-membered rings still possess considerable angle strain, but puckering reduces the internal bond angles in larger rings, thus reducing angle strain.

Many polycyclic hydrocarbons are of interest to chemists. For example, **dodecahedrane,** containing 12 five-membered rings bonded together, is one member of a family of three hydrocarbons that contain several rings of one size joined together. The two other members of this family are **tetrahedrane,** consisting of four three-membered rings, and **cubane,** consisting of six four-membered rings. These compounds are the simplest regular polyhedra whose structures resemble three of the highly symmetrical Platonic solids: the tetrahedron, the cube, and the dodecahedron.

How stable are these compounds? Tetrahedrane (with internal 60° bond angles) is so strained that all attempts to prepare it have been thus far unsuccessful. Although cubane is also highly strained because of its 90° bond angles, it was first synthesized in 1964 and is a stable molecule at room temperature. Finally, dodecahedrane is very stable because it has bond angles very close to the tetrahedral bond angle (108° versus 109.5°). Its synthesis eluded chemists for years not because of its strain or inherent instability, but because of the enormous challenge of joining 12 five-membered rings together to form a sphere.


4.12 Cyclohexane

Let's now examine the conformation of **cyclohexane,** the most common ring size in naturally occurring compounds.


4.12A The Chair Conformation

A planar cyclohexane ring would experience angle strain, because the internal bond angle between the carbon atoms would be 120°, and torsional strain, because all of the hydrogens on adjacent carbon atoms would be eclipsed.

If a cyclohexane ring were flat…



The internal bond angle is >109.5°.
angle strain




All H's are aligned.
torsional strain




Visualizing the chair. If the cyclohexane chair conformation is tipped downward, we can more easily view it as a chair with a back, seat, and foot support.

In reality, cyclohexane adopts a puckered conformation, called the **chair** form, which is more stable than any other possible conformation.



The chair conformation is so stable because it eliminates angle strain **(all C−C−C bond angles are 109.5°)** and torsional strain (all hydrogens on adjacent carbon atoms are **staggered,** not eclipsed).




All H's are **staggered.**

- **In cyclohexane, three C atoms pucker up and three C atoms pucker down, alternating around the ring. These C atoms are called *up* C's and *down* C's.**

Each cyclohexane carbon atom has one axial and one equatorial hydrogen.


Each carbon in cyclohexane has two different kinds of hydrogens.

- ***Axial* hydrogens are located above and below the ring (along a perpendicular axis).**
- ***Equatorial* hydrogens are located in the plane of the ring (around the equator).**



A three-dimensional representation of the chair form is shown in Figure 4.12.


Figure 4.12
A three-dimensional model of the chair form of cyclohexane with all H atoms drawn



- Cyclohexane has **six axial H's** and **six equatorial H's.**


How To Draw the Chair Form of Cyclohexane

Step [1] Draw the carbon skeleton.




- Draw three parts of the chair: **a wedge, a set of parallel lines,** and **another wedge.**
- Then, join them together.
- The bottom 3 C's come out of the page, and for this reason, bonds to them are sometimes highlighted in bold.

Step [2] Label the *up* C's and *down* C's on the ring.



- There are 3 *up* and 3 *down* C's, and they alternate around the ring.


Step [3] Draw in the axial H atoms.




- On an *up* C the axial H is *up.*
- On a *down* C the axial H is *down.*

Step [4] Draw in the equatorial H atoms.


- The axial H is **down** on a down C, so the equatorial H must be up.
- The axial H is **up** on an up C, so the equatorial H must be down.




All equatorial H's drawn in.




All H's drawn in.



Problem 4.21 Classify the ring carbons as *up* C's or *down* C's. Identify the bonds highlighted in bold as axial or equatorial.



Problem 4.22 Using the cyclohexane with the C's numbered as shown, draw a chair form that fits each description.



a. The ring has an axial CH_3 group at C1 and an equatorial OH on C2.
b. The ring has an equatorial CH_3 group on C6 and an axial OH group on C4.
c. The ring has equatorial OH groups on C1, C2, and C5.

4.12B Ring-Flipping

Like acyclic alkanes, **cyclohexane does not remain in a single conformation.** The bonds twist and bend, resulting in new arrangements, but the movement is more restricted. One conformational change involves **ring-flipping,** which can be viewed as a two-step process.




- **A *down* carbon flips up.** This forms a new conformation of cyclohexane called a **boat.** The boat form has two carbons oriented above a plane containing the other four carbons.
- The boat form can flip in two possible ways. The carbon labeled with a red circle can flip down, re-forming the initial conformation; or the **second *up* carbon,** labeled with a blue circle, **can flip down. This forms a second chair conformation.**

Because of ring-flipping, the ***up* carbons become *down* carbons and the *down* carbons become *up* carbons.** Thus, cyclohexane exists as two different chair conformations of equal stability, which rapidly interconvert at room temperature.

The process of ring-flipping also affects the orientation of cyclohexane's hydrogen atoms.


- Axial and equatorial H atoms are interconverted during a ring flip. Axial H atoms become equatorial H atoms, and equatorial H atoms become axial H atoms (Figure 4.13).

Figure 4.13
Ring-flipping interconverts axial and equatorial hydrogens in cyclohexane.



The chair forms of cyclohexane are 30 kJ/mol more stable than the boat forms. The boat conformation is destabilized by torsional strain because the hydrogens on the four carbon atoms in the plane are eclipsed. Additionally, there is steric strain because two hydrogens at either end of the boat—the **flagpole hydrogens**—are forced close to each other, as shown in Figure 4.14.

Figure 4.14
Two views of the boat conformation of cyclohexane



The boat form of cyclohexane is less stable than the chair forms for two reasons.

- Eclipsing interactions between H's cause torsional strain.
- The proximity of the flagpole H's causes steric strain.

4.13 Substituted Cycloalkanes

What happens when one hydrogen on cyclohexane is replaced by a larger substituent? Is there a difference in the stability of the two cyclohexane conformations? To answer these questions, remember one rule.

- **The equatorial position has more room than the axial position, so larger substituents are more stable in the equatorial position.**


4.13A Cyclohexane with One Substituent

There are two possible chair conformations of a monosubstituted cyclohexane, such as methylcyclohexane, as shown in the following *How To*.


How To Draw the Two Conformations for a Substituted Cyclohexane

Step [1] Draw one chair form and add the substituents.

- Arbitrarily pick a ring carbon, classify it as an *up* or *down* carbon, and draw the bonds. Each C has one axial and one equatorial bond.
- Add the substituents, in this case H and CH_3, arbitrarily placing one axial and one equatorial. In this example, the CH_3 group is drawn equatorial.
- This forms one of the two possible chair conformations, labeled **A.**




Step [2] Ring-flip the cyclohexane ring.



- Convert *up* C's to *down* C's and vice versa. The chosen *up* C now puckers down.

Step [3] Add the substituents to the second conformation.


- Draw axial and equatorial bonds. On a *down* C the axial bond is *down.*
- Ring-flipping converts axial bonds to equatorial bonds, and vice versa. The equatorial methyl becomes axial.
- This forms the other possible chair conformation, labeled **B.**




Although the CH_3 group flips from equatorial to axial, it starts on a down bond, and stays on a down bond. It *never* flips from below the ring to above the ring.

- **A substituent always stays on the same side of the ring—either below or above—during the process of ring-flipping.**

Each carbon atom has one *up* and one *down* bond. An *up* bond can be either axial or equatorial, depending on the carbon to which it is attached. On an *up* C, the axial bond is *up,* but on a *down* C, the equatorial bond is *up.*




The two conformations of methylcyclohexane are different, so they are not equally stable. In fact, **A,** which places the larger methyl group in the roomier equatorial position, is considerably more stable than **B,** which places it axial.



Why is a substituted cyclohexane ring more stable with a larger group in the equatorial position? Figure 4.15 shows that with an equatorial CH_3 group, steric interactions with nearby groups are minimized. An axial CH_3 group, however, is close to two other axial H atoms, creating two destabilizing steric interactions called **1,3-diaxial interactions.** Each unfavorable H,CH_3 interaction destabilizes the conformation by 3.8 kJ/mol, so **B** is 7.6 kJ/mol less stable than **A.**


Figure 4.15
Three-dimensional representations for the two conformations of methylcyclohexane



- **Larger axial substituents create unfavorable 1,3-diaxial interactions, destabilizing a cyclohexane conformation.**

The larger the substituent on the six-membered ring, the higher the percentage of the conformation containing the equatorial substituent at equilibrium. With a very large substituent like *tert*-butyl [$(CH_3)_3C-$], essentially none of the conformation containing an axial *tert*-butyl group is present at room temperature, so **the ring is essentially anchored in a single conformation having an equatorial *tert*-butyl group.** This is illustrated in Figure 4.16.

Figure 4.16
The two conformations of *tert*-butylcyclohexane



- The large *tert*-butyl group anchors the cyclohexane ring in conformation **D.**

Problem 4.23 Draw a second chair conformation for each cyclohexane. Then decide which conformation is present in higher concentration at equilibrium.

a. 

b. 

c. 

Problem 4.24 Draw both conformations for 1-ethyl-1-methylcyclohexane and decide which conformation (if any) is more stable.

4.13B A Disubstituted Cycloalkane

Rotation around the C−C bonds in the ring of a cycloalkane is restricted, so **a group on one side of the ring can *never* rotate to the other side of the ring.** As a result, there are two different 1,2-dimethylcyclopentanes—one having two CH_3 groups on the **same side** of the ring and one having them on **opposite sides** of the ring.

Wedges indicate bonds in front of the plane of the ring and dashed wedges indicate bonds behind. For a review of this convention, see Section 1.7B. If a ring carbon is bonded to a CH_3 group in **front** of the ring (on a wedge), it is *assumed* that the other atom bonded to this carbon is hydrogen, located **behind** the ring (on a dashed wedge).



2 CH_3's above the ring
A
cis isomer



1 CH_3 above and 1 CH_3 below
B
trans isomer

A and **B** are **isomers,** because they are different compounds with the same molecular formula, but they represent the second major class of isomers called **stereoisomers.**

- ***Stereoisomers*** **are isomers that differ *only* in the way the atoms are oriented in space.**

Cis and **trans** isomers are named by adding the prefixes *cis* and *trans* to the name of the cycloalkane. Thus, **A** is *cis*-1,2-dimethylcyclopentane, and **B** is *trans*-1,2-dimethylcyclopentane.

The prefixes **cis** and **trans** are used to distinguish these stereoisomers.


- **The cis isomer has two groups on the *same side* of the ring.**
- **The trans isomer has two groups on *opposite sides* of the ring.**

Problem 4.25 Draw the structure for each compound using wedges and dashed wedges.
a. *cis*-1,2-dimethylcyclopropane b. *trans*-1-ethyl-2-methylcyclopentane


Problem 4.26 For *cis*-1,3-diethylcyclobutane, draw (a) a stereoisomer; (b) a constitutional isomer.

4.13C A Disubstituted Cyclohexane

A disubstituted cyclohexane like 1,4-dimethylcyclohexane also has cis and trans stereoisomers. In addition, each of these stereoisomers has two possible chair conformations.



trans-1,4-dimethylcyclohexane



cis-1,4-dimethylcyclohexane


All disubstituted cycloalkanes with two groups bonded to *different* atoms have cis and trans isomers.

To draw both conformations for each stereoisomer, follow the procedure in Section 4.13A for a monosubstituted cyclohexane, keeping in mind that two substituents must now be added to the ring.

How To Draw Two Conformations for a Disubstituted Cyclohexane

Step [1] Draw one chair form and add the substituents.

- For *trans*-1,4-dimethylcyclohexane, arbitrarily pick two C's located 1,4- to each other, classify them as *up* or *down* C's, and draw in the substituents.
- The trans isomer must have one group *above* the ring (on an *up* bond) and one group *below* the ring (on a *down* bond). The substituents can be either axial or equatorial, as long as one is up and one is down. The easiest trans isomer to visualize has two axial CH_3 groups. This arrangement is said to be **diaxial.**
- This forms one of the two possible chair conformations, labeled **A.**




Step [2] Ring-flip the cyclohexane ring.



- The *up* C flips down, and the *down* C flips up.


Step [3] Add the substituents to the second conformation.



two equatorial CH_3 groups
B


- Ring-flipping converts axial bonds to equatorial bonds, and vice versa. The diaxial CH_3 groups become diequatorial. This trans conformation is less obvious to visualize. It is still trans, because one CH_3 group is above the ring (on an *up* bond), and one is below (on a *down* bond).

Conformations **A** and **B** are not equally stable. **Because B has both larger CH_3 groups in the roomier equatorial position, B is lower in energy.**



The cis isomer of 1,4-dimethylcyclohexane also has two conformations, as shown in Figure 4.17. Because each conformation has one CH_3 group axial and one equatorial, they are **identical in energy.** At room temperature, therefore, the two conformations exist in a 50:50 mixture at equilibrium.

Figure 4.17
The two conformations of *cis*-1,4-dimethylcyclohexane



- A cis isomer has two groups on the same side of the ring, either both *up* or both *down*. In this example, Conformations **C** and **D** have two CH_3 groups drawn up.
- Both conformations have one CH_3 group axial and one equatorial, making them equally stable.


The relative stability of the two conformations of any disubstituted cyclohexane can be analyzed using this procedure.

- **A cis isomer has two substituents on the same side, either both on *up* bonds or both on *down* bonds.**
- **A trans isomer has two substituents on opposite sides, one *up* and one *down.***
- **Whether substituents are axial or equatorial depends on the relative location of the two substituents (on carbons 1,2-, 1,3-, or 1,4-).**

Sample Problem 4.3 Draw both chair conformations for *trans*-1,3-dimethylcyclohexane.


Solution

Step [1] Draw one chair form and add substituents.




- Pick two C's 1,3- to each other.
- The trans isomer has two groups on **opposite sides.** In Conformation **A,** one CH_3 is axial (on an *up* bond), and one group is equatorial (on a *down* bond).

Steps [2–3] Ring-flip and add substituents.



- The two *up* C's flip down.
- The axial CH_3 flips equatorial (still an *up* bond) and the equatorial CH_3 flips axial (still a *down* bond). Conformation **B** is trans because the two CH_3's are still on opposite sides.
- **Conformations A and B are equally stable** because each has one CH_3 equatorial and one axial.

Problem 4.27 Label each compound as cis or trans. Then draw the second chair conformation.



Problem 4.28 Consider 1,2-dimethylcyclohexane.

a. Draw structures for the cis and trans isomers using a hexagon for the six-membered ring.
b. Draw the two possible chair conformations for the cis isomer. Which conformation, if either, is more stable?
c. Draw the two possible chair conformations for the trans isomer. Which conformation, if either, is more stable?
d. Which isomer, cis or trans, is more stable and why?

Problem 4.29 Draw a chair conformation of cyclohexane with one CH_3CH_2 group and one CH_3 group that fits each description.

a. a 1,1-disubstituted cyclohexane with an axial CH_3CH_2 group
b. a cis-1,2-disubstituted cyclohexane with an axial CH_3 group
c. a trans-1,3-disubstituted cyclohexane with an equatorial CH_3 group
d. a trans-1,4-disubstituted cyclohexane with an equatorial CH_3CH_2 group

4.14 Oxidation of Alkanes

In Chapter 3 we learned that a functional group contains a heteroatom or π bond and constitutes **the reactive part of a molecule.** Alkanes are the only family of organic molecules that has no functional group, and therefore, **alkanes undergo few reactions.** In fact, alkanes are inert to reaction unless forcing conditions are used.

In Chapter 4, we consider only one reaction of alkanes—**combustion.** Combustion is an **oxidation–reduction** reaction.

4.14A Oxidation and Reduction Reactions

Compounds that contain many C–H bonds and few C–Z bonds are said to be in a ***reduced state,*** whereas those that contain few C–H bonds and more C–Z bonds are in a ***more oxidized state.*** CH_4 is highly reduced, while CO_2 is highly oxidized.

- *Oxidation* is the *loss* of electrons.
- *Reduction* is the *gain* of electrons.

Oxidation and reduction are opposite processes. As in acid–base reactions, there are always two components in these reactions. **One component is oxidized and one is reduced.**

To determine if an organic compound undergoes oxidation or reduction, we concentrate on the carbon atoms of the starting material and product, and **compare the relative number of C–H and C–Z bonds,** where Z = an element *more electronegative* than carbon (usually O, N, or X). Oxidation and reduction are then defined in two complementary ways.


- *Oxidation* results in an *increase* in the number of C–Z bonds; *or*
- *Oxidation* results in a *decrease* in the number of C–H bonds.

Because Z is more electronegative than C, replacing C–H bonds with C–Z bonds decreases the electron density around C. Loss of electron density = oxidation.


- *Reduction* results in a *decrease* in the number of C–Z bonds; *or*
- *Reduction* results in an *increase* in the number of C–H bonds.

Figure 4.18 illustrates the oxidation of CH_4 by replacing C–H bonds with C–O bonds (from left to right). The symbol **[O]** indicates oxidation. Because reduction is the reverse of oxidation, the molecules in Figure 4.18 are progressively reduced moving from right to left, from CO_2 to CH_4. The symbol **[H]** indicates reduction.

Figure 4.18
The oxidation and reduction of a carbon compound



Sample Problem 4.4 Determine whether the organic compound is oxidized or reduced in each transformation.




Solution

a. The conversion of ethanol to acetic acid is an **oxidation** because the number of C–O bonds increases: CH_3CH_2OH has one C–O bond and CH_3COOH has three C–O bonds.

b. The conversion of cyclohexene (C_6H_{10}) to cyclohexane (C_6H_{12}) is a **reduction** because the number of C–H bonds increases: cyclohexane has two more C–H bonds than cyclohexene.


Problem 4.30 Classify each transformation as an oxidation, reduction, or neither.



4.14B Combustion of Alkanes

When an organic compound is *oxidized* by a reagent, the reagent itself is *reduced*. Similarly, when an organic compound is *reduced* by a reagent, the reagent is *oxidized*. **Organic chemists identify a reaction as an oxidation or reduction by what happens to the *organic* component of the reaction.**

Alkanes undergo **combustion**—that is, **they burn in the presence of oxygen to form carbon dioxide and water.** This is a practical example of oxidation. Every C–H and C–C bond in the starting material is converted to a C–O bond in the product. The reactions drawn show the combustion of two different alkanes. The products, **CO_2 + H_2O,** are the same, regardless of the identity of the starting material. Combustion of alkanes in the form of natural gas, gasoline, or heating oil releases energy for heating homes, powering vehicles, and cooking food.




Combustion requires a spark or a flame to initiate the reaction. Gasoline, therefore, which is composed largely of alkanes, can be safely handled and stored in the air, but the presence of a spark or match causes immediate and violent combustion.

Driving an automobile 10,000 miles at 25 miles per gallon releases ~10,000 lb of CO_2 into the atmosphere.

The combustion of alkanes and other hydrocarbons obtained from fossil fuels adds a tremendous amount of CO_2 to the atmosphere each year. Quantitatively, data show over a 20% increase in the atmospheric concentration of CO_2 in the last 56 years (from 315 parts per million in 1958 to 395 parts per million in 2014; Figure 4.19). Although the composition of the atmosphere has changed over the lifetime of the earth, this may be the first time that the actions of humankind have altered that composition significantly and so quickly.


Figure 4.19
The changing concentration of CO_2 in the atmosphere since 1958




The increasing level of atmospheric CO_2 is clearly evident on the graph. Two data points are recorded each year. The sawtooth nature of the graph is due to seasonal variation of CO_2 level with the seasonal variation in photosynthesis. (Data recorded at Mauna Loa, Hawaii)

An increased CO_2 concentration in the atmosphere may have long-range and far-reaching effects. CO_2 absorbs thermal energy that normally radiates from the earth's surface, and redirects it back to the surface. Higher levels of CO_2 may therefore contribute to an increase in the average temperature of the earth's atmosphere. The global climate change resulting from these effects may lead to melting of the polar ice caps, a rise in sea level, and many more unforeseen consequences. How great a role CO_2 plays in this process is hotly debated.

Problem 4.31 Draw the products of each combustion reaction.

a. 

b. 

4.15 Lipids—Part 1

Lipids that contain carbon–carbon double bonds are discussed in Section 10.6.


Lipids are biomolecules whose properties resemble those of alkanes and other hydrocarbons. They are unlike any other class of biomolecules, because they are defined by a **physical property,** not by the presence of a particular functional group.

- **Lipids are biomolecules that are soluble in organic solvents and insoluble in water.**

Lipids have varied sizes and shapes, and a diverse number of functional groups. Fat-soluble vitamins like vitamin A and the phospholipids that comprise cell membranes are two examples of lipids that were presented in Sections 3.5 and 3.7. Other examples are shown in Figure 4.20. One unifying feature accounts for their solubility.

- **Lipids are composed of many nonpolar C–H and C–C bonds, and have few polar functional groups.**

Figure 4.20
Three representative lipid molecules






Waxes are lipids having two long alkyl chains joined by a single oxygen-containing functional group. Because of their many C–C and C–H bonds, **waxes are hydrophobic.** They form a protective coating on the feathers of birds to make them water repellent, and on leaves to prevent water evaporation. Bees secrete $\mathbf{CH_3(CH_2)_{14}COO(CH_2)_{29}CH_3}$, a wax that forms the honeycomb in which they lay eggs.

$\mathbf{PGF_{2\alpha}}$ belongs to a class of lipids called **prostaglandins.** Prostaglandins contain many C–C and C–H bonds and a single COOH group (a **carboxy group**). Prostaglandins possess a wide range of biological activities. They control inflammation, affect blood-platelet aggregation, and stimulate uterine contractions. Nonsteroidal anti-inflammatory drugs such as ibuprofen operate by blocking the synthesis of prostaglandins, as discussed in Section 19.6.

Cholesterol is a member of the steroid family, a group of lipids having four rings joined together. Because it has just one polar OH group, cholesterol is insoluble in the aqueous medium of the blood. It is synthesized in the liver and transported to other cells bound to water-soluble organic molecules. Elevated cholesterol levels can lead to coronary artery disease.

Figure 4.21
Cholesterol embedded in a lipid bilayer of a cell membrane




- The nonpolar hydrocarbon skeleton of cholesterol is embedded in the nonpolar interior of the cell membrane. Its rigid carbon skeleton stiffens the fluid lipid bilayer, giving it strength.
- Cholesterol's polar OH group is oriented toward the aqueous media inside and outside the cell.

Cholesterol is a vital component of the cell membrane. Its hydrophobic carbon chain is embedded in the interior of the lipid bilayer, and its hydrophilic hydroxy group is oriented toward the aqueous exterior (Figure 4.21). Because its tetracyclic carbon skeleton is quite rigid compared to the long floppy side chains of a phospholipid, cholesterol stiffens the cell membrane somewhat, giving it more strength.

Lipids have a high energy content, meaning that much energy is released on their metabolism. Because lipids are composed mainly of C−C and C−H bonds, they are oxidized with the release of energy, just like alkanes are. In fact, lipids are the most efficient biomolecules for the storage of energy. **The combustion of alkanes provides heat for our homes, and the metabolism of lipids provides energy for our bodies.**

Problem 4.32 Explain why beeswax is insoluble in H_2O, slightly soluble in ethanol (CH_3CH_2OH), and soluble in chloroform ($CHCl_3$).

Problem 4.33 Pristane is a high molecular weight alkane present in shark liver oil, a lipid used as a folk remedy and lubricant by coastal societies in northern Europe for centuries. What is the IUPAC name for pristane?


pristane


Shark liver oil, the source of pristane and other lipids, is obtained from sharks that inhabit the cold waters of the northern Atlantic and southern Pacific Oceans. (Photo © Eric Couture, courtesy Canadian Shark Research Laboratory.)


KEY CONCEPTS

Alkanes

General Facts About Alkanes (4.1–4.3)


- Alkanes are composed of **tetrahedral, sp^3** hybridized C atoms.
- There are two types of alkanes: acyclic alkanes having molecular formula $\mathbf{C_nH_{2n+2}}$, and cycloalkanes having molecular formula $\mathbf{C_nH_{2n}}$.
- Alkanes have only **nonpolar C – C and C – H bonds** and no functional group, so they undergo few reactions.
- Alkanes are named with the suffix ***-ane.***

Names of Alkyl Groups (4.4A)



Conformations in Acyclic Alkanes (4.9, 4.10)

- Alkane conformations can be classified as **eclipsed, staggered, anti,** or **gauche** depending on the relative orientation of the groups on adjacent carbons.



- dihedral angle = 0°
- dihedral angle = 60°
- dihedral angle of two CH_3 groups = 180°
- dihedral angle of two CH_3 groups = 60°

- A staggered conformation is **lower in energy** than an eclipsed conformation.
- An anti conformation is **lower in energy** than a gauche conformation.


Types of Strain

- **Torsional strain**—an increase in energy caused by eclipsing interactions (4.9).
- **Steric strain**—an increase in energy when atoms are forced too close to each other (4.10).
- **Angle strain**—an increase in energy when tetrahedral bond angles deviate from 109.5° (4.11).

Two Types of Isomers

[1] **Constitutional isomers**—isomers that differ in the way the atoms are connected to each other (4.1A).

[2] **Stereoisomers**—isomers that differ *only* in the way the atoms are oriented in space (4.13B).



Conformations in Cyclohexane (4.12, 4.13)

- Cyclohexane exists as **two chair conformations** in rapid equilibrium at room temperature.
- Each carbon atom on a cyclohexane ring has **one axial** and **one equatorial hydrogen.** Ring-flipping converts axial H's to equatorial H's, and vice versa.



- In substituted cyclohexanes, groups larger than hydrogen are more stable in the **roomier equatorial position.**
- Disubstituted cyclohexanes with substituents on different atoms exist as two possible stereoisomers.
  - The **cis** isomer has two groups on the **same side** of the ring, either both up or both down.
  - The **trans** isomer has two groups on **opposite sides** of the ring, one up and one down.


Oxidation–Reduction Reactions (4.14)


- **Oxidation** results in an **increase in the number of C – Z bonds** or a **decrease in the number of C – H bonds.**
- **Reduction** results in a **decrease in the number of C – Z bonds** or an **increase in the number of C – H bonds.**

PROBLEMS


Problems Using Three-Dimensional Models

4.34 Name each alkane using the ball-and-stick model, and classify each carbon as 1°, 2°, 3°, or 4°.

a. 


b. 


4.35 Consider the substituted cyclohexane shown in the ball-and-stick model.




a. Label the substituents on C1, C2, and C4 as axial or equatorial.
b. Are the substituents on C1 and C2 cis or trans to each other?
c. Are the substituents on C2 and C4 cis or trans to each other?
d. Draw the second possible conformation in the chair form, and classify it as more stable or less stable than the conformation shown in the three-dimensional model.

4.36 Convert each three-dimensional model to a Newman projection around the indicated bond.

a. 

b. 

c. 


Constitutional Isomers


4.37 Draw the structure of all compounds that fit the following descriptions.


a. five constitutional isomers having the molecular formula C_4H_8
b. nine constitutional isomers having the molecular formula C_7H_{16}
c. twelve constitutional isomers having the molecular formula C_6H_{12} and containing one ring


IUPAC Nomenclature


4.38 Give the IUPAC name for each compound.


a. 


c. 


e. 


g. 


i. 

b. 

d. 

f. 

h. 

j. 

4.39 Give the structure and IUPAC name for each of the nine isomers having molecular formula C_9H_{20} that contains seven carbons in the longest chain and two methyl groups as substituents.


4.40 Draw the structure corresponding to each IUPAC name.


a. 3-ethyl-2-methylhexane
b. *sec*-butylcyclopentane
c. 4-isopropyl-2,4,5-trimethylundecane
d. cyclobutylcycloheptane
e. 3-ethyl-1,1-dimethylcyclohexane
f. 4-butyl-1,1-diethylcyclooctane
g. 6-isopropyl-2,3-dimethyldodecane
h. 2,2,6,6,7-pentamethyloctane
i. *cis*-1-ethyl-3-methylcyclopentane
j. *trans*-1-*tert*-butyl-4-ethylcyclohexane


4.41 Each of the following IUPAC names is incorrect. Explain why it is incorrect and give the correct IUPAC name.

a. 2,2-dimethyl-4-ethylheptane
b. 5-ethyl-2-methylhexane
c. 2-methyl-2-isopropylheptane
d. 1,5-dimethylcyclohexane
e. 1-ethyl-2,6-dimethylcycloheptane
f. 5,5,6-trimethyloctane
g. 3-butyl-2,2-dimethylhexane
h. 1,3-dimethylbutane

4.42 Give the IUPAC name for each compound.


a. 

b. 

c. 

Physical Properties



4.43 Rank the following alkanes in order of increasing boiling point.





4.44 The melting points and boiling points of two isomeric alkanes are as follows: $CH_3(CH_2)_6CH_3$, mp = –57 °C and bp = 126 °C; $(CH_3)_3CC(CH_3)_3$, mp = 102 °C and bp = 106 °C. (a) Explain why one isomer has a lower melting point but higher boiling point. (b) Explain why there is a small difference in the boiling points of the two compounds, but a huge difference in their melting points.


Conformation of Acyclic Alkanes

4.45 Which conformation in each pair is *higher* in energy? Calculate the energy difference between the two conformations using the values given in Table 4.3.




a.  or 

b.  or 

4.46 Considering rotation around the bond highlighted in red in each compound, draw Newman projections for the most stable and least stable conformations.

a.  b. 




4.47 Convert each structure to a Newman projection around the bond highlighted in red.

a.  b.  c. 



4.48 (a) Using Newman projections, draw all staggered and eclipsed conformations that result from rotation around the bond highlighted in red in each molecule; (b) draw a graph of energy versus dihedral angle for rotation around this bond.

[1]  [2] 

4.49 Label the sites of torsional and steric strain in each conformation.

a.  b.  c. 

4.50 Calculate the barrier to rotation for each bond highlighted in red.

a.  b. 




4.51 The eclipsed conformation of CH_3CH_2Cl is 15 kJ/mol less stable than the staggered conformation. How much is the H,Cl eclipsing interaction worth in destabilization?

4.52 (a) Draw the anti and gauche conformations for ethylene glycol ($HOCH_2CH_2OH$). (b) Ethylene glycol is unusual in that the gauche conformation is more stable than the anti conformation. Offer an explanation.

Conformations and Stereoisomers in Cycloalkanes

4.53 For each compound drawn below:

a. Label each OH, Br, and CH_3 group as axial or equatorial.
b. Classify each conformation as cis or trans.
c. Translate each structure into a representation with a hexagon for the six-membered ring, and wedges and dashed wedges for groups above and below the ring.
d. Draw the second possible chair conformation for each compound.

[1]  [2]  [3] 

4.54 Draw the two possible chair conformations for *cis*-1,3-dimethylcyclohexane. Which conformation, if either, is more stable?


4.55 For each compound drawn below:


a. Draw representations for the cis and trans isomers using a hexagon for the six-membered ring, and wedges and dashed wedges for substituents.


b. Draw the two possible chair conformations for the cis isomer. Which conformation, if either, is more stable?

c. Draw the two possible chair conformations for the trans isomer. Which conformation, if either, is more stable?


d. Which isomer, cis or trans, is more stable and why?


[1] 

[2] 


[3] 

4.56 Convert each of the following structures into its more stable chair form. One structure represents menthol and one represents isomenthol. Menthol, the more stable isomer, is used in lip balms and mouthwash. Which structure corresponds to menthol?





4.57 Answer the following questions about compound **A,** which contains a CH_3 group and OH group bonded to the carbon skeleton that consists of three six-membered rings in the conformation shown.



A

a. Are the CH_3 and OH groups oriented cis or trans to each other?


b. Is a substituent on C_a that is cis to the CH_3 group located in the axial or equatorial position?

c. Is an equatorial Br at C_b oriented cis or trans to the OH group?

d. Is the H atom on C_c located cis or trans to the OH group?

e. Is a substituent on C_d that is trans to the OH group located in the axial or equatorial position?

4.58 Glucose is a simple sugar with five substituents bonded to a six-membered ring.



glucose

a. Using a chair representation, draw the most stable arrangement of these substituents on the six-membered ring.



b. Convert this representation into one that uses a hexagon with wedges and dashed wedges.



c. Draw a constitutional isomer of glucose.



d. Draw a stereoisomer that has an axial OH group on one carbon.


Constitutional Isomers and Stereoisomers


4.59 Classify each pair of compounds as constitutional isomers, stereoisomers, identical molecules, or not isomers of each other.

a.  and 


c.  and 


b.  and 


d. 


and 

4.60 Classify each pair of compounds as constitutional isomers or identical molecules.


a. 


and 


b. 


and 

4.61 Answer the following questions about compounds **A–D.**


A


B


C



D


a. How are the compounds in each pair related? Choose from constitutional isomers, stereoisomers, or identical molecules: **A** and **B; A** and **C; B** and **D.**
b. Label each compound as a cis or trans isomer.
c. Draw **B** as a hexagon with wedges and dashed wedges to show the stereochemistry of substituents.
d. Draw a stereoisomer of **A** as a hexagon using wedges and dashed wedges to show the orientation of substituents.

4.62 Draw the three constitutional isomers having molecular formula C_7H_{14} that contain a five-membered ring and two methyl groups as substituents. For each constitutional isomer that can have cis and trans isomers, draw the two stereoisomers.


Oxidation and Reduction


4.63 Classify each reaction as oxidation, reduction, or neither.

a. 


b. 

4.64 Draw the products of combustion of each alkane.

a. 

b. 

4.65 Hydrocarbons like benzene are metabolized in the body to arene oxides, which rearrange to form phenols. This is an example of a general process in the body, in which an unwanted compound (benzene) is converted to a more water-soluble derivative called a *metabolite,* so that it can be excreted more readily from the body.




benzene arene oxide phenol

a. Classify each of these reactions as oxidation, reduction, or neither.
b. Explain why phenol is more water soluble than benzene. This means that phenol dissolves in urine, which is largely water, to a greater extent than benzene.

Lipids

4.66 Cholic acid, a compound called a **bile acid,** is converted to a **bile salt** in the body. Bile salts have properties similar to soaps, and they help transport lipids through aqueous solutions. Explain why this is so.



cholic acid
a bile acid


bile salt

4.67 Mineral oil, a mixture of high molecular weight alkanes, is sometimes used as a laxative. Why are individuals who use mineral oil for this purpose advised to avoid taking it at the same time they consume foods rich in fat-soluble vitamins such as vitamin A?

Challenge Problems

4.68 Cyclopropane and cyclobutane have similar strain energy despite the fact that the C–C–C bond angles of cyclopropane are much smaller than those of cyclobutane. Suggest an explanation for this observation, considering all sources of strain discussed in Chapter 4.


4.69 Although penicillin G has two amide functional groups, one is much more reactive than the other. Which amide is more reactive and why?



penicillin G

4.70 Haloethanes (CH_3CH_2X, X = Cl, Br, I) have similar barriers to rotation (13.4–15.5 kJ/mol) despite the fact that the size of the halogen increases, Cl → Br → I. Offer an explanation.


4.71 When two six-membered rings share a C–C bond, this bicyclic system is called a **decalin.** There are two possible arrangements: *trans*-decalin having two hydrogen atoms at the ring fusion on opposite sides of the rings, and *cis*-decalin having the two hydrogens at the ring fusion on the same side.



decalin *trans*-decalin *cis*-decalin


a. Draw *trans*- and *cis*-decalin using the chair form for the cyclohexane rings.
b. The trans isomer is more stable. Explain why.

4.72 Consider the tricyclic structure **A.** (a) Label each substituent on the rings as axial or equatorial. (b) Draw a skeletal structure for **A,** using wedges and dashed wedges to show whether the substituents are located above or below the rings.



A


4.73 Consider the tricyclic structure **B.** (a) Label each substituent on the rings as axial or equatorial. (b) Draw **B** using chair conformations for each six-membered ring. (c) Label the atoms on the ring fusions (the carbons that join each set of two rings together) as cis or trans to each other.




B


4.74 Read Appendix B on naming branched alkyl substituents, and draw all possible alkyl groups having the formula $C_5H_{11}-$. Give the IUPAC names for the eight compounds of molecular formula $C_{10}H_{20}$ that contain a cyclopentane ring with each of these alkyl groups as a substituent.

4.75 Read Appendix B on naming bicyclic compounds. Then give the IUPAC name for each of the following compounds.

a. 


b. 

c. 

d. 

5 Stereochemistry

5.1 Starch and cellulose
5.2 The two major classes of isomers
5.3 Looking glass chemistry—Chiral and achiral molecules
5.4 Stereogenic centers
5.5 Stereogenic centers in cyclic compounds
5.6 Labeling stereogenic centers with *R* or *S*
5.7 Diastereomers
5.8 Meso compounds
5.9 *R* and *S* assignments in compounds with two or more stereogenic centers
5.10 Disubstituted cycloalkanes
5.11 Isomers—A summary
5.12 Physical properties of stereoisomers
5.13 Chemical properties of enantiomers



Paclitaxel (trade name Taxol), a potent anticancer agent active against ovarian, breast, and several other cancers, was discovered in 1962 and approved for use by the Food and Drug Administration in 1992. Initial studies with paclitaxel were carried out with material isolated from the bark of the Pacific yew tree, but stripping the bark killed these magnificent trees. Paclitaxel was synthesized in the laboratory in 1994, and is now produced by a plant cell fermentation process. Like other widely used drugs, paclitaxel is biologically active because of its complex structure and the particular three-dimensional arrangement of its functional groups. In Chapter 5 we learn about the stereochemistry of molecules like paclitaxel.

Are you left-handed or right-handed? If you're right-handed, you've probably spent little time thinking about your hand preference. If you're left-handed, though, you probably learned at an early age that many objects—like scissors and baseball gloves—"fit" for righties, but are "backwards" for lefties. **Hands, like many objects in the world around us, are mirror images that are *not* identical.**


In Chapter 5 we examine the "handedness" of molecules, and ask, "How important is the three-dimensional shape of a molecule?"

5.1 Starch and Cellulose

Recall from Chapter 4 that ***stereochemistry* is the three-dimensional structure of a molecule.** How important is stereochemistry? Two biomolecules—starch and cellulose—illustrate how apparently minute differences in structure can result in vastly different properties.


Starch and **cellulose** are two polymers that belong to the family of biomolecules called **carbohydrates** (Figure 5.1). **A *polymer* is a large molecule composed of repeating smaller units—called monomers—that are covalently bonded together.**

Figure 5.1 Starch and cellulose—Two common carbohydrates




Starch is the main carbohydrate in the seeds and roots of plants. When we humans ingest wheat, rice, or potatoes, we consume starch, which is then hydrolyzed to the simple sugar **glucose,** one of the compounds our bodies use for energy. **Cellulose,** nature's most abundant organic material, gives rigidity to tree trunks and plant stems. Wood, cotton, and flax are composed largely of cellulose. Complete hydrolysis of cellulose also forms glucose, but unlike starch, humans cannot metabolize cellulose to glucose. In other words, we can digest starch but not cellulose.

Cellulose and starch are both composed of the same repeating unit—a six-membered ring containing an oxygen atom and three OH groups—joined by an oxygen atom. They differ in the position of the O atom joining the rings together.




- **In cellulose, the O atom joins two rings using two equatorial bonds.**
- **In starch, the O atom joins two rings using one equatorial and one axial bond.**



cellulose

two equatorial bonds (in red)



starch


one axial bond (in blue) and one equatorial bond (in red)

Starch and cellulose are **isomers** because they are different compounds with the same molecular formula $(C_6H_{10}O_5)_n$. They are **stereoisomers** because only the three-dimensional arrangement of atoms is different.

How the six-membered rings are joined together has an enormous effect on the shape and properties of these carbohydrate molecules. Cellulose is composed of long chains held together by intermolecular hydrogen bonds, forming sheets that stack in an extensive three-dimensional network. The axial–equatorial ring junction in starch creates chains that fold into a helix (Figure 5.2). Moreover, the human digestive system contains the enzyme necessary to hydrolyze starch by cleaving its axial C–O bond, but not an enzyme to hydrolyze the equatorial C–O bond in cellulose.


Figure 5.2
Three-dimensional structure of cellulose and starch

Cellulose



- Cellulose consists of an extensive three-dimensional network held together by hydrogen bonds.

Starch



- The starch polymer is composed of chains that wind into a helix.

Thus, an **apparently minor difference in the three-dimensional arrangement of atoms confers very different properties on starch and cellulose.**

Problem 5.1 Cellulose is water insoluble, despite its many OH groups. Considering its three-dimensional structure, why do you think this is so?

5.2 The Two Major Classes of Isomers

Because an understanding of isomers is integral to the discussion of stereochemistry, let's begin with an overview of isomers.

- **Isomers are different compounds with the same molecular formula.**

There are two major classes of isomers: **constitutional isomers** and **stereoisomers.** ***Constitutional (or structural) isomers*** **differ in the way the atoms are connected to each other.** Constitutional isomers have:

- different IUPAC names;
- the same or different functional groups;
- different physical properties, so they are separable by physical techniques such as distillation; and
- different chemical properties. They behave differently or give different products in chemical reactions.

Stereoisomers **differ *only* in the way atoms are oriented in space.** Stereoisomers have identical IUPAC names (except for a prefix like cis or trans). Because they differ only in the three-dimensional arrangement of atoms, stereoisomers always have the same functional group(s).

A particular three-dimensional arrangement is called a *configuration*. Thus, stereoisomers differ in configuration. The cis and trans isomers in Section 4.13B and the biomolecules starch and cellulose in Section 5.1 are two examples of stereoisomers.

Figure 5.3 illustrates examples of both types of isomers. Chapter 5 concentrates on the types and properties of stereoisomers.

Problem 5.2 Classify each pair of compounds as constitutional isomers or stereoisomers.





Figure 5.3 A comparison of constitutional isomers and stereoisomers




5.3 Looking Glass Chemistry—Chiral and Achiral Molecules



Despite the dominance of right-handedness over left-handedness, even identical twins can exhibit differences in hand preference. Pictured are Matthew (right-handed) and Zachary (left-handed), identical twin sons of the author.


Everything has a mirror image. What's significant is **whether a molecule is *identical* to or *different* from its mirror image.**

Some molecules are like hands. **Left and right hands are mirror images of each other, but they are *not* identical.** If you try to mentally place one hand inside the other hand, you can never superimpose either all the fingers, or the tops and palms. To *superimpose* an object on its mirror image means to align *all* parts of the object with its mirror image. With molecules, this means aligning all atoms and all bonds.



- A molecule (or object) that is *not* superimposable on its mirror image is said to be *chiral.*

Other molecules are like socks. **Two socks from a pair are mirror images that *are* superimposable.** One sock can fit inside another, aligning toes and heels, and tops and bottoms. A sock and its mirror image are *identical*.



- A molecule (or object) that *is* superimposable on its mirror image is said to be *achiral.*

Let's determine whether three molecules—H_2O, CH_2BrCl, and CHBrClF—are superimposable on their mirror images; that is, **are H_2O, CH_2BrCl, and CHBrClF chiral or achiral?**


The adjective ***chiral*** comes from the Greek *cheir,* meaning "hand." Left and right hands are **chiral:** they are mirror images that do not superimpose on each other.


To test chirality:

- Draw the molecule in three dimensions.
- Draw its mirror image.
- Try to align all bonds and atoms. To superimpose a molecule and its mirror image you can perform any rotation but **you cannot break bonds.**

Few beginning students of organic chemistry can readily visualize whether a compound and its mirror image are superimposable by looking at drawings on a two-dimensional page. Molecular models can help a great deal in this process.


Following this procedure, H_2O and CH_2BrCl are both **achiral** molecules because each molecule is superimposable on its mirror image.





CH_2BrCl is achiral.

With CHBrClF, the result is different. The molecule (labeled **A**) and its mirror image (labeled **B**) are not superimposable. No matter how you rotate **A** and **B,** all the atoms never align. **CHBrClF is thus a chiral molecule,** and **A** and **B** are different compounds.



CHBrClF is chiral.

A and **B** are **stereoisomers** because they are isomers differing only in the three-dimensional arrangement of substituents. These stereoisomers are called **enantiomers.**

- *Enantiomers* are mirror images that are not superimposable.

CHBrClF contains a carbon atom bonded to four different groups. **A carbon atom bonded to four different groups is called a tetrahedral *stereogenic center.*** Most chiral molecules contain one or more stereogenic centers.

Naming a carbon atom with four different groups is a topic that currently has no firm agreement among organic chemists. The IUPAC recommends the term *chirality center,* but the term has not gained wide acceptance among organic chemists since it was first suggested in 1996. Other terms in common use are chiral center, chiral carbon, asymmetric carbon, stereocenter, and stereogenic center, the term used in this text.

The general term *stereogenic center* refers to any site in a molecule at which the interchange of two groups forms a stereoisomer. A **carbon atom with four different groups is a *tetrahedral* stereogenic center,** because the interchange of two groups converts one enantiomer into another. We will learn about another type of stereogenic center in Section 8.2B.

We have now learned two related but different concepts, and it is necessary to distinguish between them.

- A molecule that is not superimposable on its mirror image is a *chiral molecule.*
- A carbon atom bonded to four different groups is a *stereogenic center.*


Molecules can contain zero, one, or more stereogenic centers.


- **With no stereogenic centers, a molecule generally is not chiral.** H_2O and CH_2BrCl have *no* stereogenic centers and are *achiral* molecules. (There are a few exceptions to this generalization, as we will learn in Section 17.5.)
- **With one tetrahedral stereogenic center, a molecule is *always* chiral.** CHBrClF is a *chiral* molecule containing *one* stereogenic center.
- **With two or more stereogenic centers, a molecule *may* or *may not* be chiral,** as we will learn in Section 5.8.

Problem 5.3 Draw the mirror image of each compound. Label each molecule as chiral or achiral.

a. 

b. 

c. 

d. 


When trying to distinguish between chiral and achiral compounds, keep in mind:

- **A *plane of symmetry* is a mirror plane that cuts a molecule in half, so that one half of the molecule is a reflection of the other half.**

- **Achiral molecules usually contain a plane of symmetry but chiral molecules do not.**

The achiral molecule CH_2BrCl has a plane of symmetry, but the chiral molecule CHBrClF does not.


CH_2BrCl
plane of symmetry



This molecule has
two identical halves.

CH_2BrCl is achiral.

CHBrClF
NO plane of symmetry



CHBrClF is chiral.

Figure 5.4 summarizes the main facts about chirality we have learned thus far.


Figure 5.4
The basic principles of chirality


- Everything has a mirror image. The fundamental question is whether a molecule and its mirror image are superimposable.
- If a molecule and its mirror image are *not* superimposable, the molecule and its mirror image are *chiral.*
- The terms *stereogenic center* and *chiral molecule* are related but distinct. In general, a chiral molecule must have one or more stereogenic centers.
- The presence of a *plane of symmetry* makes a molecule achiral.

Problem 5.4 Draw in a plane of symmetry for each molecule.


a. 


b. 

c. 

d. 

Problem 5.5 A molecule is achiral if it has a plane of symmetry in *any* conformation. Each of the following conformations does not have a plane of symmetry, but rotation around a carbon–carbon bond forms a conformation that does have a plane of symmetry. Draw this conformation for each molecule.

a. 

b. 



When a right-handed shell is held in the right hand with the thumb pointing towards the wider end, the opening is on the right side.


Stereochemistry may seem esoteric, but chirality pervades our very existence. On a molecular level, many biomolecules fundamental to life are chiral. On a macroscopic level, many naturally occurring objects possess handedness. Examples include chiral helical seashells shaped like right-handed screws, and plants such as honeysuckle that wind in a chiral left-handed helix. The human body is chiral, and hands, feet, and ears are not superimposable.

5.4 Stereogenic Centers


A necessary skill in the study of stereochemistry is the ability to locate and draw tetrahedral stereogenic centers.

5.4A Stereogenic Centers on Carbon Atoms That Are Not Part of a Ring

Recall from Section 5.3 that any carbon atom bonded to four different groups is a tetrahedral stereogenic center. To locate a stereogenic center, examine each *tetrahedral* carbon atom in a molecule, and look at the four ***groups***—not the four *atoms*—bonded to it. CBrClFI has one stereogenic center because its central carbon atom is bonded to four different elements. 3-Bromohexane also has one stereogenic center because one carbon is bonded to H, Br, CH_2CH_3, and $CH_2CH_2CH_3$. We consider all atoms in a group as a *whole unit,* not just the atom directly bonded to the carbon in question. Although C3 of 3-bromohexane is bonded to two carbon atoms, one is part of an ethyl group and one is part of a propyl group.



stereogenic center



3-bromohexane


The stereogenic center is bonded to:
- H
- Br
- CH_2CH_3
- $CH_2CH_2CH_3$


Ephedrine is isolated from ma huang, an herb used to treat respiratory ailments in traditional Chinese medicine. Once a popular drug to promote weight loss and enhance athletic performance, ephedrine has now been linked to episodes of sudden death, heart attack, and stroke.


Always omit from consideration all C atoms that can't be tetrahedral stereogenic centers. These include:

- CH_2 and CH_3 groups (more than one H bonded to C)
- any *sp* or sp^2 hybridized C (less than four groups around C)


Larger organic molecules can have two, three, or even hundreds of stereogenic centers. **Propoxyphene** and **ephedrine** each contain two stereogenic centers, and **fructose,** a simple carbohydrate, has three.



propoxyphene
Trade name Darvon
(analgesic)



ephedrine
(bronchodilator, decongestant)




fructose
(a simple sugar)

= stereogenic center


Sample Problem 5.1 Locate the stereogenic centers in each drug. Albuterol is a bronchodilator—that is, it widens airways—so it is used to treat asthma. Chloramphenicol is an antibiotic used extensively in developing countries because of its low cost.

a.



albuterol

b.




chloramphenicol


Heteroatoms surrounded by four different groups are also stereogenic centers. Stereogenic N atoms are discussed in Chapter 25.

Solution

Omit all CH_2 and CH_3 groups and all doubly bonded (sp^2 hybridized) C's. In albuterol, one C has three CH_3 groups bonded to it, so it can be eliminated as well. Draw in H atoms on tetrahedral C's in skeletal structures to more clearly see the groups. This leaves one C in albuterol and two C's in chloramphenicol surrounded by four different groups, making them stereogenic centers.


a. 


one stereogenic center


b. 


two stereogenic centers


Problem 5.6 Locate the stereogenic centers in each molecule. Compounds may have one or more stereogenic centers.


a. 

b. 


c. 

d. 

e. 

f. 


Problem 5.7 The facts in Section 5.4A can be used to locate stereogenic centers in any molecule, no matter how complicated. Always look for carbons surrounded by four different groups. With this in mind, locate the four stereogenic centers in aliskiren, a drug introduced in 2007 for the treatment of hypertension.



aliskiren

5.4B Drawing a Pair of Enantiomers


- **Any molecule with one tetrahedral stereogenic center is a chiral compound and exists as a pair of enantiomers.**



butan-2-ol
one stereogenic center

Butan-2-ol, for example, has one stereogenic center. To draw both enantiomers, use the typical convention for depicting a tetrahedron: **place two bonds in the plane, one in front of the plane on a wedge, and one behind the plane on a dashed wedge.** Then, to form the first enantiomer **A,** arbitrarily place the four groups—H, OH, CH_3, and CH_2CH_3—on any bond to the stereogenic center.

Draw the molecule…then the mirror image.




mirror
not superimposable
enantiomers


In Section 28.2, we will learn about Fischer projection formulas, an older convention used for drawing stereogenic centers utilized mainly in carbohydrate chemistry.

Then, draw a mirror plane and arrange the substituents in the mirror image so that they are a reflection of the groups in the first molecule, forming **B.** No matter how **A** and **B** are rotated, it is impossible to align all of their atoms. Because **A** and **B** are mirror images and not superimposable, **A** and **B** are a pair of **enantiomers.** Two other pairs of enantiomers are drawn in Figure 5.5.

Figure 5.5
Three-dimensional representations for pairs of enantiomers




Problem 5.8 Locate the stereogenic center in each compound and draw both enantiomers.




5.5 Stereogenic Centers in Cyclic Compounds

In drawing a tetrahedron using solid lines, wedges, and dashed wedges, always draw the two solid lines first; then draw the wedge and the dashed wedge on the *opposite side* of the solid lines.




Stereogenic centers may also occur at carbon atoms that are part of a ring. To find stereogenic centers on ring carbons always draw the rings as flat polygons, and look for tetrahedral carbons that are bonded to four different groups, as usual. Each ring carbon is bonded to two other atoms in the ring, as well as two substituents attached to the ring. When the two substituents on the ring are *different,* we must compare the ring atoms equidistant from the atom in question.

Does methylcyclopentane have a stereogenic center? All of the carbon atoms are bonded to two or three hydrogen atoms except for C1, the ring carbon bonded to the methyl group. Next, compare the ring atoms and bonds on both sides equidistant from C1, and continue until a point of difference is reached, or until both sides meet, either at an atom or in the middle of a bond. In this case, there is no point of difference on either side, so C1 is bonded to identical alkyl groups that happen to be part of a ring. **C1, therefore, is *not* a stereogenic center.**




With 3-methylcyclohexene, the result is different. All carbon atoms are bonded to two or three hydrogen atoms or are sp^2 hybridized except for C3, the ring carbon bonded to the methyl group. In this case, the atoms equidistant from C3 are different, so C3 is bonded to *different* alkyl groups in the ring. **C3 is therefore bonded to four different groups, making it a stereogenic center.**




Two enantiomers are *different* compounds. To convert one enantiomer to another you must **switch the position of two atoms.** This amounts to breaking bonds.

Because 3-methylcyclohexene has one tetrahedral stereogenic center it is a chiral compound and exists as a pair of enantiomers.



Many biologically active compounds contain one or more stereogenic centers on ring carbons. For example, **thalidomide,** a drug once prescribed as a sedative and anti-nausea agent for pregnant women in Great Britain and Europe, contains one stereogenic center, so it exists as a pair of enantiomers.







Today, thalidomide is prescribed under strict controls for the treatment of Hansen's disease (leprosy). Because it was once thought to be highly contagious, individuals in Hawai'i with Hansen's disease were sent to Kalaupapa, a remote and inaccessible peninsula on the north shore of the Hawaiian island of Moloka'i. Hansen's disease is now known to be a curable bacterial infection, which is treated by the sulfa drugs discussed in Section 25.16.


Unfortunately thalidomide was sold as a mixture of its two enantiomers, and each of these stereoisomers has a different biological activity. This is a property not uncommon in chiral drugs, as we will see in Section 5.13. Although one enantiomer was an effective sedative and anti-nausea drug, the other enantiomer was responsible for thousands of catastrophic birth defects in children born to women who took the drug during pregnancy. Thalidomide was never approved for use in the United States due to the diligence of Frances Oldham Kelsey, a medical reviewing officer for the Food and Drug Administration, who insisted that the safety data on thalidomide were inadequate.


Sucrose and **paclitaxel** (the chapter-opening molecule) are two useful compounds with several stereogenic centers at ring carbons. Identify the stereogenic centers in these more complicated compounds in exactly the same way, looking at one carbon at a time. **Sucrose,** with nine stereogenic centers on two rings, is the carbohydrate used as table sugar. **Paclitaxel,** with 11 stereogenic centers, is an anticancer agent active against ovarian, breast, and some lung tumors.




Problem 5.9 Locate the stereogenic centers in each compound. A molecule may have zero, one, or more stereogenic centers. Gabapentin [part (d)] is used clinically to treat seizures and certain types of chronic pain. Gabapentin enacarbil [part (e)] is a related compound that is three times more potent.


a. 

c. 


e. 

gabapentin enacarbil


b. 

d. 


gabapentin

f. 

Problem 5.10 Locate the stereogenic centers in each compound.

a. 

cholesterol

b. 

simvastatin
Trade name Zocor
(cholesterol-lowering drug)

5.6 Labeling Stereogenic Centers with *R* or *S*


Naming enantiomers with the prefixes *R* or *S* is called the Cahn–Ingold–Prelog system after the three chemists who devised it.

Because enantiomers are two different compounds, we need a method to distinguish them by name. This is done by adding the prefix *R* or *S* to the IUPAC name of the enantiomer. To designate an enantiomer as *R* or *S,* first **assign a priority** (1, 2, 3, or 4) to each group bonded to the stereogenic center, and then use these priorities to label one enantiomer *R* and one *S*.

Rules Needed to Assign Priority


Rule 1 **Assign priorities (1, 2, 3, or 4) to the atoms directly bonded to the stereogenic center in order of decreasing atomic number. The atom of *highest* atomic number gets the *highest* priority (1).**

- In CHBrClF, priorities are assigned as follows: Br (1, highest) → Cl (2) → F (3) → H (4, lowest). In many molecules the lowest priority group will be H.



Rule 2 **If two atoms on a stereogenic center are the *same,* assign priority based on the atomic number of the atoms bonded to these atoms. *One* atom of higher atomic number determines a higher priority.**

- With butan-2-ol, the O atom gets highest priority (1) and H gets lowest priority (4) using Rule 1. Butan-2-ol also has two carbon atoms bonded to the stereogenic center, one that is part of a CH_3 group and one that is part of a CH_2CH_3 group. To assign priority (either 2 or 3) to the two C atoms, look at what atoms (other than the stereogenic center) are bonded to each C.




- The CH_2CH_3 gets higher priority (2) than the CH_3 group (priority 3) because the carbon of the ethyl group is bonded to another carbon.
- The order of priority of groups in butan-2-ol is: $-OH$ **(1),** $-CH_2CH_3$ **(2),** $-CH_3$ **(3),** and $-H$ **(4).**
- If priority still cannot be assigned, continue along a chain until a point of difference is reached.

Rule 3 **If two isotopes are bonded to the stereogenic center, assign priorities in order of decreasing *mass* number.**


- In comparing two isotopes of the element hydrogen, deuterium, which has a mass number of two (one proton and one neutron), has a higher priority than hydrogen, which has a mass number of one (one proton only).

Rule 4 **To assign a priority to an atom that is part of a multiple bond, treat a multiply bonded atom as an equivalent number of singly bonded atoms.**

- The C of a C=O is considered to be bonded to two O atoms.



- Other common multiple bonds are drawn below.








Figure 5.6 gives examples of priorities assigned to stereogenic centers.

Figure 5.6
Examples of assigning priorities to stereogenic centers




- The stereogenic center is bonded to Br, Cl, C, and H.
- The stereogenic center is *not* bonded directly to I.



- $CH(CH_3)_2$ gets the highest priority because the C is bonded to 2 other C's.

[* = stereogenic center]



- OH gets the highest priority because O has the highest atomic number.
- CO_2H (three bonds to O) gets higher priority than CH_2OH (one bond to O).

Problem 5.11 Which group in each pair is assigned the *higher* priority?

a. $-CH_3$, $-CH_2CH_3$
b. $-I$, $-Br$
c. $-H$, $-D$
d. $-CH_2Br$, $-CH_2CH_2Br$
e. $-CH_2CH_2Cl$, $-CH_2CH(CH_3)_2$
f. $-CH_2OH$, $-CHO$

Problem 5.12 Rank the following groups in order of *decreasing* priority.


a. $-COOH$, $-H$, $-NH_2$, $-OH$
b. $-H$, $-CH_3$, $-Cl$, $-CH_2Cl$
c. $-CH_2CH_3$, $-CH_3$, $-H$, $-CH(CH_3)_2$
d. $-CH=CH_2$, $-CH_3$, $-C\equiv CH$, $-H$

R is derived from the Latin word *rectus* meaning "right" and *S* is from the Latin word *sinister* meaning "left."

Once priorities are assigned to the four groups around a stereogenic center, we can use three steps to designate the center as either *R* or *S*.

How To Assign *R* or *S* to a Stereogenic Center


Example **Label each enantiomer as *R* or *S*.**



two enantiomers of butan-2-ol

Step [1] **Assign priorities from 1 to 4 to each group bonded to the stereogenic center.**

- The priorities for the four groups around the stereogenic center in butan-2-ol were given in Rule 2, on page 186.





—*Continued*

How To, continued . . .


Step [2] **Orient the molecule with the lowest priority group (4) *back* (on a *dashed wedge*), and visualize the relative positions of the remaining three groups (priorities 1, 2, and 3).**

- For each enantiomer of butan-2-ol, look toward the lowest priority group, drawn behind the plane, down the C – H bond.






Step [3] **Trace a circle from priority group 1 → 2 → 3.**




clockwise
***R* isomer**




counterclockwise
***S* isomer**

- If tracing the circle goes in the **clockwise** direction—to the right from the noon position—the isomer is named ***R*.**
- If tracing the circle goes in the **counterclockwise** direction—to the left from the noon position—the isomer is named ***S*.**

- The letters *R* or *S* precede the IUPAC name of the molecule. For the enantiomers of butan-2-ol:




A
clockwise
***R* isomer**
(*R*)-butan-2-ol



B
counterclockwise
***S* isomer**
(*S*)-butan-2-ol


Sample Problem 5.2

Label the following compound as *R* or *S*.




Solution

[1] Assign priorities.



[2] Looking toward the lowest priority group (H), trace a circle, 1→2→3.




counterclockwise

Answer: ***S*** isomer


How do you assign *R* or *S* to a molecule when the lowest priority group is not oriented toward the back, on a dashed wedge? You could rotate and flip the molecule until the lowest priority group is in the back, as shown in Figure 5.7; then follow the stepwise procedure for assigning the configuration. Or, if manipulating and visualizing molecules in three dimensions is difficult for you, try the procedure suggested in Sample Problem 5.3.

Figure 5.7
Orienting the lowest priority group in back



- In rotating a molecule about a single bond, the position of three groups changes.
- In flipping a molecule 180°, the position of all four groups changes.

Sample Problem 5.3 Label the following compound as *R* or *S*.




Solution


In this problem, the lowest priority group (H) is oriented in **front** of, not behind, the page. To assign *R* or *S* in this case:

- **Switch** the position of the lowest priority group (H) with the group located *behind* the page (OH).
- Determine *R* or *S* in the usual manner.
- **Reverse the answer.** Because we switched the position of two groups on the stereogenic center to begin with, and there are only two possibilities, the answer is **opposite** to the correct answer.


[1] Assign priorities.



[2] Switch groups 4 and 1.




[3] Trace a circle, 1 → 2 → 3, and reverse the answer.




counterclockwise
It looks like an *S* isomer, but we must reverse the answer because we switched groups 1 and 4, *S* → *R*.

Answer: ***R*** isomer

Problem 5.13 Label each compound as *R* or *S*.



Problem 5.14 Draw both enantiomers of clopidogrel (trade name Plavix), a drug given to prevent the formation of blood clots in persons who have a history of stroke or coronary artery disease. Plavix is sold as a single enantiomer with the *S* configuration. Which enantiomer is Plavix?




clopidogrel



Lisinopril (trade name Zestril) is an ACE inhibitor, a drug that lowers blood pressure by decreasing the amount of angiotensin in the blood. Angiotensin is a polyamide that narrows blood vessels, thus increasing blood pressure.

Problem 5.15 (a) Locate the stereogenic centers in the ball-and-stick model of lisinopril, a drug used to treat high blood pressure. (b) Label each stereogenic center as *R* or *S*.



lisinopril

5.7 Diastereomers

We have now seen many examples of compounds containing one tetrahedral stereogenic center. The situation is more complex for compounds with two stereogenic centers, because more stereoisomers are possible. Moreover, a molecule with two or more stereogenic centers ***may* or *may not* be chiral.**


- **For *n* stereogenic centers, the maximum number of stereoisomers is 2^n.**

- When $n = 1$, $2^1 = 2$. With one stereogenic center there are always two stereoisomers and they are **enantiomers.**
- When $n = 2$, $2^2 = 4$. With two stereogenic centers, the maximum number of stereoisomers is four, although sometimes there are *fewer* than four.

Problem 5.16 What is the maximum number of stereoisomers possible for a compound with: (a) three stereogenic centers; (b) eight stereogenic centers?


Let's illustrate a stepwise procedure for finding all possible stereoisomers using 2,3-dibromopentane. Because 2,3-dibromopentane has two stereogenic centers, the maximum number of stereoisomers is four.

In testing to see if one compound is superimposable on another, rotate atoms and flip the entire molecule, but **do not break any bonds.**




How To Find and Draw All Possible Stereoisomers for a Compound with Two Stereogenic Centers

Step [1] **Draw one stereoisomer by arbitrarily arranging substituents around the stereogenic centers. Then draw its mirror image.**



- Arbitrarily add the H, Br, CH_3, and CH_2CH_3 groups to the stereogenic centers, forming **A.** Then draw the mirror image **B** so that substituents in **B** are a reflection of the substituents in **A.**
- Determine whether **A** and **B** are superimposable by flipping or rotating one molecule to see if all the atoms align.
- If you have drawn the compound and the mirror image in the described manner, you only have to do two operations to see if the atoms align. Place **B** directly on top of **A** (either in your mind or use models); and, rotate **B** 180° and place it on top of **A** to see if the atoms align.




- In this case, the atoms of **A** and **B** do not align, making **A** and **B** nonsuperimposable mirror images—**enantiomers. A** and **B** are two of the four possible stereoisomers for 2,3-dibromopentane.

—*Continued*


How To, continued . . .

Step [2] **Draw a third possible stereoisomer by switching the positions of any two groups on *one* stereogenic center only. Then draw its mirror image.**

- Switching the positions of H and Br (or any two groups) on one stereogenic center of either **A** or **B** forms a new stereoisomer (labeled **C** in this example), which is different from both **A** and **B.** Then draw the mirror image of **C,** labeled **D. C** and **D** are nonsuperimposable mirror images—**enantiomers.** We have now drawn four stereoisomers for 2,3-dibromopentane, the maximum number possible.




With models…



There are only two types of stereoisomers: ***Enantiomers*** **are stereoisomers that *are* mirror images. *Diastereomers* are stereoisomers that are *not* mirror images.**

There are four stereoisomers for 2,3-dibromopentane: enantiomers **A** and **B,** and enantiomers **C** and **D.** What is the relationship between two stereoisomers like **A** and **C? A** and **C** represent the second class of stereoisomers, called **diastereomers.** ***Diastereomers*** **are stereoisomers that are *not* mirror images of each other. A** and **B** are diastereomers of **C** and **D,** and vice versa. Figure 5.8 summarizes the relationships between the stereoisomers of 2,3-dibromopentane.

Problem 5.17 Label the two stereogenic centers in each compound and draw all possible stereoisomers.



Problem 5.18 Compounds **E** and **F** are two isomers of 2,3-dibromopentane drawn in staggered conformations. Which compounds (**A**–**D**) in Figure 5.8 are identical to **E** and **F?**






Figure 5.8
The four stereoisomers of 2,3-dibromopentane



- Pairs of enantiomers: **A** and **B; C** and **D.**
- Pairs of diastereomers: **A** and **C; A** and **D; B** and **C; B** and **D.**


5.8 Meso Compounds

Whereas 2,3-dibromopentane has two stereogenic centers and the maximum of four stereoisomers, **2,3-dibromobutane** has two stereogenic centers but fewer than the maximum number of stereoisomers.




2,3-dibromobutane
= stereogenic center

To find and draw all the stereoisomers of 2,3-dibromobutane, follow the same stepwise procedure outlined in Section 5.7. Arbitrarily add the H, Br, and CH_3 groups to the stereogenic centers, forming one stereoisomer **A,** and then draw its mirror image **B. A** and **B** are nonsuperimposable mirror images—**enantiomers.**




enantiomers

To find the other two stereoisomers (if they exist), switch the position of two groups on *one* stereogenic center of *one* enantiomer only. In this case, switching the positions of H and Br on one stereogenic center of **A** forms **C,** which is different from both **A** and **B** and is thus a new stereoisomer.




identical
C = D



However, the mirror image of **C,** labeled **D,** is superimposable on **C,** so **C** and **D** are *identical.* Thus, **C** is achiral, even though it has two stereogenic centers. **C** is a **meso compound.**

- A *meso compound* is an achiral compound that contains tetrahedral stereogenic centers.


C contains a **plane of symmetry. Meso compounds generally have a plane of symmetry,** so they possess two identical halves.



two identical halves

Because one stereoisomer of 2,3-dibromobutane is superimposable on its mirror image, there are only three stereoisomers and not four, as summarized in Figure 5.9.


Figure 5.9
The three stereoisomers of 2,3-dibromobutane




A and **B** are diastereomers of **C.**

- Pair of enantiomers: **A** and **B.**
- Pairs of diastereomers: **A** and **C; B** and **C.**


Problem 5.19 Draw all the possible stereoisomers for each compound and label pairs of enantiomers and diastereomers.



Problem 5.20 Which compounds are meso compounds?




Problem 5.21 Draw a meso compound for each of the following molecules.



5.9 *R* and *S* Assignments in Compounds with Two or More Stereogenic Centers

When a compound has more than one stereogenic center, the *R* or *S* configuration must be assigned to each of them. In the stereoisomer of 2,3-dibromopentane drawn here, C2 has the *S* configuration and C3 has the *R,* so the complete name of the compound is (2*S*,3*R*)-2,3-dibromopentane.



(2*S*,3*R*)-2,3-dibromopentane

R,S configurations can be used to determine whether two compounds are identical, enantiomers, or diastereomers.

- Identical compounds have the *same R,S* designations at every tetrahedral stereogenic center.
- Enantiomers have exactly *opposite R,S* designations.
- Diastereomers have the *same R,S* designation for at least one stereogenic center and the *opposite* for at least one of the other stereogenic centers.


For example, if a compound has two stereogenic centers, both with the *R* configuration, then its enantiomer is *S,S* and the diastereomers are either *R,S* or *S,R*.

Problem 5.22 If the two stereogenic centers of a compound are *R,S* in configuration, what are the *R,S* assignments for its enantiomer and two diastereomers?


Problem 5.23 Without drawing out the structures, label each pair of compounds as enantiomers or diastereomers.

a. (2*R*,3*S*)-hexane-2,3-diol and (2*R*,3*R*)-hexane-2,3-diol
b. (2*R*,3*R*)-hexane-2,3-diol and (2*S*,3*S*)-hexane-2,3-diol
c. (2*R*,3*S*,4*R*)-hexane-2,3,4-triol and (2*S*,3*R*,4*R*)-hexane-2,3,4-triol


Problem 5.24 (a) Label the four stereogenic centers in sorbitol as *R* or *S*. (b) How are sorbitol and **A** related? (c) How are sorbitol and **B** related?



sorbitol



A




B



Sorbitol (Problem 5.24) occurs naturally in some berries and fruits. It is used as a substitute sweetener in sugar-free—that is, sucrose-free—candy and gum.

5.10 Disubstituted Cycloalkanes


Let us now turn our attention to disubstituted cycloalkanes, and draw all possible stereoisomers for **1,3-dibromocyclopentane.** Because 1,3-dibromocyclopentane has two stereogenic centers (labeled in blue), it has a maximum of four stereoisomers.




1,3-dibromocyclopentane

In determining chirality in substituted cycloalkanes, always draw the rings as **flat polygons.** This is especially true for cyclohexane derivatives, where having two chair forms that interconvert can make analysis especially difficult.

To draw all possible stereoisomers, remember that a disubstituted cycloalkane can have two substituents on the same side of the ring (**cis isomer,** labeled **A**) or on opposite sides of the ring (**trans isomer,** labeled **B**). These compounds are **stereoisomers but not mirror images of each other,** making them **diastereomers. A** and **B** are two of the four possible stereoisomers.



A
cis isomer




B
trans isomer

A and **B** are diastereomers.


cis-1,3-Dibromocyclopentane contains a plane of symmetry.

plane of symmetry




two identical halves


To find the other two stereoisomers (if they exist), draw the mirror image of each compound and determine whether the compound and its mirror image are superimposable.



=




A
cis isomer




identical to **A**

=



- The cis isomer is superimposable on its mirror image, making them *identical.* Thus, **A** is an **achiral meso compound.**




B and **C** are enantiomers.


- The trans isomer **B** is *not* superimposable on its mirror image, labeled **C,** making **B** and **C** different compounds. Thus, **B** and **C** are **enantiomers.**

Because one stereoisomer of 1,3-dibromocyclopentane is superimposable on its mirror image, there are only three stereoisomers, not four. **A** is an achiral meso compound and **B** and **C** are a pair of chiral enantiomers. **A** and **B** are diastereomers, as are **A** and **C.**

Problem 5.25 Which of the following cyclic molecules are *meso* compounds?



Problem 5.26 Draw all possible stereoisomers for each compound. Label pairs of enantiomers and diastereomers.



5.11 Isomers—A Summary

Before moving on to other aspects of stereochemistry, take the time to review Figures 5.10 and 5.11. Keep in mind the following facts, and use Figure 5.10 to summarize the types of isomers.

- **There are two major classes of isomers: constitutional isomers and stereoisomers.**
- **There are only two kinds of stereoisomers: enantiomers and diastereomers.**

Then, to determine the relationship between two nonidentical molecules, refer to the flowchart in Figure 5.11.

Figure 5.10
Summary—Types of isomers







Figure 5.11
Determining the relationship between two nonidentical molecules






Problem 5.27 State how each pair of compounds is related. Are they enantiomers, diastereomers, constitutional isomers, or identical?


a. 


and 

c. 

and 

b. 
and 

d. 

and 

5.12 Physical Properties of Stereoisomers


Recall from Section 5.2 that constitutional isomers have different physical and chemical properties. How, then, do the physical and chemical properties of enantiomers compare?

- **The chemical and physical properties of two enantiomers are *identical* except in their interaction with *chiral* substances.**

5.12A Optical Activity


Two enantiomers have identical physical properties—melting point, boiling point, solubility—except for how they interact with plane-polarized light.

What is plane-polarized light? Ordinary light consists of electromagnetic waves that oscillate in all planes perpendicular to the direction in which the light travels. Passing light through a polarizer allows light in only one plane to come through, resulting in **plane-polarized light** (or simply **polarized light**). Plane-polarized light has an electric vector that oscillates in a single plane.




A **polarimeter** is an instrument that allows plane-polarized light to travel through a sample tube containing an organic compound. After the light exits the sample tube, an analyzer slit is rotated to determine the direction of the plane of the exiting polarized light.

With **achiral compounds,** the light exits the sample tube *unchanged,* and the plane of the polarized light is in the same position it was before entering the sample tube. **A compound that does not change the plane of polarized light is said to be *optically inactive.***




With **chiral compounds,** the plane of the polarized light is rotated through an angle α. The angle α, measured in degrees (°), is called the **observed rotation. A compound that rotates the plane of polarized light is said to be *optically active.***



The achiral compound CH_2BrCl is optically inactive, whereas a single enantiomer of CHBrClF, a chiral compound, is optically active.

The rotation of polarized light can be in the **clockwise** or **counterclockwise** direction.


- If the rotation is *clockwise* (to the right from the noon position), the compound is called *dextrorotatory.* The rotation is labeled *d* or (+).
- If the rotation is *counterclockwise* (to the left from noon), the compound is called *levorotatory.* The rotation is labeled *l* or (–).



(*S*)-(–)-glyceraldehyde

No relationship exists between the *R* and *S* prefixes that designate configuration and the (+) and (–) designations indicating optical rotation. For example, the *S* enantiomer of lactic acid is dextrorotatory (+), whereas the *S* enantiomer of glyceraldehyde is levorotatory (–).

How does the rotation of two enantiomers compare?



(*S*)-(+)-lactic acid

- Two enantiomers rotate plane-polarized light *to an equal extent* but in the *opposite* direction.

Thus, if enantiomer **A** rotates polarized light +5°, then the same concentration of enantiomer **B** rotates it –5°.

5.12B Racemic Mixtures

What is the observed rotation of an equal amount of two enantiomers? Because **two enantiomers rotate plane-polarized light to an equal extent but in opposite directions, the rotations cancel,** and no rotation is observed.

- An equal amount of two enantiomers is called a *racemic mixture* or a *racemate.* A racemic mixture is optically inactive.

Besides optical rotation, other physical properties of a racemate are not readily predicted. The melting point and boiling point of a racemic mixture are not necessarily the same as either pure enantiomer, and this fact is not easily explained. The physical properties of two enantiomers and their racemic mixture are summarized in Table 5.1.

Table 5.1 The Physical Properties of Enantiomers **A** and **B** Compared

Property	A alone	B alone	Racemic A + B
Melting point	identical to **B**	identical to **A**	may be different from **A** and **B**
Boiling point	identical to **B**	identical to **A**	may be different from **A** and **B**
Optical rotation	equal in magnitude but opposite in sign to **B**	equal in magnitude but opposite in sign to **A**	0°

5.12C Specific Rotation

The observed rotation depends on the number of chiral molecules that interact with polarized light. This in turn depends on the concentration of the sample and the length of the sample tube. To standardize optical rotation data, the quantity **specific rotation** ([α]) is defined using a specific sample tube length (usually 1 dm), concentration, temperature (25 °C), and wavelength (589 nm, the D line emitted by a sodium lamp).


$$\textbf{specific rotation} = [\alpha] = \frac{\alpha}{l \times c}$$

α = observed rotation (°)
l = length of sample tube (dm)
c = concentration (g/mL)

[dm = decimeter
1 dm = 10 cm]

Specific rotations are physical constants just like melting points or boiling points, and are reported in chemical reference books for a wide variety of compounds.

Problem 5.28 The amino acid (*S*)-alanine has the physical characteristics listed under the structure.



(*S*)-alanine
[α] = +8.5
mp = 297 °C

a. What is the melting point of (*R*)-alanine?
b. How does the melting point of a racemic mixture of (*R*)- and (*S*)-alanine compare to the melting point of (*S*)-alanine?
c. What is the specific rotation of (*R*)-alanine, recorded under the same conditions as the reported rotation of (*S*)-alanine?
d. What is the optical rotation of a racemic mixture of (*R*)- and (*S*)-alanine?
e. Label each of the following as optically active or inactive: a solution of pure (*S*)-alanine; an equal mixture of (*R*)- and (*S*)-alanine; a solution that contains 75% (*S*)- and 25% (*R*)-alanine.

Problem 5.29 A natural product was isolated in the laboratory, and its observed rotation was +10° when measured in a 1 dm sample tube containing 1.0 g of compound in 10 mL of water. What is the specific rotation of this compound?

5.12D Enantiomeric Excess

Sometimes in the laboratory we have neither a pure enantiomer nor a racemic mixture, but rather a mixture of two enantiomers in which one enantiomer is present in excess of the other. The **enantiomeric excess (*ee*),** also called the **optical purity,** tells how much more there is of one enantiomer.

- **Enantiomeric excess = *ee* = % of one enantiomer – % of the other enantiomer.**

Enantiomeric excess tells how much one enantiomer is present in excess of the racemic mixture. For example, if a mixture contains 75% of one enantiomer and 25% of the other, the enantiomeric excess is 75% – 25% = 50%. There is a 50% excess of one enantiomer over the racemic mixture.

Problem 5.30 What is the *ee* for each of the following mixtures of enantiomers **A** and **B?**

a. 95% **A** and 5% **B** b. 85% **A** and 15% **B**

Knowing the *ee* of a mixture makes it possible to calculate the amount of each enantiomer present, as shown in Sample Problem 5.4.

Sample Problem 5.4 If the enantiomeric excess is 95%, how much of each enantiomer is present?

Solution

Label the two enantiomers **A** and **B** and assume that **A** is in excess. A 95% *ee* means that the solution contains an excess of 95% of **A,** and 5% of the racemic mixture of **A** and **B.** Because a racemic mixture is an equal amount of both enantiomers, it has 2.5% of **A** and 2.5% of **B.**

- Total amount of **A** = 95% + 2.5% = 97.5%
- Total amount of **B** = 2.5% (or 100% – 97.5%)

Problem 5.31 For the given *ee* values, calculate the percentage of each enantiomer present.

a. 90% *ee* b. 99% *ee* c. 60% *ee*

The enantiomeric excess can also be calculated if two quantities are known—the specific rotation [α] of a mixture and the specific rotation [α] of a pure enantiomer.

$$ee = \frac{[\alpha] \text{ mixture}}{[\alpha] \text{ pure enantiomer}} \times 100\%$$

Sample Problem 5.5 Pure cholesterol has a specific rotation of –32. A sample of cholesterol prepared in the lab had a specific rotation of –16. What is the enantiomeric excess of this sample of cholesterol?


Solution

Calculate the *ee* of the mixture using the given formula.

$$ee = \frac{[\alpha] \text{ mixture}}{[\alpha] \text{ pure enantiomer}} \times 100\% = \frac{-16}{-32} \times 100\% = 50\%\ ee$$

Answer

Problem 5.32 Pure MSG, a common flavor enhancer, exhibits a specific rotation of +24. (a) Calculate the *ee* of a solution whose [α] is +10. (b) If the *ee* of a solution of MSG is 80%, what is [α] for this solution?




MSG
monosodium glutamate

Problem 5.33 (*S*)-Lactic acid has a specific rotation of +3.8. (a) If the *ee* of a solution of lactic acid is 60%, what is [α] for this solution? (b) How much of the dextrorotatory and levorotatory isomers does the solution contain?

5.12E The Physical Properties of Diastereomers

Diastereomers are not mirror images of each other, and as such, **their physical properties are different, including optical rotation.** Figure 5.12 compares the physical properties of the three stereoisomers of tartaric acid, consisting of a meso compound that is a diastereomer of a pair of enantiomers.

Figure 5.12
The physical properties of the three stereoisomers of tartaric acid



- **A** and **B** are enantiomers.
- **A** and **B** are diastereomers of **C.**

Property	A	B	C	A + B (1:1)
melting point (°C)	171	171	146	206
solubility (g/100 mL H_2O)	139	139	125	139
[α]	+13	–13	0	0
R,S designation	*R,R*	*S,S*	*R,S*	—
d,l designation	*d*	*l*	none	*d,l*


- The physical properties of **A** and **B** differ from their diastereomer **C.**
- The physical properties of a racemic mixture of **A** and **B** (last column) can also differ from either enantiomer and diastereomer **C.**
- **C** is an achiral meso compound, so it is optically inactive; [α] = 0.

Whether the physical properties of a set of compounds are the same or different has practical applications in the lab. Physical properties characterize a compound's physical state, and two compounds can usually be separated only if their physical properties are different.

Two enantiomers can be separated by the process of **resolution,** as described in Section 29.3.

- **Because two enantiomers have identical physical properties, they cannot be separated by common physical techniques like distillation.**
- **Diastereomers and constitutional isomers have different physical properties, and therefore they can be separated by common physical techniques.**

Problem 5.34 Compare the physical properties of the three stereoisomers of 1,3-dimethylcyclopentane.



three stereoisomers of 1,3-dimethylcyclopentane

a. How do the boiling points of **A** and **B** compare? What about **A** and **C?**
b. Characterize a solution of each of the following as optically active or optically inactive: pure **A;** pure **B;** pure **C;** an equal mixture of **A** and **B;** an equal mixture of **A** and **C.**
c. A reaction forms a 1:1:1 mixture of **A, B,** and **C.** If this mixture is distilled, how many fractions would be obtained? Which fractions would be optically active and which would be optically inactive?

5.13 Chemical Properties of Enantiomers

When two enantiomers react with an achiral reagent, they react at the same rate, but when they react with a chiral, non-racemic reagent, they react at different rates.

- **Two enantiomers have exactly the same chemical properties except for their reaction with chiral, non-racemic reagents.**

For an everyday analogy, consider what happens when you are handed an achiral object like a pen and a chiral object like a right-handed glove. Your left and right hands are enantiomers, but they can both hold the achiral pen in the same way. With the glove, however, only your right hand can fit inside it, not your left.


We will examine specific reactions of chiral molecules with both chiral and achiral reagents later in this text. Here, we examine two more general applications.

5.13A Chiral Drugs


A living organism is a sea of chiral molecules. Many drugs are chiral, and often they must interact with a chiral receptor or a chiral enzyme to be effective. One enantiomer of a drug may treat a disease whereas its mirror image may be ineffective. Alternatively, one enantiomer may trigger one biochemical response and its mirror image may elicit a totally different response.

Although (*R*)-ibuprofen shows no anti-inflammatory activity itself, it is slowly converted to the *S* enantiomer in vivo.

The drugs ibuprofen and fluoxetine each contain one stereogenic center, and thus exist as a pair of enantiomers, only one of which exhibits biological activity. **(*S*)-Ibuprofen** is the active component of the anti-inflammatory agents Motrin and Advil, and **(*R*)-fluoxetine** is the active component in the antidepressant Prozac.



(*S*)-ibuprofen
anti-inflammatory agent




(*R*)-fluoxetine
antidepressant



(*S*)-Naproxen is the active ingredient in the widely used pain relievers Naprosyn and Aleve.

For more examples of two enantiomers that exhibit very different biochemical properties, see *Journal of Chemical Education*, **1996,** *73,* 481–484.

Changing the orientation of two substituents to form a mirror image can also alter biological activity to produce an undesirable side effect in the other enantiomer. The *S* enantiomer of **naproxen** is an active anti-inflammatory agent, but the *R* enantiomer is a harmful liver toxin.



(*S*)-naproxen
anti-inflammatory agent


(*R*)-naproxen
liver toxin

If a chiral drug could be sold as a single active enantiomer, it should be possible to use smaller doses with fewer side effects. Many chiral drugs continue to be sold as racemic mixtures, however, because it is more difficult and therefore more costly to obtain a single enantiomer. An enantiomer is not easily separated from a racemic mixture because the two enantiomers have the same physical properties. In Chapter 12 we will study a reaction that can form a single active enantiomer, an important development in making chiral drugs more readily available.

5.13B Enantiomers and the Sense of Smell

Research suggests that the odor of a particular molecule is determined more by its shape than by the presence of a particular functional group. For example, hexachloroethane (Cl_3CCCl_3) and cyclooctane have no obvious structural similarities, but they both have a camphor-like odor, a fact attributed to their similar spherical shape. Each molecule binds to spherically shaped olfactory receptors present on the nerve endings in the nasal passage, resulting in similar odors (Figure 5.13).

Figure 5.13
The shape of molecules and the sense of smell




Cyclooctane and other molecules similar in shape bind to a particular olfactory receptor on the nerve cells that lie at the top of the nasal passage. Binding results in a nerve impulse that travels to the brain, which interprets impulses from particular receptors as specific odors.


Because enantiomers interact with chiral smell receptors, some enantiomers have different odors. There are a few well-characterized examples of this phenomenon in nature. For example,

(*S*)-carvone is responsible for the odor of caraway, whereas (*R*)-carvone is responsible for the odor of spearmint.


caraway seeds



(*S*)-carvone




(*R*)-carvone


spearmint leaves

These examples demonstrate that understanding the three-dimensional structure of a molecule is very important in organic chemistry.

Problem 5.35 Like carvone, the two enantiomers of celery ketone smell differently. The *R* enantiomer smells like celery leaves, whereas the *S* enantiomer smells like licorice. Draw each enantiomer and assign its odor.



celery ketone


(*R*)-Celery ketone (Problem 5.35) has an odor reminiscent of celery leaves.

KEY CONCEPTS

Stereochemistry

Isomers Are Different Compounds with the Same Molecular Formula (5.2, 5.11).


[1] **Constitutional isomers**—isomers that differ in the way the atoms are connected to each other. They have:
- different IUPAC names
- the same or different functional groups
- different physical and chemical properties

[2] **Stereoisomers**—isomers that differ only in the way atoms are oriented in space**.** They have the same functional group and the same IUPAC name except for prefixes such as cis, trans, *R,* and *S*.
- **Enantiomers**—stereoisomers that are nonsuperimposable mirror images of each other (5.4).
- **Diastereomers**—stereoisomers that are not mirror images of each other (5.7).


Some Basic Principles

- When a compound and its mirror image are **superimposable,** they are **identical achiral compounds.** When a compound has a plane of symmetry in one conformation, the compound is achiral (5.3).
- When a compound and its mirror image are **not superimposable,** they are **different chiral compounds** called **enantiomers.** A chiral compound has no plane of symmetry in any conformation (5.3).
- A **tetrahedral stereogenic center** is a carbon atom bonded to four different groups (5.4, 5.5).
- For ***n* stereogenic centers,** the maximum number of stereoisomers is $\mathbf{2^n}$ (5.7).


plane of symmetry



no stereogenic centers
achiral



one stereogenic center
chiral



two stereogenic centers
chiral



two stereogenic centers
achiral

Optical Activity Is the Ability of a Compound to Rotate Plane-Polarized Light (5.12).

- An optically active solution contains a chiral compound.
- An optically inactive solution contains one of the following:
  - an achiral compound with no stereogenic centers
  - a meso compound—an achiral compound with two or more stereogenic centers
  - a racemic mixture—an equal amount of two enantiomers

The Prefixes *R* and *S* Compared with *d* and *l*

The prefixes *R* and *S* are labels used in nomenclature. Rules on assigning *R,S* are found in Section 5.6.

- An enantiomer has every stereogenic center opposite in configuration. If a compound with two stereogenic centers has the *R,R* configuration, its enantiomer has the *S,S* configuration.
- A diastereomer of this same compound has either the *R,S* or *S,R* configuration; one stereogenic center has the same configuration and one is opposite.

The prefixes *d* (or +) and *l* (or –) tell the direction a compound rotates plane-polarized light (5.12).

- Dextrorotatory (*d* or +) compounds rotate polarized light clockwise.
- Levorotatory (*l* or –) compounds rotate polarized light counterclockwise.
- There is no relation between whether a compound is *R* or *S* and whether it is *d* or *l*.

The Physical Properties of Isomers Compared (5.12)

Type of isomer	Physical properties
Constitutional isomers	Different
Enantiomers	Identical except for the direction polarized light is rotated
Diastereomers	Different
Racemic mixture	Possibly different from either enantiomer

Equations

- Specific rotation (5.12C):

$$\text{specific rotation} = [\alpha] = \frac{\alpha}{l \times c}$$

α = observed rotation (°)
l = length of sample tube (dm)
c = concentration (g/mL)

[dm = decimeter, 1 dm = 10 cm]


- Enantiomeric excess (5.12D):

$$ee = \%\text{ of one enantiomer} - \%\text{ of the other enantiomer}$$
$$= \frac{[\alpha]\text{ mixture}}{[\alpha]\text{ pure enantiomer}} \times 100\%$$

PROBLEMS

Problems Using Three-Dimensional Models

5.36 (a) Locate the stereogenic centers in the ball-and-stick model of ezetimibe (trade name Zetia), a cholesterol-lowering drug. (b) Label each stereogenic center as *R* or *S*.



ezetimibe


5.37 Consider the ball-and-stick models **A–D.** How is each pair of compounds related: (a) **A** and **B;** (b) **A** and **C;** (c) **A** and **D;** (d) **C** and **D?** Choose from identical molecules, enantiomers, or diastereomers.



A **B** **C** **D**


Constitutional Isomers versus Stereoisomers

5.38 Label each pair of compounds as constitutional isomers, stereoisomers, or not isomers of each other.




Mirror Images and Chirality

5.39 Label each compound as chiral or achiral.




5.40 Determine if each compound is identical to or an enantiomer of **A.**




A

5.41 Indicate a plane of symmetry for each molecule that contains one. A molecule may require rotation around a carbon–carbon bond to see the plane of symmetry.




Finding and Drawing Stereogenic Centers


5.42 Locate the tetrahedral stereogenic center(s) in each compound. A molecule may have one or more stereogenic centers.




5.43 Locate the stereogenic centers in each drug.

a. 

amoxicillin
(an antibiotic)


b. 

norethindrone
(oral contraceptive component)


c. 

heroin
(an opiate)

5.44 Draw both enantiomers for each biologically active compound.

a. 

amphetamine
(a powerful central nervous stimulant)

b. 

ketoprofen
(analgesic and anti-inflammatory agent)

Nomenclature


5.45 Which group in each pair is assigned the higher priority in *R,S* nomenclature?


a. $-CD_3$, $-CH_3$
b. $-CH(CH_3)_2$, $-CH_2OH$
c. $-CH_2Cl$, $-CH_2CH_2CH_2Br$
d. $-CH_2NH_2$, $-NHCH_3$


5.46 Rank the following groups in order of decreasing priority.


a. $-F$, $-NH_2$, $-CH_3$, $-OH$
b. $-CH_3$, $-CH_2CH_3$, $-CH_2CH_2CH_3$, $-(CH_2)_3CH_3$
c. $-NH_2$, $-CH_2NH_2$, $-CH_3$, $-CH_2NHCH_3$
d. $-COOH$, $-CH_2OH$, $-H$, $-CHO$
e. $-Cl$, $-CH_3$, $-SH$, $-OH$
f. $-C\equiv CH$, $-CH(CH_3)_2$, $-CH_2CH_3$, $-CH=CH_2$


5.47 Label each stereogenic center as *R* or *S*.


a. 


c. 


e. 

g. 


b. 


d. 

f. 


h. 

5.48 Locate the stereogenic centers in each Newman projection and label each center as *R* or *S*.

a. 


b. 

5.49 Draw the structure of (*S,S*)-ethambutol, a drug used to treat tuberculosis that is 10 times more potent than any of its other stereoisomers.



ethambutol

5.50 Draw the structure of (*S*)-citalopram, a drug used to treat depression and anxiety that is much more potent than its *R* enantiomer.





citalopram


5.51 Draw the structure for each compound.

a. (*R*)-3-methylhexane
b. (4*R*,5*S*)-4,5-diethyloctane
c. (3*R*,5*S*,6*R*)-5-ethyl-3,6-dimethylnonane
d. (3*S*,6*S*)-6-isopropyl-3-methyldecane


5.52 Give the IUPAC name for each compound, including the *R,S* designation for each stereogenic center.

a. 

b. 

c. 


5.53 Label the stereogenic centers in paclitaxel, the chapter-opening anticancer drug, as *R* or *S*.





paclitaxel

Compounds with More Than One Stereogenic Center


5.54 What is the maximum number of stereoisomers possible for each compound?

a. 

b. 

c. 


5.55 The shrub ma huang (Section 5.4A) contains two biologically active stereoisomers—ephedrine and pseudoephedrine—with two stereogenic centers as shown in the given structure. Ephedrine is one component of a once popular combination drug used by body builders to increase energy and alertness, while pseudoephedrine is a nasal decongestant.





isolated from ma huang


a. Draw the structure of naturally occurring (–)-ephedrine, which has the 1*R*,2*S* configuration.
b. Draw the structure of naturally occurring (+)-pseudoephedrine, which has the 1*S*,2*S* configuration.
c. How are ephedrine and pseudoephedrine related?
d. Draw all other stereoisomers of (–)-ephedrine and (+)-pseudoephedrine and give the *R,S* designation for all stereogenic centers.
e. How is each compound drawn in part (d) related to (–)-ephedrine?

5.56 Draw all possible stereoisomers for each compound. Label pairs of enantiomers and diastereomers. Label any meso compound.


a. 


b. 


c. 

d. 

5.57 Draw the enantiomer and a diastereomer for each compound.


a. 

b. 

c. 

5.58 Draw all possible constitutional and stereoisomers for a compound of molecular formula C_6H_{12} having a cyclobutane ring and two methyl groups as substituents. Label each compound as chiral or achiral.

5.59 Explain each statement by referring to compounds **A–E.**




A **B** **C** **D** **E**

a. **A** has a mirror image but no enantiomer.
b. **B** has an enantiomer and no diastereomer.
c. **C** has both an enantiomer and a diastereomer.
d. **D** has a diastereomer but no enantiomer.
e. **E** has a diastereomer but no enantiomer.


Comparing Compounds: Enantiomers, Diastereomers, and Constitutional Isomers

5.60 How is each compound related to the simple sugar D-erythrose? Is it an enantiomer, diastereomer, or identical?




D-erythrose

5.61 Consider Newman projections (**A–D**) for four-carbon carbohydrates. How is each pair of compounds related: (a) **A** and **B;** (b) **A** and **C;** (c) **A** and **D;** (d) **C** and **D?** Choose from identical molecules, enantiomers, or diastereomers.





A **B** **C** **D**



5.62 How is compound **A** related to compounds **B–E?** Choose from enantiomers, diastereomers, constitutional isomers, or identical molecules.





A **B** **C** **D** **E**



5.63 How are the compounds in each pair related to each other? Are they identical, enantiomers, diastereomers, constitutional isomers, or not isomers of each other?



a.  and 

b.  and 

c.  and 

d.  and 

e.  and 

f.  and 

Physical Properties of Isomers


5.64 Drawn are four isomeric dimethylcyclopropanes.



A **B** **C** **D**

a. How are the compounds in each pair related (enantiomers, diastereomers, constitutional isomers): **A** and **B; A** and **C; B** and **C; C** and **D?**
b. Label each compound as chiral or achiral.
c. Which compounds, alone, would be optically active?
d. Which compounds have a plane of symmetry?
e. How do the boiling points of the compounds in each pair compare: **A** and **B; B** and **C; C** and **D?**
f. Which of the compounds are meso compounds?
g. Would an equal mixture of compounds **C** and **D** be optically active? What about an equal mixture of **B** and **C?**


5.65 The [α] of pure quinine, an antimalarial drug, is –165.



quinine
(antimalarial drug)

a. Calculate the *ee* of a solution with the following [α] values: –50, –83, and –120.
b. For each *ee,* calculate the percent of each enantiomer present.
c. What is [α] for the enantiomer of quinine?
d. If a solution contains 80% quinine and 20% of its enantiomer, what is the *ee* of the solution?
e. What is [α] for the solution described in part (d)?

5.66 Amygdalin, a compound isolated from the pits of apricots, peaches, and wild cherries, has been used as an unsanctioned anticancer drug both within and outside of the United States. One hydrolysis product formed from amygdalin is mandelic acid, used in treating common skin problems caused by photo-aging and acne.




amygdalin

mandelic acid


a. How many stereogenic centers are present in amygdalin? What is the maximum number of stereoisomers possible?
b. Draw both enantiomers of mandelic acid and label each stereogenic center as *R* or *S*.
c. Pure (*R*)-mandelic acid has a specific rotation of –154. If a sample contains 60% of the *R* isomer and 40% of its enantiomer, what is [α] of this solution?
d. Calculate the *ee* of a solution of mandelic acid having [α] = +50. What is the percentage of each enantiomer present?

General Problems

5.67 Artemisinin and mefloquine are widely used antimalarial drugs. A ball-and-stick model of artemisinin appears on the cover of this text.




artemisinin



mefloquine

a. Locate the stereogenic centers in both drugs.
b. Label each stereogenic center in mefloquine as *R* or *S*.
c. What is the maximum number of stereoisomers possible for artemisinin?
d. How are the N atoms in mefloquine hybridized?
e. Can two molecules of artemisinin intermolecularly hydrogen bond to each other?
f. What product is formed when mefloquine is treated with HCl?

5.68 Saquinavir (trade name Invirase) is a protease inhibitor, used to treat HIV (human immunodeficiency virus).




saquinavir
Trade name Invirase


a. Locate all stereogenic centers in saquinavir, and label each stereogenic center as *R* or *S*.
b. Draw the enantiomer of saquinavir.
c. Draw a diastereomer of saquinavir.
d. Draw a constitutional isomer that contains at least one different functional group.

Challenge Problems


5.69 A limited number of chiral compounds having no stereogenic centers exist. For example, although **A** is achiral, constitutional isomer **B** is chiral. Make models and explain this observation. Compounds containing two double bonds that share a single carbon atom are called *allenes.* Locate the allene in the antibiotic mycomycin and decide whether mycomycin is chiral or achiral.



achiral
A




chiral
B



mycomycin

5.70 A sulfonium ion (R_3S^+) is a stereogenic center if three different alkyl groups are bonded to sulfur because sulfur is surrounded by four different groups, including its lone pair. In assigning an *R* or *S* designation to sulfur, the lone pair is always assigned the lowest priority (4). SAM, *S*-adenosylmethionine, is a biologically active sulfonium ion that we will learn about in Section 7.16. Locate all the stereogenic centers in SAM, and assign an *R,S* designation to each center.




SAM

5.71 a. Locate all the tetrahedral stereogenic centers in discodermolide, a tumor inhibitor isolated from the Caribbean marine sponge *Discodermia dissoluta.*


b. Certain carbon–carbon double bonds can also be stereogenic centers. With reference to the definition in Section 5.3, explain how this can occur, and then locate the three additional stereogenic centers in discodermolide.

c. Considering all stereogenic centers, what is the maximum number of stereoisomers possible for discodermolide?



discodermolide

5.72 Label each compound as chiral or achiral. Compounds that contain a single carbon common to two rings are called spiro compounds. Because carbon is tetrahedral, the two rings are perpendicular to each other.



5.73 An acid–base reaction of (*R*)-*sec*-butylamine with a racemic mixture of 2-phenylpropanoic acid forms two products having different melting points and somewhat different solubilities. Draw the structure of these two products. Assign *R* and *S* to any stereogenic centers in the products. How are the two products related? Choose from enantiomers, diastereomers, constitutional isomers, or not isomers.



2-phenylpropanoic acid (racemic mixture)

(*R*)-*sec*-butylamine

Understanding Organic Reactions

6



6.1 Writing equations for organic reactions
6.2 Kinds of organic reactions
6.3 Bond breaking and bond making
6.4 Bond dissociation energy
6.5 Thermodynamics
6.6 Enthalpy and entropy
6.7 Energy diagrams
6.8 Energy diagram for a two-step reaction mechanism
6.9 Kinetics
6.10 Catalysts
6.11 Enzymes

Glucose, the most abundant simple carbohydrate, is the building block for starch and cellulose and a major sweet-tasting component of honey. Glucose is used as an energy source by most organisms. In humans, when glucose levels are high after a meal is digested, the body stores glucose as glycogen, which is then hydrolyzed when glucose levels fall and energy demands increase. Glucose is transported in the bloodstream and metabolized aerobically to carbon dioxide and water and a great deal of energy. In Chapter 6, we learn about energy changes that accompany chemical reactions.

Why do certain reactions occur when two compounds are mixed together whereas others do not? To answer this question we must learn how and why organic compounds react.

Reactions are at the heart of organic chemistry. An understanding of chemical processes has made possible the conversion of natural substances into new compounds with different, and sometimes superior, properties. Aspirin, ibuprofen, nylon, and polyethylene are all products of chemical reactions between substances derived from petroleum.

Reactions are difficult to learn when each reaction is considered a unique and isolated event. *Avoid this tendency.* Virtually all chemical reactions are woven together by a few basic themes. After we learn the general principles, specific reactions then fit neatly into a general pattern.

In our study of organic reactions we will begin with the functional groups, looking for electron-rich and electron-deficient sites, and bonds that might be broken easily. These reactive sites give us a clue as to the general type of reaction a particular class of compound undergoes. Finally, we will learn about how a reaction occurs. Does it occur in one step or in a series of steps? Understanding the details of an organic reaction allows us to determine when it might be used in preparing interesting and useful organic compounds.

6.1 Writing Equations for Organic Reactions


Like other reactions, equations for organic reactions are usually drawn with a single reaction arrow (→) between the starting material and product, but other conventions make these equations look different from those encountered in general chemistry.

Often the solvent and temperature of a reaction are omitted from chemical equations, to further focus attention on the main substances involved in the reaction.

The **reagent,** the chemical substance with which an organic compound reacts, is sometimes drawn on the left side of the equation with the other reactants. At other times, the reagent is drawn above or below the reaction arrow itself, to focus attention on the organic starting material by itself on the left side. The solvent and temperature of a reaction may be added above or below the arrow. **The symbols "*hν*" and "Δ" are used for reactions that require *light* or *heat,* respectively.** Figure 6.1 presents an organic reaction in different ways.


Most organic reactions take place in a **liquid solvent.** Solvents solubilize key reaction components and serve as heat reservoirs to maintain a given temperature. Chapter 7 presents the two major types of reaction solvents and how they affect substitution reactions.

When two sequential reactions are carried out without drawing any intermediate compound, the steps are usually numbered above or below the reaction arrow. This convention signifies that the first step occurs *before* the second, and the reagents are added *in sequence,* not at the same time.



In this equation only the organic product is drawn on the right side of the arrow. Although the reagent CH_3MgBr contains both Mg and Br, these elements do not appear in the organic product, and they are often omitted on the product side of the equation. These elements have not disappeared. They are part of an inorganic by-product (HOMgBr in this case), and are often of little interest to an organic chemist.

Figure 6.1
Different ways of writing organic reactions



• The reagent (Br_2) can be on the left side or above the arrow.


• Other reaction parameters can be indicated.

6.2 Kinds of Organic Reactions

Like other compounds, organic molecules undergo acid–base and oxidation–reduction reactions, as discussed in Chapters 2 and 4. Organic molecules also undergo **substitution, elimination,** and **addition** reactions.


6.2A Substitution Reactions

- *Substitution* is a reaction in which an atom or a group of atoms is *replaced* by another atom or group of atoms.



Z = H or a heteroatom

In a general substitution reaction, Y *replaces* Z on a carbon atom. **Substitution reactions involve σ bonds: one σ bond breaks and another forms at the same carbon atom.** The most common examples of substitution occur when Z is hydrogen or a heteroatom that is more electronegative than carbon.



6.2B Elimination Reactions

- *Elimination* is a reaction in which elements of the starting material are "lost" and a π bond is formed.




Two σ bonds are broken. A π bond is formed.

In an elimination reaction, two groups X and Y are removed from a starting material. **Two σ bonds are broken, and a π bond is formed between adjacent atoms.** The most common examples of elimination occur when X = H and Y is a heteroatom more electronegative than carbon.




loss of HBr new π bond




loss of H_2O new π bond


6.2C Addition Reactions

- *Addition* is a reaction in which elements are added to a starting material.




In an addition reaction, new groups X and Y are added to a starting material. **A π bond is broken and two σ bonds are formed.**






A summary of the general types of organic reactions is given in Appendix G.

Addition and elimination reactions are exactly opposite. A π bond is *formed* in elimination reactions, whereas a π bond is *broken* in addition reactions.



Problem 6.1


Classify each transformation as substitution, elimination, or addition.






Capsaicin is responsible for the characteristic spicy flavor of jalapeño and habañero peppers.

To determine whether a reaction is a substitution, elimination, or addition with a complex starting material, concentrate on the functional groups that *change*. The conversion of amine **X** and acid chloride **Y** to the naturally occurring compound capsaicin is a substitution reaction, because the N atom of the amine *replaces* the Cl of the acid chloride.



X **Y** capsaicin

Problem 6.2 Classify the conversion of **A** to **B** as a substitution, elimination, or addition. **B** can be converted to the female sex hormone estrone in two steps.



6.3 Bond Breaking and Bond Making

Having now learned how to write and identify some common kinds of organic reactions, we can turn to a discussion of **reaction mechanism.**

- A *reaction mechanism* is a detailed description of how bonds are broken and formed as a starting material is converted to a product.


A reaction mechanism describes the relative order and rate of bond cleavage and formation. It explains all the known facts about a reaction and accounts for all products formed, and it is subject to modification or refinement as new details are discovered.

A reaction can occur either in one step or in a series of steps.

- **A one-step reaction is called a *concerted reaction.*** No matter how many bonds are broken or formed, a starting material is converted *directly* to a product.




- **A stepwise reaction** involves more than one step. A starting material is first converted to an unstable intermediate, called a **reactive intermediate,** which then goes on to form the product.




6.3A Bond Cleavage

Bonds are broken and formed in all chemical reactions. When a bond is broken, the electrons in the bond can be divided **equally** or **unequally** between the two atoms of the bond.

- Breaking a bond by **equally dividing the electrons** between the two atoms in the bond is called **homolysis** or **homolytic cleavage.**



- Breaking a bond by **unequally dividing the electrons** between the two atoms in the bond is called **heterolysis** or **heterolytic cleavage.**



Heterolysis of a bond between **A** and **B** can give either **A** or **B** the two electrons in the bond. When **A** and **B** have different electronegativities, the ***electrons normally end up on the more electronegative atom*.**

Homolysis and heterolysis require energy. Both processes generate reactive intermediates, but the products are different in each case.


- **Homolysis generates uncharged reactive intermediates with *unpaired* electrons.**
- **Heterolysis generates *charged* intermediates.**

Each of these reactive intermediates has a very short lifetime and reacts quickly to form a stable organic product.

6.3B Radicals, Carbocations, and Carbanions

The curved arrow notation first discussed in Section 1.6 works fine for heterolytic bond cleavage because it illustrates the movement of an **electron pair.** For homolytic cleavage, however, one electron moves to one atom in the bond and one electron moves to the other, so a different kind of curved arrow is needed.

- **To illustrate the movement of a single electron, use a half-headed curved arrow, sometimes called a *fishhook.***



- Two **half-headed** curved arrows are needed for two **single** electrons.
- One **full-headed** curved arrows is needed for one electron **pair.**

A full-headed curved arrow ( ) shows the movement of an electron *pair.* **A half-headed curved arrow** ( ) shows the movement of a *single* electron.


Figure 6.2 illustrates homolysis and two different heterolysis reactions for a carbon compound using curved arrows. Three different reactive intermediates are formed.

Homolysis of the C−Z bond generates two uncharged products with unpaired electrons.

- **A reactive intermediate with a single unpaired electron is called a *radical.***

Most radicals are highly unstable because they contain an atom that does not have an octet of electrons. Radicals typically have **no charge. They are intermediates in a group of reactions called *radical reactions,*** which are discussed in detail in Chapter 15.

Figure 6.2
Three reactive intermediates resulting from homolysis and heterolysis of a C−Z bond



- Radicals are intermediates in **radical** reactions.
- **Ionic intermediates** are seen in **polar** reactions.


Heterolysis of the C–Z bond can generate a **carbocation** or a **carbanion.**

- **Giving two electrons to Z and none to carbon generates a positively charged carbon intermediate called a *carbocation.***
- **Giving two electrons to C and none to Z generates a negatively charged carbon species called a *carbanion.***

Both carbocations and carbanions are unstable reactive intermediates: A carbocation contains a carbon atom surrounded by only six electrons. A carbanion has a negative charge on carbon, which is not a very electronegative atom. **Carbocations (electrophiles)** and **carbanions (nucleophiles)** can be intermediates in ***polar reactions*—reactions in which a nucleophile reacts with an electrophile.**

The chemistry of **carbenes,** another type of organic reactive intermediate, is discussed in Section 26.4.


Thus, homolysis and heterolysis generate radicals, carbocations, and carbanions, the three most common reactive intermediates in organic chemistry.



radical	carbocation	carbanion
C is surrounded by seven electrons.	C is surrounded by six electrons.	C has a lone pair.


- **Radicals and carbocations are electrophiles because they contain a carbon with no octet.**
- **Carbanions are nucleophiles because they contain a carbon with a lone pair.**

Problem 6.3 By taking into account electronegativity differences, draw the products formed by heterolysis of the carbon–heteroatom bond in each molecule. Classify the organic reactive intermediate as a carbocation or a carbanion.



6.3C Bond Formation

Like bond cleavage, bond formation occurs in two different ways. Two radicals can each donate **one electron** to form a two-electron bond. Alternatively, two ions with unlike charges can come together, with the negatively charged ion donating **both electrons** to form the resulting two-electron bond. **Bond formation always releases energy.**



6.3D All Kinds of Arrows


Table 6.1 summarizes the many kinds of arrows used in describing organic reactions. Curved arrows are especially important because they explicitly show what electrons are involved in a reaction, how these electrons move in forming and breaking bonds, and if a reaction proceeds via a radical or polar pathway.


A more complete summary of the arrows used in organic chemistry is given in the table Common Abbreviations, Arrows, and Symbols, located on the inside back cover.

Table 6.1 A Summary of Arrow Types in Chemical Reactions

Arrow	Name	Use
	Reaction arrow	Drawn between the starting materials and products in an equation (6.1)
	Double reaction arrows (equilibrium arrows)	Drawn between the starting materials and products in an equilibrium equation (2.2)
	Double-headed arrow	Drawn between resonance structures (1.6)
	Full-headed curved arrow	Shows movement of an electron pair (1.6, 2.2)
	Half-headed curved arrow (fishhook)	Shows movement of a single electron (6.3)

Sample Problem 6.1 Use full-headed or half-headed curved arrows to show the movement of electrons in each equation.


a. 

b. 


Solution

Look for all bonds that are broken or formed. Use full-headed curved arrows for electron pairs and half-headed curved arrows for single electrons.

a. In this reaction, the C – O bond is broken heterolytically. Because only one electron pair is involved, **one full-headed curved arrow** is needed. The electron pair in the C – O bond ends up on O.




b. This reaction involves radicals, so half-headed curved arrows are needed to show the movement of single electrons. One new two-electron bond is formed between H and Cl, and an unpaired electron is left on C. Because a total of three electrons are involved, three half-headed curved arrows are needed.




An electron remains on C.

Two electrons form a bond.

Problem 6.4 Use curved arrows to show the movement of electrons in each equation.




Sample Problem 6.2 Follow the curved arrows and draw the products of the following reaction.




Solution

Three full-headed curved arrows are drawn, so three electron *pairs* take part in the reaction. Arrow **1** shows that a lone pair on $^-$OH forms a new bond to H, forming H_2O. Arrow **2** indicates that the electron pair in the C – H bond forms a carbon–carbon double bond. Arrow **3** shows that the electron pair in the C – Cl bond ends up on Cl, forming Cl^-. After breaking and making bonds, formal charges on the atoms involved in the reaction are adjusted when necessary.




Problem 6.5 Follow the curved arrows and draw the products of the following reaction.



6.4 Bond Dissociation Energy

Bond breaking can be quantified using the bond dissociation energy.

- **The *bond dissociation energy* is the energy needed to homolytically cleave a covalent bond.**



The energy absorbed or released in any reaction, symbolized by ΔH°, is called the **enthalpy change** or **heat of reaction.**

The superscript (°) means that values are determined under standard conditions (pure compounds in their most stable state at 25 °C and 1 atm pressure).

- **When ΔH° is positive (+), energy is absorbed and the reaction is *endothermic.***
- **When ΔH° is negative (–), energy is released and the reaction is *exothermic.***

Additional bond dissociation energies for C – C multiple bonds are given in Table 1.6.

A bond dissociation energy is the $\Delta H°$ for a specific kind of reaction—the homolysis of a covalent bond to form two radicals. Because bond breaking requires energy, **bond dissociation energies are always *positive* numbers,** and homolysis is always **endothermic.** Conversely, **bond formation always *releases* energy,** so this reaction is always **exothermic.** The H – H bond requires +435 kJ/mol to cleave and releases –435 kJ/mol when formed. Table 6.2 contains a representative list of bond dissociation energies for many common bonds.

A table of bond dissociation energies also appears in Appendix C.

$$H{-}H \xrightarrow[\text{endothermic reaction}]{\Delta H = +435 \text{ kJ/mol}} H\cdot + \cdot H$$

$$H\cdot + \cdot H \xrightarrow[\text{exothermic reaction}]{\Delta H = -435 \text{ kJ/mol}} H{-}H$$

Table 6.2 Bond Dissociation Energies for Some Common Bonds [A–B → A• + •B]


Bond	$\Delta H°$ kJ/mol	(kcal/mol)
H–Z bonds		
H–F	569	(136)
H–Cl	431	(103)
H–Br	368	(88)
H–I	297	(71)
H–OH	498	(119)
Z–Z bonds		
H–H	435	(104)
F–F	159	(38)
Cl–Cl	242	(58)
Br–Br	192	(46)
I–I	151	(36)
HO–OH	213	(51)
R–H bonds		
CH_3–H	435	(104)
CH_3CH_2–H	410	(98)
$CH_3CH_2CH_2$–H	410	(98)
$(CH_3)_2CH$–H	397	(95)
$(CH_3)_3C$–H	381	(91)
$CH_2{=}CH$–H	435	(104)
HC≡C–H	523	(125)
$CH_2{=}CHCH_2$–H	364	(87)
C_6H_5–H	460	(110)
$C_6H_5CH_2$–H	356	(85)
R–R bonds		
CH_3–CH_3	368	(88)
CH_3–CH_2CH_3	356	(85)
CH_3–$CH{=}CH_2$	385	(92)
CH_3–C≡CH	489	(117)

Bond	$\Delta H°$ kJ/mol	(kcal/mol)
R–X bonds		
CH_3–F	456	(109)
CH_3–Cl	351	(84)
CH_3–Br	293	(70)
CH_3–I	234	(56)
CH_3CH_2–F	448	(107)
CH_3CH_2–Cl	339	(81)
CH_3CH_2–Br	285	(68)
CH_3CH_2–I	222	(53)
$(CH_3)_2CH$–F	444	(106)
$(CH_3)_2CH$–Cl	335	(80)
$(CH_3)_2CH$–Br	285	(68)
$(CH_3)_2CH$–I	222	(53)
$(CH_3)_3C$–F	444	(106)
$(CH_3)_3C$–Cl	331	(79)
$(CH_3)_3C$–Br	272	(65)
$(CH_3)_3C$–I	209	(50)
R–OH bonds		
CH_3–OH	389	(93)
CH_3CH_2–OH	393	(94)
$CH_3CH_2CH_2$–OH	385	(92)
$(CH_3)_2CH$–OH	401	(96)
$(CH_3)_3C$–OH	401	(96)

Comparing bond dissociation energies is equivalent to comparing **bond strength.**


- The *stronger* the bond, the *higher* its bond dissociation energy.

For example, the H−H bond is stronger than the Cl−Cl bond because its bond dissociation energy is higher [Table 6.2: 435 kJ/mol (H_2) versus 242 kJ/mol (Cl_2)]. The data in Table 6.2 demonstrate that **bond dissociation energies *decrease* down a column of the periodic table as the valence electrons used in bonding are farther from the nucleus.** Bond dissociation energies for a group of methyl–halogen bonds exemplify this trend.



Because bond length increases down a column of the periodic table, bond dissociation energies are a quantitative measure of the general phenomenon noted in Chapter 1—***shorter* bonds are *stronger* bonds.**

Problem 6.6 Which bond in each pair has the higher bond dissociation energy?



Bond dissociation energies are also used to calculate the enthalpy change ($\Delta H°$) in a reaction in which several bonds are broken and formed. **$\Delta H°$ indicates the relative strength of bonds broken and formed in a reaction.**

- When $\Delta H°$ is positive, more energy is needed to break bonds than is released in forming bonds. The bonds broken in the starting material are *stronger* than the bonds formed in the product.

- When $\Delta H°$ is negative, more energy is released in forming bonds than is needed to break bonds. The bonds formed in the product are *stronger* than the bonds broken in the starting material.

To determine the overall $\Delta H°$ for a reaction:

[1] Beginning with a *balanced* equation, add the bond dissociation energies for all bonds broken in the starting materials. This (+) value represents the **energy needed** to break bonds.

[2] Add the bond dissociation energies for all bonds formed in the products. This (–) value represents the **energy released** in forming bonds.

[3] **The overall $\Delta H°$ is the sum in Step [1] *plus* the sum in Step [2].**

$\Delta H°$ overall enthalpy change	=	sum of $\Delta H°$ of bonds broken	+	(–) sum of $\Delta H°$ of bonds formed

Sample Problem 6.3 Use the values in Table 6.2 to determine $\Delta H°$ for the following reaction.



Solution

[1] Bonds broken

	$\Delta H°$ (kJ/mol)
$(CH_3)_3C-Cl$	+331
H–OH	+498
Total	+829 kJ/mol

Energy needed to break bonds.

[2] Bonds formed

	$\Delta H°$ (kJ/mol)
$(CH_3)_3C-OH$	–401
H–Cl	–431
Total	–832 kJ/mol

Energy released in forming bonds.

[3] Overall $\Delta H°$ =

sum in Step [1]
+
sum in Step [2]


+829 kJ/mol
–832 kJ/mol

Answer: –3 kJ/mol


Because $\Delta H°$ is a negative value, this reaction is **exothermic** and energy is released. **The bonds broken in the starting material are weaker than the bonds formed in the product.**


Problem 6.7 Use the values in Table 6.2 to calculate $\Delta H°$ for each reaction. Classify each reaction as endothermic or exothermic.

a. 

b. 

The oxidation of both isooctane and glucose, the molecule that introduced Chapter 6, forms CO_2 and H_2O.







isooctane

+ (25/2) O_2 ⟶ 8 CO_2 + 9 H_2O

$\Delta H°$ = –5447 kJ/mol

Energy is released.





glucose

+ 6 O_2 ⟶ 6 CO_2 + 6 H_2O

$\Delta H°$ = –2872 kJ/mol

$\Delta H°$ is negative for both oxidations, so both reactions are exothermic. **Both isooctane and glucose release energy on oxidation because the bonds in the products are stronger than the bonds in the reactants.**

Bond dissociation energies have two important limitations. They present *overall* energy changes only. They reveal nothing about the reaction mechanism or how fast a reaction proceeds. Moreover, bond dissociation energies are determined for reactions in the gas phase, whereas most organic reactions are carried out in a liquid solvent where solvation energy contributes to the overall enthalpy of a reaction. As such, bond dissociation energies are imperfect indicators of energy changes in a reaction. Despite these limitations, using bond dissociation energies to calculate $\Delta H°$ gives a useful approximation of the energy changes that occur when bonds are broken and formed in a reaction.

Problem 6.8 Calculate $\Delta H°$ for each oxidation reaction. Each equation is balanced as written; remember to take into account the coefficients in determining the number of bonds broken or formed.
[$\Delta H°$ for O_2 = 497 kJ/mol; $\Delta H°$ for one C=O in CO_2 = 535 kJ/mol]

a. $CH_4 + 2\ O_2 \longrightarrow CO_2 + 2\ H_2O$ b. $2\ CH_3CH_3 + 7\ O_2 \longrightarrow 4\ CO_2 + 6\ H_2O$

6.5 Thermodynamics

For a reaction to be practical, the equilibrium must favor the products, *and* the reaction rate must be fast enough to form them in a reasonable time. These two conditions depend on the **thermodynamics** and the **kinetics** of a reaction, respectively.

- ***Thermodynamics*** **describes energy and equilibrium. How do the *energies* of the reactants and the products compare? What are the relative *amounts* of reactants and products at equilibrium?**

Reaction kinetics are discussed in Section 6.9.

- ***Kinetics*** **describes reaction rates. How *fast* are reactants converted to products?**

6.5A Equilibrium Constant and Free Energy Changes

K_{eq} was first defined in Section 2.3 for acid–base reactions.

The **equilibrium constant, K_{eq},** is a mathematical expression that relates the amount of starting material and product at equilibrium. For example, when starting materials **A** and **B** react to form products **C** and **D,** the equilibrium constant is given by the following expression.

$$\mathbf{A} + \mathbf{B} \rightleftharpoons \mathbf{C} + \mathbf{D}$$

$$K_{eq} = \frac{[\text{products}]}{[\text{starting materials}]} = \frac{[C][D]}{[A][B]}$$

The size of K_{eq} tells about the position of equilibrium; that is, it expresses whether the starting materials or products predominate once equilibrium has been reached.

- When $K_{eq} > 1$, equilibrium favors the products (**C** and **D**) and the equilibrium lies to the right as the equation is written.
- When $K_{eq} < 1$, equilibrium favors the starting materials (**A** and **B**) and the equilibrium lies to the left as the equation is written.

- **For a reaction to be useful, the equilibrium must favor the products, and $K_{eq} > 1$.**

What determines whether equilibrium favors the products in a given reaction? **The position of equilibrium is determined by the relative energies of the reactants and products.** The free energy of a molecule, also called its **Gibbs free energy,** is symbolized by $G°$**.** The **change in free energy** between reactants and products, symbolized by $\Delta G°$, determines whether the starting materials or products are favored at equilibrium.

- **$\Delta G°$ is the overall energy difference between reactants and products.**

$$\Delta G° = G°_{\text{products}} - G°_{\text{reactants}}$$

$G°_{\text{products}}$: free energy of the products; $G°_{\text{reactants}}$: free energy of the reactants

$\Delta G°$ is related to the equilibrium constant K_{eq} by the following equation:

$$\Delta G° = -2.303RT \log K_{eq}$$

R = 8.314 J/(K•mol), the gas constant
T = Kelvin temperature (K)

At 25 °C, $2.303RT = 5.7$ kJ/mol; thus, $\Delta G° = -5.7 \log K_{eq}$.


$K_{eq} > 1$ when $\Delta G° < 0$, and equilibrium favors the products. $K_{eq} < 1$ when $\Delta G° > 0$, and equilibrium favors the starting materials.

Using this expression we can determine the relationship between the equilibrium constant and the free energy change between reactants and products.

- **When $K_{eq} > 1$, $\log K_{eq}$ is positive, making $\Delta G°$ negative, and energy is released. Thus, equilibrium favors the products when the energy of the products is *lower* than the energy of the reactants.**
- **When $K_{eq} < 1$, $\log K_{eq}$ is negative, making $\Delta G°$ positive, and energy is absorbed. Thus, equilibrium favors the reactants when the energy of the products is *higher* than the energy of the reactants.**

Compounds that are lower in energy have increased stability. Thus, **equilibrium favors the products when they are more stable (lower in energy) than the starting materials of a reaction.** This is summarized in Figure 6.3.

Figure 6.3
Summary of the relationship between $\Delta G°$ and K_{eq}



The symbol ~ means *approximately.*

Because $\Delta G°$ depends on the logarithm of K_{eq}, a **small change in energy corresponds to a large difference in the relative amount of starting material and product at equilibrium.** Several values of $\Delta G°$ and K_{eq} are given in Table 6.3. For example, a difference in energy of only ~6 kJ/mol means that there is 10 times as much of the more stable species at equilibrium. A difference in energy of ~18 kJ/mol means that there is essentially only one compound, either starting material or product, at equilibrium.

Table 6.3 Representative Values for $\Delta G°$ and K_{eq} at 25 °C, for a Reaction A → B

$\Delta G°$ (kJ/mol)	K_{eq}	Relative amount of A and B at equilibrium
+18	**10^{-3}**	**Essentially all A (99.9%)**
+12	10^{-2}	100 times as much **A** as **B**
+6	10^{-1}	10 times as much **A** as **B**
0	1	Equal amounts of **A** and **B**
–6	10^{1}	10 times as much **B** as **A**
–12	10^{2}	100 times as much **B** as **A**
–18	**10^{3}**	**Essentially all B (99.9%)**

Increasing [product]

Problem 6.9 a. Which K_{eq} corresponds to a negative value of $\Delta G°$, $K_{eq} = 1000$ or $K_{eq} = .001$?
b. Which K_{eq} corresponds to a lower value of $\Delta G°$, $K_{eq} = 10^{-2}$ or $K_{eq} = 10^{-5}$?


Problem 6.10 Given each of the following values, is the starting material or product favored at equilibrium?
a. $K_{eq} = 5.5$ b. $\Delta G° = 40$ kJ/mol

Problem 6.11 Given each of the following values, is the starting material or product lower in energy?
a. $\Delta G° = 8.0$ kJ/mol b. $K_{eq} = 10$ c. $\Delta G° = -12$ kJ/mol d. $K_{eq} = 10^{-3}$

6.5B Energy Changes and Conformational Analysis

These equations can be used for any process with two states in equilibrium. As an example, monosubstituted cyclohexanes exist as two different chair conformations that rapidly interconvert at room temperature, with the conformation having the substituent in the roomier equatorial position favored (Section 4.13). Knowing the energy difference between the two conformations allows us to calculate the amount of each at equilibrium.

For example, the energy difference between the two chair conformations of phenylcyclohexane is –12.1 kJ/mol, as shown in the accompanying equation. Using the values in Table 6.3, this corresponds to an equilibrium constant of ~100, meaning that there is approximately 100 times more **B** (equatorial phenyl group) than **A** (axial phenyl group) at equilibrium.




$\Delta G° = -2.303RT \log K_{eq}$

$-12.1 \text{ kJ/mol} \longrightarrow K_{eq} \approx 100$

There is ~100 times more **B** than **A** at equilibrium.

Problem 6.12 The equilibrium constant for the conversion of the axial to the equatorial conformation of methoxycyclohexane is 2.7.



a. Given these data, which conformation is present in the larger amount at equilibrium?
b. Is $\Delta G°$ for this process positive or negative?
c. From the values in Table 6.3, approximate the size of $\Delta G°$.

6.6 Enthalpy and Entropy

The **free energy change ($\Delta G°$)** depends on the **enthalpy change ($\Delta H°$)** and the **entropy change ($\Delta S°$).** $\Delta H°$ indicates relative bond strength, but what does $\Delta S°$ measure?

Entropy is a rather intangible concept that comes up again and again in chemistry courses. One way to remember the relation between entropy and disorder is to consider a handful of chopsticks. Dropped on the floor, they are arranged randomly (a state of high entropy). Placed end-to-end in a straight line, they are arranged intentionally (a state of low entropy). The more disordered, random arrangement is favored and easier to achieve energetically.

***Entropy* ($S°$) is a measure of the randomness in a system.** The more freedom of motion or the more disorder present, the higher the entropy. Gas molecules move more freely than liquid molecules and are higher in entropy. Cyclic molecules have more restricted bond rotation than similar acyclic molecules and are lower in entropy.

The *entropy change* ($\Delta S°$) is the change in the amount of disorder between reactants and products. $\Delta S°$ is positive (+) when the products are more disordered than the reactants. $\Delta S°$ is negative (–) when the products are less disordered (more ordered) than the reactants.


- Reactions resulting in an increase in entropy are favored.

$\Delta G°$ is related to $\Delta H°$ and $\Delta S°$ by the following equation:

$$\underset{\text{total energy change}}{\Delta G°} = \underset{\text{change in bonding energy}}{\Delta H°} - \underset{\text{change in disorder}}{T\Delta S°} \qquad [T = \text{Kelvin temperature}]$$

This equation tells us that the total energy change in a reaction is due to two factors: the change in the **bonding energy** and the change in **disorder.** The change in bonding energy can be calculated from bond dissociation energies (Section 6.4). Entropy changes, on the other hand, are more difficult to assess, but they are important when the number of molecules of starting material *differs* from the number of molecules of product in the balanced chemical equation. The entropy of a system also changes when an acyclic molecule is *cyclized* to a cyclic one, or a cyclic molecule is converted to an acyclic one.

For example, **when a single starting material forms two products,** as in the homolytic cleavage of a bond to form two radicals, **entropy increases** and favors formation of the products. In contrast, **entropy decreases when an acyclic compound forms a ring,** because a ring has fewer degrees of freedom. In this case, therefore, entropy does *not* favor formation of the product.



The metabolism of glucose (Section 6.4) is favored by entropy because the number of molecules of products formed (6 CO_2 and 6 H_2O) is greater than the number of molecules of reactants ($C_6H_{12}O_6$ and 6 O_2). Moreover, a cyclic reactant is cleaved to form 12 acyclic product molecules.

In most other reactions that are not carried out at high temperature, the entropy term ($T\Delta S°$) is small compared to the enthalpy term ($\Delta H°$) and it can be neglected. Thus, **we will often approximate the overall free energy change of a reaction by the change in the bonding energy only.** Keep in mind that this is an approximation, but it gives us a starting point from which to decide if the reaction is energetically favorable.

Recall from Section 6.4 that a reaction is endothermic when $\Delta H°$ is positive and exothermic when $\Delta H°$ is negative. A reaction is **endergonic when $\Delta G°$ is positive** and **exergonic when $\Delta G°$ is negative.** $\Delta G°$ is usually approximated by $\Delta H°$ in this text, so the terms endergonic and exergonic are rarely used.

$$\Delta G° \approx \Delta H°$$

According to this approximation:

- **The product is favored when $\Delta H°$ is a *negative* value; that is, the bonds in the product are *stronger* than the bonds in the starting material.**

- **The starting material is favored when $\Delta H°$ is a *positive* value; that is, the bonds in the starting material are *stronger* than the bonds in the product.**

Problem 6.13 Considering each of the following values and neglecting entropy, tell whether the starting material or product is favored at equilibrium: (a) $\Delta H°$ = 80 kJ/mol; (b) $\Delta H°$ = –40 kJ/mol.

Problem 6.14 For a reaction with $\Delta H°$ = 40 kJ/mol, decide which of the following statements is (are) true. Correct any false statement to make it true. (a) The reaction is exothermic; (b) $\Delta G°$ for the reaction is positive; (c) K_{eq} is greater than 1; (d) the bonds in the starting materials are stronger than the bonds in the product; and (e) the product is favored at equilibrium.

Problem 6.15 Answer Problem 6.14 for a reaction with $\Delta H°$ = –20 kJ/mol.


6.7 Energy Diagrams

An **energy diagram** is a schematic representation of the energy changes that take place as reactants are converted to products. An energy diagram indicates how readily a reaction proceeds, how many steps are involved, and how the energies of the reactants, products, and intermediates compare.

Consider a concerted reaction between molecule **A**–**B** with anion **C:**$^-$ to form products **A:**$^-$ and **B**–**C.** If the reaction occurs in a single step, the bond between **A** and **B** is broken *as* the bond between **B** and **C** is formed. Let's assume that the products are lower in energy than the reactants in this hypothetical reaction.



An energy diagram plots **energy on the *y* axis** versus the progress of reaction, often labeled the **reaction coordinate,** on the ***x* axis.** As the starting materials **A**–**B** and **C:**$^-$ approach one another, their electron clouds feel some repulsion, causing an increase in energy, until a maximum value is reached. This unstable energy maximum is called the **transition state.** In the transition state the bond between **A** and **B** is partially broken, and the bond between **B** and **C** is partially formed. Because it is at the top of an energy "hill," **a transition state can never be isolated.**



At the transition state, the bond between **A** and **B** can re-form to regenerate starting material, *or* the bond between **B** and **C** can form to generate product. As the bond forms between **B** and **C** the energy decreases until some stable energy minimum of the products is reached.

- The energy difference between the reactants and products is $\Delta H°$. Because the products are at lower energy than the reactants, this reaction is exothermic and energy is released.

- The energy difference between the transition state and the starting material is called the *energy of activation,* symbolized by E_a.


The *energy of activation* is the minimum amount of energy needed to break bonds in the reactants. It represents an **energy barrier** that must be overcome for a reaction to occur. The size of E_a tells us about the reaction rate.

A slow reaction has a large E_a. A fast reaction has a low E_a.

- The larger the E_a, the greater the amount of energy that is needed to break bonds, and the slower the reaction rate.

How can we draw the structure of the unstable transition state? The structure of the transition state is somewhere in between the structures of the starting material and product. Any bond that is partially broken or formed is drawn with a dashed line. Any atom that gains or loses a charge contains a partial charge in the transition state. Transition states are drawn in brackets, with a superscript double dagger (‡).

In the hypothetical reaction between **A**–**B** and **C:**$^-$ to form **A:**$^-$ and **B**–**C,** the bond between **A** and **B** is partially broken, and the bond between **B** and **C** is partially formed. Because **A** gains a negative charge and **C** loses a charge in the course of the reaction, each atom bears a partial negative charge in the transition state.




Several energy diagrams are drawn in Figure 6.4. For any energy diagram:

- E_a determines the height of the energy barrier.
- $\Delta H°$ determines the relative position of the reactants and products.

Figure 6.4
Some representative energy diagrams


Example [1]

- **Large E_a ⟶ slow reaction**
- **(+) $\Delta H°$ ⟶ endothermic reaction**




Example [3]

- **Low E_a ⟶ fast reaction**
- **(+) $\Delta H°$ ⟶ endothermic reaction**




Example [2]

- **Large E_a ⟶ slow reaction**
- **(–) $\Delta H°$ ⟶ exothermic reaction**



Example [4]


- **Low E_a ⟶ fast reaction**
- **(–) $\Delta H°$ ⟶ exothermic reaction**



The two variables, E_a **and** $\Delta H°$**, are independent** of each other. Two reactions can have identical values for $\Delta H°$ but very different E_a values. For two exothermic reactions with the same negative value of $\Delta H°$ but different E_a values, the reaction with the lower E_a is faster.


Problem 6.16 Draw an energy diagram for a reaction in which the products are higher in energy than the starting materials and E_a is large. Clearly label all of the following on the diagram: the axes, the starting materials, the products, the transition state, $\Delta H°$, and E_a.

Problem 6.17 Draw the structure for the transition state in each reaction.

a. 

b. 

Problem 6.18 Compound **A** can be converted to either **B** or **C.** The energy diagrams for both processes are drawn on the graph below.




a. Label each reaction as endothermic or exothermic.
b. Which reaction is faster?
c. Which reaction generates the product lower in energy?
d. Which points on the graphs correspond to transition states?
e. Label the energy of activation for each reaction.
f. Label the $\Delta H°$ for each reaction.

6.8 Energy Diagram for a Two-Step Reaction Mechanism

Although the hypothetical reaction in Section 6.7 is concerted, many reactions involve more than one step with formation of a reactive intermediate. Consider the same overall reaction, **A−B + C:⁻** to form products **A:⁻ + B−C,** but in this case begin with the assumption that the reaction occurs by a *stepwise* pathway—that is, bond breaking occurs *before* bond making. Once again, assume that the overall process is exothermic.




One possible stepwise mechanism involves heterolysis of the **A−B** bond to form two ions **A:⁻** and **B⁺,** followed by reaction of **B⁺** with anion **C:⁻** to form product **B−C,** as outlined in the accompanying equations. Species **B⁺** is a **reactive intermediate. B⁺** is a product in Step [1] that reacts with **C:⁻** in Step [2].




To draw an energy diagram for a two-step mechanism, we must draw an energy diagram for each step, and then combine them together. Each step has its own energy barrier, with a transition state at the energy maximum.

Step [1] is endothermic because energy is needed to cleave the **A−B** bond, making $\Delta H°$ a positive value and placing the products of Step [1] at higher energy than the starting materials. In the transition state, the **A−B** bond is partially broken.



Step [2] is exothermic because energy is released in forming the **B−C** bond, making $\Delta H°$ a negative value and placing the products of Step [2] at lower energy than the starting materials of Step [2]. In the transition state, the **B−C** bond is partially formed.




The overall process is shown in Figure 6.5 as a single energy diagram that combines both steps. Because the reaction has two steps, there are two transition states, each corresponding to an energy barrier. The transition states are separated by an energy minimum, at which the reactive intermediate $\mathbf{B^+}$ is located. Because we made the assumption that the overall two-step process is exothermic, the overall energy difference between the reactants and products, labeled $\Delta H°_{\text{overall}}$, has a negative value, and the final products are at a lower energy than the starting materials.

The energy barrier for Step [1], labeled E_a[1], is higher than the energy barrier for Step [2], labeled E_a[2]. This is because bond cleavage (Step [1]) is more difficult (requires more energy) than bond formation (Step [2]). A higher energy transition state for Step [1] makes it the slower step of the mechanism.

- **In a multistep mechanism, the step with the highest energy transition state is called the rate-determining step.**


In this reaction, the rate-determining step is Step [1].

Figure 6.5 Complete energy diagram for the two-step conversion of A–B + C:⁻ → A:⁻ + B–C



- **The transition states are located at energy maxima, while the reactive intermediate B^+ is located at an energy minimum.**
- Each step has its own value of ΔH° and E_a.
- The overall energy difference between starting material and products is called $\Delta H^\circ_{overall}$. In this example, the products of the two-step sequence are at lower energy than the starting materials.
- Since Step [1] has the higher energy transition state, it is the **rate-determining step.**

Problem 6.19 Consider the following energy diagram.



a. How many steps are involved in this reaction?
b. Label ΔH° and E_a for each step, and label $\Delta H^\circ_{overall}$.
c. Label each transition state.
d. Which point on the graph corresponds to a reactive intermediate?
e. Which step is rate-determining?
f. Is the overall reaction endothermic or exothermic?

Problem 6.20 Draw an energy diagram for a two-step reaction, A → B → C, where the relative energy of these compounds is C < A < B, and the conversion of B → C is rate-determining.

6.9 Kinetics

We now turn to a more detailed discussion of **reaction rate**—that is, how fast a particular reaction proceeds. **The study of reaction rates is called *kinetics*.**


The rate of chemical processes affects many facets of our lives. Aspirin is an effective anti-inflammatory agent because it rapidly inhibits the synthesis of prostaglandins (Section 19.6). Butter turns rancid with time because its lipids are only slowly oxidized by oxygen in the air to undesirable by-products (Section 15.11). DDT (Section 7.4) is a persistent environmental pollutant because it does not react appreciably with water, oxygen, or any other chemical with which it comes into contact. All of these processes occur at different rates, resulting in beneficial or harmful effects.



Some reactions have a very favorable equilibrium constant (K_{eq} >> 1), but the rate is very slow. Gasoline can be safely handled in the air because its reaction with O_2 is slow unless there is a spark to provide energy to initiate the reaction.

6.9A Energy of Activation

As we learned in Section 6.7, the energy of activation, E_a, is the energy difference between the reactants and the transition state. It is the **energy barrier** that must be exceeded for reactants to be converted to products.



- **The *larger* the E_a, the *slower* the reaction.**



Practically, the effect of temperature on reaction rate is used to an advantage in the kitchen. Food is stored in a cold refrigerator to slow the reactions that cause spoilage.

Concentration and temperature also affect reaction rate.

- **The *higher* the concentration, the *faster* the rate. Increasing concentration increases the number of collisions between reacting molecules, which in turn increases the rate.**

- **The *higher* the temperature, the *faster* the rate. Increasing temperature increases the average kinetic energy of the reacting molecules. Because the kinetic energy of colliding molecules is used for bond cleavage, increasing the average kinetic energy increases the rate.**

The E_a values of most organic reactions are 40–150 kJ/mol. When E_a < 80 kJ/mol, the reaction occurs readily at or below room temperature. When E_a > 80 kJ/mol, higher temperatures are needed. As a rule of thumb, increasing the temperature by 10 °C doubles the reaction rate. Thus, reactions in the lab are often heated to increase their rates so they occur in a reasonable amount of time.

Keep in mind that certain **reaction quantities have *no effect* on reaction rate.**

- **$\Delta G°$, $\Delta H°$, and K_{eq} do *not* determine the rate of a reaction. These quantities indicate the direction of equilibrium and the relative energy of reactants and products.**

Problem 6.21 Which value (if any) corresponds to a faster reaction: (a) E_a = 40 kJ/mol or E_a = 4 kJ/mol; (b) a reaction temperature of 0 °C or a reaction temperature of 25 °C; (c) K_{eq} = 10 or K_{eq} = 100; (d) $\Delta H°$ = –10 kJ/mol or $\Delta H°$ = 10 kJ/mol?

Problem 6.22 For a reaction with K_{eq} = 0.8 and E_a = 80 kJ/mol, decide which of the following statements is (are) true. Correct any false statement to make it true. Ignore entropy considerations. (a) The reaction is faster than a reaction with K_{eq} = 8 and E_a = 80 kJ/mol. (b) The reaction is faster than a reaction with K_{eq} = 0.8 and E_a = 40 kJ/mol. (c) $\Delta G°$ for the reaction is a positive value. (d) The starting materials are lower in energy than the products of the reaction. (e) The reaction is exothermic.

6.9B Rate Equations

The rate of a chemical reaction is determined by measuring the decrease in the concentration of the reactants over time, or the increase in the concentration of the products over time. A **rate law** (or **rate equation**) is an equation that shows the relationship between the rate of a reaction and the concentration of the reactants. A rate law is determined *experimentally,* and it depends on the mechanism of the reaction.

A rate law has two important terms: the **rate constant symbolized by *k*,** and the **concentration of the reactants.** Not all reactant concentrations may appear in the rate equation, as we shall soon see.

$$\text{rate} = k[\text{reactants}]$$

k = the rate constant

A rate constant k and the energy of activation E_a are inversely related. **A high E_a corresponds to a small k.**

A rate constant k is a fundamental characteristic of a reaction. It is a complex mathematical term that takes into account the dependence of a reaction rate on temperature and the energy of activation.

- **Fast reactions have large rate constants.**
- **Slow reactions have small rate constants.**

What concentration terms appear in the rate equation? That depends on the mechanism. For the organic reactions we will encounter:

- **A rate equation contains concentration terms for *all* reactants involved in a *one-step* mechanism.**
- **A rate equation contains concentration terms for *only* the reactants involved in the *rate-determining step* in a multistep reaction.**

In the one-step reaction of **A−B** + **C:**$^-$ to form **A:**$^-$ + **B−C,** *both* reactants appear in the transition state of the only step of the mechanism. The **concentration of *both* reactants affects the reaction rate** and *both* terms appear in the rate equation. This type of reaction involving two reactants is said to be **bimolecular.**



rate = *k* [AB][C:$^-$]

Both reactants are involved in the only step.
Both reactants determine the rate.


sum of the exponents = **2**

Second-order rate equation

The ***order* of a rate equation equals the sum of the exponents of the concentration terms** in the rate equation. In the rate equation for the concerted reaction of **A−B** + **C:**$^-$**,** there are two concentration terms, each with an exponent of one. Thus, the sum of the exponents is two and the **rate equation is *second order*** (the reaction follows second-order kinetics).

Because the rate of the reaction depends on the concentration of both reactants, doubling the concentration of *either* **A−B** or **C:**$^-$ doubles the rate of the reaction. Doubling the concentration of *both* **A−B** and **C:**$^-$ increases the reaction rate by a factor of *four.*

The situation is different in the stepwise conversion of **A−B** + **C:**$^-$ to form **A:**$^-$ + **B−C.** The mechanism shown in Section 6.8 has two steps: a slow step (the **rate-determining** step) in which the **A−B** bond is broken, and a fast step in which the **B−C** bond is formed.



Step [1]
rate-determining

Step [2]

rate = *k* [AB]

only one concentration term

Only AB is involved in the rate-determining step.
Only [AB] determines the rate.

First-order rate equation

In a multistep mechanism, a reaction can occur no faster than its rate-determining step. **Only the concentrations of the reactants in the rate-determining step appear in the rate equation.** In this example, the rate depends on the concentration of **A−B** *only,* because only **A−B** appears in the rate-determining step. A reaction involving only one reactant is said to be **unimolecular.** Because there is only one concentration term (raised to the first power), the **rate equation is *first order*** (the reaction follows first-order kinetics).

Because the rate of the reaction depends on the concentration of only *one* reactant, doubling the concentration of **A−B** doubles the rate of the reaction, but **doubling the concentration of C:⁻ has *no effect* on the reaction rate.**

This might seem like a puzzling result. If **C:⁻** is involved in the reaction, why doesn't it affect the overall rate of the reaction? Not only can you change the concentration of **C:⁻** and not affect the rate, but you also can replace it by a different anion without affecting the rate. How can this be? **C:⁻ is not involved in the slow step of the reaction, so neither its concentration nor its identity affects the reaction rate.**

The following analogy is useful. Let's say three students must make 20 peanut butter and jelly sandwiches for a class field trip. Student (**1**) spreads the peanut butter on the bread. Student (**2**) spreads on the jelly, and student (**3**) cuts the sandwiches in half. Suppose student (**2**) is very slow in spreading the jelly. It doesn't matter how fast students (**1**) and (**3**) are; they can't finish making sandwiches any faster than student (**2**) can add the jelly. Five more students can spread on the peanut butter, or an entirely different individual can replace student (**3**), and this doesn't speed up the process. How fast the sandwiches are made is determined entirely by the rate-determining step—that is, spreading the jelly.

Rate equations provide very important information about the mechanism of a reaction. Rate laws for new reactions with unknown mechanisms are determined by a set of experiments that measure how a reaction's rate changes with concentration. Then, a mechanism is suggested based on which reactants affect the rate.

Problem 6.23 For each rate equation, what effect does the indicated concentration change have on the overall rate of the reaction?


[1] rate = $k[CH_3CH_2Br][^-OH]$


a. tripling the concentration of CH_3CH_2Br only
b. tripling the concentration of ^-OH only
c. tripling the concentration of both CH_3CH_2Br and ^-OH

[2] rate = $k[(CH_3)_3COH]$

a. doubling the concentration of $(CH_3)_3COH$
b. increasing the concentration of $(CH_3)_3COH$ by a factor of 10

Problem 6.24 Write a rate equation for each reaction, given the indicated mechanism.

a. 


b. 

6.10 Catalysts

Some reactions do not occur in a reasonable time unless a **catalyst** is added.

- A *catalyst* is a substance that speeds up the rate of a reaction. A catalyst is recovered unchanged in a reaction, and it does not appear in the product.

Common catalysts in organic reactions are **acids** and **metals.** Two examples are shown with the catalyst drawn in red.




The reaction of acetic acid with ethanol to yield ethyl acetate and water occurs in the presence of an acid catalyst. The acid catalyst is written over or under the arrow to emphasize that it is not part of the starting materials or the products. The details of this reaction are discussed in Chapter 22.

The reaction of cyclohexene with hydrogen to form cyclohexane occurs only in the presence of a metal catalyst such as palladium, platinum, or nickel. The metal provides a surface that binds both the cyclohexene and the hydrogen, and in doing so, facilitates the reaction. We return to this mechanism in Chapter 12.

Catalysts accelerate a reaction by lowering the energy of activation (Figure 6.6). They have no effect on the equilibrium constant, so they do not change the amount of reactant and product at equilibrium. Thus, catalysts affect how *quickly* equilibrium is achieved, but not the relative amounts of reactants and products at equilibrium. If a catalyst is somehow used up in one step of a reaction sequence, it must be regenerated in another step. Because only a small amount of a catalyst is needed relative to starting material, it is said to be present in a **catalytic amount.**

Figure 6.6
The effect of a catalyst on a reaction



- The catalyst lowers the energy of activation, thus **increasing the rate of the catalyzed reaction.**
- The energy of the reactants and products is the same in both the uncatalyzed and catalyzed reactions, so the **position of equilibrium is unaffected.**

Problem 6.25 Identify the catalyst in each equation.

a. $CH_2{=}CH_2 \xrightarrow[H_2SO_4]{H_2O} CH_3CH_2OH$ b. $CH_3Cl \xrightarrow[^-OH]{I^-} CH_3OH + Cl^-$

6.11 Enzymes


The catalysts that synthesize and break down biomolecules in living organisms are governed by the same principles as the acids and metals in organic reactions. The catalysts in living organisms, however, are usually protein molecules called **enzymes.**

- *Enzymes* are biochemical catalysts composed of amino acids held together in a very specific three-dimensional shape.

An enzyme contains a region called its **active site,** which binds an organic reactant, called a **substrate.** When bound, this unit is called the **enzyme–substrate complex,** as shown schematically in Figure 6.7 for the enzyme lactase, the enzyme that binds lactose, the principal carbohydrate in milk. Once bound, the organic substrate undergoes a very specific reaction at an enhanced rate. In this example, lactose is converted into two simpler sugars, glucose and galactose. When individuals lack adequate amounts of lactase, they are unable to digest lactose, causing abdominal cramping and diarrhea.

An enzyme speeds up a biological reaction in a variety of ways. It may hold reactants in the proper conformation to facilitate reaction, or it may provide an acidic site needed for a particular transformation. Once the reaction is completed, the enzyme releases the substrate and it is then able to catalyze another reaction.

Figure 6.7
Lactase, an example of a biological catalyst



The enzyme lactase binds the carbohydrate lactose ($C_{12}H_{22}O_{11}$) in its active site in Step [1]. Lactose then reacts with water to break a bond and form two simpler sugars, galactose and glucose, in Step [2]. This process is the first step in digesting lactose, the principal carbohydrate in milk.

KEY CONCEPTS

Understanding Organic Reactions

Writing Equations for Organic Reactions (6.1)

- Use curved arrows to show the movement of electrons. Full-headed arrows are used for electron pairs and half-headed arrows are used for single electrons.
- Reagents can be drawn either on the left side of an equation or over the reaction arrow. Catalysts are drawn over or under the reaction arrow.

Types of Reactions (6.2)

[1] Substitution



Z = H or a heteroatom

[2] Elimination



Two σ bonds are broken. A π bond is formed.

[3] Addition



A π bond is broken. Two σ bonds are formed.

Important Trends


Values compared	Trend
Bond dissociation energy and **bond strength**	The ***higher*** the bond dissociation energy, the ***stronger*** the bond (6.4).
Energy and **stability**	The ***higher*** the energy, the ***less stable*** the species (6.5A).
E_a and **reaction rate**	The ***larger*** the energy of activation, the ***slower*** the reaction (6.9A).
E_a and **rate constant**	The ***larger*** the energy of activation, the ***smaller*** the rate constant (6.9B).

Reactive Intermediates (6.3)

- Breaking bonds generates reactive intermediates.
- Homolysis generates radicals with unpaired electrons.
- Heterolysis generates ions.

Reactive intermediate	General structure	Reactive feature	Reactivity
Radical		Unpaired electron	Electrophilic
Carbocation		Positive charge; only six electrons around C	Electrophilic
Carbanion		Net negative charge; lone electron pair on C	Nucleophilic

Energy Diagrams (6.7, 6.8)



- E_a determines the rate.
- $\Delta H°$ is the difference in bonding energy between the reactants and products.

Conditions Favoring Product Formation (6.5, 6.6)

Variable	Value	Meaning
K_{eq}	$K_{eq} > 1$	More products than reactants are present at equilibrium.
$\Delta G°$	$\Delta G° < 0$	The free energy of the products is **lower** than the energy of the reactants.
$\Delta H°$	$\Delta H° < 0$	Bonds in the products are **stronger** than bonds in the reactants.
$\Delta S°$	$\Delta S° > 0$	The products are **more disordered** than the reactants.

Equations (6.5, 6.6)

$$\Delta G° = -2.303RT \log K_{eq}$$

K_{eq} depends on the energy difference between reactants and products.

R = 8.314 J/(K•mol), the gas constant
T = Kelvin temperature (K)

$$\Delta G° = \Delta H° - T\Delta S°$$

free energy change | change in bonding energy | change in disorder

T = Kelvin temperature (K)


Factors Affecting Reaction Rate (6.9)


Factor	Effect
Energy of activation	Larger E_a ----→ slower reaction
Concentration	Higher concentration ----→ faster reaction
Temperature	Higher temperature ----→ faster reaction

PROBLEMS

Problems Using Three-Dimensional Models

6.26 Draw the products of homolysis or heterolysis of each indicated bond. Use electronegativity differences to decide on the location of charges in the heterolysis reaction. Classify each carbon reactive intermediate as a radical, carbocation, or carbanion.

a. 

b. 


6.27 Explain why the bond dissociation energy for bond (a) is lower than the bond dissociation energy for bond (b).



$\Delta H°$ = 356 kJ/mol $\Delta H°$ = 385 kJ/mol


Types of Reactions

6.28 Classify each transformation as substitution, elimination, or addition.




Curved Arrows


6.29 Use full-headed or half-headed curved arrows to show the movement of electrons in each reaction.




6.30 Draw the products of each reaction by following the curved arrows.




6.31 (a) Add curved arrows for each step to show how **A** is converted to the epoxy ketone **C.** (b) Classify the conversion of **A** to **C** as a substitution, elimination, or addition. (c) Draw one additional resonance structure for **B.**



6.32 (a) Draw in the curved arrows to show how **A** is converted to **B** in Step [1]. (b) Identify **X,** using the curved arrows drawn for Step [2].




6.33 $PGF_{2\alpha}$ (Section 4.15) is synthesized in cells using a cyclooxygenase enzyme that catalyzes a multistep radical pathway. Two steps in the pathway are depicted in the accompanying equations. (a) Draw in curved arrows to illustrate how **C** is converted to **D** in Step [1]. (b) Identify **Y,** the product of Step [2], using the curved arrows that are drawn on compound **D.**



C D $PGF_{2\alpha}$

Bond Dissociation Energy and Calculating $\Delta H°$

6.34 Rank the indicated bonds in order of increasing bond dissociation energy.




6.35 Calculate $\Delta H°$ for each reaction.

a. $HO\cdot + CH_4 \longrightarrow \cdot CH_3 + H_2O$

b. $CH_3OH + HBr \longrightarrow CH_3Br + H_2O$

6.36 Homolysis of the indicated C – H bond in propene forms a resonance-stabilized radical.



propene

a. Draw the two possible resonance structures for this radical.
b. Use half-headed curved arrows to illustrate how one resonance structure can be converted to the other.
c. Draw a structure for the resonance hybrid.

6.37 As we learned in Chapter 4, propane ($CH_3CH_2CH_3$) has both 1° and 2° hydrogens.

a. Draw the carbon radical formed by homolysis of each type of C – H bond.
b. Use the values in Table 6.2 to determine which C – H bond is stronger.
c. Explain how this information can be used to determine the relative stability of the two radicals formed. Which radical formed from propane is more stable?

Thermodynamics, $\Delta G°$, $\Delta H°$, $\Delta S°$, and K_{eq}

6.38 Given each value, determine whether the starting material or product is favored at equilibrium.


a. $K_{eq} = 0.5$
b. $\Delta G° = -100$ kJ/mol
c. $\Delta H° = 8.0$ kJ/mol
d. $K_{eq} = 16$
e. $\Delta G° = 2.0$ kJ/mol
f. $\Delta H° = 200$ kJ/mol
g. $\Delta S° = 8$ J/(K•mol)
h. $\Delta S° = -8$ J/(K•mol)

6.39 a. Which value corresponds to a negative value of $\Delta G°$: $K_{eq} = 10^{-2}$ or $K_{eq} = 10^{2}$?

b. In a unimolecular reaction with five times as much starting material as product at equilibrium, what is the value of K_{eq}? Is $\Delta G°$ positive or negative?


c. Which value corresponds to a larger K_{eq}: $\Delta G° = -8$ kJ/mol or $\Delta G° = 20$ kJ/mol?


6.40 As we learned in Chapter 4, monosubstituted cyclohexanes exist as an equilibrium mixture of two conformations having either an axial or equatorial substituent. When R = CH_2CH_3, K_{eq} for this process is 23. When R = $C(CH_3)_3$, K_{eq} for this process is 4000.



a. When R = CH_2CH_3, which conformation is present in higher concentration?
b. Which R shows the higher percentage of equatorial conformation at equilibrium?
c. Which R shows the higher percentage of axial conformation at equilibrium?
d. For which R is $\Delta G°$ more negative?
e. How is the size of R related to the amount of axial and equatorial conformations at equilibrium?


6.41 For which of the following reactions is $\Delta S°$ a positive value?

a. 

b. 

Energy Diagrams and Transition States

6.42 Draw the transition state for each reaction.

a. 

b. 

6.43 Draw an energy diagram for each reaction. Label the axes, the starting material, product, transition state, $\Delta H°$, and E_a.

a. a concerted reaction with $\Delta H° = -80$ kJ/mol and $E_a = 16$ kJ/mol

b. a two-step reaction, **A** → **B** → **C,** in which the relative energy of the compounds is **A** < **C** < **B,** and the step **A** → **B** is rate-determining

6.44 Consider the following reaction: $CH_4 + Cl\cdot \rightarrow \cdot CH_3 + HCl$.


a. Use curved arrows to show the movement of electrons.

b. Calculate $\Delta H°$ using the bond dissociation energies in Table 6.2.

c. Draw an energy diagram assuming that $E_a = 16$ kJ/mol.

d. What is E_a for the reverse reaction ($\cdot CH_3 + HCl \rightarrow CH_4 + Cl\cdot$)?

6.45 Consider the following energy diagram for the conversion of **A** → **G.**



a. Which points on the graph correspond to transition states?

b. Which points on the graph correspond to reactive intermediates?


c. How many steps are present in the reaction mechanism?

d. Label each step of the mechanism as endothermic or exothermic.

e. Label the overall reaction as endothermic or exothermic.

6.46 Draw an energy diagram for the Brønsted–Lowry acid–base reaction of CH_3CO_2H with $^-OC(CH_3)_3$ to form $CH_3CO_2^-$ and $(CH_3)_3COH$. Label the axes, starting materials, products, $\Delta H°$, and E_a. Draw the structure of the transition state.

6.47 Consider the following two-step reaction:



a. How many bonds are broken and formed in Step [1]? Would you predict $\Delta H°$ of Step [1] to be positive or negative?

b. How many bonds are broken and formed in Step [2]? Would you predict the $\Delta H°$ of Step [2] to be positive or negative?

c. Which step is rate-determining?

d. Draw the structure for the transition state in both steps of the mechanism.

e. If $\Delta H°_{overall}$ is negative for this two-step reaction, draw an energy diagram illustrating all of the information in parts (a)–(d).

Kinetics and Rate Laws


6.48 Indicate which factors affect the rate of a reaction.

a. $\Delta G°$ c. E_a e. concentration g. k
b. $\Delta H°$ d. temperature f. K_{eq} h. catalysts

6.49 The following is a concerted, bimolecular reaction: $CH_3Br + NaCN \rightarrow CH_3CN + NaBr$.

a. What is the rate equation for this reaction?
b. What happens to the rate of the reaction if $[CH_3Br]$ is doubled?
c. What happens to the rate of the reaction if [NaCN] is halved?
d. What happens to the rate of the reaction if $[CH_3Br]$ and [NaCN] are both increased by a factor of five?

6.50 The conversion of acetyl chloride to methyl acetate occurs via the following two-step mechanism:



acetyl chloride methyl acetate


a. Add curved arrows to show the movement of the electrons in each step.
b. Write the rate equation for this reaction, assuming the first step is rate-determining.
c. If the concentration of $^-OCH_3$ were increased 10 times, what would happen to the rate of the reaction?
d. If the concentrations of both CH_3COCl and $^-OCH_3$ were increased 10 times, what would happen to the rate of the reaction?
e. Classify the conversion of acetyl chloride to methyl acetate as an addition, elimination, or substitution.

6.51 Label each statement as true or false. Correct any false statement to make it true.

a. Increasing temperature increases reaction rate.
b. If a reaction is fast, it has a large rate constant.
c. A fast reaction has a large negative $\Delta G°$ value.
d. When E_a is large, the rate constant k is also large.
e. Fast reactions have equilibrium constants > 1.
f. Increasing the concentration of a reactant always increases the rate of a reaction.

General Problems


6.52 The Diels–Alder reaction, a powerful reaction discussed in Chapter 16, occurs when a 1,3-diene such as **A** reacts with an alkene such as **B** to form the six-membered ring in **C.**



A **B** **C**

a. Draw curved arrows to show how **A** and **B** react to form **C.**
b. What bonds are broken and formed in this reaction?
c. Would you expect this reaction to be endothermic or exothermic?
d. Does entropy favor the reactants or products?
e. Is the Diels–Alder reaction a substitution, elimination, or addition?

6.53 The conversion of $(CH_3)_3CI$ to $(CH_3)_2C=CH_2$ can occur by either a one-step or a two-step mechanism, as shown in Equations [1] and [2].




a. What rate equation would be observed for the mechanism in Equation [1]?
b. What rate equation would be observed for the mechanism in Equation [2]?
c. What is the order of each rate equation (i.e., first, second, and so forth)?
d. How can these rate equations be used to show which mechanism is the right one for this reaction?
e. Assume Equation [1] represents an endothermic reaction and draw an energy diagram for the reaction. Label the axes, reactants, products, E_a, and $\Delta H°$. Draw the structure for the transition state.
f. Assume Equation [2] represents an endothermic reaction and that the product of the rate-determining step is higher in energy than the reactants or products. Draw an energy diagram for this two-step reaction. Label the axes, reactants and products for each step, and the E_a and $\Delta H°$ for each step. Label $\Delta H°_{overall}$. Draw the structure for both transition states.

Challenge Problems


6.54 Explain why HC≡CH is more acidic than CH_3CH_3, even though the C–H bond in HC≡CH has a higher bond dissociation energy than the C–H bond in CH_3CH_3.

6.55 The use of curved arrows is a powerful tool that illustrates even complex reactions.


a. Add curved arrows to show how carbocation **A** is converted to carbocation **B.** Label each new σ bond formed. Similar reactions have been used in elegant syntheses of steroids.



b. Draw the product by following the curved arrows. This reaction is an example of a [3,3] sigmatropic rearrangement, as we will learn in Chapter 27.




6.56



propylbenzene

a. What carbon radical is formed by homolysis of the C–H_a bond in propylbenzene? Draw all reasonable resonance structures for this radical.
b. What carbon radical is formed by homolysis of the C–H_b bond in propylbenzene? Draw all reasonable resonance structures for this radical.
c. The bond dissociation energy of one of the C–H bonds is considerably less than the bond dissociation energy of the other. Which C–H bond is weaker? Offer an explanation.

6.57 Esterification is the reaction of a carboxylic acid (RCOOH) with an alcohol (R'OH) to form an ester (RCOOR') with loss of water. Equation [1] is an example of an *intermolecular* esterification reaction. Equation [2] is an example of an *intramolecular* esterification reaction; that is, the carboxylic acid and alcohol are contained in the same starting material, forming a cyclic ester as product. The equilibrium constants for both reactions are given. Explain why K_{eq} is different for these two apparently similar reactions.




6.58 Although K_{eq} of Equation [1] in Problem 6.57 does not greatly favor formation of the product, it is sometimes possible to use Le Châtelier's principle to increase the yield of ethyl acetate. Le Châtelier's principle states that if an equilibrium is disturbed, a system will react to counteract this disturbance. How can Le Châtelier's principle be used to drive the equilibrium to increase the yield of ethyl acetate? Another example of Le Châtelier's principle is given in Section 9.8.

6.59 As we will learn in Section 15.12, many antioxidants—compounds that prevent unwanted radical oxidation reactions from occurring—are phenols, compounds that contain an OH group bonded directly to a benzene ring.

a. Explain why homolysis of the O – H bond in phenol requires considerably less energy than homolysis of the O – H bond in ethanol (362 kJ/mol vs. 438 kJ/mol).

b. Why is the C – O bond in phenol shorter than the C – O bond in ethanol?



phenol ethanol

Alkyl Halides and Nucleophilic Substitution

7



Fluticasone is a prominent member of a growing number of useful drugs that contain halogens. Sold under the trade name Flonase, fluticasone is a synthetic steroid that is used as an aerosol inhalation for the treatment of seasonal and recurrent nasal allergies. Because it acts by decreasing inflammation, fluticasone is also used to combat the chronic lung inflammation associated with asthma. In Chapter 7 we learn about alkyl halides like fluticasone and one of their characteristic reactions, nucleophilic substitution.

7.1 Introduction to alkyl halides
7.2 Nomenclature
7.3 Physical properties
7.4 Interesting alkyl halides
7.5 The polar carbon–halogen bond
7.6 General features of nucleophilic substitution
7.7 The leaving group
7.8 The nucleophile
7.9 Possible mechanisms for nucleophilic substitution
7.10 Two mechanisms for nucleophilic substitution
7.11 The S_N2 mechanism
7.12 The S_N1 mechanism
7.13 Carbocation stability
7.14 The Hammond postulate
7.15 When is the mechanism S_N1 or S_N2?
7.16 Biological nucleophilic substitution
7.17 Vinyl halides and aryl halides
7.18 Organic synthesis

This is the first of three chapters dealing with an in-depth study of the organic reactions of compounds containing C–Z σ bonds, where Z is an element more electronegative than carbon. In Chapter 7 we learn about **alkyl halides** and one of their characteristic reactions, **nucleophilic substitution**. In Chapter 8, we look at **elimination**, a second general reaction of alkyl halides. We conclude this discussion in Chapter 9 by examining other molecules that also undergo nucleophilic substitution and elimination reactions.

7.1 Introduction to Alkyl Halides

Alkyl halides are organic molecules containing a halogen atom X bonded to an sp^3 hybridized carbon atom. As we learned in Section 3.2, alkyl halides are classified as **primary (1°), secondary (2°),** or **tertiary (3°)** depending on the number of carbons bonded to the carbon with the halogen. Whether an alkyl halide is 1°, 2°, or 3° is the *most important factor* in determining the course of its chemical reactions.


Alkyl halides have the general molecular formula $\mathbf{C_nH_{2n+1}X}$, and are formally derived from an alkane by replacing a hydrogen atom with a halogen.



alkyl halide

X = F, Cl, Br, I

C is sp^3 hybridized.



Four types of organic halides having the halogen atom in close proximity to a π bond are illustrated in Figure 7.1. **Vinyl halides** have a halogen atom bonded to a carbon–carbon double bond, and **aryl halides** have a halogen atom bonded to a benzene ring. These two types of organic halides with X bonded directly to an sp^2 hybridized carbon atom do *not* undergo the reactions presented in Chapter 7, as discussed in Section 7.17.

Figure 7.1
Four types of organic halides (RX) having X near a π bond



vinyl halide — aryl halide


- C bonded to X is sp^2 hybridized.
- These organic halides are **unreactive** in the reactions discussed in Chapter 7.

allylic halide — benzylic halide


- C bonded to X is sp^3 hybridized.
- These organic halides do participate in the reactions discussed in Chapter 7.

Allylic halides and benzylic halides have halogen atoms bonded to sp^3 hybridized carbon atoms and *do* undergo the reactions described in Chapter 7. **Allylic halides** have X bonded to the carbon atom *adjacent* to a carbon–carbon double bond, and **benzylic halides** have X bonded to the carbon atom *adjacent* to a benzene ring. The synthesis of allylic and benzylic halides is discussed in Sections 15.10 and 18.14, respectively.

Problem 7.1 Telfairine, a naturally occurring insecticide, and halomon, an antitumor agent, are two polyhalogenated compounds isolated from red algae. (a) Classify each halide bonded to an sp^3 hybridized carbon as 1°, 2°, or 3°. (b) Label each halide as vinyl, allylic, or neither.



telfairine



halomon

7.2 Nomenclature


The systematic (IUPAC) method for naming alkyl halides follows from the basic rules described in Chapter 4. Common names are also discussed in Section 7.2B, because many low molecular weight alkyl halides are often referred to by their common names.

7.2A IUPAC System


An alkyl halide is named as an alkane with a halogen substituent—that is, as a ***halo alkane.*** To name a halogen substituent, change the ***-ine*** ending of the name of the halogen to the suffix ***-o*** (chlor*ine* → chlor*o*).

How To Name an Alkyl Halide Using the IUPAC System

Example Give the IUPAC name of the following alkyl halide:



Step [1] Find the parent carbon chain and name it as an alkane.




7 C's in the longest chain

7 C's ----→ **heptane**

- Name the parent chain as an ***alkane,*** with the halogen as a substituent bonded to the longest chain.


Step [2] Apply all other rules of nomenclature.

a. **Number** the chain.



- Begin at the end nearest the first substituent, either alkyl or halogen.

b. **Name** and **number** the substituents.



c. **Alphabetize: c** for **c**hloro, then **m** for **m**ethyl.

Answer: 2-chloro-5-methylheptane

7.2B Common Names

Common names for alkyl halides are used only for simple alkyl halides. To assign a common name:

- **Name all the carbon atoms of the molecule as a single alkyl group.**
- **Name the halogen bonded to the alkyl group. To name the halogen, change the *-ine* ending of the halogen name to the suffix *-ide;* for example, brom*ine* → brom*ide*.**
- **Combine the names of the alkyl group and halide, separating the words with a space.**



tert-butyl group

***tert*-butyl iodide**




ethyl group


ethyl chloride

Other examples of alkyl halide nomenclature are given in Figure 7.2.

Figure 7.2
Examples: Nomenclature of alkyl halides




IUPAC: **1-chloro-2-methylpropane**
Common: isobutyl chloride





- **e**thyl group at **C1**
- **f**luoro group at **C2**


IUPAC: **1-ethyl-2-fluorocyclopentane**
earlier letter – – ➤ lower number
[too complex to use a common name]

Problem 7.2 Give the IUPAC name for each compound.

a. 

b. 

c. 


d. 

Problem 7.3 Give the structure corresponding to each name.

a. 3-chloro-2-methylhexane
b. 4-ethyl-5-iodo-2,2-dimethyloctane
c. *cis*-1,3-dichlorocyclopentane
d. 1,1,3-tribromocyclohexane
e. *sec*-butyl bromide
f. 6-ethyl-3-iodo-3,5-dimethylnonane



7.3 Physical Properties

Alkyl halides are weakly polar molecules. They exhibit **dipole–dipole** interactions because of their polar C–X bond, but because the rest of the molecule contains only C–C and C–H bonds they are incapable of intermolecular hydrogen bonding. How this affects their physical properties is summarized in Table 7.1.




Opposite ends of the dipoles interact.

Table 7.1 Physical Properties of Alkyl Halides

Property	Observation
Boiling point and melting point	• Alkyl halides have higher bp's and mp's than alkanes having the same number of carbons. 
	• Bp's and mp's increase as the size of R increases.  larger surface area higher mp and bp
	• Bp's and mp's increase as the size of X increases.  more polarizable halogen higher mp and bp
Solubility	• RX is soluble in organic solvents. • RX is insoluble in water.


Problem 7.4 An sp^3 hybridized C – Cl bond is more polar than an sp^2 hybridized C – Cl bond. (a) Explain why this phenomenon arises. (b) Rank the following compounds in order of increasing boiling point.



7.4 Interesting Alkyl Halides

Many simple alkyl halides make excellent solvents because they are not flammable and dissolve a wide variety of organic compounds. Compounds in this category include **$CHCl_3$** (chloroform or trichloromethane) and **CCl_4** (carbon tetrachloride or tetrachloromethane). Large quantities of these solvents are produced industrially each year, but like many chlorinated organic compounds, both chloroform and carbon tetrachloride are toxic if inhaled or ingested. Other simple alkyl halides are shown in Figure 7.3.

Figure 7.3
Some simple alkyl halides




CH_3Cl

- **Chloromethane (CH_3Cl)** is produced by giant kelp and algae and also found in emissions from volcanoes such as Hawai'i's Kilauea. Almost all of the atmospheric chloromethane results from these natural sources.



CH_2Cl_2


- **Dichloromethane (or methylene chloride, CH_2Cl_2)** is an important solvent, once used to decaffeinate coffee. Coffee is now decaffeinated by using supercritical CO_2 due to concerns over the possible ill effects of trace amounts of residual CH_2Cl_2 in the coffee. Subsequent studies on rats have shown, however, that no cancers occurred when animals ingested the equivalent of over 100,000 cups of decaffeinated coffee per day.




$CF_3CHClBr$


- **Halothane ($CF_3CHClBr$)** is a safe general anesthetic compared to other organic anesthetics such as $CHCl_3$, which causes liver and kidney damage, and $CH_3CH_2OCH_2CH_3$ (diethyl ether), which is very flammable.


Synthetic organic halides are also used in insulating materials, plastic wrap, and coatings. Two such compounds are **Teflon** and **poly(vinyl chloride) (PVC).**





Teflon
(nonstick coating)






poly(vinyl chloride) (PVC)
(plastic used in films, pipes, and insulation)

Thousands of organic halides have now been isolated from a variety of marine organisms. The characteristic smell of the ocean is due in part to simple alkyl halides found in seaweed. While many sponges and corals use these compounds as a chemical defense against predators and parasites, recent research has shown that some may have novel medicinal properties that offer




Hundreds of organic halides with diverse structures and biological activities have been isolated from red algae of the genus *Laurencia,* seaweed that grows in shallow water at the edges of reefs.

treatment for cancer and other diseases. Two halogenated compounds are ma'ilione, isolated from the Hawaiian red algae *Laurencia cartilaginea,* and plocoralide B, which kills esophageal cancer cells.




ma'ilione




plocoralide B

Although the beneficial effects of many organic halides are undisputed, certain synthetic chlorinated organics such as the **chlorofluorocarbons** and the pesticide **DDT** have caused lasting harm to the environment.




$CFCl_3$
CFC 11
Freon 11



DDT

=




Chlorofluorocarbons (CFCs) have the general molecular structure $\mathbf{CF_xCl_{4-x}}$. Trichlorofluoromethane [$CFCl_3$, CFC 11, or Freon 11 (trade name)] is an example of these easily vaporized compounds, having been extensively used as a refrigerant and an aerosol propellant. CFCs slowly rise to the stratosphere, where sunlight catalyzes their decomposition, a process that contributes to the destruction of the ozone layer, the thin layer of atmosphere that shields the earth's surface from harmful ultraviolet radiation (Section 15.9). Although it is now easy to second-guess the extensive use of CFCs, it is also easy to see why they were used so widely. **CFCs made refrigeration available to the general public.** Would you call your refrigerator a comfort or a necessity?

DDT, a nonbiodegradable pesticide, has been labeled both a "miraculous" discovery by Winston Churchill in 1945 and the "elixir of death" by Rachel Carson in her 1962 book *Silent Spring.* DDT use was banned in the United States in 1973, but because of its effectiveness and low cost, it is still widely used to control insect populations in developing countries.

The story of the insecticide **DDT** (**d**ichloro**d**iphenyl**t**richloroethane) follows the same theme: DDT is an organic molecule with valuable short-term effects that has caused long-term problems. DDT kills insects that spread diseases such as malaria and typhus, and in controlling insect populations, DDT has saved millions of lives worldwide. DDT is a weakly polar organic compound that persists in the environment for years. Because DDT is soluble in organic media, it accumulates in fatty tissues. Most adults in the United States have low concentrations of DDT (or a degradation product of DDT) in their bodies. DDT is acutely toxic to many types of marine life (crayfish, sea shrimp, and some fish), but the long-term effect on humans is not known.

Problem 7.5 Chondrocole A is a marine natural product isolated from red seaweed that grows in regions of heavy surf in the Pacific Ocean. (a) Predict the solubility of chondrocole A in water and CH_2Cl_2. (b) Locate the stereogenic centers and label each as *R* or *S*. (c) Draw a stereoisomer and a constitutional isomer of chondrocole A.



chondrocole A

7.5 The Polar Carbon–Halogen Bond

The properties of alkyl halides dictate their reactivity. The electrostatic potential maps of four simple alkyl halides in Figure 7.4 illustrate that the electronegative halogen X creates a polar C−X bond, making the carbon atom electron deficient. **The chemistry of alkyl halides is determined by this polar C−X bond.**





Figure 7.4 Electrostatic potential maps of four halomethanes (CH_3X)

- The polar C−X bond makes the carbon atom *electron deficient* in each CH_3X molecule.


What kind of reactions do alkyl halides undergo? **The characteristic reactions of alkyl halides are substitution and elimination.** Because alkyl halides contain an electrophilic carbon, they react with electron-rich reagents—Lewis bases (nucleophiles) and Brønsted–Lowry bases.

- **Alkyl halides undergo substitution reactions with nucleophiles.**



In a substitution reaction of an alkyl halide, **the halogen X is replaced by an electron-rich nucleophile :Nu⁻.** The C−X σ bond is broken and the C−Nu σ bond is formed.

- **Alkyl halides undergo elimination reactions with Brønsted–Lowry bases.**



In an elimination reaction of an alkyl halide, the **elements of HX are removed by a Brønsted–Lowry base :B.**

The remainder of Chapter 7 is devoted to a discussion of the substitution reactions of alkyl halides. Elimination reactions are discussed in Chapter 8.

7.6 General Features of Nucleophilic Substitution

Three components are necessary in any substitution reaction.




[1] An alkyl group containing an ***sp*³** hybridized carbon bonded to X.

[2] **X**—An atom X (or a group of atoms) called **a leaving group,** which is able to accept the electron density in the C−X bond. The most common leaving groups are halide anions (X^-), but H_2O (from ROH_2^+) and N_2 (from RN_2^+) are also encountered.


[3] **:Nu⁻**—A **nucleophile.** Nucleophiles contain a lone pair or a π bond but not necessarily a negative charge.

Because these substitution reactions involve electron-rich nucleophiles, they are called ***nucleophilic* substitution reactions.** Examples are shown in Equations [1]–[3]. **Nucleophilic**


substitutions are Lewis acid–base reactions. The nucleophile donates its electron pair, the alkyl halide (Lewis acid) accepts it, and the C – X bond is heterolytically cleaved. Curved arrow notation can be used to show the movement of electron pairs, as shown in Equation [3].



Negatively charged nucleophiles like $^-$OH and $^-$SH are used as **salts** with Li^+, Na^+, or K^+ counterions to balance charge. The identity of the cation is usually inconsequential, and therefore it is often omitted from the chemical equation.




When a neutral nucleophile is used, the substitution product bears a positive charge. **All atoms originally bonded to the nucleophile stay bonded to it after substitution occurs.** All three CH_3 groups stay bonded to the N atom in the given example.



The reaction of alkyl halides with NH_3 to form amines (RNH_2) is discussed in Chapter 25.


Furthermore, when the substitution product bears a positive charge and also contains a *proton* bonded to O or N, the initial substitution product readily loses a proton in a Brønsted–Lowry acid–base reaction, forming a neutral product.




All of these reactions are nucleophilic substitutions and have the same overall result—**replacement of the leaving group by the nucleophile,** regardless of the identity or charge of the nucleophile. To draw any nucleophilic substitution product:

- Find the sp^3 hybridized carbon with the leaving group.
- Identify the nucleophile, the species with a lone pair or π bond.
- Substitute the nucleophile for the leaving group and assign charges (if necessary) to any atom that is involved in bond breaking or bond formation.

Problem 7.6 Identify the nucleophile and leaving group and draw the products of each substitution reaction.




Problem 7.7 Draw the product of nucleophilic substitution with each neutral nucleophile. When the initial substitution product can lose a proton to form a neutral product, draw the product after proton transfer.





Cepacol throat lozenges and Crest Pro-Health Mouth Rinse contain the antiseptic CPC, which is prepared by nucleophilic substitution (Problem 7.8).

Problem 7.8 CPC (cetylpyridinium chloride), an antiseptic found in throat lozenges and mouthwash, is synthesized by the following reaction. Draw the structure of CPC.



Problem 7.9 What neutral nucleophile is needed to convert dihalide **A** to ticlopidine, an antiplatelet drug used to reduce the risk of strokes?



7.7 The Leaving Group

Nucleophilic substitution is a general reaction of organic compounds. Why, then, are alkyl halides the most common substrates, and halide anions the most common leaving groups? To answer this question, we must understand leaving group ability. **What makes a good leaving group?**

In a nucleophilic substitution reaction of R−X, the C−X bond is heterolytically cleaved, and the leaving group departs with the electron pair in that bond, forming $X:^-$. **The more stable the leaving group $X:^-$, the better able it is to accept an electron pair,** giving rise to the following generalization:


- **In comparing two leaving groups, the better leaving group is the weaker base.**




Good leaving groups are weak bases.

For example, H_2O is a better leaving group than ^{-}OH because H_2O is a weaker base. Moreover, the periodic trends in basicity can now be used to identify **periodic trends in leaving group ability:**

- **Left-to-right across a row of the periodic table, basicity *decreases* so leaving group ability *increases.***



- Down a column of the periodic table, basicity *decreases* so leaving group ability *increases.*



All good leaving groups are weak bases with strong conjugate acids having low pK_a values. Thus, all halide anions except F^- are good leaving groups because their conjugate acids (HCl, HBr, and HI) have low pK_a values. Tables 7.2 and 7.3 list good and poor leaving groups for nucleophilic substitution reactions, respectively. Nucleophilic substitution does not occur with any of the leaving groups in Table 7.3 because these leaving groups are strong bases.

Table 7.2 Good Leaving Groups for Nucleophilic Substitution

Starting material	Leaving group	Conjugate acid	pK_a
R—Cl	$:\ddot{Cl}:^-$	HCl	–7
R—Br	$:\ddot{Br}:^-$	HBr	–9
R—I	$:\ddot{I}:^-$	HI	–10
R—OH_2^+	$H_2\ddot{O}:$	H_3O^+	–1.7





Table 7.3 Poor Leaving Groups for Nucleophilic Substitution

Starting material	Leaving group	Conjugate acid	pK_a
R—F	$:\ddot{F}:^-$	HF	3.2
R—OH	$:\ddot{O}H^-$	H_2O	15.7
R—NH_2	$:\ddot{N}H_2^-$	NH_3	38
R—H	$H:^-$	H_2	35
R—R	$R:^-$	RH	50

Problem 7.10 Which is the better leaving group in each pair?

a. Cl^-, I^- b. NH_3, $^-NH_2$ c. H_2O, H_2S

Problem 7.11 Which molecules contain good leaving groups?


a.  b.  c.  d. 

Given a particular nucleophile and leaving group, how can we determine whether the equilibrium will favor products in a nucleophilic substitution? We can often correctly predict the direction of equilibrium by comparing the basicity of the nucleophile and the leaving group.

- **Equilibrium favors the products of nucleophilic substitution when the leaving group is a weaker base than the nucleophile.**

Sample Problem 7.1 illustrates how to apply this general rule.

Sample Problem 7.1 Will the following substitution reaction favor formation of the products?




Solution

Determine the basicity of the nucleophile ($^-$OH) and the leaving group (Cl^-) by comparing the pK_a values of their conjugate acids. **The stronger the conjugate acid, the weaker the base, and the better the leaving group.**

			conjugate acids	
nucleophile	$^-$OH	⟶	H_2O	pK_a = 15.7
leaving group	Cl^-	⟶	HCl	pK_a = −7
	weaker base		**stronger acid**	

Because Cl^-, the leaving group, is a weaker base than $^-$OH, the nucleophile, the reaction favors the products.

Problem 7.12 Does the equilibrium favor the reactants or products in each substitution reaction?




7.8 The Nucleophile

We use the word *base* to mean *Brønsted–Lowry* base and the word *nucleophile* to mean a *Lewis base* that reacts with electrophiles *other than protons*.

Nucleophiles and bases are structurally similar: both have a lone pair or a π bond. They differ in what they attack.

- **Bases attack protons. Nucleophiles attack other electron-deficient atoms (usually carbons).**



7.8A Nucleophilicity Versus Basicity

How is **nucleophilicity** (nucleophile strength) related to basicity? Although it is generally true that **a strong base is a strong nucleophile,** nucleophile size and steric factors can sometimes change this relationship.

Nucleophilicity parallels basicity in three instances.


[1] For two nucleophiles with the same nucleophilic atom, the stronger base is the stronger nucleophile.

- The relative nucleophilicity of $^-$OH and $CH_3CO_2^-$, two oxygen nucleophiles, is determined by comparing the pK_a values of their conjugate acids (H_2O and CH_3CO_2H). CH_3CO_2H (pK_a = 4.8) is a stronger acid than H_2O (pK_a = 15.7), so $^-$OH is a stronger base and stronger nucleophile than $CH_3CO_2^-$.

[2] A negatively charged nucleophile is always stronger than its conjugate acid.

- ^{-}OH is a stronger base and stronger nucleophile than H_2O, its conjugate acid.

[3] Right-to-left across a row of the periodic table, nucleophilicity increases as basicity increases.



Problem 7.13 Identify the stronger nucleophile in each pair.


a. NH_3, $^{-}NH_2$ b. CH_3NH_2, CH_3OH c. $CH_3CO_2^{-}$, $CH_3CH_2O^{-}$

7.8B Steric Effects and Nucleophilicity

All steric effects arise because two atoms cannot occupy the same space. In Chapter 4, for example, we learned that **steric strain** is an increase in energy when big groups (occupying a large volume) are forced close to each other.

Nucleophilicity does not parallel basicity when **steric hindrance** becomes important. ***Steric hindrance* is a decrease in reactivity resulting from the presence of bulky groups at the site of a reaction.**

For example, although pK_a tables indicate that *tert*-butoxide [$(CH_3)_3CO^{-}$] is a stronger base than ethoxide ($CH_3CH_2O^{-}$), **ethoxide is the *stronger* nucleophile.** The three CH_3 groups around the O atom of *tert*-butoxide create steric hindrance, making it more difficult for this big, bulky base to attack a tetravalent carbon atom.



Steric hindrance decreases nucleophilicity but *not* basicity. Because bases pull off small, easily accessible protons, they are unaffected by steric hindrance. Nucleophiles, on the other hand, must attack a crowded tetrahedral carbon, so bulky groups decrease reactivity.

Sterically hindered bases that are poor nucleophiles are called *nonnucleophilic bases*. Potassium *tert*-butoxide [$K^{+}\,^{-}OC(CH_3)_3$] is a strong, nonnucleophilic base.

7.8C Comparing Nucleophiles of Different Size—Solvent Effects

Atoms vary greatly in size down a column of the periodic table, and in this case, **nucleophilicity depends on the solvent used in a substitution reaction.** Although solvent has thus far been ignored, most organic reactions take place in a liquid solvent that dissolves all reactants to some

extent. Because substitution reactions involve polar starting materials, polar solvents are used to dissolve them. There are two main kinds of polar solvents—**polar *protic* solvents and polar *aprotic* solvents.**

Polar Protic Solvents


In addition to dipole–dipole interactions, **polar *protic* solvents are capable of intermolecular hydrogen bonding,** because they contain an O–H or N–H bond. The most common polar protic solvents are water and alcohols (ROH) (Figure 7.5). **Polar protic solvents solvate *both* cations and anions well.**

- Cations are solvated by ion–dipole interactions.
- Anions are solvated by hydrogen bonding.

Figure 7.5
Polar protic solvents


H_2O	CH_3OH methanol	CH_3CH_2OH ethanol	$(CH_3)_3COH$ *tert*-butanol	CH_3CO_2H acetic acid

For example, if the salt NaBr is used as a source of the nucleophile Br^- in H_2O, the Na^+ cations are solvated by ion–dipole interactions with H_2O molecules, and the Br^- anions are solvated by strong hydrogen bonding interactions.



How do polar protic solvents affect nucleophilicity? **In polar protic solvents, nucleophilicity *increases* down a column of the periodic table as the size of the anion increases. This is *opposite* to basicity.** A small electronegative anion like F^- is very well solvated by hydrogen bonding, effectively shielding it from reaction. On the other hand, a large, less electronegative anion like I^- does not hold onto solvent molecules as tightly. The solvent does not "hide" a large nucleophile as well, and the nucleophile is much more able to donate its electron pairs in a reaction. Thus, **nucleophilicity increases down a column** even though basicity decreases, giving rise to the following trend in polar protic solvents:

I^- is a weak base but a strong nucleophile in polar protic solvents.




Polar Aprotic Solvents

Polar *aprotic* solvents also exhibit dipole–dipole interactions, but they have no O−H or N−H bond so they are **incapable of hydrogen bonding.** Examples of polar aprotic solvents are shown in Figure 7.6. **Polar aprotic solvents solvate only cations well.**


- Cations are solvated by ion–dipole interactions.
- Anions are not well solvated because the solvent cannot hydrogen bond to them.

Figure 7.6
Polar aprotic solvents

Abbreviations are often used in organic chemistry, instead of a compound's complete name. A list of common abbreviations is given on the inside back cover.




When the salt NaBr is dissolved in acetone, $(CH_3)_2C{=}O$, the Na^+ cations are solvated by ion–dipole interactions with the acetone molecules, but, with no possibility for hydrogen bonding, the Br^- anions are not well solvated. Often these anions are called **naked anions** because they are not bound by tight interactions with solvent.





Br^- anions are surrounded by solvent but not well solvated by the $(CH_3)_2C{=}O$ molecules.


How do polar aprotic solvents affect nucleophilicity? Because anions are not well solvated in polar aprotic solvents, there is no need to consider whether solvent molecules more effectively hide one anion than another. Nucleophilicity parallels basicity and **the stronger base is the stronger nucleophile.** Because basicity decreases with size down a column, nucleophilicity decreases as well:



Problem 7.14 Classify each solvent as protic or aprotic.

a. 

b. 

c. 

Problem 7.15 Identify the stronger nucleophile in each pair of anions.

a. Br^- or Cl^- in a polar protic solvent

b. HO^- or Cl^- in a polar aprotic solvent

c. HS^- or F^- in a polar protic solvent

7.8D Summary

This discussion of nucleophilicity has brought together many new concepts, such as steric hindrance and solvent effects, both of which we will meet again in our study of organic chemistry. Keep in mind the central relationship between nucleophilicity and basicity in comparing two nucleophiles.

- **It is generally true that the *stronger* base is the *stronger* nucleophile.**
- **In polar protic solvents, however, nucleophilicity increases with increasing size of an anion (opposite to basicity).**
- **Steric hindrance decreases nucleophilicity without decreasing basicity, making $(CH_3)_3CO^-$ a stronger base but a weaker nucleophile than $CH_3CH_2O^-$.**

Table 7.4 lists some common nucleophiles used in nucleophilic substitution reactions.


Problem 7.16 Rank the nucleophiles in each group in order of increasing nucleophilicity.


a. ^-OH, $^-NH_2$, H_2O


b. ^-OH, Br^-, F^- (polar aprotic solvent)

c. H_2O, ^-OH, $CH_3CO_2^-$

Problem 7.17 What nucleophile is needed to convert $(CH_3)_2CHCH_2CH_2Br$ to each product?

a. 

b. 

c. 


d. 

Table 7.4 Common Nucleophiles in Organic Chemistry

	Negatively charged nucleophiles			Neutral nucleophiles	
Oxygen	^-OH	^-OR	$CH_3CO_2^-$	H_2O	ROH
Nitrogen	N_3^-			NH_3	RNH_2
Carbon	^-CN	$HC\equiv C^-$			
Halogen	Cl^-	Br^-	I^-		
Sulfur	HS^-	RS^-		H_2S	RSH


7.9 Possible Mechanisms for Nucleophilic Substitution

Now that you know something about the general features of nucleophilic substitution, you can begin to understand the mechanism.




Nucleophilic substitution at an sp^3 hybridized carbon involves two σ bonds: the bond to the leaving group is broken and the bond to the nucleophile is formed. To understand the mechanism of this reaction, though, we must know the timing of these two events; that is, **what is the order of bond breaking and bond making?** Do they happen at the same time, or does one event precede the other? There are three possibilities.

[1] **The mechanism has one step, and bond breaking and bond making occur at the *same* time.**




- If the C−X bond is broken ***as*** the C−Nu bond is formed, the mechanism has **one step.** As we learned in Section 6.9, the rate of such a bimolecular reaction depends on the concentration of both reactants; that is, the rate equation is **second order,** and **rate = *k*[RX][:Nu⁻].**

[2] **The mechanism has two steps, and bond breaking occurs *before* bond making.**



- If the C−X bond is broken ***first*** and then the C−Nu bond is formed, the mechanism has **two steps** and a **carbocation** is formed as an intermediate. Because the first step is rate-determining, the rate depends on the concentration of RX only; that is, the rate equation is **first order,** and **rate = *k*[RX].**

[3] **The mechanism has two steps and bond making occurs *before* bond breaking.**



- If the C−Nu bond is formed ***first*** and then the C−X bond is broken, the mechanism has **two steps,** but this mechanism has an inherent problem. The intermediate generated in the first step has 10 electrons around carbon, violating the octet rule, so this last possibility can be disregarded.

Thus, there are two possible mechanisms for nucleophilic substitution: **a one-step mechanism in which bond breaking and bond making are simultaneous, and a two-step mechanism in which bond breaking comes before bond making.** In Section 7.10 we look at data for two specific nucleophilic substitution reactions and see if those data fit either of these proposed mechanisms.

7.10 Two Mechanisms for Nucleophilic Substitution


Rate equations for two different reactions give us insight into the possible mechanism for nucleophilic substitution.

Reaction of bromomethane (CH_3Br) with acetate ($CH_3CO_2^-$) affords the substitution product methyl acetate with loss of Br^- as the leaving group (Equation [1]). Kinetic data show that the

reaction rate depends on the concentration of *both* reactants; that is, the rate equation is **second order.** This suggests a **bimolecular reaction with a one-step mechanism** in which the C–X bond is broken *as* the C–Nu bond is formed.



Equation [2] illustrates a similar nucleophilic substitution reaction with a different alkyl halide, $(CH_3)_3CBr$, which also leads to substitution of Br^- by $CH_3CO_2^-$. Kinetic data show that this reaction rate depends on the concentration of only *one* reactant, the alkyl halide; that is, the rate equation is **first order.** This suggests a **two-step mechanism in which the rate-determining step involves the alkyl halide only.**



The numbers **1** and **2** in the names S_N1 and S_N2 refer to the kinetic order of the reactions. For example, S_N**2** means that the kinetics are **second** order. The number *2* does *not* refer to the number of steps in the mechanism.


How can these two different results be explained? Although these two reactions have the same nucleophile and leaving group, **there must be two different mechanisms** because there are two different rate equations. These equations are specific examples of two well known mechanisms for nucleophilic substitution at an sp^3 hybridized carbon:

- **S_N2 mechanism (substitution nucleophilic bimolecular).**
- **S_N1 mechanism (substitution nucleophilic unimolecular).**

The reaction in Equation [1] illustrates an S_N2 mechanism, whereas the reaction in Equation [2] illustrates an S_N1 mechanism.

7.11 The S_N2 Mechanism

The reaction of CH_3Br with $CH_3CO_2^-$ is an example of an **S_N2 reaction.** What are the general features of this mechanism?



7.11A Kinetics

An S_N2 reaction exhibits **second-order kinetics;** that is, the reaction is **bimolecular** and both the alkyl halide and the nucleophile appear in the rate equation.

- rate = $k[CH_3Br][CH_3CO_2^-]$

Changing the concentration of *either* reactant affects the rate. For example, doubling the concentration of *either* the nucleophile or the alkyl halide doubles the rate. Doubling the concentration of *both* reactants increases the rate by a factor of four.

Problem 7.18 What happens to the rate of an S_N2 reaction under each of the following conditions?


a. [RX] is tripled, and $[:Nu^-]$ stays the same.
b. Both [RX] and $[:Nu^-]$ are tripled.
c. [RX] is halved, and $[:Nu^-]$ stays the same.
d. [RX] is halved, and $[:Nu^-]$ is doubled.

7.11B A One-Step Mechanism

The most straightforward explanation for the observed second-order kinetics is a **concerted reaction—bond breaking and bond making occur at the *same* time,** as shown in Mechanism 7.1.

Mechanism 7.1 The S_N2 Mechanism

One step The C – Br bond breaks as the C – O bond forms.




An energy diagram for the reaction of $CH_3Br + CH_3CO_2^-$ is shown in Figure 7.7. The reaction has one step, so there is one energy barrier between reactants and products. Because the equilibrium for this S_N2 reaction favors the products, the products are drawn at lower energy than the starting materials.

Problem 7.19 Draw an energy diagram for the following S_N2 reaction. Label the axes, the starting materials, and the product. Draw the structure of the transition state.



Figure 7.7
An energy diagram for the S_N2 reaction: $CH_3Br + CH_3CO_2^- \rightarrow CH_3CO_2CH_3 + Br^-$



- In the transition state, the C—Br bond is partially broken, the C—O bond is partially formed, and both the attacking nucleophile and the departing leaving group bear a partial negative charge.

7.11C Stereochemistry of the S_N2 Reaction


From what direction does the nucleophile approach the substrate in an S_N2 reaction? There are two possibilities.

- **Frontside attack:** The nucleophile approaches from the ***same*** side as the leaving group.
- **Backside attack:** The nucleophile approaches from the side ***opposite*** the leaving group.


The results of frontside and backside attack of a nucleophile are illustrated with $CH_3CH(D)Br$ as substrate and the general nucleophile $:Nu^-$. This substrate has the leaving group bonded to a stereogenic center, thus allowing us to see the structural difference that results when the nucleophile attacks from two different directions.

In **frontside attack,** the nucleophile approaches from the **same** side as the leaving group, forming **A.** In this example, the leaving group was drawn on the right, so the nucleophile attacks from the right, and all other groups remain in their original positions. Because the nucleophile and leaving group are in the same position relative to the other three groups on carbon, frontside attack results in **retention of configuration** around the stereogenic center.


Recall from Section 1.1 that D stands for the isotope deuterium (^2H).



In **backside attack,** the nucleophile approaches from the **opposite** side to the leaving group, forming **B.** In this example, the leaving group was drawn on the right, so the nucleophile attacks from the left. Because the nucleophile and leaving group are in the opposite position relative to the other three groups on carbon, backside attack results in **inversion of configuration** around the stereogenic center.



The products of frontside and backside attack are *different* compounds. **A** and **B** are stereoisomers that are nonsuperimposable—they are **enantiomers.**



Inversion of configuration in an S_N2 reaction is often called **Walden inversion,** after Latvian chemist Dr. Paul Walden, who first observed this process in 1896.

Which product is formed in an S_N2 reaction? When the stereochemistry of the product is determined, **only B, the product of backside attack, is formed.**


Backside attack occurs in all S_N2 reactions, but we can observe this change only when the leaving group is bonded to a stereogenic center.

- **All S_N2 reactions proceed with backside attack of the nucleophile, resulting in *inversion* of configuration at a stereogenic center.**

One explanation for backside attack is based on an electronic argument. Both the nucleophile and leaving group are electron rich and these like charges repel each other. Backside attack

keeps these two groups as far away from each other as possible. In the transition state, the nucleophile and leaving group are 180° away from each other, and the other three groups around carbon occupy a plane, as illustrated in Figure 7.8.


Figure 7.8
Stereochemistry of the S_N2 reaction



• :Nu⁻ and Br⁻ are 180° away from each other, on either side of a plane containing R, H, and D.


Two additional examples of inversion of configuration in S_N2 reactions are given in Figure 7.9.

Figure 7.9
Two examples of inversion of configuration in the S_N2 reaction




- The bond to the nucleophile in the product is always on the **opposite side** compared to the bond to the leaving group in the starting material. If the leaving group is drawn to the left, the nucleophile approaches from the right. If the leaving group is drawn in front of the plane (on a wedge), the nucleophile approaches from the back and ends up on a dashed wedge.

Sample Problem 7.2 Label the nucleophile and leaving group, and draw the product (including stereochemistry) of the following S_N2 reaction.



Solution

Br^- is the leaving group and ^-CN is the nucleophile. Because S_N2 reactions proceed with **inversion** of configuration and the leaving group is drawn above the ring (on a wedge), the nucleophile must come in from below (ending up on a dashed wedge).



- **Inversion** of configuration occurs at the C – Br bond.
- Backside attack converts the **cis** starting material to a **trans** product because the nucleophile (^-CN) attacks from below the plane of the ring.


Problem 7.20 Draw the product of each S_N2 reaction and indicate stereochemistry.



7.11D The Identity of the R Group

How does the rate of an S_N2 reaction change as the alkyl group in the substrate alkyl halide changes from CH_3 ⟶ 1° ⟶ 2° ⟶ 3°?


- **As the number of R groups on the carbon with the leaving group *increases,* the rate of an S_N2 reaction *decreases.***



- **Methyl and 1° alkyl halides undergo S_N2 reactions with ease.**
- **2° Alkyl halides react more slowly.**
- **3° Alkyl halides *do not* undergo S_N2 reactions.**

This order of reactivity can be explained by steric effects. As small H atoms are replaced by larger alkyl groups, **steric hindrance caused by bulky R groups makes nucleophilic attack from the back side more difficult,** slowing the reaction rate. Figure 7.10 illustrates the effect of increasing steric hindrance in a series of alkyl halides.

Figure 7.10
Steric effects in the S_N2 reaction




- **The S_N2 reaction is fastest with unhindered halides.**

Table 7.5 summarizes what we have learned thus far about an S_N2 mechanism.

Table 7.5 Characteristics of the S_N2 Mechanism


Characteristic	Result
Kinetics	• Second-order kinetics; rate = k[RX][:Nu^-]
Mechanism	• One step
Stereochemistry	• Backside attack of the nucleophile • Inversion of configuration at a stereogenic center
Identity of R	• Unhindered halides react fastest. • Rate: $CH_3X > RCH_2X > R_2CHX > R_3CX$

Problem 7.21 Which compound in each pair undergoes a faster S_N2 reaction?




The S_N2 reaction is a key step in the laboratory synthesis of many drugs including **ethambutol** (trade name Myambutol), used in the treatment of tuberculosis, and **fluoxetine** (trade name Prozac), an antidepressant, as illustrated in Figure 7.11.

Figure 7.11 Nucleophilic substitution in the synthesis of two useful drugs




- In both examples, the initial substitution product bears a positive charge and goes on to lose a proton to form the product drawn.
- The NH_2 group serves as a neutral nucleophile to displace halogen in each synthesis. The new bonds formed by nucleophilic substitution are drawn in red in the products.

Problem 7.22 What neutral nucleophile is needed to convert **C** to **D?** Imatinib, an effective treatment for certain cancers, is prepared in one step from **D.**



7.12 The S_N1 Mechanism

The reaction of $(CH_3)_3CBr$ with $CH_3CO_2^-$ is an example of the second mechanism for nucleophilic substitution, the **S_N1 mechanism.** What are the general features of this mechanism?



7.12A Kinetics

The S_N1 reaction exhibits **first-order kinetics.**

- rate = $k[(CH_3)_3CBr]$

As we learned in Section 7.10, the kinetics suggest that the S_N1 mechanism involves **more than one step,** and that the slow step is **unimolecular,** involving *only* the alkyl halide. **The identity and concentration of the nucleophile have no effect on the reaction rate.** Doubling the concentration of $(CH_3)_3CBr$ doubles the rate, but doubling the concentration of the nucleophile has no effect.


Problem 7.23 What happens to the rate of an S_N1 reaction under each of the following conditions?

a. [RX] is tripled, and [:Nu^-] stays the same.
b. Both [RX] and [:Nu^-] are tripled.
c. [RX] is halved, and [:Nu^-] stays the same.
d. [RX] is halved, and [:Nu^-] is doubled.

7.12B A Two-Step Mechanism

The most straightforward explanation for the observed first-order kinetics is a **two-step mechanism** in which **bond breaking occurs *before* bond making,** as shown in Mechanism 7.2.

Mechanism 7.2 The S_N1 Mechanism



1. Heterolysis of the C–Br bond forms a carbocation in the rate-determining step.
2. Nucleophilic attack of acetate (a Lewis base) on the carbocation (a Lewis acid) forms the new C–O bond.


The key features of the S_N1 mechanism are:

- The mechanism has two steps.
- Carbocations are formed as reactive intermediates.

An energy diagram for the reaction of $(CH_3)_3CBr + CH_3CO_2^-$ is shown in Figure 7.12. Each step has its own energy barrier, with a transition state at each energy maximum. Because the transition state for Step [1] is at higher energy, Step [1] is rate-determining. $\Delta H°$ for Step [1] has a positive value because only bond breaking occurs, whereas $\Delta H°$ of Step [2] has a negative value because only bond making occurs. The overall reaction is assumed to be exothermic, so the final product is drawn at lower energy than the initial starting material.

Figure 7.12


An energy diagram for the S_N1 reaction: $(CH_3)_3CBr + CH_3CO_2^- \rightarrow (CH_3)_3COCOCH_3 + Br^-$



- The S_N1 mechanism has two steps, so there are two energy barriers.
- $E_a[1] > E_a[2]$ because Step [1] involves bond breaking and Step [2] involves bond formation.
- In each step only one bond is broken or formed, so the transition state for each step has one partial bond.


7.12C Stereochemistry of the S_N1 Reaction

To understand the stereochemistry of the S_N1 reaction, we must examine the geometry of the carbocation intermediate.



- **A carbocation (with three groups around C) is sp^2 hybridized and trigonal planar, and contains a vacant *p* orbital extending above and below the plane.**

To illustrate the consequences of having a trigonal planar carbocation formed as a reactive intermediate, we examine the S_N1 reaction of a 3° alkyl halide **A** having the leaving group bonded to a stereogenic center.




Loss of the leaving group in Step [1] generates a planar carbocation that is now achiral. Attack of the nucleophile in Step [2] can occur from either the front or the back to afford two products, **B** and **C.** These two products are *different* compounds containing one stereogenic center. **B** and **C** are stereoisomers that are not superimposable—they are **enantiomers.** Because there is no preference for nucleophilic attack from either direction, an equal amount of the two enantiomers is formed—a **racemic mixture.** We say that ***racemization*** has occurred.

Nucleophilic attack from both sides of a planar carbocation occurs in S_N1 reactions, but we see the result of this phenomenon only when the leaving group is bonded to a stereogenic center.

- ***Racemization*** **is the formation of equal amounts of two enantiomeric products from a single starting material.**
- **S_N1 reactions proceed with *racemization* at a single stereogenic center.**


Two additional examples of racemization in S_N1 reactions are given in Figure 7.13.

Figure 7.13
Two examples of racemization in the S_N1 reaction




- Nucleophilic substitution of each starting material by an S_N1 mechanism forms a **racemic mixture** of two products.
- With H_2O, a neutral nucleophile, the initial product of nucleophilic substitution (ROH_2^+) loses a proton to form the final neutral product, ROH (Section 7.6).

Sample Problem 7.3 Label the nucleophile and leaving group, and draw the products (including stereochemistry) of the following S_N1 reaction.




Solution

Br^- is the leaving group and H_2O is the nucleophile. Loss of the leaving group generates a trigonal planar carbocation, which can react with the nucleophile from either direction to form two products.




In this example, the initial products of nucleophilic substitution bear a positive charge. They readily lose a proton to form neutral products. The overall process with a neutral nucleophile thus has **three steps:** the first two constitute the **two-step S_N1 mechanism** (loss of the leaving group and attack of the nucleophile), and the third is a **Brønsted–Lowry acid–base reaction** leading to a neutral organic product.



The two products in this reaction are nonsuperimposable mirror images—**enantiomers.** Because nucleophilic attack on the trigonal planar carbocation occurs with equal frequency from both directions, a **racemic mixture is formed.**


Problem 7.24 Draw the products of each S_N1 reaction and indicate the stereochemistry of any stereogenic centers.



7.12D The Identity of the R Group

How does the rate of an S_N1 reaction change as the alkyl group in the substrate alkyl halide changes from CH_3 --→ 1° --→ 2° --→ 3°?

- **As the number of R groups on the carbon with the leaving group *increases,* the rate of an S_N1 reaction *increases.***



- 3° Alkyl halides undergo S_N1 reactions rapidly.
- 2° Alkyl halides react more slowly.
- Methyl and 1° alkyl halides do *not* undergo S_N1 reactions.


This trend is exactly opposite to that observed for the S_N2 mechanism. To explain this result, we must examine the rate-determining step, the formation of the carbocation, and learn about the effect of alkyl groups on **carbocation stability.** Table 7.6 summarizes the characteristics of the S_N1 mechanism.

Table 7.6 Characteristics of the S_N1 Mechanism

Characteristic	Result
Kinetics	• First-order kinetics; rate = k[RX]
Mechanism	• Two steps
Stereochemistry	• Trigonal planar carbocation intermediate • Racemization at a single stereogenic center
Identity of R	• More substituted halides react fastest. • Rate: $R_3CX > R_2CHX > RCH_2X > CH_3X$




7.13 Carbocation Stability

Carbocations are classified as **primary (1°), secondary (2°), or tertiary (3°)** by the number of R groups bonded to the charged carbon atom. As the number of R groups on the positively charged carbon atom increases, the stability of the carbocation **increases.**



When we speak of carbocation stability, we really mean *relative* stability. Tertiary carbocations are too unstable to isolate, but they are more stable than secondary carbocations. We will examine the reason for this order of stability by invoking two different principles: **inductive effects** and **hyperconjugation.**

Problem 7.25 Classify each carbocation as 1°, 2°, or 3°.

a. 
b. 
c. 
d. 


Problem 7.26 Draw the structure of a 1°, 2°, and 3° carbocation, each having molecular formula $C_4H_9^+$. Rank the three carbocations in order of increasing stability.

7.13A Inductive Effects

Inductive effects are electronic effects that occur through σ bonds. In Section 2.5B, for example, we learned that more electronegative atoms stabilize a negative charge by an **electron-withdrawing inductive effect.**


Electron-donor groups (Z) stabilize a (+) charge; Z→Y^+. Electron-withdrawing groups (W) stabilize a (–) charge; W←Y^-.

To stabilize a positive charge, **electron-donating groups** are needed. **Alkyl groups are electron-donor groups that stabilize a positive charge.** An alkyl group with several σ bonds is more polarizable than a hydrogen atom, and more able to donate electron density. Thus, as R groups successively replace the H atoms in CH_3^+, **the positive charge is more dispersed on the electron donor R groups, and the carbocation is more stabilized.**




Electrostatic potential maps for four carbocations in Figure 7.14 illustrate the effect of increasing alkyl substitution on the positive charge of the carbocation.

Figure 7.14
Electrostatic potential maps for different carbocations



- Dark blue areas in electrostatic potential plots indicate regions low in electron density. As alkyl substitution increases, the region of positive charge is less concentrated on carbon.

Problem 7.27 Rank the following carbocations in order of increasing stability.




7.13B Hyperconjugation

A second explanation for the observed trend in carbocation stability is based on orbital overlap. A 3° carbocation is more stable than a 2°, 1°, or methyl carbocation because the positive charge is delocalized over more than one atom.

- **Spreading out charge by the overlap of an empty *p* orbital with an adjacent σ bond is called *hyperconjugation.***

For example, CH_3^+ cannot be stabilized by hyperconjugation, but $(CH_3)_2CH^+$ can:




Both carbocations contain an sp^2 hybridized carbon, so both are trigonal planar with a vacant p orbital extending above and below the plane. There are no adjacent C−H σ bonds with which the p orbital can overlap in CH_3^+, but there *are* adjacent C−H σ bonds in $(CH_3)_2CH^+$. This overlap (the **hyperconjugation**) delocalizes the positive charge on the carbocation, spreading it over a larger volume, and this stabilizes the carbocation.

The larger the number of alkyl groups on the adjacent carbons, the greater the possibility for hyperconjugation, and the larger the stabilization. Hyperconjugation thus provides an alternate way of explaining why **carbocations with a larger number of R groups are more stabilized**.


7.14 The Hammond Postulate

The rate of an S_N1 reaction depends on the rate of formation of the carbocation (the product of the rate-determining step) via heterolysis of the C−X bond.

- The rate of an S_N1 reaction *increases* as the number of R groups on the carbon with the leaving group *increases.*
- The stability of a carbocation *increases* as the number of R groups on the positively charged carbon *increases.*



- Thus, the rate of an S_N1 reaction *increases* as the stability of the carbocation *increases.*




The rate of a reaction depends on the magnitude of E_a, and the stability of a product depends on $\Delta G°$. The **Hammond postulate,** first proposed in 1955, **relates rate to stability.**

7.14A The General Features of the Hammond Postulate

The Hammond postulate provides a qualitative estimate of the energy of a transition state. Because the energy of the transition state determines the energy of activation and therefore the reaction rate, predicting the relative energy of two transition states allows us to determine the relative rates of two reactions.

According to the Hammond postulate, the transition state of a reaction resembles the structure of the species (reactant or product) to which it is closer in energy. In **endothermic reactions,** the transition state is closer in energy to the **products.** In **exothermic reactions,** the transition state is closer in energy to the **reactants.**




- Transition states in endothermic reactions resemble the products.
- Transition states in exothermic reactions resemble the reactants.

What happens to the reaction rate if the energy of the product is lowered? In an **endothermic reaction,** the transition state resembles the products, so anything that stabilizes the product stabilizes the transition state, too. **Lowering the energy of the transition state *decreases* the energy of activation (E_a), which *increases* the reaction rate.**

Suppose there are two possible products of an endothermic reaction, but one is more stable (lower in energy) than the other (Figure 7.15). According to the Hammond postulate, **the transition state to form the more stable product is lower in energy, so this reaction should occur faster.**

Figure 7.15
An endothermic reaction—how the energies of the transition state and products are related




• The lower energy transition state leads to the lower energy product.

- In an endothermic reaction, the more stable product forms faster.

What happens to the reaction rate of an **exothermic reaction** if the energy of the product is lowered? The transition state resembles the reactants, so **lowering the energy of the products has little or no effect on the energy of the transition state.** If E_a is unaffected, then the reaction rate is unaffected, too, as shown in Figure 7.16.

Figure 7.16
An exothermic reaction—how the energies of the transition state and products are related



- Decreasing the energy of the product often has little effect on the energy of the transition state.

- **In an exothermic reaction, the more stable product may or may not form faster because E_a is similar for both products.**

7.14B The Hammond Postulate and the S_N1 Reaction

In the S_N1 reaction, the rate-determining step is the formation of the carbocation, an *endothermic* reaction. According to the Hammond postulate, the **stability of the carbocation determines the rate of its formation.**

For example, heterolysis of the C−Cl bond in $(CH_3)_2CHCl$ affords a less stable 2° carbocation, $(CH_3)_2CH^+$ (Equation [1]), whereas heterolysis of the C−Cl bond in $(CH_3)_3CCl$ affords a more stable 3° carbocation, $(CH_3)_3C^+$ (Equation [2]). The Hammond postulate states that Reaction [2] is faster than Reaction [1], because the transition state to form the more stable 3° carbocation is lower in energy. Figure 7.17 depicts an energy diagram comparing these two endothermic reactions.






Figure 7.17
Energy diagram for carbocation formation in two different S_N1 reactions





- $(CH_3)_2CH^+$ is less stable than $(CH_3)_3C^+$, so $E_a[1] > E_a[2]$, and Reaction [1] is slower.

In conclusion, the Hammond postulate can be used to predict the relative rates of two reactions. In the S_N1 reaction the rate-determining step is endothermic, so the more stable carbocation is formed faster.

Problem 7.28 Which alkyl halide in each pair reacts faster in an S_N1 reaction?

a.  or 

b.  or 

7.15 When Is the Mechanism S_N1 or S_N2?


Given a particular starting material and nucleophile, how do we know whether a reaction occurs by the S_N1 or S_N2 mechanism? Four factors are examined:

- **The alkyl halide—CH_3X, RCH_2X, R_2CHX, or R_3CX**
- **The nucleophile—strong or weak**
- **The leaving group—good or poor**
- **The solvent—protic or aprotic**

7.15A The Alkyl Halide—The Most Important Factor

The most important factor in determining whether a reaction follows the S_N1 or S_N2 mechanism is the *identity of the alkyl halide.*


- Increasing alkyl substitution favors S_N1.
- Decreasing alkyl substitution favors S_N2.



- Methyl and 1° halides (CH_3X and RCH_2X) undergo S_N2 reactions only.
- 3° Alkyl halides (R_3CX) undergo S_N1 reactions only.
- 2° Alkyl halides (R_2CHX) undergo both S_N1 and S_N2 reactions. Other factors determine the mechanism.


Examples are given in Figure 7.18.


Figure 7.18
The identity of RX and the mechanism of nucleophilic substitution





1° halide	2° halide	3° halide
S_N2	Both **S_N2** and **S_N1** are possible.	**S_N1**

Problem 7.29 What is the likely mechanism of nucleophilic substitution for each alkyl halide?

a. 

b. 

c. 

d. 


7.15B The Nucleophile

How does the strength of the nucleophile affect an S_N1 or S_N2 mechanism? The rate of the S_N1 reaction is unaffected by the identity of the nucleophile because the nucleophile does not appear in the rate equation (rate = k[RX]). The identity of the nucleophile *is* important for the S_N2 reaction, however, because the nucleophile does appear in the rate equation for this mechanism (rate = k[RX][:Nu^-]).

- Strong nucleophiles present in high concentration favor S_N2 reactions.
- Weak nucleophiles favor S_N1 reactions by decreasing the rate of any competing S_N2 reaction.


The most common nucleophiles in S_N2 reactions bear a net negative charge. The most common nucleophiles in S_N1 reactions are weak nucleophiles such as H_2O and ROH. The identity of the nucleophile is especially important in determining the mechanism and therefore the stereochemistry of nucleophilic substitution when 2° alkyl halides are starting materials.

Let's compare the substitution products formed when the 2° alkyl halide **A** (*cis*-1-bromo-4-methylcyclohexane) is treated with either the strong nucleophile ^-OH or the weak nucleophile H_2O. Because a 2° alkyl halide can react by either mechanism, the strength of the nucleophile determines which mechanism takes place.




cis-1-bromo-4-methylcyclohexane
A

The **strong nucleophile $^{-}$OH favors an S_N2 reaction,** which occurs with **backside** attack of the nucleophile, resulting in **inversion of configuration.** Because the leaving group Br^- is above the plane of the ring, the nucleophile attacks from below, and a single product **B** is formed.





The **weak nucleophile H_2O favors an S_N1 reaction,** which occurs by way of an intermediate carbocation. Loss of the leaving group in **A** forms the carbocation, which undergoes nucleophilic attack from both above and below the plane of the ring to afford two products, **C** and **D.** Loss of a proton by proton transfer forms the final products, **B** and **E. B** and **E** are diastereomers of each other (**B** is a trans isomer and **E** is a cis isomer).





Thus, the mechanism of nucleophilic substitution determines the stereochemistry of the products formed.

Problem 7.30 For each alkyl halide and nucleophile: [1] Draw the product of nucleophilic substitution; [2] determine the likely mechanism (S_N1 or S_N2) for each reaction.


a.  + CH_3OH


b.  + ^{-}SH

c.  + $CH_3CH_2O^-$

d.  + CH_3OH

Problem 7.31 Draw the products (including stereochemistry) for each reaction.

a.  + H_2O ⟶

b.  + $^{-}:C{\equiv}C-H$ ⟶


7.15C The Leaving Group

How does the identity of the leaving group affect an S_N1 or S_N2 reaction?


- A better leaving group increases the rate of *both* S_N1 and S_N2 reactions.


Because the bond to the leaving group is partially broken in the transition state of the only step of the S_N2 mechanism and the slow step of the S_N1 mechanism, **a better leaving group increases the rate of both reactions.** The better the leaving group, the more willing it is to accept the electron pair in the C−X bond, and the faster the reaction.


For alkyl halides, the following order of reactivity is observed for the S_N1 and the S_N2 mechanisms:




Problem 7.32 Which compound in each pair reacts faster in nucleophilic substitution?

a. 

c. 

b. 

d. 

7.15D The Solvent

Polar protic solvents and polar aprotic solvents affect the rates of S_N1 and S_N2 reactions differently.

- **Polar *protic* solvents are especially good for S_N1 reactions.**
- **Polar *aprotic* solvents are especially good for S_N2 reactions.**

Summary of solvent effects:
- Polar protic solvents favor S_N1 reactions because the ionic intermediates are stabilized by solvation.
- Polar aprotic solvents favor S_N2 reactions because nucleophiles are not well solvated, and therefore are more nucleophilic.

Polar protic solvents like H_2O and ROH solvate both cations and anions well, and this characteristic is important for the S_N1 mechanism, in which two ions (a carbocation and a leaving group) are formed by heterolysis of the C–X bond. The carbocation is solvated by ion–dipole interactions with the polar solvent, and the leaving group is solvated by hydrogen bonding, in much the same way that Na^+ and Br^- are solvated in Section 7.8C. These interactions stabilize the reactive intermediate. In fact, a polar protic solvent is generally needed for an S_N1 reaction.


Polar aprotic solvents exhibit dipole–dipole interactions but not hydrogen bonding, and as a result, they do not solvate anions well. This has a pronounced effect on the nucleophilicity of anionic nucleophiles. Because these nucleophiles are not "hidden" by strong interactions with the solvent, they are **more nucleophilic.** Because stronger nucleophiles favor S_N2 reactions, **polar aprotic solvents are especially good for S_N2 reactions.**

Problem 7.33 Which solvents favor S_N1 reactions and which favor S_N2 reactions?


a. 


b. CH_3CN


c. 

d. 

Problem 7.34 For each reaction, use the identity of the alkyl halide and nucleophile to determine which substitution mechanism occurs. Then determine which solvent affords the faster reaction.

a. 

b. 

c. 

7.15E Summary of Factors That Determine Whether the S_N1 or S_N2 Mechanism Occurs


Table 7.7 summarizes the factors that determine whether a reaction occurs by the S_N1 or S_N2 mechanism. Sample Problems 7.4 and 7.5 illustrate how these factors are used to determine the mechanism of a given reaction.


Table 7.7 Summary of Factors That Determine the S_N1 or S_N2 Mechanism

Alkyl halide	Mechanism	Other factors
CH_3X RCH_2X (1°)	**S_N2**	Favored by • **strong nucleophiles** (usually a net negative charge) • polar **aprotic** solvents
R_3CX (3°)	**S_N1**	Favored by • **weak nucleophiles** (usually neutral) • polar **protic** solvents
R_2CHX (2°)	**S_N1 or S_N2**	The mechanism depends on the conditions. • **Strong nucleophiles favor the S_N2 mechanism over the S_N1 mechanism.** RO^- is a stronger nucleophile than ROH, so RO^- favors the S_N2 reaction and ROH favors the S_N1 reaction. • **Protic solvents favor the S_N1 mechanism and aprotic solvents favor the S_N2 mechanism.** H_2O and CH_3OH are polar protic solvents that favor the S_N1 mechanism, whereas acetone $[(CH_3)_2C{=}O]$ and DMSO $[(CH_3)_2S{=}O]$ are polar aprotic solvents that favor the S_N2 mechanism.

Sample Problem 7.4


Determine the mechanism of nucleophilic substitution for each reaction and draw the products.

a. 


b. 

Solution

a. The alkyl halide is 1°, so it must react by an S_N2 mechanism with the nucleophile $^-:C{\equiv}CH$.





b. The alkyl halide is 2°, so it can react by either the S_N1 or S_N2 mechanism. The strong nucleophile (^-CN) favors the S_N2 mechanism.



Sample Problem 7.5


Determine the mechanism of nucleophilic substitution for each reaction and draw the products, including stereochemistry.

a. 


b. 

Solution


a. The 2° alkyl halide can react by either the S_N1 or S_N2 mechanism. The strong nucleophile ($^-OCH_3$) favors the S_N2 mechanism, as does the polar aprotic solvent (DMSO). S_N2 reactions proceed with inversion of configuration.



b. The alkyl halide is 3°, so it reacts by an S_N1 mechanism with the weak nucleophile CH_3OH. S_N1 reactions proceed with racemization at a single stereogenic center, so two products are formed.



Problem 7.35 Determine the mechanism and draw the products of each reaction. Include the stereochemistry at all stereogenic centers.




7.16 Biological Nucleophilic Substitution

Nucleophilic substitution occurs in a wide variety of biological reactions.


7.16A Leaving Groups Derived from Phosphorus

In contrast to nucleophilic substitutions run in the laboratory that use alkyl halides as substrates and halide anions as leaving groups, biological substitutions often occur with phosphorus leaving groups, such as phosphate (PO_4^{3-}, abbreviated as P_i for inorganic phosphate), diphosphate ($P_2O_7^{4-}$, abbreviated as PP_i), and triphosphate ($P_3O_{10}^{5-}$, abbreviated as PPP_i). These anions are excellent leaving groups because they are weak, resonance-stabilized bases.




phosphate P_i | diphosphate PP_i | triphosphate PPP_i

When an organic compound contains a carbon bonded to one of these leaving groups, the compound is called an organic monophosphate, diphosphate, or triphosphate.



organic monophosphate R–OP | organic diphosphate R–OPP | organic triphosphate R–OPPP


Adenosine triphosphate (ATP) is an organic triphosphate.




adenosine triphosphate

ATP

Nucleophilic substitutions with these substrates may proceed by either an S_N2 or S_N1 pathway, as shown with the general diphosphate R_2CHOPP.



The final step in the biosynthesis of geraniol, a component of rose oil used in perfumery, is an S_N1 reaction of geranyl diphosphate with water. This reaction occurs by way of a resonance-stabilized carbocation. We will learn more about reactions of diphosphates in Chapter 16.




geranyl diphosphate



SAM, a nutritional supplement sold under the name SAMe (pronounced sammy), has been used in Europe to treat depression and arthritis for over 20 years. In cells, SAM is used in nucleophilic substitutions that synthesize key amino acids, hormones, and neurotransmitters.

7.16B *S*-Adenosylmethionine

A common nucleophilic substitution occurs with *S*-adenosylmethionine, or **SAM.** SAM is the cell's equivalent of CH_3I. The many polar functional groups in SAM make it soluble in the aqueous environment in the cell.




S-adenosylmethionine

SAM


The CH_3 group in SAM [abbreviated as $(CH_3SR_2)^+$] is part of a sulfonium salt, a positively charged sulfur species that contains a good leaving group. Nucleophilic attack at the CH_3 group

of SAM displaces R_2S, a good neutral leaving group. This reaction is called **methylation,** because a CH_3 group is transferred from one compound (SAM) to another ($:Nu^-$).




For example, **adrenaline** (epinephrine) is a hormone synthesized in the adrenal glands from noradrenaline (norepinephrine) by nucleophilic substitution using SAM (Figure 7.19). When an individual senses danger or is confronted by stress, the hypothalamus region of the brain signals the adrenal glands to synthesize and release adrenaline, which enters the bloodstream and then

Figure 7.19
Adrenaline synthesis from noradrenaline in response to stress



stimulates a response in many organs. Stored carbohydrates are metabolized in the liver to form glucose, which is further metabolized to provide an energy boost. Heart rate and blood pressure increase, and lung passages are dilated. These physiological changes result from the "rush of adrenaline," and prepare an individual for "fight or flight."

Problem 7.36 Nicotine, a toxic and addictive component of tobacco, is synthesized from **A** using SAM. Write out the reaction that converts **A** into nicotine.



A nicotine

7.17 Vinyl Halides and Aryl Halides

S_N1 and S_N2 reactions occur only at sp^3 hybridized carbon atoms. Now that we have learned about the mechanisms for nucleophilic substitution we can understand why **vinyl halides** and **aryl halides,** which have a halogen atom bonded to an sp^2 hybridized C, do not undergo nucleophilic substitution by either the S_N1 or S_N2 mechanism. The discussion here centers on vinyl halides, but similar arguments hold for aryl halides as well.




vinyl halide aryl halide

C bonded to X is sp^2 hybridized.




Vinyl halides do not undergo S_N2 reactions in part because of the percent *s*-character in the hybrid orbital of the carbon atom in the C−X bond. The higher percent *s*-character in the sp^2 hybrid orbital of the vinyl halide compared to the sp^3 hybrid orbital of the alkyl halide (33% vs. 25%) makes the bond shorter and stronger.

Vinyl halides do not undergo S_N1 reactions because heterolysis of the C−X bond would form a **highly unstable vinyl carbocation.** Because this carbocation has only two groups around the positively charged carbon, it is ***sp*** hybridized. These carbocations are even less stable than 1° carbocations, so the S_N1 reaction does not take place.



vinyl carbocation
highly unstable



Problem 7.37 Rank the following carbocations in order of increasing stability.

a.  b.  c. 

7.18 Organic Synthesis

Thus far we have concentrated on the starting material in nucleophilic substitution—the alkyl halide—and have not paid much attention to the product formed. Nucleophilic substitution reactions, and in particular S_N2 reactions, introduce a wide variety of different functional groups in molecules, depending on the nucleophile. For example, when $^-$OH, $^-$OR, and $^-$CN are used as nucleophiles, the products are alcohols (ROH), ethers (ROR), and nitriles (RCN), respectively. Table 7.8 lists some functional groups readily introduced using nucleophilic substitution.

Table 7.8 Molecules Synthesized from R–X by the S_N2 Reaction

	Nucleophile (:Nu⁻)	Product	Name
Oxygen compounds	^-OH	R–OH	alcohol
	$^-OR'$	R–OR'	ether
			ester
Carbon compounds	^-CN	R–CN	nitrile
	$^-:C\equiv C-H$	R–C≡C–H	alkyne
Nitrogen compounds	N_3^-	$R-N_3$	azide
	$:NH_3$	$R-NH_2$	amine
Sulfur compounds	^-SH	R–SH	thiol
	$^-SR'$	R–SR'	sulfide

By thinking of **nucleophilic substitution as a reaction that *makes* a particular kind of organic compound,** we begin to think about ***synthesis*.**

- **Organic synthesis is the systematic preparation of a compound from a readily available starting material by one or many steps.**



Aspirin is synthesized by a two-step procedure from simple, cheap starting materials.


Phenol, the starting material for the aspirin synthesis, is a petroleum product, like most of the starting materials used in large quantities in industrial syntheses. A shortage of petroleum reserves thus affects the availability not only of fuels for transportation, but also of raw materials needed for most chemical synthesis.

7.18A Background on Organic Synthesis

Chemists synthesize molecules for many reasons. Sometimes a **natural product,** a compound isolated from natural sources, has useful medicinal properties, but is produced by an organism in only minute quantities. Synthetic chemists then prepare this molecule from simpler starting materials, so that it can be made available to a large number of people.

Sometimes, chemists prepare molecules that do not occur in nature (although they may be similar to those in nature), because these molecules have superior properties to their naturally occurring relatives. **Aspirin, or acetylsalicylic acid** (Section 2.7), is a well known example. Acetylsalicylic acid is prepared from phenol, a product of the petroleum industry, by a two-step procedure (Figure 7.20). Aspirin has become one of the most popular and widely used drugs in the world because it has excellent analgesic and anti-inflammatory properties, *and* it is cheap and readily available.


Figure 7.20 Synthesis of aspirin




7.18B Nucleophilic Substitution and Organic Synthesis

To carry out synthesis we must think backwards. We examine a compound and ask: **What starting material and reagent are needed to make it?** If we are using nucleophilic substitution, we must determine what alkyl halide and what nucleophile can be used to form a specific product. This is the simplest type of synthesis because it involves only one step. In Chapter 11 we will learn about multistep syntheses.

Suppose, for example, that we are asked to prepare $(CH_3)_2CHCH_2OH$ (2-methylpropan-1-ol) from an alkyl halide and any required reagents. To accomplish this synthesis, we must "fill in the boxes" for the starting material and reagent in the accompanying equation.






To determine the two components needed for the synthesis, remember that the carbon atoms come from the organic starting material, in this case a 1° alkyl halide [$(CH_3)_2CHCH_2Br$]. The functional group comes from the nucleophile, ^-OH in this case. With these two components, we can "fill in the boxes" to complete the synthesis.



After any synthesis is proposed, check to see if it is reasonable, given what we know about reactions. Will the reaction written give a high yield of product? The synthesis of $(CH_3)_2CHCH_2OH$ is reasonable, because the starting material is a 1° alkyl halide and the nucleophile (^-OH) is strong, and both facts contribute to a successful S_N2 reaction.

Problem 7.38 What alkyl halide and nucleophile are needed to prepare each compound?

a. 
b. 
c. 
d. 

Problem 7.39 The ether, $CH_3OCH_2CH_3$, can be prepared by two different nucleophilic substitution reactions, one using CH_3O^- as nucleophile and the other using $CH_3CH_2O^-$ as nucleophile. Draw both routes.

KEY CONCEPTS

Alkyl Halides and Nucleophilic Substitution


General Facts about Alkyl Halides

- Alkyl halides contain a halogen atom X bonded to an sp^3 hybridized carbon (7.1).
- Alkyl halides are named as halo alkanes, with the halogen as a substituent (7.2).
- Alkyl halides have a polar C – X bond, so they exhibit dipole–dipole interactions but are incapable of intermolecular hydrogen bonding (7.3).

- The polar C – X bond containing an electrophilic carbon makes alkyl halides reactive towards nucleophiles and bases (7.5).

The Central Theme (7.6)

- Nucleophilic substitution is one of the two main reactions of alkyl halides. A nucleophile replaces a leaving group on an sp^3 hybridized carbon.



- One σ bond is broken and one σ bond is formed.
- There are two possible mechanisms: S_N1 and S_N2.

S_N1 and S_N2 Mechanisms Compared

	S_N2 mechanism	S_N1 mechanism
[1] Mechanism	• One step (7.11B)	• Two steps (7.12B)
[2] Alkyl halide	• Order of reactivity: $CH_3X > RCH_2X > R_2CHX > R_3CX$ (7.11D)	• Order of reactivity: $R_3CX > R_2CHX > RCH_2X > CH_3X$ (7.12D)
[3] Rate equation	• Rate = k[RX][:Nu⁻]	• Rate = k[RX]
	• Second-order kinetics (7.11A)	• First-order kinetics (7.12A)
[4] Stereochemistry	• Backside attack of the nucleophile (7.11C)	• Trigonal planar carbocation intermediate (7.12C)
	• Inversion of configuration at a stereogenic center	• Racemization at a single stereogenic center
[5] Nucleophile	• Favored by stronger nucleophiles (7.15B)	• Favored by weaker nucleophiles (7.15B)
[6] Leaving group	• Better leaving group ⇢ faster reaction (7.15C)	• Better leaving group ⇢ faster reaction (7.15C)
[7] Solvent	• Favored by polar aprotic solvents (7.15D)	• Favored by polar protic solvents (7.15D)

Important Trends

- The best leaving group is the weakest base. Leaving group ability increases left-to-right across a row and down a column of the periodic table (7.7).
- Nucleophilicity decreases left-to-right across a row of the periodic table (7.8A).
- Nucleophilicity decreases down a column of the periodic table in polar aprotic solvents (7.8C).
- Nucleophilicity increases down a column of the periodic table in polar protic solvents (7.8C).
- The stability of a carbocation increases as the number of R groups bonded to the positively charged carbon increases (7.13).


Important Principles


Principle	Example
• Electron-donating groups (such as R groups) stabilize a positive charge (7.13A).	• 3° Carbocations (R_3C^+) are more stable than 2° carbocations (R_2CH^+), which are more stable than 1° carbocations (RCH_2^+).
• Steric hindrance decreases nucleophilicity but not basicity (7.8B).	• $(CH_3)_3CO^-$ is a stronger base but a weaker nucleophile than $CH_3CH_2O^-$.
• Hammond postulate: In an endothermic reaction, the more stable product is formed faster. In an exothermic reaction, this is not necessarily true (7.14).	• S_N1 reactions are faster when more stable (more substituted) carbocations are formed, because the rate-determining step is endothermic.
• Planar, sp^2 hybridized atoms react with reagents from both sides of the plane (7.12C).	• A trigonal planar carbocation reacts with nucleophiles from both sides of the plane.

PROBLEMS

Problems Using Three-Dimensional Models


7.40 Give the IUPAC name for each compound, including any *R,S* designation.

a. 

b. 


7.41 Draw the products formed when each alkyl halide is treated with NaCN.


a. 


b. 


Nomenclature


7.42 Give the IUPAC name for each compound.


a. 

b. 

c. 

d. 

e. 

f. 

Include the *R,S* designation in the name.

7.43 Give the structure corresponding to each name.


a. 3-bromo-4-ethylheptane
b. 1,1-dichloro-2-methylcyclohexane
c. 1-bromo-4-ethyl-3-fluorooctane
d. (*S*)-3-iodo-2-methylnonane
e. (1*R*,2*R*)-*trans*-1-bromo-2-chlorocyclohexane
f. (*R*)-4,4,5-trichloro-3,3-dimethyldecane


7.44 Draw the eight constitutional isomers having the molecular formula $C_5H_{11}Cl$.

a. Give the IUPAC name for each compound (ignoring *R* and *S* designations).
b. Classify each alkyl halide as 1°, 2°, or 3°.
c. Label any stereogenic centers.
d. For each constitutional isomer that contains a stereogenic center, draw all possible stereoisomers, and label each stereogenic center as *R* or *S*.

Physical Properties


7.45 Which compound in each pair has the higher boiling point?

a. 


b. 

General Nucleophilic Substitution, Leaving Groups, and Nucleophiles

7.46 Draw the products of each nucleophilic substitution reaction.




7.47 Which of the following molecules contain a good leaving group?



7.48 Rank the species in each group in order of increasing leaving group ability.

a. Br^-, Cl^-, I^- b. NH_3, H_2S, H_2O

7.49 Which of the following nucleophilic substitution reactions will take place?



7.50 Rank the species in each group in order of increasing nucleophilicity.

a. $CH_3CH_2S^-$, $CH_3CH_2O^-$, $CH_3CO_2^-$ in CH_3OH
b. CH_3NH_2, CH_3SH, CH_3OH in acetone
c. ^-OH, F^-, Cl^- in acetone
d. HS^-, F^-, Cl^- in CH_3OH


7.51 Classify each solvent as protic or aprotic.

a. $(CH_3)_2CHOH$
b. CH_3NO_2
c. CH_2Cl_2
d. NH_3
e. $N(CH_3)_3$
f. $HCONH_2$

7.52 Why is the amine N atom more nucleophilic than the amide N atom in $CH_3CONHCH_2CH_2CH_2NHCH_3$?

The S_N2 Reaction

7.53 Consider the following S_N2 reaction.




a. Draw a mechanism using curved arrows.
b. Draw an energy diagram. Label the axes, the reactants, products, E_a, and $\Delta H°$. Assume that the reaction is exothermic.
c. Draw the structure of the transition state.
d. What is the rate equation?
e. What happens to the reaction rate in each of the following instances? [1] The leaving group is changed from Br^- to I^-; [2] The solvent is changed from acetone to CH_3CH_2OH; [3] The alkyl halide is changed from $CH_3(CH_2)_4Br$ to $CH_3CH_2CH_2CH(Br)CH_3$; [4] The concentration of ^-CN is increased by a factor of five; and [5] The concentrations of both the alkyl halide and ^-CN are increased by a factor of five.


7.54 Pick the reactant or solvent in each part that gives the faster S_N2 reaction.

a. reaction of ^-OH with CH_3CH_2Br or CH_3CH_2Cl
b. reaction of $CH_3CH_2CH_2Cl$ with NaOH or $NaOCOCH_3$
c. reaction of $CH_3CH_2CH_2I$ with $^-OCH_3$ in CH_3OH or DMSO


7.55 Draw the products of each S_N2 reaction and indicate the stereochemistry where appropriate.

a. 

+ $^-OCH_3$ ⟶

b. 


+ $^-OCH_2CH_3$ ⟶


c. 

+ ^-CN ⟶


Carbocations

7.56 Classify the carbocations as 1°, 2°, or 3°, and rank the carbocations in each group in order of increasing stability.


a. 

b. 

7.57 Which of the following carbocations (**A** or **B**) is more stable? Explain your choice.



A



B

The S_N1 Reaction


7.58 Consider the following S_N1 reaction.




a. Draw a mechanism for this reaction using curved arrows.
b. Draw an energy diagram. Label the axes, starting material, product, E_a, and $\Delta H°$. Assume that the starting material and product are equal in energy.
c. Draw the structure of any transition states.
d. What is the rate equation for this reaction?
e. What happens to the reaction rate in each of the following instances? [1] The leaving group is changed from I^- to Cl^-; [2] The solvent is changed from H_2O to DMF; [3] The alkyl halide is changed from $(CH_3)_2C(I)CH_2CH_3$ to $(CH_3)_2CHCH(I)CH_3$; [4] The concentration of H_2O is increased by a factor of five; and [5] The concentrations of both the alkyl halide and H_2O are increased by a factor of five.

7.59 Pick the reactant or solvent in each part that gives the faster S_N1 reaction.
a. reaction of H_2O with $(CH_3)_3CCl$ or $(CH_3)_3CI$
b. reaction of CH_3OH with $(CH_3)_3CBr$ or $(CH_3)_2CHCH_2Br$
c. reaction of $CH_3CH_2CH(I)CH_3$ with CH_3CH_2OH in water or DMSO

7.60 Draw the products of each S_N1 reaction and indicate the stereochemistry when necessary.


a. 

+ CH_3CH_2OH ⟶

b. 

+ H_2O ⟶


7.61 Draw a stepwise mechanism for the following reaction that illustrates how two substitution products are formed. Explain why 1-bromohex-2-ene reacts rapidly with a weak nucleophile (CH_3OH) under S_N1 reaction conditions, even though it is a 1° alkyl halide.



1-bromohex-2-ene


S_N1 and S_N2 Reactions

7.62 (a) Rank **A, B,** and **C** in order of increasing S_N2 reactivity. (b) Rank **A, B,** and **C** in order of increasing S_N1 reactivity.




A **B** **C**

7.63 Determine the mechanism of nucleophilic substitution of each reaction and draw the products, including stereochemistry.




7.64 Fluticasone, the chapter-opening molecule, can be prepared by the following reaction. Draw a stepwise mechanism for this reaction.




A fluticasone

7.65 Uridine monophosphate (UMP) is one of the four nucleotides that compose RNA, the nucleic acid that translates the genetic information of DNA into proteins needed by cells for proper function and development. A key step in the synthesis of UMP is the S_N1 reaction of **A** with **B** to form **C,** which is then converted to UMP in one step. Draw a stepwise mechanism for this S_N1 reaction.




A **B** **C** + HO—PP UMP


7.66 Diphenhydramine, the antihistamine in Benadryl, can be prepared by the following two-step sequence. What is the structure of diphenhydramine?




7.67 Draw a stepwise, detailed mechanism for the following reaction. Use curved arrows to show the movement of electrons.




7.68 When a single compound contains both a nucleophile and a leaving group, an **intramolecular** reaction may occur. With this in mind, draw the product of the following reaction.




7.69 Nicotine can be made when the following ammonium salt is treated with Na_2CO_3. Draw a stepwise mechanism for this reaction.



7.70 Quinapril (trade name Accupril) is used to treat high blood pressure and congestive heart failure. One step in the synthesis of quinapril involves reaction of the racemic alkyl bromide **A** with a single enantiomer of the amino ester **B.** (a) What two products are formed in this reaction? (b) Given the structure of quinapril, which one of these two products is needed to synthesize the drug?




7.71 Draw a stepwise, detailed mechanism for the following reaction.




7.72 When (*R*)-6-bromo-2,6-dimethylnonane is dissolved in CH_3OH, nucleophilic substitution yields an optically inactive solution. When the isomeric halide (*R*)-2-bromo-2,5-dimethylnonane is dissolved in CH_3OH under the same conditions, nucleophilic substitution forms an optically active solution. Draw the products formed in each reaction, and explain why the difference in optical activity is observed.

Synthesis


7.73 Fill in the appropriate reagent or starting material in each of the following reactions.



7.74 Devise a synthesis of each compound from an alkyl halide using any other organic or inorganic reagents.




7.75 Suppose you have compounds **A**–**D** at your disposal. Using these compounds, devise two different ways to make **E.** Which one of these methods is preferred, and why?




A **B** **C** **D** **E**

7.76 Muscalure, the sex pheromone of the common housefly, can be prepared by a reaction sequence that uses two nucleophilic substitutions. Identify compounds **A**–**D** in the following synthesis of muscalure.



Challenge Problems

7.77 We will return often to nucleophilic substitution, in particular the S_N2 reaction, in subsequent chapters and concentrate on the *nucleophile,* rather than the alkyl halide. By using different nucleophiles, nucleophilic substitution allows the synthesis of a wide variety of organic compounds with many different functional groups. With this in mind, draw the products of each two-step sequence. (Hint: Step [1] in each part involves an acid–base reaction that removes the most acidic hydrogen from the starting material.)




7.78 Explain why quinuclidine is a much more reactive nucleophile than triethylamine, even though both compounds have N atoms surrounded by three R groups.




quinuclidine triethylamine

7.79 Draw a stepwise mechanism for the following reaction sequence.




major product minor product


7.80 As we will learn in Chapter 9, an epoxide is an ether with an oxygen atom in a three-membered ring. Epoxides can be made by intramolecular S_N2 reactions of intermediates that contain a nucleophile and a leaving group on adjacent carbons, as shown.



Assume that each of the following starting materials can be converted to an epoxide by this reaction. Draw the product formed (including stereochemistry) from each starting material. Why might some of these reactions be more difficult than others in yielding nucleophilic substitution products?




7.81 When trichloride **J** is treated with CH_3OH, nucleophilic substitution forms the dihalide **K.** Draw a mechanism for this reaction and explain why one Cl is much more reactive than the other two Cl's so that a single substitution product is formed.



J **K**

7.82 In some nucleophilic substitutions under S_N1 conditions, complete racemization does not occur and a small excess of one enantiomer is present. For example, treatment of optically pure 1-bromo-1-phenylpropane with water forms 1-phenylpropan-1-ol. (a) Calculate how much of each enantiomer is present using the given optical rotation data. (b) Which product predominates—the product of inversion or the product of retention of configuration? (c) Suggest an explanation for this phenomenon.




1-bromo-1-phenylpropane

1-phenylpropan-1-ol
observed [α] = +5.0
optically pure *S* isomer, [α] = –48

Alkyl Halides and Elimination Reactions

8



DDE, dichlorodiphenyl**d**ichloro**e**thylene, is formed by the elimination of HCl from the pesticide DDT. DDE and DDT accumulate in the fatty tissues of predator birds such as osprey that feed on fish contaminated with DDT. When DDE and DDT concentration is high, mother osprey produce eggs with very thin shells that are easily crushed, so fewer osprey chicks hatch. In Chapter 8, we learn about **elimination reactions,** the second general reaction of alkyl halides, which form alkenes like DDE.

8.1 General features of elimination
8.2 Alkenes—The products of elimination reactions
8.3 The mechanisms of elimination
8.4 The E2 mechanism
8.5 The Zaitsev rule
8.6 The E1 mechanism
8.7 S_N1 and E1 reactions
8.8 Stereochemistry of the E2 reaction
8.9 When is the mechanism E1 or E2?
8.10 E2 reactions and alkyne synthesis
8.11 When is the reaction S_N1, S_N2, E1, or E2?


Elimination reactions introduce π bonds into organic compounds, so they can be used to synthesize **alkenes** and **alkynes**—hydrocarbons that contain one and two π bonds, respectively. Like nucleophilic substitution, elimination reactions can occur by two different pathways, depending on the conditions. By the end of Chapter 8, therefore, you will have learned four different reaction mechanisms, two for nucleophilic substitution (S_N1 and S_N2) and two for elimination (E1 and E2).

The biggest challenge with this material is learning how to sort out two different reactions that follow four different mechanisms. **Will a particular alkyl halide undergo substitution or elimination with a given reagent, and by which of the four possible mechanisms?** To answer this question, we conclude Chapter 8 with a summary that allows you to predict which reaction and mechanism are likely for a given substrate.


8.1 General Features of Elimination

All **elimination reactions** involve loss of elements from the starting material to form a new π bond in the product.


- **Alkyl halides undergo elimination reactions with Brønsted–Lowry bases. The elements of HX are lost and an alkene is formed.**



Equations [1] and [2] illustrate examples of elimination reactions. In both reactions a base removes the elements of an acid, HBr or HCl, from the organic starting material.



Removal of the elements of HX, called **dehydrohalogenation,** is one of the most common methods to introduce a π bond and prepare an alkene. Dehydrohalogenation is an example of **β elimination,** because it involves loss of elements from two adjacent atoms: the **α carbon** bonded to the leaving group X, and the **β carbon** adjacent to it. Three curved arrows illustrate how four bonds are broken or formed in the process.



- The base (B:) removes a proton on the β carbon, thus forming H−B+.
- The electron pair in the β C−H bond forms the new π bond between the α and β carbons.
- The electron pair in the C−X bond ends up on halogen, forming the leaving group :X−.

The most common bases used in elimination reactions are negatively charged oxygen compounds such as −OH and its alkyl derivatives, −OR, called **alkoxides,** listed in Table 8.1. Potassium *tert*-butoxide, $K^+ {}^-OC(CH_3)_3$, a bulky nonnucleophilic base, is especially useful (Section 7.8B).


Table 8.1
Common Bases Used in Dehydrohalogenation

Structure	Name
$Na^+ {}^-OH$	Sodium hydroxide
$K^+ {}^-OH$	Potassium hydroxide
$Na^+ {}^-OCH_3$	Sodium methoxide
$Na^+ {}^-OCH_2CH_3$	Sodium ethoxide
$K^+ {}^-OC(CH_3)_3$	Potassium *tert*-butoxide


To draw any product of dehydrohalogenation:

- **Find the α carbon—the sp^3 hybridized carbon bonded to the leaving group.**
- **Identify all β carbons with H atoms.**
- **Remove the elements of H and X from the α and β carbons and form a π bond.**

For example, 2-bromo-2-methylpropane has three β carbons (three CH_3 groups), but because all three are *identical,* only *one* alkene is formed upon elimination of HBr. In contrast, 2-bromobutane has two *different* β carbons (labeled β_1 and β_2), so elimination affords *two* constitutional isomers by loss of HBr across either the α and β_1 carbons, or the α and β_2 carbons. We learn about which product predominates and why in Section 8.5.




2-bromo-2-methylpropane one alkene




2-bromobutane two constitutional isomers

An elimination reaction is the first step in the slow degradation of the **pesticide DDT** (Chapter 8 opening paragraph and Section 7.4). Elimination of HCl from DDT forms the degradation product **DDE** (dichlorodiphenyldichloroethylene). This stable alkene is found in minute concentration in the fatty tissues of most adults in the United States.



DDT DDE

Problem 8.1 Label the α and β carbons in each alkyl halide. Draw all possible elimination products formed when each alkyl halide is treated with $K^{+-}OC(CH_3)_3$.




8.2 Alkenes—The Products of Elimination Reactions

Because elimination reactions of alkyl halides form alkenes, let's review earlier material on alkene structure and learn some additional facts as well.

8.2A Bonding in a Carbon–Carbon Double Bond

Recall from Section 1.10B that alkenes are hydrocarbons containing a carbon–carbon double bond. Each carbon of the double bond is sp^2 hybridized and trigonal planar, and all bond angles are 120°.




ethylene



Ethylene, the simplest alkene, is a hormone that regulates plant growth and fruit ripening. A ripe banana speeds up the ripening of green tomatoes because the banana gives off ethylene.

The double bond of an alkene consists of a σ bond and a π bond.



- **The σ bond, formed by end-on overlap of the two sp^2 hybrid orbitals, lies in the plane of the molecule.**
- **The π bond, formed by side-by-side overlap of two 2*p* orbitals, lies perpendicular to the plane of the molecule. The π bond is formed during elimination.**

Alkenes are classified according to the number of carbon atoms bonded to the carbons of the double bond. A **monosubstituted alkene** has one carbon atom bonded to the carbons of the double bond. A **disubstituted alkene** has two carbon atoms bonded to the carbons of the double bond, and so forth.






Figure 8.1 shows several alkenes and how they are classified. You must be able to classify alkenes in this way to determine the major and minor products of elimination reactions, when a mixture of alkenes is formed.

Figure 8.1
Classifying alkenes by the number of R groups bonded to the double bond



- Carbon atoms bonded to the double bond are screened in blue.


Problem 8.2 Classify each alkene in the following vitamins by the number of carbon substituents bonded to the double bond.




8.2B Restricted Rotation

Figure 8.2 shows that there is free rotation about the carbon–carbon single bonds of butane, but *not* around the carbon–carbon double bond of but-2-ene. Because of restricted rotation, two stereoisomers of but-2-ene are possible.

Figure 8.2
Rotation around C–C and C=C compared



These conformations **interconvert** by rotation.
They represent the **same** molecule.



These molecules **do not interconvert** by rotation.
They are **different** molecules.


The concept of cis and trans isomers was first introduced for disubstituted cycloalkanes in Chapter 4. In both cases, a ring or a double bond restricts motion, preventing the rotation of a group from one side of the ring or double bond to the other.

- **The cis isomer has two groups on the *same side* of the double bond.**
- **The trans isomer has two groups on *opposite sides* of the double bond.**




cis-But-2-ene and *trans*-but-2-ene are stereoisomers, but not mirror images of each other, so they are **diastereomers.**

The cis and trans isomers of but-2-ene are a specific example of a general type of stereoisomer occurring at carbon–carbon double bonds. **Whenever the two groups on *each* end of a carbon–carbon double bond are *different from each other,* two diastereomers are possible.**




Problem 8.3 For which double bonds are stereoisomers possible?







The characteristic fragrance of lilac is a mixture of (*E*)-ocimene and other volatile ethers, aldehydes, and alcohols.



Problem 8.4 (a) Which double bonds in (*E*)-ocimene, a major component of the odor of lilac flowers, can exhibit stereoisomerism? (b) Draw a diastereomer of (*E*)-ocimene.






(*E*)-ocimene

Problem 8.5 Label each pair of alkenes as constitutional isomers, stereoisomers, or identical.

a.  and 

c.  and 


b.  and 

d.  and 

8.2C Stability of Alkenes


Some alkenes are more stable than others. For example, **trans alkenes are generally more stable than cis alkenes** because the larger groups bonded to the double bond carbons are farther apart, reducing steric interactions.

The trans isomer has the CH_3 groups farther away from each other.



more stable

Steric interactions of the CH_3 groups destabilize the cis isomer.



less stable


The stability of an alkene increases, moreover, as the **number of R groups bonded to the double bond carbons increases.**

least stable — most stable

$CH_2{=}CH_2 < RCH{=}CH_2 < R_2C{=}CH_2 \sim RCH{=}CHR < R_2C{=}CHR < R_2C{=}CR_2$


Increasing number of R groups
Increasing stability

R groups increase the stability of an alkene because R groups are sp^3 hybridized, whereas the carbon atoms of the double bond are sp^2 hybridized. Recall from Sections 1.11B and 2.5D that the percent *s*-character of a hybrid orbital increases from 25% to 33% in going from sp^3 to sp^2. The higher the percent *s*-character, the more readily an atom accepts electron density. Thus, ***sp*^{2} hybridized carbon atoms are more able to *accept* electron density** and ***sp*^{3} hybridized carbon atoms are more able to *donate* electron density.**



- **As a result, increasing the number of electron-donating R groups on a carbon atom able to accept electron density makes the alkene more stable.**


Thus, *trans*-but-2-ene (a disubstituted alkene) is more stable than *cis*-but-2-ene (another disubstituted alkene), but both are more stable than but-1-ene (a monosubstituted alkene).






In summary:

- **Trans alkenes are more stable than cis alkenes because they have fewer steric interactions.**
- **Increasing alkyl substitution stabilizes an alkene by an electron-donating inductive effect.**


Problem 8.6 Which alkene in each pair is more stable?

a. or 


b.  or 

c.  or 

Problem 8.7 Several factors can affect alkene stability. Explain why alkene **A** is more stable than alkene **B** even though both contain disubstituted carbon–carbon double bonds.



A



B

8.3 The Mechanisms of Elimination

What is the mechanism for elimination? What is the order of bond breaking and bond making? Is the reaction a one-step process or does it occur in many steps?

There are two mechanisms for elimination—E2 and E1—just as there are two mechanisms for nucleophilic substitution—S_N2 and S_N1.

- **The E2 mechanism (bimolecular elimination)**
- **The E1 mechanism (unimolecular elimination)**

The E2 and E1 mechanisms differ in the timing of bond cleavage and bond formation, analogous to the S_N2 and S_N1 mechanisms. In fact, E2 and S_N2 reactions have some features in common, as do E1 and S_N1 reactions.

8.4 The E2 Mechanism

The most common mechanism for dehydrohalogenation is the E2 mechanism. For example, $(CH_3)_3CBr$ reacts with ^-OH to form $(CH_3)_2C{=}CH_2$ via an E2 mechanism.



8.4A Kinetics


An E2 reaction exhibits **second-order kinetics;** that is, the reaction is **bimolecular** and both the alkyl halide and the base appear in the rate equation.

- **rate = k[$(CH_3)_3CBr$][^-OH]**

8.4B A One-Step Mechanism

The most straightforward explanation for the second-order kinetics is a **concerted reaction: all bonds are broken and formed in a single step,** as shown in Mechanism 8.1.


Mechanism 8.1 The E2 Mechanism



- The base $^-$OH removes a proton from the β carbon, forming H_2O (a by-product).
- The electron pair in the β C–H bond forms the new π bond.
- The leaving group Br^- comes off with the electron pair in the C–Br bond.


An energy diagram for the reaction of $(CH_3)_3CBr$ with $^-$OH is shown in Figure 8.3. The reaction has one step, so there is one energy barrier between reactants and products. Two bonds are broken (C–H and C–Br) and two bonds are formed (H–OH and the π bond) in a single step, so the transition state contains four partial bonds, with the negative charge distributed over the base and the leaving group. Entropy favors the products of an E2 reaction because two molecules of starting material form three molecules of product.

Figure 8.3
An energy diagram for an E2 reaction: $(CH_3)_3CBr + {}^-OH \rightarrow (CH_3)_2C{=}CH_2 + H_2O + Br^-$



- In the transition state, the C–H and C–Br bonds are partially broken, the O–H and π bonds are partially formed, and both the base and the departing leaving group bear a partial negative charge.

Problem 8.8 Use curved arrows to show the movement of electrons in the following E2 mechanism. Draw the structure of the transition state.



There are close parallels between the E2 and S_N2 mechanisms in how the identity of the base, the leaving group, and the solvent affect the rate.

The Base

- **The base appears in the rate equation, so the rate of the E2 reaction increases as the strength of the base increases.**

The IUPAC names for **DBN** and **DBU** are rarely used because the names are complex. **DBN** stands for 1,5-**d**iaza**b**icyclo[4.3.0]-**n**on-5-ene, and **DBU** stands for 1,8-**d**iaza**b**icyclo[5.4.0]-**u**ndec-7-ene.

E2 reactions are generally run with strong, negatively charged bases like $^-$OH and $^-$OR. Two strong, sterically hindered nitrogen bases, called **DBN** and **DBU,** are also sometimes used. An example of an E2 reaction with DBN is shown in Figure 8.4.






Figure 8.4 An E2 elimination with DBN used as the base



The Leaving Group

- **Because the bond to the leaving group is partially broken in the transition state, the better the leaving group the faster the E2 reaction.**



The Solvent

- **Polar aprotic solvents increase the rate of E2 reactions.**


Because **polar aprotic solvents** like $(CH_3)_2C{=}O$ do not solvate anions well, a negatively charged base is not "hidden" by strong interactions with the solvent (Section 7.17D), and the base is stronger. **A stronger base increases the reaction rate.**

Problem 8.9 Consider an E2 reaction between CH_3CH_2Br and $KOC(CH_3)_3$. What effect does each of the following changes have on the rate of elimination? (a) The base is changed to KOH. (b) The alkyl halide is changed to CH_3CH_2Cl.

8.4C The Identity of the Alkyl Halide


The S_N2 and E2 mechanisms differ in how the R group affects the reaction rate.

- **As the number of R groups on the carbon with the leaving group increases, the rate of the E2 reaction increases.**



This trend is exactly *opposite* to the reactivity of alkyl halides in S_N2 reactions, where increasing alkyl substitution decreases the rate of reaction (Section 7.11D).


Why does increasing alkyl substitution increase the rate of an E2 reaction? In the transition state, the double bond is partially formed, so increasing the stability of the double bond with alkyl substituents stabilizes the transition state (i.e., it lowers E_a), which increases the rate of the reaction.



The double bond is partially formed.

- **Increasing the number of R groups on the carbon with the leaving group forms more highly substituted, more stable alkenes in E2 reactions.**

For example, the E2 reaction of a 1° alkyl halide (1-bromobutane) forms a monosubstituted alkene, whereas the E2 reaction of a 3° alkyl halide (2-bromo-2-methylpropane) forms a disubstituted alkene. The disubstituted alkene is more stable, so the 3° alkyl halide reacts faster than the 1° alkyl halide.



Problem 8.10 Rank the alkyl halides in each group in order of increasing reactivity in an E2 reaction.




Table 8.2 summarizes the characteristics of the E2 mechanism, and Figure 8.5 illustrates two examples of E2 reactions used in the synthesis of **quinine** and **estradiol,** two naturally occurring compounds.


Table 8.2 Characteristics of the E2 Mechanism

Characteristic	Result
Kinetics	• Second order
Mechanism	• One step
Identity of R	• More substituted halides react faster. • Rate: $R_3CX > R_2CHX > RCH_2X$
Base	• Favored by strong bases
Leaving group	• Better leaving group ⟶ faster reaction
Solvent	• Favored by polar aprotic solvents

Figure 8.5
Two examples of the E2 reaction used in organic synthesis



Quinine, a natural product isolated from the bark of the cinchona tree native to the Andes Mountains, is a powerful antipyretic—that is, it reduces fever—and for centuries, it was the only effective treatment for malaria.




- Bonds and atoms in quinine and estradiol that originate in the alkene intermediate are shown in red.

Problem 8.11 How does each of the following changes affect the rate of an E2 reaction?

a. tripling [RX]
b. halving [B:]
c. changing the solvent from CH_3OH to DMSO
d. changing the leaving group from I^- to Br^-
e. changing the base from ^-OH to H_2O
f. changing the alkyl halide from CH_3CH_2Br to $(CH_3)_2CHBr$

8.5 The Zaitsev Rule

Recall from Section 8.1 that a mixture of alkenes can form from the dehydrohalogenation of alkyl halides having two or more different β carbon atoms. When this occurs, one of the products usually predominates. The **major product is the more stable product—the one with the more substituted double bond.** For example, elimination of the elements of H and I from 1-iodo-1-methylcyclohexane yields two constitutional isomers: the trisubstituted alkene **A** (the major product) and the disubstituted alkene **B** (the minor product).



1-iodo-1-methylcyclohexane

A
major product
trisubstituted alkene


B
minor product
disubstituted alkene

This phenomenon is called the **Zaitsev rule** (also called the **Saytzeff rule,** depending on the translation) for the Russian chemist who first noted this trend.

- **The Zaitsev rule: The major product in β elimination has the more substituted double bond.**

A reaction is *regioselective* when it yields predominantly or exclusively one constitutional isomer when more than one is possible. The E2 reaction is **regioselective** because the more substituted alkene predominates.

The Zaitsev rule results because the double bond is partially formed in the transition state for the E2 reaction. Thus, increasing the stability of the double bond by adding R groups lowers the energy of the transition state, which increases the reaction rate. E2 elimination of HBr from 2-bromo-2-methylbutane yields alkenes **C** and **D. D,** having the more substituted double bond, is the major product, because the transition state leading to its formation is lower in energy.




less stable
transition state

C
minor product
disubstituted alkene

more stable
transition state

D
major product
trisubstituted alkene

When a mixture of stereoisomers is possible from dehydrohalogenation, the **major product is the more stable stereoisomer.** Dehydrohalogenation of alkyl halide **X** forms a mixture of trans and cis alkenes, **Y** and **Z.** The trans alkene **Y** is the major product because it is more stable.




X

Y
trans alkene
major product

Z
cis alkene
minor product


A reaction is *stereoselective* when it forms predominantly or exclusively one stereoisomer when two or more are possible. The **E2 reaction is stereoselective** because one stereoisomer is formed preferentially.

Sample Problem 8.1 Predict the major product in the following E2 reaction.




Solution

The alkyl halide has two different β C atoms (labeled β_1 and β_2), so two different alkenes are possible: one formed by removal of HCl across the α and β_1 carbons, and one formed by removal of HCl across the α and β_2 carbons. Using the Zaitsev rule, the major product should be **A,** because it has the more substituted double bond.





A
trisubstituted alkene
major product





B
disubstituted alkene
minor product

Problem 8.12 What alkenes are formed from each alkyl halide by an E2 reaction? Use the Zaitsev rule to predict the major product.

a. 

b. 

c. 

d. 

8.6 The E1 Mechanism

The dehydrohalogenation of $(CH_3)_3CI$ with H_2O to form $(CH_3)_2C{=}CH_2$ can be used to illustrate the second general mechanism of elimination, the **E1 mechanism.**



8.6A Kinetics

An E1 reaction exhibits first-order kinetics.


- rate = $k[(CH_3)_3CI]$

Like the S_N1 mechanism, the kinetics suggest that the reaction mechanism involves more than one step, and that the slow step is **unimolecular,** involving *only* the alkyl halide.

8.6B A Two-Step Mechanism

The most straightforward explanation for the observed first-order kinetics is a **two-step reaction: the bond to the leaving group breaks first *before* the π bond is formed,** as shown in Mechanism 8.2.

Mechanism 8.2 The E1 Mechanism




1. Heterolysis of the C–I bond forms a carbocation in the rate-determining step.
2. A base (either H_2O or I^-) removes a proton from a carbon adjacent to the carbocation, and the electron pair in the C–H bond forms the π bond.

The E1 and E2 mechanisms both involve the same number of bonds broken and formed. **The only difference is the timing.**

- **In an E1 reaction, the leaving group comes off *before* the β proton is removed, and the reaction occurs in two steps.**
- **In an E2 reaction, the leaving group comes off *as* the β proton is removed, and the reaction occurs in one step.**


An energy diagram for the reaction of $(CH_3)_3CI + H_2O$ is shown in Figure 8.6. Each step has its own energy barrier, with a transition state at each energy maximum. Because its transition state is higher in energy, Step [1] is rate-determining. $\Delta H°$ for Step [1] is positive because only bond breaking occurs, whereas $\Delta H°$ of Step [2] is negative because two bonds are formed and only one is broken.

Figure 8.6
Energy diagram for an E1 reaction: $(CH_3)_3CI + H_2O \rightarrow (CH_3)_2C{=}CH_2 + H_3O^+ + I^-$



- The E1 mechanism has two steps, so there are two energy barriers.
- Step [1] is rate-determining.

Problem 8.13 Draw an E1 mechanism for the following reaction. Draw the structure of the transition state for each step.




8.6C Other Characteristics of E1 Reactions

Three other features of E1 reactions are worthy of note.

[1] The rate of an E1 reaction increases as the number of R groups on the carbon with the leaving group increases.

Increasing alkyl substitution has the same effect on the rate of *both* an E1 and E2 reaction; increasing rate of the E1 and E2 reactions: RCH_2X (1°) < R_2CHX (2°) < R_3CX (3°).



Like an S_N1 reaction, more substituted alkyl halides yield more substituted (and more stable) carbocations in the rate-determining step. Increasing the stability of a carbocation, in turn, decreases E_a for the slow step, which increases the rate of the E1 reaction according to the Hammond postulate.


[2] Because the base does not appear in the rate equation, weak bases favor E1 reactions.

The strength of the base usually determines whether a reaction follows the E1 or E2 mechanism.

- **Strong bases like $^-$OH and $^-$OR favor E2 reactions, whereas weaker bases like H_2O and ROH favor E1 reactions.**

[3] E1 reactions are regioselective, favoring formation of the more substituted, more stable alkene.

The Zaitsev rule applies to E1 reactions, too. For example, E1 elimination of HBr from 1-bromo-1-methylcyclopentane yields alkenes **A** and **B. A,** having the more substituted double bond, is the major product.



1-bromo-1-methylcyclopentane

A
trisubstituted alkene
major product


B
disubstituted alkene
minor product


Table 8.3 summarizes the characteristics of E1 reactions.

Table 8.3 Characteristics of the E1 Mechanism

Characteristic	Result
Kinetics	• First order
Mechanism	• Two steps
Identity of R	• More substituted halides react faster. • Rate: $R_3CX > R_2CHX > RCH_2X$
Base	• Favored by weaker bases such as H_2O and ROH
Leaving group	• A better leaving group makes the reaction faster because the bond to the leaving group is partially broken in the rate-determining step.
Solvent	• Polar protic solvents that solvate the ionic intermediates are needed.

Problem 8.14 What alkenes are formed from each alkyl halide by an E1 reaction? Use the Zaitsev rule to predict the major product.

a. 


b. 

Problem 8.15 How does each of the following changes affect the rate of an E1 reaction?

a. doubling [RX]
b. doubling [B:]
c. changing the halide from $(CH_3)_3CBr$ to $CH_3CH_2CH_2Br$
d. changing the leaving group from Cl^- to Br^-
e. changing the solvent from DMSO to CH_3OH

8.7 S_N1 and E1 Reactions

S_N1 and E1 reactions have exactly the same first step—formation of a carbocation. They differ in what happens to the carbocation.



- In an S_N1 reaction, a nucleophile attacks the carbocation, forming a substitution product.
- In an E1 reaction, a base removes a proton, forming a new π bond.

The same conditions that favor substitution by an S_N1 mechanism also favor elimination by an E1 mechanism: a 3° alkyl halide as substrate, a weak nucleophile or base as reagent, and a polar protic solvent. As a result, both reactions usually occur in the same reaction mixture to afford a mixture of products, as illustrated in Sample Problem 8.2.


Sample Problem 8.2 Draw the S_N1 and E1 products formed in the reaction of $(CH_3)_3CBr$ with H_2O.

Solution

The first step in both reactions is heterolysis of the C–Br bond to form a **carbocation.**




Reaction of the carbocation with H_2O as a nucleophile affords the substitution product (Reaction [1]). Alternatively, H_2O acts as a base to remove a proton, affording the elimination product (Reaction [2]). **Two products are formed.**



Because E1 reactions often occur with a competing S_N1 reaction, **E1 reactions of alkyl halides are *much less useful* than E2 reactions.**

Problem 8.16 Draw both the S_N1 and E1 products of each reaction.




8.8 Stereochemistry of the E2 Reaction


Although the E2 reaction does not produce products with tetrahedral stereogenic centers, its transition state consists of four atoms that react at the same time, and they react only if they possess a particular stereochemical arrangement.

8.8A General Stereochemical Features

The transition state of an E2 reaction consists of **four atoms** from the alkyl halide—one hydrogen atom, two carbon atoms, and the leaving group (X)—**all aligned in a plane.** There are two ways for the C−H and C−X bonds to be coplanar.



syn periplanar



anti periplanar


The dihedral angle for the C−H and C−X bonds equals 0° for the syn periplanar arrangement and 180° for the anti periplanar arrangement.

- **The H and X atoms can be oriented on the same side of the molecule. This geometry is called *syn periplanar.***
- **The H and X atoms can be oriented on opposite sides of the molecule. This geometry is called *anti periplanar.***

All evidence suggests that **E2 elimination occurs most often in the anti periplanar geometry.** This arrangement allows the molecule to react in the lower energy *staggered* conformation. It also allows two electron-rich species, the incoming base and the departing leaving group, to be farther away from each other, as illustrated in Figure 8.7.

Figure 8.7
Two possible geometries for the E2 reaction


a.



preferred geometry

- An **anti periplanar** arrangement has a **staggered** conformation.
- The two electron-rich groups are far apart.

b.




- A **syn periplanar** arrangement has an **eclipsed** conformation.
- The two electron-rich groups are close.


Anti periplanar geometry is the preferred arrangement for any alkyl halide undergoing E2 elimination, regardless of whether it is cyclic or acyclic. This stereochemical requirement has important consequences for compounds containing six-membered rings.

Problem 8.17 Given that an E2 reaction proceeds with anti periplanar stereochemistry, draw the products of each elimination. The alkyl halides in (a) and (b) are diastereomers of each other. How are the products of these two reactions related? Recall from Section 3.2A that C_6H_5- is a phenyl group, a benzene ring bonded to another group.

a.




b.




8.8B Anti Periplanar Geometry and Halocyclohexanes

Recall from Section 4.13 that cyclohexane exists as two chair conformations that rapidly interconvert, and that substituted cyclohexanes are more stable with substituents in the roomier equatorial position. Chlorocyclohexane exists as two chair conformations, but **X** is preferred because the Cl group is equatorial.




For E2 elimination, **the C–Cl bond must be anti periplanar to a C–H bond on a β carbon,** and this occurs only when the H and Cl atoms are both in the **axial** position. This requirement for **trans diaxial geometry** means that E2 elimination must occur from the less stable conformation **Y,** as shown in Figure 8.8.

Figure 8.8
The trans diaxial geometry for the E2 elimination in chlorocyclohexane




- In conformation **X** (**equatorial** Cl group), a β C–H bond and a C–Cl bond are never anti periplanar; therefore, no E2 elimination can occur. β Carbons are highlighted in blue.
- In conformation **Y** (**axial** Cl group), two β C–H bonds and the C–Cl bond are **trans diaxial;** therefore, E2 elimination occurs. Axial H's on β carbons that can react are shown in red.


Sometimes this rigid stereochemical requirement affects the regioselectivity of the E2 reaction of substituted cyclohexanes. Dehydrohalogenation of *cis*- and *trans*-1-chloro-2-methylcyclohexane via an E2 mechanism illustrates this phenomenon.




The **cis isomer** exists as two conformations (**A** and **B**), each of which has one group axial and one group equatorial. E2 reaction must occur from conformation **B,** which contains an axial Cl atom.




Because conformation **B** has two different axial β H atoms, labeled H_a and H_b, E2 reaction occurs in two different directions to afford two alkenes. **The major product contains the more stable trisubstituted double bond, as predicted by the Zaitsev rule.**



The **trans isomer** exists as two conformations, **C,** having two equatorial substituents, and **D,** having two axial substituents. E2 reaction must occur from conformation **D,** which contains an axial Cl atom.




Because conformation **D** has **only one axial β H,** E2 reaction occurs in only one direction to afford a **single product,** having the disubstituted double bond. This is *not* predicted by the Zaitsev rule. E2 reaction requires H and Cl to be trans and diaxial, and with the trans isomer, this is possible only when the less stable alkene is formed as product.




- **In conclusion, with substituted cyclohexanes, E2 elimination must occur with a trans diaxial arrangement of H and X, and as a result of this requirement, the more substituted alkene is not necessarily the major product.**

Sample Problem 8.3 Draw the major E2 elimination product formed from the following alkyl halide.




Solution

To draw the elimination products, locate the β carbons and look for H atoms that are trans to the leaving group. The given alkyl chloride has two different β carbons, labeled β_1 and β_2. Elimination can occur only when the leaving group (Cl) and a H atom on the β carbon are **trans.**



The β_1 C has a H atom trans to Cl, so E2 elimination occurs to form a disubstituted alkene. Because there is no trans H on the β_2 C, E2 elimination cannot occur in this direction, and the more stable trisubstituted alkene is *not* formed. Although this result is not predicted by the Zaitsev rule, it is consistent with the requirement that the H and X atoms in an E2 elimination must be located trans to each other.

Problem 8.18 Draw the major E2 elimination products from each of the following alkyl halides.



Problem 8.19 Explain why *cis*-1-chloro-2-methylcyclohexane undergoes E2 elimination much faster than its trans isomer.

8.9 When Is the Mechanism E1 or E2?

Given a particular starting material and base, how do we know whether a reaction occurs by the E1 or E2 mechanism?

Because the rate of *both* the E1 and E2 reactions increases as the number of R groups on the carbon with the leaving group increases, you cannot use the identity of the alkyl halide to decide which elimination mechanism occurs.


- **The strength of the base is the most important factor in determining the mechanism for elimination. Strong bases favor the E2 mechanism. Weak bases favor the E1 mechanism.**

Table 8.4 compares the E1 and E2 mechanisms.

Table 8.4 A Comparison of the E1 and E2 Mechanisms

Mechanism	Comment
E2 mechanism	• Much more common and useful • Favored by strong, negatively charged bases, especially $^-$OH and $^-$OR • The reaction occurs with 1°, 2°, and 3° alkyl halides. Order of reactivity: $R_3CX > R_2CHX > RCH_2X$.
E1 mechanism	• Much less useful because a mixture of S_N1 and E1 products usually results • Favored by weaker, neutral bases, such as H_2O and ROH • This mechanism does not occur with 1° RX because they form highly unstable 1° carbocations.

Problem 8.20 Which mechanism, E1 or E2, will occur in each reaction?



8.10 E2 Reactions and Alkyne Synthesis

Recall from Section 1.10C that the carbon–carbon triple bond of alkynes consists of one σ and two π bonds.

A single elimination reaction produces the π bond of an alkene. Two consecutive elimination reactions produce the two π bonds of an alkyne.



alkene
one π bond


One elimination reaction is needed.

alkyne
two π bonds

Two elimination reactions are needed.

- **Alkynes are prepared by two successive dehydrohalogenation reactions.**

Two elimination reactions are needed to remove two moles of HX from a **dihalide** as substrate. Two different starting materials can be used.




vicinal dihalide

geminal dihalide

- **A vicinal dihalide has two X atoms on *adjacent* carbon atoms.**
- **A geminal dihalide has two X atoms on the *same* carbon atom.**

The word *geminal* comes from the Latin *geminus,* meaning *twin.*

Equations [1] and [2] illustrate how two moles of HX can be removed from these dihalides with base. Two equivalents of strong base are used and each step follows an E2 mechanism.




The relative strength of C–H bonds depends on the hybridization of the carbon atom: $sp > sp^2 > sp^3$. For more information, review Section 1.11B.


Stronger bases are needed to synthesize alkynes by dehydrohalogenation than are needed to synthesize alkenes. The typical base is **amide ($^-NH_2$),** used as the sodium salt $NaNH_2$ (sodium amide). $KOC(CH_3)_3$ can also be used with DMSO as solvent. Because DMSO is a polar aprotic solvent, the anionic base is not well solvated, thus **increasing its basicity** and making it strong enough to remove two equivalents of HX. Examples are given in Figure 8.9.

The strongly basic conditions needed for alkyne synthesis result from the difficulty of removing the second equivalent of HX from the intermediate vinyl halide, RCH=C(R)X. Because H and X are both bonded to sp^2 hybridized carbons, these bonds are shorter and stronger than the sp^3 hybridized C–H and C–X bonds of an alkyl halide, necessitating the use of a stronger base.

Figure 8.9
Examples of dehydrohalogenation of dihalides to afford alkynes



Problem 8.21 Draw the alkynes formed when each dihalide is treated with excess base.




8.11 When Is the Reaction S_N1, S_N2, E1, or E2?

We have now considered two different kinds of reactions (substitution and elimination) and four different mechanisms (S_N1, S_N2, E1, and E2) that begin with one class of compounds (alkyl halides). How do we know if a given alkyl halide will undergo substitution or elimination with a given base or nucleophile, and by what mechanism?

Unfortunately, there is no easy answer, and often mixtures of products result. Two generalizations help to determine whether substitution or elimination occurs.


[1] Good nucleophiles that are weak bases favor substitution over elimination.

Certain anions generally give products of substitution because they are good nucleophiles but weak bases. These include: **I^-, Br^-, HS^-, ^-CN,** and **$CH_3CO_2^-$.**



[2] Bulky, nonnucleophilic bases favor elimination over substitution.

$KOC(CH_3)_3$, DBU, and **DBN** are too sterically hindered to attack a tetravalent carbon, but are able to remove a small proton, favoring elimination over substitution.



Most often, however, we will have to rely on other criteria to predict the outcome of these reactions. To determine the product of a reaction with an alkyl halide:

[1] Classify the alkyl halide as 1°, 2°, or 3°.

[2] Classify the base or nucleophile as strong, weak, or bulky.

Predicting the substitution and elimination products of a reaction can then be organized by the type of alkyl halide, as summarized in Table 8.5. The explanation that follows the table is organized with 2° alkyl halides last, because their reactions can follow any of the four mechanisms and product mixtures often result.


Table 8.5 Summary of Alkyl Halides and S_N1, S_N2, E1, and E2 Mechanisms

Alkyl halide type	Reaction with	Mechanism
1° RCH_2X	• Strong nucleophile	S_N2
	• Strong bulky base	E2
2° R_2CHX	• Strong base and nucleophile	S_N2 and E2
	• Strong bulky base	E2
	• Weak base and nucleophile	S_N1 and E1
3° R_3CX	• Weak base and nucleophile	S_N1 and E1
	• Strong base	E2

8.11A Tertiary Alkyl Halides


Tertiary alkyl halides react by all mechanisms *except* S_N2.

- **With strong bases, elimination occurs by an E2 mechanism.**



A strong base or nucleophile favors an S_N2 or E2 mechanism, but 3° halides are too sterically hindered to undergo an S_N2 reaction, so only E2 elimination occurs.

- **With weak nucleophiles or bases, a mixture of S_N1 and E1 products results.**



A weak base or nucleophile favors S_N1 and E1 mechanisms and both occur.

8.11B Primary Alkyl Halides


Primary alkyl halides react by S_N2 and E2 mechanisms.

- With strong nucleophiles, substitution occurs by an S_N2 mechanism.



A strong base or nucleophile favors S_N2 or E2, but 1° halides are the least reactive halide type in elimination, so only S_N2 reaction occurs.

- With strong, bulky bases, elimination occurs by an E2 mechanism.




A strong, bulky base cannot act as a nucleophile, so elimination occurs and the mechanism is E2.

8.11C Secondary Alkyl Halides


Secondary alkyl halides react by *all* mechanisms.

- With strong bases and nucleophiles, a mixture of S_N2 and E2 products results.



A strong base that is also a strong nucleophile gives a mixture of S_N2 and E2 products.

- With strong, bulky bases, elimination occurs by an E2 mechanism.



A strong, bulky base cannot act as a nucleophile, so elimination occurs and the mechanism is E2.


- With weak nucleophiles or bases, a mixture of S_N1 and E1 products results.



A weak base or nucleophile favors S_N1 and E1 mechanisms and both occur.


Sample Problems 8.4–8.6 illustrate how to apply the information in Table 8.5 to specific alkyl halides.

Sample Problem 8.4 Draw the products of the following reaction.



Solution


- Classify the halide as 1°, 2°, or 3° and the reagent as a strong or weak base (and nucleophile) to determine the mechanism. In this case, the alkyl halide is 3° and the reagent (H_2O) is a weak base and nucleophile, so products of both **S_N1** and **E1** mechanisms are formed.
- To draw the S_N1 product, substitute the nucleophile (H_2O) for the leaving group (Br^-), and draw the neutral product after loss of a proton.
- To draw the E1 product, remove the elements of H and Br from the α and β carbons. There are two identical β C atoms with H atoms, so only one elimination product is possible.



S_N1 product **E1 product**


Sample Problem 8.5

Draw the products of the following reaction.




Solution

- Classify the halide as 1°, 2°, or 3° and the reagent as a strong or weak base (and nucleophile) to determine the mechanism. In this case, the alkyl halide is 2° and the reagent (CH_3O^-) is a strong base and nucleophile, so products of both **S_N2** and **E2** mechanisms are formed.
- To draw the S_N2 product, substitute the nucleophile (CH_3O^-) for the leaving group (Br^-).
- To draw the E2 product, remove the elements of H and Br from the α and β carbons. There are two identical β C atoms with H atoms, so only one elimination product is possible.




S_N2 product



E2 product

Sample Problem 8.6


Draw the products of the following reaction, and include the mechanism showing how each product is formed.



Solution


[1] Classify the halide as 1°, 2°, or 3° and the reagent as a strong or weak base (and nucleophile) to determine the mechanism. In this case, the alkyl halide is 3° and the reagent (CH_3OH) is a weak base and nucleophile, so products of both **S_N1** and **E1** mechanisms are formed.

[2] Draw the steps of the mechanisms to give the products. Both mechanisms begin with the same first step: loss of the leaving group to form a carbocation.




carbocation

- **For S_N1:** The carbocation reacts with a nucleophile. Nucleophilic attack of CH_3OH on the carbocation generates a positively charged intermediate that loses a proton to afford the neutral S_N1 product.




- **For E1:** The carbocation reacts with a base (CH_3OH or Br^-). Two different products of elimination can form because the carbocation has two different β carbons.




In this problem, three products are formed: one from an S_N1 reaction and two from E1 reactions.

Problem 8.22 Draw the products in each reaction.



Problem 8.23 Draw a stepwise mechanism for the following reaction.




KEY CONCEPTS


Alkyl Halides and Elimination Reactions

A Comparison Between Nucleophilic Substitution and β Elimination

Nucleophilic substitution—A nucleophile attacks a carbon atom (7.6).



β Elimination—A base attacks a proton (8.1).




Similarities	Differences
• In both reactions RX acts as an electrophile, reacting with an electron-rich reagent. • Both reactions require a **good leaving group X:⁻** that can accept the electron density in the C – X bond.	• In substitution, a nucleophile attacks a single carbon atom. • In elimination, a Brønsted–Lowry base removes a proton to form a π bond, and two carbons are involved in the reaction.

The Importance of the Base in E2 and E1 Reactions (8.9)

The strength of the base determines the mechanism of elimination.

- Strong bases favor E2 reactions.
- Weak bases favor E1 reactions.
- The same organic product can form in two eliminations by different mechanisms.



E1 and E2 Mechanisms Compared

	E2 mechanism	E1 mechanism
Mechanism	• One step (8.4B)	• Two steps (8.6B)
Alkyl halide	• Rate: $R_3CX > R_2CHX > RCH_2X$ (8.4C)	• Rate: $R_3CX > R_2CHX > RCH_2X$ (8.6C)
Rate equation	• Rate = k[RX][B:]	• Rate = k[RX]
	• Second-order kinetics (8.4A)	• First-order kinetics (8.6A)
Stereochemistry	• Anti periplanar arrangement of H and X (8.8)	• Trigonal planar carbocation intermediate (8.6B)
Base	• Favored by strong bases (8.4B)	• Favored by weak bases (8.6C)
Leaving group	• Better leaving group ⇢ faster reaction (8.4B)	• Better leaving group ⇢ faster reaction (Table 8.3)
Solvent	• Favored by polar aprotic solvents (8.4B)	• Favored by polar protic solvents (Table 8.3)
Product	• More substituted alkene favored (Zaitsev rule, 8.5)	• More substituted alkene favored (Zaitsev rule, 8.6C)


Summary Chart on the Four Mechanisms: S_N1, S_N2, E1, or E2 (8.11)

Alkyl halide type	Conditions		Mechanism
1° RCH_2X	strong nucleophile	⇢	**S_N2**
	strong bulky base	⇢	**E2**
2° R_2CHX	strong base and nucleophile	⇢	**S_N2 + E2**
	strong bulky base	⇢	**E2**
	weak base and nucleophile	⇢	**S_N1 + E1**
3° R_3CX	weak base and nucleophile	⇢	**S_N1 + E1**
	strong base	⇢	**E2**


PROBLEMS

Problems Using Three-Dimensional Models


8.24 Rank the alkenes shown in the ball-and-stick models (**A**–**C**) in order of increasing stability.



A




B



C

8.25 Name each compound and decide which stereoisomer will react faster in an E2 elimination reaction. Explain your choice.




D




E

8.26 What is the major E2 elimination product formed from each alkyl halide?

a.




b.




General Elimination

8.27 Draw all possible constitutional isomers formed by dehydrohalogenation of each alkyl halide.


a.




b.



c.




d.




8.28 What alkyl halide forms each of the following alkenes as the *only* product in an elimination reaction?


a.




b.



c.




d.



Alkenes


8.29 Which double bonds in the following natural products can exhibit stereoisomerism? Nerolidol is isolated from the angel's trumpet plant, caryophyllene is present in hemp, and humulene comes from hops.

a.




nerolidol

b.





caryophyllene


c.







humulene

8.30 Label each pair of alkenes as constitutional isomers, stereoisomers, or identical.


a.  and 

c.  and 


b.  and 

d.  and 


8.31 Rank the following alkenes in order of increasing stability:



A




B




C

8.32 $\Delta H°$ values obtained for a series of similar reactions are one set of experimental data used to determine the relative stability of alkenes. Explain how the following data suggest that *cis*-but-2-ene is more stable than but-1-ene (Section 12.3A).




but-1-ene





cis-but-2-ene


E2 Reaction

8.33 Draw all constitutional isomers formed in each E2 reaction and predict the major product using the Zaitsev rule.


a. 


c. 


b. 

d. 


8.34 For each of the following alkenes, draw the structure of two different alkyl halides that yield the given alkene as the only product of dehydrohalogenation.

a. 

b. 

c. 


8.35 Consider the following E2 reaction.



a. Draw the by-products of the reaction and use curved arrows to show the movement of electrons.


b. What happens to the reaction rate with each of the following changes? [1] The solvent is changed to DMF. [2] The concentration of $^{-}OC(CH_3)_3$ is decreased. [3] The base is changed to ^{-}OH. [4] The halide is changed to $CH_3CH_2CH_2CH_2CH(Br)CH_3$. [5] The leaving group is changed to I^{-}.

8.36 Dehydrohalogenation of 1-chloro-1-methylcyclopropane affords two alkenes (**A** and **B**) as products. Explain why **A** is the major product despite the fact that it contains the less substituted double bond.




1-chloro-1-methyl-cyclopropane

8.37 What is the major stereoisomer formed when each alkyl halide is treated with $KOC(CH_3)_3$?




E1 Reaction

8.38 What alkene is the major product formed from each alkyl halide in an E1 reaction?




E1 and E2

8.39 Draw all constitutional isomers formed in each elimination reaction. Label the mechanism as E2 or E1.



8.40 Rank the alkyl halides in order of increasing E2 reactivity. Then do the same for E1 reactivity.



A **B** **C**

8.41 Pick the reactant or solvent in each part that gives the faster elimination reaction.


a. reaction of ^-OH with 1-chloro-1-methylcyclohexane or 1-chloro-3-methylcyclohexane
b. reaction of H_2O with $CH_3CH(Cl)CH_2CH_3$ or $(CH_3)_2C(Cl)CH_2CH_3$
c. reaction of $(CH_3)_3CCl$ with ^-OH in H_2O or DMSO

8.42 In the dehydrohalogenation of bromocyclodecane, the major product is *cis*-cyclodecene rather than *trans*-cyclodecene. Offer an explanation.


8.43 Elimination of HBr from 2-bromobutane affords a mixture of but-1-ene and but-2-ene. With sodium ethoxide as base, but-2-ene constitutes 81% of the alkene products, but with potassium *tert*-butoxide, but-2-ene constitutes only 67% of the alkene products. Offer an explanation for this difference.


Stereochemistry and the E2 Reaction

8.44 What is the major E2 elimination product formed from each halide?



8.45 Taking into account anti periplanar geometry, predict the major E2 product formed from each starting material.


a. 

b. 

8.46 Does *cis*- or *trans*-1-bromo-4-*tert*-butylcylohexane react faster in an E2 reaction?


8.47 a. Draw three-dimensional representations for all stereoisomers of 2-chloro-3-methylpentane, and label pairs of enantiomers.
b. Considering dehydrohalogenation across C2 and C3 only, draw the E2 product that results from each of these alkyl halides. How many different products have you drawn?
c. How are these products related to each other?

8.48 Explain why compound **A** does not undergo an E2 elimination with strong base.




A


8.49 What alkyl chloride affords the following alkene exclusively under E2 reaction conditions?





Alkynes

8.50 Draw the products formed when each dihalide is treated with excess $NaNH_2$.

a. 


b. 


c. 

d. 

8.51 Draw the structure of a dihalide that could be used to prepare each alkyne. There may be more than one possible dihalide.

a.

b. 

c. 


8.52 Under certain reaction conditions, 2,3-dibromobutane reacts with two equivalents of base to give three products, each of which contains two new π bonds. Product **A** has two *sp* hybridized carbon atoms, product **B** has one *sp* hybridized carbon atom, and product **C** has none. What are the structures of **A, B,** and **C?**

S_N1, S_N2, E1, and E2 Mechanisms


8.53 For which reaction mechanisms—S_N1, S_N2, E1, or E2—are each of the following statements true? A statement may be true for one or more mechanisms.
a. The mechanism involves carbocation intermediates.
b. The mechanism has two steps.
c. The reaction rate increases with better leaving groups.
d. The reaction rate increases when the solvent is changed from CH_3OH to $(CH_3)_2SO$.
e. The reaction rate depends on the concentration of the alkyl halide only.
f. The mechanism is concerted.

g. The reaction of CH_3CH_2Br with NaOH occurs by this mechanism.
h. Racemization at a stereogenic center occurs.
i. Tertiary (3°) alkyl halides react faster than 2° or 1° alkyl halides.
j. The reaction follows a second-order rate equation.


8.54 Draw the organic products formed in each reaction.




8.55 What is the major product formed when each alkyl halide is treated with each of the following reagents: [1] $NaOCOCH_3$; [2] $NaOCH_3$; [3] $KOC(CH_3)_3$? If it is not possible to predict the major product, identify the products in the mixture and the mechanism by which each is formed.




8.56 Draw all products, including stereoisomers, in each reaction.




8.57 

Draw all of the substitution and elimination products formed from the given alkyl halide with each reagent: (a) CH_3OH; (b) KOH. Indicate the stereochemistry around the stereogenic centers present in the products, as well as the mechanism by which each product is formed.

8.58 The following reactions do not afford the major product that is given. Explain why this is so, and draw the structure of the major product actually formed.




8.59 Draw a stepwise, detailed mechanism for each reaction.



8.60 Draw the major product formed when (*R*)-1-chloro-3-methylpentane is treated with each reagent: (a) $NaOCH_2CH_3$; (b) KCN; (c) DBU.


8.61 Draw a stepwise, detailed mechanism for the following reaction.



8.62 Explain why the reaction of 2-bromopropane with $NaOCOCH_3$ gives $(CH_3)_2CHOCOCH_3$ exclusively as product, but the reaction of 2-bromopropane with $NaOCH_2CH_3$ gives a mixture of $(CH_3)_2CHOCH_2CH_3$ (20%) and $CH_3CH=CH_2$ (80%).


Challenge Problems

8.63 Draw a stepwise detailed mechanism that illustrates how four organic products are formed in the following reaction.




8.64 Although there are nine stereoisomers of 1,2,3,4,5,6-hexachlorocyclohexane, one stereoisomer reacts 7000 times more slowly than any of the others in an E2 elimination. Draw the structure of this isomer and explain why this is so.


8.65 Explain the selectivity observed in the following reactions.




8.66 Draw a stepwise mechanism for the following reaction. The four-membered ring in the starting material and product is called a β-lactam. This functional group confers biological activity on penicillin and many related antibiotics, as is discussed in Chapter 22. (Hint: The mechanism begins with β elimination and involves only two steps.)



8.67 Although dehydrohalogenation occurs with anti periplanar geometry, some eliminations have syn periplanar geometry. Examine the starting material and product of each elimination, and state whether the elimination occurs with syn or anti periplanar geometry.



8.68 (a) Draw all products formed by treatment of each dibromide (**A** and **B**) with one equivalent of $NaNH_2$. (b) Label pairs of diastereomers and constitutional isomers.



A **B**

Alcohols, Ethers, and Related Compounds

9



Linalool is a 10-carbon alcohol found in a wide variety of scented herbs, laurels, and citrus fruits. The *R* enantiomer is one of the two main components of lavender oil, while the *S* enantiomer is found in coriander and sweet orange flowers. Because of its pleasant odor, linalool is used commercially in scented soaps and lotions, and as an insecticide for controlling fleas and cockroaches. In Chapter 9, we learn about alcohols like linalool, as well as related oxygen- and sulfur-containing functional groups.


9.1 Introduction
9.2 Structure and bonding
9.3 Nomenclature
9.4 Physical properties
9.5 Interesting alcohols, ethers, and epoxides
9.6 Preparation of alcohols, ethers, and epoxides
9.7 General features—Reactions of alcohols, ethers, and epoxides
9.8 Dehydration of alcohols to alkenes
9.9 Carbocation rearrangements
9.10 Dehydration using $POCl_3$ and pyridine
9.11 Conversion of alcohols to alkyl halides with HX
9.12 Conversion of alcohols to alkyl halides with $SOCl_2$ and PBr_3
9.13 Tosylate—Another good leaving group
9.14 Reaction of ethers with strong acid
9.15 Thiols and sulfides
9.16 Reactions of epoxides
9.17 Application: Epoxides, leukotrienes, and asthma
9.18 Application: Benzo[*a*]-pyrene, epoxides, and cancer

In Chapter 9, we take the principles learned in Chapters 7 and 8 about leaving groups, nucleophiles, and bases, and apply them to **alcohols, ethers,** and **epoxides,** three new functional groups that contain polar C–O bonds. In the process, you will discover that all of the reactions in Chapter 9 follow one of the four mechanisms introduced in Chapters 7 and 8—S_N1, S_N2, E1, or E2—so there are **no new general mechanisms to learn.**

Later in the chapter, we will also examine **thiols (RSH)** and **sulfides (R_2S),** sulfur analogues of alcohols and ethers, respectively. These functional groups play a key role in the chemistry of biomolecules, especially the proteins discussed in Chapter 29.


9.1 Introduction

Alcohols, ethers, and **epoxides** are three functional groups that contain carbon–oxygen σ bonds.




alcohol **ether** **epoxide**

Alcohols **contain a hydroxy group** (**OH** group) bonded to an sp^3 hybridized carbon atom. As we learned in Section 3.2, alcohols are classified as **primary (1°), secondary (2°),** or **tertiary (3°)** based on the number of carbon atoms bonded to the carbon with the OH group.



alcohol

C is sp^3 hybridized.



cortisol

anti-inflammatory steroid


Compounds having a hydroxy group on an sp^2 hybridized carbon atom—**enols** and **phenols**—undergo different reactions than alcohols and are discussed in Chapters 11 and 19, respectively. **Enols** have an OH group on a carbon of a C−C double bond. **Phenols** have an OH group on a benzene ring.



enol phenol

• C bonded to OH is sp^2 hybridized.

Ethers **have two alkyl groups bonded to an oxygen atom.** An ether is **symmetrical** if the two alkyl groups are the same, and **unsymmetrical** if they are different. Both alcohols and ethers are organic derivatives of H_2O, formed by replacing one or both of the hydrogens on the oxygen atom by R groups, respectively.




ether symmetrical ether identical R groups unsymmetrical ether different R groups

***Epoxides* are ethers having the oxygen atom in a three-membered ring.** Epoxides are also called **oxiranes.**



epoxide or oxirane

Problem 9.1 Label each ether and alcohol in brevenal, a marine natural product. Classify each alcohol as 1°, 2°, or 3°.




brevenal




Brevenal is a nontoxic marine polyether produced by *Karenia brevis,* a single-celled organism that proliferates during red tides, vast algal blooms that turn the ocean water red, brown, or green.

9.2 Structure and Bonding

Alcohols, ethers, and epoxides each contain an oxygen atom surrounded by two atoms and two nonbonded electron pairs, making the O atom **tetrahedral** and sp^3 hybridized. Because only two of the four groups around O are atoms, alcohols and ethers have a **bent** shape like H_2O.




The bond angle around the O atom in an alcohol or ether is similar to the tetrahedral bond angle of **109.5°.** In contrast, the C−O−C bond angle of an epoxide must be **60°,** a considerable deviation from the tetrahedral bond angle. For this reason, **epoxides have angle strain,** making them much more reactive than other ethers.



a strained, three-membered ring

Because oxygen is much more electronegative than carbon or hydrogen, the C−O and O−H bonds are all polar, with the O atom electron rich and the C and H atoms electron poor. The electrostatic potential maps in Figure 9.1 show these polar bonds for all three functional groups.

Figure 9.1
Electrostatic potential maps for a simple alcohol, ether, and epoxide



- Electron-rich regions are shown by the red around the O atoms.

9.3 Nomenclature


To name an alcohol, ether, or epoxide using the IUPAC system, we must learn how to name the functional group either as a substituent or by using a suffix added to the parent name.

9.3A Naming Alcohols


- **In the IUPAC system, alcohols are identified by the suffix *-ol.***

How To Name an Alcohol Using the IUPAC System

Example Give the IUPAC name of the following alcohol:



Step [1] Find the longest carbon chain containing the carbon bonded to the OH group.




6 C's in the longest chain

6 C's → hexane → hexanol

- Change the ***-e*** ending of the parent alkane to the suffix ***-ol.***

Step [2] Number the carbon chain to give the OH group the lower number, and apply all other rules of nomenclature.

a. **Number** the chain.




- Number the chain to put the OH group at C3, not C4.

hexan-3-ol

b. **Name** and **number** the substituents.

methyl at C5




Answer: 5-methylhexan-3-ol

$CH_3CH_2CH_2CH_2OH$ is named as 1-butanol using the 1979 IUPAC recommendations and butan-1-ol using the 1993 IUPAC recommendations.


When an OH group is bonded to a ring, the **ring is numbered beginning with the OH group.** Because the functional group is always at C1, the "1" is usually omitted from the name. The ring is then numbered in a clockwise or counterclockwise fashion to give the next substituent the lower number. Representative examples are given in Figure 9.2.

Figure 9.2
Examples: Naming cyclic alcohols



3-methylcyclohexanol

[The OH group is at C1; the second substituent (CH_3) gets the lower number.]




2,5,5-trimethylcyclohexanol

[The OH group is at C1; the second substituent (CH_3) gets the lower number.]


Common names are often used for simple alcohols. To assign a common name:

- Name all the carbon atoms of the molecule as a single **alkyl group.**
- Add the word **alcohol,** separating the words with a space.




isopropyl alcohol


Compounds with two hydroxy groups are called **diols** (using the IUPAC system) or **glycols.** Compounds with three hydroxy groups are called **triols,** and so forth. To name a diol, for example, the suffix ***-diol*** is added to the name of the parent alkane, and numbers are used to indicate the location of the two OH groups.



ethylene glycol
(ethane-1,2-diol)



glycerol
(propane-1,2,3-triol)





***trans*-cyclopentane-1,2-diol**


Common names are usually used for these simple compounds.

Numbers are needed to show the location of **two** OH groups.

Problem 9.2 Give the IUPAC name for each compound.

a. 

b. 


c. 

Problem 9.3 Give the structure corresponding to each name.

a. 7,7-dimethyloctan-4-ol
b. 5-methyl-4-propylheptan-3-ol
c. 2-*tert*-butyl-3-methylcyclohexanol
d. *trans*-cyclohexane-1,2-diol


9.3B Naming Ethers

Simple ethers are usually assigned common names. To do so, **name both alkyl groups** bonded to the oxygen, arrange these names alphabetically, and add the word **ether.** For symmetrical ethers, name the alkyl group and add the prefix **di-.**



***sec*-butyl methyl ether**


[Alphabetize the **b** of **b**utyl before the **m** of **m**ethyl.]




diethyl ether

More complex ethers are named using the IUPAC system. One alkyl group is named as a hydrocarbon chain, and the other is named as part of a substituent bonded to that chain.

- Name the simpler alkyl group + O atom as an **alkoxy** substituent by changing the ***-yl*** ending of the alkyl group to ***-oxy.***
- Name the remaining alkyl group as an alkane, with the alkoxy group as a substituent bonded to this chain.




Sample Problem 9.1 Give the IUPAC name for the following ether.




Solution

[1] Name the longer chain as an alkane and the shorter chain as an alkoxy group.




[2] Apply the other nomenclature rules to complete the name.



Answer: 4-ethoxyoctane

Problem 9.4 Name each of the following ethers.





tetrahydrofuran
THF


Cyclic ethers have an O atom in a ring. A common cyclic ether is **tetrahydrofuran (THF),** a polar aprotic solvent used in nucleophilic substitution (Section 7.8C) and many other organic reactions.

9.3C Naming Epoxides

Any cyclic compound containing a heteroatom is called a *heterocycle.*

Epoxides are named in three different ways—**epoxyalkanes, oxiranes,** or **alkene oxides.**

To name an epoxide as an **epoxyalkane,** first name the alkane chain or ring to which the oxygen is attached, and use the prefix ***epoxy*** to name the epoxide as a substituent. Use two numbers to designate the location of the atoms to which the O's are bonded.



1,2-epoxycyclohexane **1,2-epoxy-2-methylpropane** ***cis*-2,3-epoxypentane**

Epoxides bonded to a chain of carbon atoms can also be named as derivatives of **oxirane,** the simplest epoxide having two carbons and one oxygen atom in a ring. The oxirane ring is numbered to put the O atom at position "1," and the first substituent at position "2." No number is used for a substituent in a monosubstituted oxirane.



oxirane **2,2-dimethyloxirane**


Epoxides are also named as **alkene oxides,** because they are often prepared by adding an O atom to an alkene (Chapter 12). To name an epoxide this way, mentally replace the epoxide oxygen by a double bond, name the alkene (Section 10.3), and then add the word ***oxide.*** For example, the common name for oxirane is ethylene oxide, because it is an epoxide derived from the alkene ethylene. We will use this method of naming epoxides after the details of alkene nomenclature are presented in Chapter 10.




ethylene **ethylene oxide** oxirane

Problem 9.5 Name each epoxide.

a.  (two ways)


b. 

c. 

(two ways)

9.4 Physical Properties

Alcohols, ethers, and epoxides exhibit dipole–dipole interactions because they have a bent structure with two polar bonds. **Alcohols are also capable of intermolecular hydrogen bonding,** because they possess a hydrogen atom on an oxygen, making alcohols much more polar than ethers and epoxides.



hydrogen bond

Steric factors affect the extent of hydrogen bonding. Although all alcohols can hydrogen bond, increasing the number of R groups around the carbon atom bearing the OH group decreases the extent of hydrogen bonding. Thus, 3° alcohols are least able to hydrogen bond, whereas 1° alcohols are most able to.







Table 9.1 Physical Properties of Alcohols, Ethers, and Epoxides

Property	Observation
Boiling point and melting point	• For compounds of comparable molecular weight, the **stronger** the intermolecular forces, the **higher** the bp or mp. 
	• Bp's **increase** as the extent of hydrogen bonding **increases.** 
Solubility	• Alcohols, ethers, and epoxides having ≤ 5 C's are H_2O soluble because they each have an oxygen atom capable of hydrogen bonding to H_2O (Section 3.4C). • Alcohols, ethers, and epoxides having > 5 C's are H_2O insoluble because the nonpolar alkyl portion is too large to dissolve in H_2O. • Alcohols, ethers, and epoxides of any size are soluble in organic solvents.

Key: VDW = van der Waals forces; DD = dipole–dipole; HB = hydrogen bonding

How these factors affect the physical properties of alcohols, ethers, and epoxides is summarized in Table 9.1.

Problem 9.6 Rank the following compounds in order of increasing boiling point.




9.5 Interesting Alcohols, Ethers, and Epoxides

A large number of alcohols, ethers, and epoxides have interesting and useful properties.

9.5A Interesting Alcohols


The structure and properties of three simple alcohols—methanol, propan-2-ol, and ethylene glycol—are given in Figure 9.3. **Ethanol** (CH_3CH_2OH), formed by the fermentation of the carbohydrates in grains, grapes, and potatoes, is the alcohol present in alcoholic beverages. It is perhaps the first organic compound synthesized by humans, because alcohol production has been known for at least 4000 years. Ethanol depresses the central nervous system, increases the production of stomach acid, and dilates blood vessels, producing a flushed appearance. Ethanol is also a common laboratory solvent, which is sometimes made unfit to ingest by adding small amounts of benzene or methanol (both of which are toxic).

Figure 9.3
Some simple alcohols




CH_3OH

- **Methanol (CH_3OH)** is also called wood alcohol, because it can be obtained by heating wood at high temperatures in the absence of air. Methanol is extremely toxic because of the oxidation products formed when it is metabolized in the liver (Section 12.14). Ingestion of as little as 15 mL causes blindness, and 100 mL causes death.



$(CH_3)_2CHOH$

- **Propan-2-ol [$(CH_3)_2CHOH$]** is the major component of rubbing alcohol. When rubbed on the skin it evaporates readily, producing a pleasant cooling sensation. Because it has weak antibacterial properties, propan-2-ol is used to clean skin before minor surgery and to sterilize medical instruments.




$HOCH_2CH_2OH$

- **Ethylene glycol ($HOCH_2CH_2OH$)** is the major component of antifreeze. It is readily prepared from ethylene oxide by reactions discussed in Section 9.16. It is sweet tasting but toxic.

Ethanol is a common gasoline additive, widely touted as an environmentally friendly fuel source. Two common gasoline–ethanol fuels are gasohol, which contains 10% ethanol, and E-85, which contains 85% ethanol. Ethanol is now routinely prepared from the carbohydrates in corn (Figure 9.4). Starch, a complex carbohydrate polymer, can be hydrolyzed to the simple sugar glucose, which forms ethanol by the process of fermentation. Combining ethanol with gasoline forms a usable fuel, which combusts to form CO_2, H_2O, and a great deal of energy.

Since green plants use sunlight to convert CO_2 and H_2O to carbohydrates during photosynthesis, next year's corn crop removes CO_2 from the atmosphere to make new molecules of starch as the corn grows. While in this way ethanol is a renewable fuel source, the need for large-scale farm equipment and the heavy reliance on fertilizers and herbicides make ethanol expensive to produce. Moreover, many criticize the use of valuable farmland for an energy-producing crop rather than for food production. As a result, discussion continues on ethanol as an alternative to fossil fuels.

Figure 9.4 Ethanol from corn, a renewable fuel source




- Hydrolysis of amylose (one form of starch) and fermentation of the resulting simple sugars (Step [1]) yield ethanol, which is mixed with hydrocarbons from petroleum refining (Step [2]) to form usable fuels.
- Combustion of this ethanol–hydrocarbon fuel forms CO_2 and releases a great deal of energy (Step [3]).
- Photosynthesis converts atmospheric CO_2 back to plant carbohydrates in Step [4], and the cycle continues.

9.5B Interesting Ethers

The discovery that **diethyl ether** ($CH_3CH_2OCH_2CH_3$) is a general anesthetic revolutionized surgery in the nineteenth century. For years, a heated controversy existed over who first discovered diethyl ether's anesthetic properties and recognized the enormous benefit in its use. Early experiments were performed by a dentist, Dr. William Morton, resulting in a public demonstration of diethyl ether as an anesthetic in Boston in 1846. In fact, Dr. Crawford Long, a Georgia physician, had been using diethyl ether in surgery and obstetrics for several years, but had not presented his findings to a broader audience.


Diethyl ether is an imperfect anesthetic, but considering the alternatives in the nineteenth century, it was a miracle drug. It is safe, easy to administer, and causes little patient mortality, but it is highly flammable and causes nausea in many patients. For these reasons, it has largely been replaced by sevoflurane and other halogenated ethers, which are non-flammable and cause little patient discomfort.



sevoflurane

Recall from Section 3.7B that crown ethers are named as ***x*-crown-*y***, where ***x*** is the total number of atoms in the ring and ***y*** is the number of O atoms.

Recall from Section 3.7B that some cyclic **polyethers**—compounds with two or more ether linkages—contain cavities that can complex specific-sized cations. For example, 18-crown-6 binds K^+, whereas 12-crown-4 binds Li^+.




18-crown-6 complex with K^+ complex with Li^+ 12-crown-4

- **A crown ether–cation complex is called a *host–guest* complex. The crown ether is the *host* and the cation is the *guest.***
- **The ability of a host molecule to bind specific guests is called *molecular recognition.***

The ability of crown ethers to complex cations can be exploited in nucleophilic substitution reactions, as shown in Figure 9.5. Nucleophilic substitution reactions are usually run in polar solvents to dissolve both the polar organic substrate and the ionic nucleophile. With a crown ether, though, the reaction can be run in a nonpolar solvent under conditions that enhance nucleophilicity.

Figure 9.5 The use of crown ethers in nucleophilic substitution reactions

KCN is insoluble in nonpolar solvents alone, but with 18-crown-6:



18-crown-6 host–guest complex soluble in nonpolar solvents


A rapid nucleophilic substitution reaction occurs in nonpolar solvents when a crown ether is added.

When 18-crown-6 is added to the reaction of CH_3CH_2Br with KCN, for example, the crown ether forms a tight complex with K^+ that has nonpolar C–H bonds on the outside, making the complex soluble in nonpolar solvents like benzene (C_6H_6) or hexane. When the crown ether/K^+ complex dissolves in the nonpolar solvent, it carries the ^-CN along with it to maintain electrical neutrality. The result is a solution of tightly complexed cation and relatively unsolvated anion (nucleophile). The anion, therefore, is extremely nucleophilic because it is not hidden from the substrate by solvent molecules.


Problem 9.7 Which mechanism is favored by the use of crown ethers in nonpolar solvents, S_N1 or S_N2?

9.5C Interesting Epoxides

Although epoxides occur less widely in natural products than alcohols or ethers, interesting and useful epoxides are also known. Examples include two recently introduced drugs, eplerenone and tiotropium bromide. Eplerenone (trade name Inspra) is prescribed to reduce cardiovascular risk in patients who have already had a heart attack. Tiotropium bromide (trade name Spiriva) is a long-acting bronchodilator used to treat the chronic obstructive pulmonary disease of smokers and those routinely exposed to secondhand smoke.




eplerenone



tiotropium bromide


9.6 Preparation of Alcohols, Ethers, and Epoxides

Alcohols and ethers are both common products of nucleophilic substitution. They are synthesized from alkyl halides by S_N2 reactions using strong nucleophiles. As in all S_N2 reactions, highest yields of products are obtained with unhindered methyl and 1° alkyl halides.




The preparation of ethers by this method is called the **Williamson ether synthesis,** and, although it was first reported in the 1800s, it is still the most general method to prepare an ether. Unsymmetrical ethers can be synthesized in two different ways, but often one path is preferred.


For example, ethyl isopropyl ether can be prepared from $CH_3CH_2O^-$ and 2-bromopropane (Path [a]), or from $(CH_3)_2CHO^-$ and bromoethane (Path [b]). Because the mechanism is S_N2, **the preferred path uses the less sterically hindered halide,** CH_3CH_2Br—Path [b].



Problem 9.8 Draw the organic product of each reaction.



Problem 9.9 A key step in the synthesis of the antidepressant paroxetine (trade name Paxil) involves a Williamson ether synthesis of the acyclic ether in **X.** Draw two different routes to this ether and state which is preferred.




A **hydroxide** nucleophile is needed to synthesize an alcohol, and salts such as **NaOH** and **KOH** are inexpensive and commercially available. An **alkoxide** salt is needed to make an ether. Simple alkoxides such as sodium methoxide ($NaOCH_3$) can be purchased, but others are prepared from alcohols by a Brønsted–Lowry acid–base reaction. For example, **sodium ethoxide ($NaOCH_2CH_3$)** is prepared by treating ethanol with NaH.

NaH is an especially good base for forming an alkoxide, because the by-product of the reaction, H_2, is a gas that just bubbles out of the reaction mixture.



alkoxide nucleophile


Sample Problem 9.2 Draw the product of the following two-step reaction sequence.



Solution


[1] The base removes a proton from the OH group, forming an alkoxide.

[2] The alkoxide acts as a nucleophile in an S_N2 reaction, forming an ether.




- This two-step sequence converts an alcohol to an ether.

When an organic compound contains both a hydroxy group and a halogen atom on adjacent carbon atoms, an *intramolecular* version of this reaction forms an epoxide. The starting material for this two-step sequence, a **halohydrin,** is prepared from an alkene, as we will learn in Chapter 10.



Problem 9.10 Draw the products of each reaction.




9.7 General Features—Reactions of Alcohols, Ethers, and Epoxides

We begin our discussion of the chemical reactions of alcohols, ethers, and epoxides with a look at the general reactive features of each functional group.


9.7A Alcohols

Unlike many families of molecules, the reactions of alcohols do *not* fit neatly into a single reaction class. In Chapter 9, we discuss only the substitution and β elimination reactions of alcohols. Alcohols are also key starting materials in oxidation reactions (Chapter 12), and their polar O−H bond makes them more acidic than many other organic compounds, a feature we will explore in Chapter 19.

Alcohols are similar to alkyl halides in that both contain an electronegative element bonded to an sp^3 hybridized carbon atom. Alkyl halides contain a good leaving group (X^-), however, whereas alcohols do not. Nucleophilic substitution with ROH as starting material would displace $^-$**OH, a strong base and therefore a poor leaving group.**




For an alcohol to undergo a nucleophilic substitution or elimination reaction, the **OH group must be converted into a better leaving group.** This can be done by reaction with acid. Treatment of an alcohol with a strong acid like HCl or H_2SO_4 protonates the O atom via an acid–base reaction. This transforms the $^-$OH leaving group into **H_2O, a weak base and therefore a good leaving group.**




If the OH group of an alcohol is made into a good leaving group, alcohols *can* undergo β elimination and nucleophilic substitution, as described in Sections 9.8–9.12.

Because the pK_a of $(ROH_2)^+$ is ~–2, protonation of an alcohol occurs only with very strong acids—namely, those having a p$K_a \leq -2$.




9.7B Ethers and Epoxides

Like alcohols, **ethers do not contain a good leaving group,** which means that nucleophilic substitution and β elimination do not occur directly. Ethers undergo fewer useful reactions than alcohols.




Epoxides don't have a good leaving group either, but they have one characteristic that neither alcohols nor ethers have: **the "leaving group" is contained in a strained three-membered ring.** Nucleophilic attack opens the three-membered ring and relieves angle strain, making nucleophilic attack a favorable process that occurs even with the poor leaving group. Specific examples are presented in Section 9.16.



9.8 Dehydration of Alcohols to Alkenes

The dehydrohalogenation of alkyl halides, discussed in Chapter 8, is one way to introduce a π bond into a molecule. Another way is to eliminate water from an alcohol in a **dehydration** reaction.

- **Dehydration is a β elimination reaction in which the elements of OH and H are removed from the α and β carbon atoms, respectively.**




Dehydration is typically carried out using H_2SO_4 and other strong acids, or phosphorus oxychloride ($POCl_3$) in the presence of an amine base. We consider dehydration in acid first, followed by dehydration with $POCl_3$ in Section 9.10.

9.8A General Features of Dehydration in Acid


Alcohols undergo dehydration in the presence of strong acid to afford alkenes, as illustrated in Equations [1] and [2]. Typical acids used for this conversion are H_2SO_4 or *p*-toluenesulfonic acid (abbreviated as TsOH).

Recall from Section 2.6 that *p*-toluenesulfonic acid is a strong organic acid ($pK_a = -7$).




p-toluenesulfonic acid


TsOH



More substituted alcohols dehydrate more readily, giving rise to the following order of reactivity:



When an alcohol has two or three different β carbons, dehydration is regioselective and follows the Zaitsev rule. **The more substituted alkene is the major product when a mixture of constitutional isomers is possible.** For example, elimination of H and OH from 2-methylbutan-2-ol yields two constitutional isomers: the trisubstituted alkene **A** as major product and the disubstituted alkene **B** as minor product.





2-methylbutan-2-ol


A
major product
trisubstituted alkene

B
minor product
disubstituted alkene


Problem 9.11 Draw the products formed when each alcohol undergoes dehydration with TsOH, and label the major product when a mixture results.

a. 

b. 

c. 

Problem 9.12 Rank the alcohols in order of increasing reactivity when dehydrated with H_2SO_4.




9.8B The E1 Mechanism for the Dehydration of 2° and 3° Alcohols

The mechanism of dehydration depends on the structure of the alcohol: **2° and 3° alcohols react by an E1 mechanism, whereas 1° alcohols react by an E2 mechanism.** Regardless of the type of alcohol, however, strong acid is *always* needed to protonate the O atom to form a good leaving group.

The E1 dehydration of 2° and 3° alcohols is illustrated with $(CH_3)_3COH$ (a 3° alcohol) as starting material to form $(CH_3)_2C{=}CH_2$ as product (Mechanism 9.1). The mechanism consists of **three steps.**

Mechanism 9.1 Dehydration of 2° and 3° ROH—An E1 Mechanism



1. Protonation of the oxygen atom converts the poor leaving group (^-OH) into a good leaving group (H_2O).
2. Heterolysis of the C–O bond forms a carbocation in the rate-determining step.
3. A base (such as HSO_4^- or H_2O) removes a proton from a carbon adjacent to the carbocation to form the new π bond.


Thus, **dehydration of 2° and 3° alcohols occurs via an E1 mechanism with an added first step.** Step [1] protonates the OH group to make a good leaving group. Steps [2] and [3] are the two steps of an E1 mechanism: loss of a leaving group (H_2O in this case) to form a carbocation, followed by removal of a β proton to form a π bond. The acid used to protonate the alcohol in Step [1] is regenerated upon removal of the proton in Step [3], so dehydration is **acid-catalyzed.**

The E1 dehydration of 2° and 3° alcohols with acid gives clean elimination products without by-products formed from an S_N1 reaction. This makes the E1 dehydration of alcohols much more synthetically useful than the E1 dehydrohalogenation of alkyl halides (Section 8.7). Clean elimination takes place because the reaction mixture contains no good nucleophile to react with the intermediate carbocation, so no competing S_N1 reaction occurs.

9.8C The E2 Mechanism for the Dehydration of 1° Alcohols

Because 1° carbocations are highly unstable, the dehydration of 1° alcohols cannot occur by an E1 mechanism involving a carbocation intermediate. With 1° alcohols, therefore, **dehydration follows an E2 mechanism.** The two-step process for the conversion of $CH_3CH_2CH_2OH$ (a 1° alcohol) to $CH_3CH{=}CH_2$ with H_2SO_4 as acid catalyst is shown in Mechanism 9.2.

Mechanism 9.2 Dehydration of a 1° ROH—An E2 Mechanism



1. Protonation of the oxygen atom converts the poor leaving group (^-OH) into a good leaving group (H_2O).
2. Two bonds are broken and two bonds are formed. The base (HSO_4^- or H_2O) removes a proton from the β carbon; the electron pair in the β C–H bond forms the new π bond and the leaving group (H_2O) departs.

The dehydration of a 1° alcohol begins with the protonation of the OH group to form a good leaving group, just as in the dehydration of a 2° or 3° alcohol. With 1° alcohols, however, loss of the leaving group and removal of a β proton occur at the *same* time, so that no highly unstable 1° carbocation is generated.


Problem 9.13 Draw the structure of each transition state in the two-step mechanism for the reaction, $CH_3CH_2CH_2OH + H_2SO_4 \rightarrow CH_3CH{=}CH_2 + H_2O$.

9.8D Le Châtelier's Principle

Although entropy favors product formation in dehydration (one molecule of reactant forms two molecules of products), enthalpy does not, because the two σ bonds broken in the reactant are

Figure 9.6

The dehydration of CH_3CH_2OH to $CH_2=CH_2$—An endothermic reaction



ΔH° calculation:

[1] Bonds broken

	ΔH° (kJ/mol)
CH_3CH_2–OH	+393
$HOCH_2CH_2$–H	+410
Total	+803 kJ/mol

Energy needed to break bonds.

[2] Bonds formed

	ΔH° (kJ/mol)
CH_2=CH_2 π bond	–267
H–OH	–498
Total	–765 kJ/mol

Energy released in forming bonds.

[3] Overall ΔH° =

sum in Step [1]
+
sum in Step [2]

+803 kJ/mol
–765 kJ/mol

ΔH° = +38 kJ/mol

The reaction is endothermic.

[Values taken from Appendix C.]


stronger than the σ and π bonds formed in the products. For example, $\Delta H°$ for the dehydration of CH_3CH_2OH to $CH_2=CH_2$ is +38 kJ/mol (Figure 9.6).

According to **Le Châtelier's principle, a system at equilibrium will react to counteract any disturbance to the equilibrium.** Thus, removing a product from a reaction mixture as it is formed drives the equilibrium to the right, forming more product.

Le Châtelier's principle can be used to favor products in dehydration reactions because the alkene product has a lower boiling point than the alcohol reactant. Thus, the alkene can be distilled from the reaction mixture as it is formed, leaving the alcohol and acid to react further, forming more product.

9.9 Carbocation Rearrangements


Sometimes "unexpected" products are formed in dehydration; that is, the carbon skeletons of the starting material and product might be different, or the double bond might be in an unexpected location. For example, the dehydration of 3,3-dimethylbutan-2-ol yields two alkenes, whose carbon skeletons do not match the carbon framework of the starting material.



3,3-dimethylbutan-2-ol

This phenomenon sometimes occurs when carbocations are reactive intermediates. **A less stable carbocation can rearrange to a more stable carbocation by shift of a hydrogen atom or an alkyl group.** These **1,2-shifts** involve migration of an alkyl group or hydrogen atom from one carbon to an adjacent carbon atom. The migrating group moves with the two electrons that bonded it to the carbon skeleton.

Because the migrating group in a 1,2-shift moves with two bonding electrons, the carbon it leaves behind now has only three bonds (six electrons), giving it a net positive (+) charge.




- **Movement of a hydrogen atom is called a 1,2-hydride shift.**
- **Movement of an alkyl group is called a 1,2-alkyl shift.**

The dehydration of 3,3-dimethylbutan-2-ol illustrates the rearrangement of a 2° to a 3° carbocation by a **1,2-methyl shift,** as shown in Mechanism 9.3. The carbocation rearrangement occurs in Step [3] of the four-step mechanism.

Mechanism 9.3 A 1,2-Methyl Shift—Carbocation Rearrangement During Dehydration

Part [1] Formation of a 2° carbocation and rearrangement




1 Protonation of the oxygen atom converts the poor leaving group ($^-$OH) into a good leaving group (H_2O).

2 Heterolysis of the C–O bond forms a 2° carbocation.

3 1,2-Shift of a CH_3 group converts a 2° carbocation to a more stable 3° carbocation.

Part [2] Loss of a proton to form the π bond




4 Loss of a proton from a β carbon (β_1 or β_2) forms two different alkenes.

Steps [1], [2], and [4] in the mechanism for the dehydration of 3,3-dimethylbutan-2-ol are exactly the same steps previously seen in dehydration: protonation, loss of H_2O, and loss of a proton. Only Step [3], rearrangement of the less stable 2° carbocation to the more stable 3° carbocation, is new.


- **1,2-Shifts convert a less stable carbocation to a more stable carbocation.**

For example, 2° carbocation **A** rearranges to the more stable 3° carbocation by a 1,2-hydride shift, whereas carbocation **B** does not rearrange because it is 3° to begin with.




Sample Problem 9.3 illustrates a dehydration reaction that occurs with a **1,2-hydride** shift.

Sample Problem 9.3 Show how the dehydration of alcohol **X** forms alkene **Y** using a 1,2-hydride shift.




Solution

Steps [1] and [2] Protonation of **X** and loss of H_2O form a 2° carbocation.




Steps [3] and [4] Rearrangement of the 2° carbocation by a 1,2-hydride shift forms a more stable 3° carbocation. Loss of a proton from a β carbon forms alkene **Y**.




Problem 9.14 What other alkene is also formed along with **Y** in Sample Problem 9.3? What alkenes would form from **X** if no carbocation rearrangement occurred?

Rearrangements are not unique to dehydration reactions. **Rearrangements can occur whenever a carbocation is formed as reactive intermediate,** meaning any S_N1 or E1 reaction. In fact, the formation of rearranged products often indicates the presence of a carbocation intermediate.

Problem 9.15 Show how a 1,2-shift forms a more stable carbocation from each intermediate.



Problem 9.16 Explain why two substitution products are formed in the following reaction.




9.10 Dehydration Using $POCl_3$ and Pyridine

Because some organic compounds decompose in the presence of strong acid, other methods that avoid strong acid have been developed to convert alcohols to alkenes. A common method uses phosphorus oxychloride ($POCl_3$) and pyridine (an amine base) in place of H_2SO_4 or TsOH. For example, the treatment of cyclohexanol with $POCl_3$ and pyridine forms cyclohexene in good yield.




pyridine



cyclohexanol cyclohexene

$POCl_3$ serves much the same role as strong acid does in acid-catalyzed dehydration. **It converts a poor leaving group ($^-$OH) into a good leaving group.** Dehydration then proceeds by an **E2 mechanism,** as shown in Mechanism 9.4. Pyridine is the base that removes a β proton during elimination.

Mechanism 9.4 Dehydration Using $POCl_3$ + Pyridine—An E2 Mechanism



1–2 Reaction of the OH with $POCl_3$ followed by loss of a proton converts a poor leaving group ($^-$OH) into a good leaving group ($^-OPOCl_2$).


3 Two bonds are broken and two bonds are formed. The base (pyridine) removes a proton; the electron pair in the β C–H bond forms the π bond, and the leaving group ($^-OPOCl_2$) departs.

No rearrangements occur during dehydration with $POCl_3$, suggesting that carbocations are *not* formed as intermediates in this reaction. Steps [1] and [2] of the mechanism convert the OH group to a good leaving group. In Step [3], the C−H and C−O bonds are broken and the π bond is formed.

We have now learned about two different reagents for alcohol dehydration—strong acid (H_2SO_4 or TsOH) and $POCl_3$ + pyridine. The best dehydration method for a given alcohol is often hard to know ahead of time, and this is why organic chemists develop more than one method for a given

Figure 9.7

Dehydration reactions in the synthesis of two natural products




- New double bonds formed by dehydration are shown in red.
- **Patchouli alcohol,** obtained from the patchouli plant native to Malaysia, has been used in perfumery because of its exotic fragrance. In the 1800s, shawls imported from India were often packed with patchouli leaves to ward off insects, thus permeating the clothing with the distinctive odor.


type of transformation. Two examples of dehydration reactions used in the synthesis of natural products are given in Figure 9.7.

9.11 Conversion of Alcohols to Alkyl Halides with HX


Alcohols undergo nucleophilic substitution reactions only if the OH group is converted into a better leaving group before nucleophilic attack. Thus, substitution does *not* occur when an alcohol is treated with X^- because ^-OH is a poor leaving group (Reaction [1]), but substitution *does* occur on treatment of an alcohol with HX because H_2O is now the leaving group (Reaction [2]).




- **The reaction of alcohols with HX (X = Cl, Br, I) is a general method to prepare 1°, 2°, and 3° alkyl halides.**



More substituted alcohols usually react more rapidly with HX:




Problem 9.17 Draw the products of each reaction.



9.11A Two Mechanisms for the Reaction of ROH with HX

How does the reaction of ROH with HX occur? Acid–base reactions are very fast, so the strong acid HX protonates the OH group of the alcohol, forming a **good leaving group** (H_2O) and a **good nucleophile** (the conjugate base, X^-). Both components are needed for nucleophilic substitution. The mechanism of substitution of X^- for H_2O then depends on the structure of the R group.


When there is an oxygen-containing reactant and a strong acid, generally the first step in the mechanism is protonation of the oxygen atom.



- **Methyl and 1° ROH form RX by an S_N2 mechanism.**
- **Secondary (2°) and 3° ROH form RX by an S_N1 mechanism.**

The reaction of CH_3CH_2OH with HBr illustrates the S_N2 mechanism of a 1° alcohol (Mechanism 9.5). Nucleophilic attack on the protonated alcohol occurs in one step: **the bond to the nucleophile X^- is formed *as* the bond to the leaving group (H_2O) is broken.**

Mechanism 9.5 Reaction of a 1° ROH with HX—An S_N2 Mechanism




1 Protonation of the OH group forms a good leaving group (H_2O).

2 The bond to the nucleophile forms as the leaving group departs.

The reaction of $(CH_3)_3COH$ with HBr illustrates the S_N1 mechanism of a 3° alcohol (Mechanism 9.6). Nucleophilic attack on the protonated alcohol occurs in two steps: **the bond to the leaving group (H_2O) is broken *before* the bond to the nucleophile X^- is formed.**


Mechanism 9.6 Reaction of 2° and 3° ROH with HX—An S_N1 Mechanism




1. Protonation of the OH group forms a good leaving group (H_2O).
2. Loss of the leaving group forms a carbocation.
3. Nucleophilic attack of Br^- forms the substitution product.

Both mechanisms begin with the same first step—protonation of the O atom to form a good leaving group—and both mechanisms give an alkyl halide (RX) as product. The mechanisms differ only in the timing of bond breaking and bond making.


The reactivity of hydrogen halides increases with increasing acidity:




Because Cl^- is a poorer nucleophile than Br^- or I^-, the reaction of 1° alcohols with HCl occurs only when an additional Lewis acid catalyst, usually **$ZnCl_2$,** is added. $ZnCl_2$ complexes with the O atom of the alcohol in a Lewis acid–base reaction, making a leaving group and facilitating the S_N2 reaction.




Knowing the mechanism allows us to predict the stereochemistry of the products when reaction occurs at a stereogenic center.





- The 1° alcohol **A** reacts with HBr via an S_N2 mechanism to yield the alkyl bromide **B** with **inversion** of stereochemistry at the stereogenic center.
- The 3° alcohol **C** reacts with HCl via an S_N1 mechanism to yield a **racemic mixture** of alkyl chlorides **D** and **E,** because a trigonal planar carbocation intermediate is formed.


Problem 9.18 Draw the products of each reaction, indicating the stereochemistry around any stereogenic centers.



9.11B Carbocation Rearrangement in the S_N1 Reaction

Because carbocations are formed in the S_N1 reaction of 2° and 3° alcohols with HX, **carbocation rearrangements are possible,** as illustrated in Sample Problem 9.4.


Sample Problem 9.4 Draw a stepwise mechanism for the following reaction.




Solution

A 2° alcohol reacts with HBr by an S_N1 mechanism. Because substitution converts a 2° alcohol to a 3° alkyl halide in this example, a carbocation rearrangement must occur.


Steps [1] and [2] Protonation of the O atom and then loss of H_2O form a 2° carbocation.



Steps [3] and [4] **Rearrangement of the 2° carbocation by a 1,2-hydride shift forms a more stable 3° carbocation.** Nucleophilic attack forms the substitution product.



Problem 9.19 What is the major product formed when each alcohol is treated with HCl?



9.12 Conversion of Alcohols to Alkyl Halides with $SOCl_2$ and PBr_3


Primary (1°) and 2° alcohols can be converted to alkyl halides using $SOCl_2$ and PBr_3.

- **$SOCl_2$ (thionyl chloride) converts alcohols into alkyl chlorides.**
- **PBr_3 (phosphorus tribromide) converts alcohols into alkyl bromides.**

Both reagents convert $^-$OH into a good leaving group *in situ*—that is, directly in the reaction mixture—as well as provide the **nucleophile,** either Cl^- or Br^-, to displace the leaving group.


9.12A Reaction of ROH with $SOCl_2$

The treatment of a 1° or 2° alcohol with thionyl chloride, $SOCl_2$, and pyridine forms an alkyl chloride, with SO_2 and HCl as by-products.



The mechanism for this reaction consists of two parts: **conversion of the OH group into a better leaving group, and nucleophilic attack by Cl^- via an S_N2 reaction,** as shown in Mechanism 9.7.

Mechanism 9.7 Reaction of ROH with $SOCl_2$ + Pyridine—An S_N2 Mechanism




1–2 Reaction of the alcohol with $SOCl_2$ and loss of a proton convert the OH group to OSOCl, a good leaving group.

3 Nucleophilic attack of chloride and loss of the leaving group (SO_2 and Cl^-) form RCl in a single step.

Problem 9.20 If the reaction of an alcohol with $SOCl_2$ and pyridine follows an S_N2 mechanism, what is the stereochemistry of the alkyl chloride formed from (*R*)-butan-2-ol?


9.12B Reaction of ROH with PBr_3

In a similar fashion, the treatment of a 1° or 2° alcohol with phosphorus tribromide, PBr_3, forms an alkyl bromide.



The mechanism for this reaction also consists of two parts: **conversion of the OH group into a better leaving group, and nucleophilic attack by Br^- via an S_N2 reaction,** as shown in Mechanism 9.8.

Mechanism 9.8 Reaction of ROH with PBr_3—An S_N2 Mechanism



1. Reaction of the alcohol with PBr_3 converts the OH group to $OPBr_2$, a good leaving group, and generates the nucleophile, Br^-.
2. Nucleophilic attack of bromide and loss of the leaving group form RBr in a single step.


Table 9.2 summarizes the methods for converting an alcohol to an alkyl halide presented in Sections 9.11 and 9.12.

Table 9.2 Summary of Methods for ROH → RX

Overall reaction	Reagent	Comment
ROH → RCl	HCl	• Useful for all ROH • An S_N1 mechanism for 2° and 3° ROH; an S_N2 mechanism for CH_3OH and 1° ROH
	$SOCl_2$	• Best for CH_3OH, and 1° and 2° ROH • An S_N2 mechanism
ROH → RBr	HBr	• Useful for all ROH • An S_N1 mechanism for 2° and 3° ROH; an S_N2 mechanism for CH_3OH and 1° ROH
	PBr_3	• Best for CH_3OH, and 1° and 2° ROH • An S_N2 mechanism
ROH → RI	HI	• Useful for all ROH • An S_N1 mechanism for 2° and 3° ROH; an S_N2 mechanism for CH_3OH and 1° ROH

Problem 9.21 If the reaction of an alcohol with PBr_3 follows an S_N2 mechanism, what is the stereochemistry of the alkyl bromide formed from (*R*)-butan-2-ol?

Problem 9.22 Draw the organic products formed in each reaction, and indicate the stereochemistry of products that contain stereogenic centers.




9.12C The Importance of Making RX from ROH

We have now learned two methods to prepare an alkyl chloride and two methods to prepare an alkyl bromide from an alcohol. If there is one good way to carry out a reaction, why search for more? A particular reagent might work well for one starting material, but not so well for another, so, organic chemists try to devise several different ways to perform the same overall reaction.


Why are there so many ways to convert an alcohol to an alkyl halide? Alkyl halides are versatile starting materials in organic synthesis, as shown in Sample Problem 9.5.

Sample Problem 9.5 Convert propan-1-ol to butanenitrile (**A**).



Solution


Direct conversion of propan-1-ol to **A** using ^-CN as a nucleophile is not possible because ^-OH is a poor leaving group. However, conversion of the OH group to a Br atom forms a good leaving group, which can then readily undergo an S_N2 reaction with ^-CN to yield **A**. The overall result of this two-step sequence is the substitution of ^-OH by ^-CN.



Problem 9.23 Draw two steps to convert $(CH_3)_2CHOH$ into each of the following compounds: $(CH_3)_2CHN_3$ and $(CH_3)_2CHOCH_2CH_3$.

9.13 Tosylate—Another Good Leaving Group


We have now learned two methods to convert the OH group of an alcohol to a better leaving group: treatment with strong acids (Section 9.8), and conversion to an alkyl halide (Sections 9.11–9.12). Alcohols can also be converted to **alkyl tosylates.**



Recall from Section 1.5 that a third-row element like sulfur can have 10 or 12 electrons around it in a valid Lewis structure.

An alkyl tosylate is often called simply a **tosylate.**


An **alkyl tosylate** is composed of two parts: the **alkyl group R,** derived from an alcohol; and the **tosylate** (short for ***p*-toluenesulfonate**), which is a good leaving group. A tosyl group, $CH_3C_6H_4SO_2-$, is abbreviated as **Ts,** so an alkyl tosylate becomes **ROTs.**




9.13A Conversion of Alcohols to Alkyl Tosylates

A tosylate (TsO^-) is similar to I^- in leaving group ability.


Alcohols are converted to alkyl tosylates by treatment with *p*-toluenesulfonyl chloride (TsCl) in the presence of pyridine. This overall process converts a poor leaving group (^-OH) into a good one (^-OTs). A tosylate is a good leaving group because its conjugate acid, *p*-toluenesulfonic acid ($CH_3C_6H_4SO_3H$, TsOH), is a strong acid ($pK_a = -7$, Section 2.6).



(*S*)-Butan-2-ol is converted to its tosylate with **retention of configuration** at the stereogenic center. Thus, the C–O bond of the alcohol must *not* be broken when the tosylate is formed.




Problem 9.24 Draw the products of each reaction, and indicate the stereochemistry at any stereogenic center.



9.13B Reactions of Alkyl Tosylates

Because alkyl tosylates have good leaving groups, **they undergo both nucleophilic substitution and β elimination,** exactly as alkyl halides do. Generally, alkyl tosylates are treated with strong nucleophiles and bases, so that the mechanism of substitution is **S_N2** and the mechanism of elimination is **E2.**

For example, propyl tosylate, which has the leaving group on a 1° carbon, reacts with $NaOCH_3$ to yield methyl propyl ether, the product of nucleophilic substitution by an S_N2 mechanism. Propyl tosylate reacts with $KOC(CH_3)_3$, a strong bulky base, to yield propene by an E2 mechanism.




Because substitution occurs via an **S_N2** mechanism, **inversion of configuration** results when the leaving group is bonded to a stereogenic center.




Inversion of configuration

Sample Problem 9.6 Draw the product of the following reaction, including stereochemistry.




Solution

The 1° alkyl tosylate and the strong nucleophile both favor substitution by an **S_N2** mechanism, which proceeds by backside attack, resulting in **inversion** of configuration at the stereogenic center.




Problem 9.25 Draw the products of each reaction, and include the stereochemistry at any stereogenic center in the products.



9.13C The Two-Step Conversion of an Alcohol to a Substitution Product


We now have another **two-step method to convert an alcohol to a substitution product:** reaction of an alcohol with TsCl and pyridine to form an alkyl tosylate (Step [1]), followed by nucleophilic attack on the tosylate (Step [2]).



Let's look at the stereochemistry of this two-step process.

- Step [1], formation of the tosylate, proceeds with **retention** of configuration at a stereogenic center because the C—O bond remains intact.
- Step [2] is an S_N2 reaction, so it proceeds with **inversion of configuration** because the nucleophile attacks from the back side.
- Overall there is a **net inversion of configuration** at a stereogenic center.

For example, the treatment of *cis*-3-methylcyclohexanol with *p*-toluenesulfonyl chloride and pyridine forms a cis tosylate **A,** which undergoes backside attack by the nucleophile $^{-}OCH_3$ to yield the trans ether **B.**



Problem 9.26 Draw the products formed when (*S*)-butan-2-ol is treated with TsCl and pyridine, followed by NaOH. Label the stereogenic center in each compound as *R* or *S*. What is the stereochemical relationship between the starting alcohol and the final product?

9.13D A Summary of Substitution and Elimination Reactions of Alcohols

The reactions of alcohols in Sections 9.8–9.13C share two similarities:


- **The OH group is converted into a better leaving group by treatment with acid or another reagent.**
- **The resulting product undergoes either elimination or substitution, depending on the reaction conditions.**

Figure 9.8 summarizes these reactions with cyclohexanol as starting material.

Problem 9.27 Draw the product formed when $(CH_3)_2CHOH$ is treated with each reagent.

a. $SOCl_2$, pyridine
b. TsCl, pyridine
c. H_2SO_4
d. HBr
e. PBr_3, then NaCN
f. $POCl_3$, pyridine


Figure 9.8
Summary: Nucleophilic substitution and β elimination reactions of alcohols




9.14 Reaction of Ethers with Strong Acid

Because ethers are so unreactive, diethyl ether and tetrahydrofuran (THF) are often used as solvents for organic reactions.

Recall from Section 9.7B that ethers have a poor leaving group, so they cannot undergo nucleophilic substitution or β elimination reactions directly. Instead, they must first be converted to a good leaving group by reaction with strong acids. Only **HBr** and **HI** can be used, though, because they are strong acids that are also sources of good nucleophiles (Br^- and I^-, respectively). **When ethers react with HBr or HI, both C−O bonds are cleaved and two alkyl halides are formed as products.**




HBr or HI serves as a strong acid that both protonates the O atom of the ether and is the source of a good nucleophile (Br^- or I^-). Because both C−O bonds in the ether are broken, **two successive nucleophilic substitution reactions occur.**




- **The mechanism of ether cleavage is S_N1 or S_N2, depending on the identity of R.**
- **With 2° or 3° alkyl groups bonded to the ether oxygen, the C−O bond is cleaved by an S_N1 mechanism involving a carbocation; with methyl or 1° R groups, the C−O bond is cleaved by an S_N2 mechanism.**

For example, cleavage of $(CH_3)_3COCH_3$ with HI occurs at two bonds, as shown in Mechanism 9.9. The 3° alkyl group undergoes nucleophilic substitution by an $\mathbf{S_N1}$ mechanism, resulting in the cleavage of one C−O bond. The methyl group undergoes nucleophilic substitution by an $\mathbf{S_N2}$ mechanism, resulting in the cleavage of the second C−O bond.




Mechanism 9.9 Mechanism of Ether Cleavage in Strong Acid—$(CH_3)_3COCH_3 + HI \rightarrow (CH_3)_3CI + CH_3I + H_2O$

Part [1] Cleavage of the 3° C – O bond by an S_N1 mechanism



1. Protonation of the ether O atom forms a good leaving group.
2. Cleavage of the C–O bond to the 3° carbon forms a 3° carbocation and CH_3OH.
3. Nucleophilic attack of I^- forms the substitution product.

Part [2] Cleavage of the CH_3 – O bond by an S_N2 mechanism




4. Protonation of the OH group forms a good leaving group (H_2O).
5. Nucleophilic attack of iodide forms the second alkyl halide, CH_3I. Because the mechanism is S_N2, the C–O bond is broken as the C–I bond is formed.


The mechanism illustrates the central role of HX in the reaction:

- **HX protonates the ether oxygen, thus making a good leaving group.**
- **HX provides a source of X^- for nucleophilic attack.**

Problem 9.28 What alkyl halides are formed when each ether is treated with HBr?




Problem 9.29 Explain why the treatment of anisole with HBr yields phenol and CH_3Br, but not bromobenzene.




9.15 Thiols and Sulfides

Thiols and sulfides are sulfur analogues of alcohols and ethers, respectively.




thiol




sulfide

9.15A Thiols


Thiols, also called mercaptans, contain a mercapto group (SH) bonded to a carbon atom. Because sulfur is below oxygen in the periodic table, the sulfur atom is surrounded by two atoms and two lone pairs, giving thiols a bent shape. Unlike alcohols, however, thiols are incapable of intermolecular hydrogen bonds, so thiols have lower boiling points and melting points than alcohols with a similar number of carbons.



thiol




ethanethiol
bp 35 °C




ethanol
bp 78 °C


Many simple thiols have pungent and disagreeable odors. Skunks, onions, and human sweat all contain thiols.



propane-1-thiol
onion odor



3-methylbutane-1-thiol
skunk odor




(*S*)-3-methyl-3-sulfanylhexan-1-ol
onion-like odor in human sweat

Thiols are named in a similar method to alcohols, using the suffix ***-thiol*** instead of the suffix ***-ol***. To name a thiol in the IUPAC system:


- **Name the parent carbon chain and add the suffix *-thiol*.**
- **Number the carbon chain to give the SH group the lower number and apply the other rules of nomenclature.**

Examples of thiol nomenclature are given in Figure 9.9.


Figure 9.9
Naming thiols



pentane-3-thiol




4-methylhexane-2-thiol




2-methylcyclohexanethiol


Problem 9.30 Name each thiol.



Thiols are prepared by S_N2 reactions of alkyl halides with ^-SH, a good nucleophile.





Thiols are easily oxidized with Br_2 or I_2 to disulfides (RSSR), compounds that contain a sulfur–sulfur bond. This reaction is an oxidation (Section 4.14) because H atoms are removed from the thiol in forming the disulfide. Disulfides are reduced to thiols with Zn and acid.



Disulfide formation is especially important in determining the shape and properties of some proteins that contain the amino acid cysteine, as we will learn in Chapter 29.

Problem 9.31 Draw the product of each reaction.






The potent odor of grapefruit mercaptan (Problem 9.31c) contributes to the characteristic aroma of grapefruit.

9.15B Sulfides

Sulfides contain two alkyl groups bonded to a sulfur atom. Sulfides are named with the same rules used to name ethers. The suffix ***sulfide*** is used instead of *ether* for simple compounds.




sulfide

sec-butyl ethyl sulfide


diethyl sulfide

To name more complex sulfides using the IUPAC system, one alkyl group is named as a parent chain and the other is named as part of a substituent bonded to that chain.

- Name the simpler alkyl group + S atom as an *alkylthio* substituent.
- Name the remaining alkyl group as an alkane with an alkylthio substituent using the usual rules of nomenclature.





methylthiocyclohexane




3-ethylthio-5-methyloctane

Problem 9.32 Give the IUPAC name for each sulfide.

a. 

b. 

Sulfides are prepared from thiols by an S_N2 reaction that is analogous to the Williamson ether synthesis.




propane-1-thiol

sulfur nucleophile + H_2


ethyl propyl sulfide

Sulfides contain a nucleophilic sulfur atom that reacts readily with unhindered alkyl halides to form sulfonium ions.



sulfonium ion

S-Adenosylmethionine (SAM), a biological sulfonium ion that was introduced in Section 7.16, is synthesized from the amino acid methionine, which contains a nucleophilic sulfide, and adenosine triphosphate (ATP), which contains a triphosphate leaving group (Section 7.16).




methionine


adenosine triphosphate
ATP

S-adenosylmethionine

triphosphate
leaving group


Problem 9.33 Draw the product of each reaction.

a. 


b. 

9.16 Reactions of Epoxides

Although epoxides do not contain a good leaving group, they contain a strained three-membered ring with two polar bonds. **Nucleophilic attack opens the strained three-membered ring,** making it a favorable process even with the poor leaving group.




This reaction occurs readily with strong nucleophiles like ⁻CN, and with acids like HZ, where Z is a nucleophilic atom.



9.16A Opening of Epoxide Rings with Strong Nucleophiles


Virtually all strong nucleophiles open an epoxide ring by a two-step reaction sequence.



- **Step [1]:** The nucleophile attacks an electron-deficient carbon of the epoxide, cleaving a C−O bond and relieving the strain of the three-membered ring.
- **Step [2]:** Protonation of the alkoxide with water generates a neutral product with two functional groups on adjacent atoms.


Common nucleophiles that open epoxide rings include ⁻OH, ⁻OR, ⁻CN, ⁻SR, and NH_3. With these strong nucleophiles, the reaction occurs via an **S_N2 mechanism,** resulting in two consequences:

- **The nucleophile opens the epoxide ring from the back side.**




Other examples of the nucleophilic opening of epoxide rings are presented in Sections 12.6 and 20.14.


- **In an unsymmetrical epoxide, the nucleophile attacks at the less substituted carbon atom.**




Problem 9.34 Draw the product of each reaction, indicating the stereochemistry at any stereogenic center.



1,2-Epoxycyclohexane, an achiral epoxide with a plane of symmetry, reacts with $^-OCH_3$ to yield two *trans*-1,2-disubstituted cyclohexanes, **A** and **B,** which are **enantiomers;** each has two stereogenic centers.



Nucleophilic attack of $^-OCH_3$ occurs from the back side at *either* C–O bond, because both ends are equally substituted. Because attack at either side occurs with equal probability, an equal amount of the two enantiomers is formed—**a racemic mixture.** This is a specific example of a general rule concerning the stereochemistry of products obtained from an achiral reactant.



- **Whenever an achiral reactant yields a product with stereogenic centers, the product must be achiral (meso) or racemic.**

This general rule can be restated in terms of optical activity. Recall from Section 5.12 that achiral compounds and racemic mixtures are optically inactive.

- **Optically inactive starting materials give optically inactive products.**

Problem 9.35 The cis and trans isomers of 2,3-dimethyloxirane both react with $^-$OH to give butane-2,3-diol. One stereoisomer gives a single achiral product, and one gives two chiral enantiomers. Which epoxide gives one product and which gives two?




cis-2,3-dimethyloxirane

one enantiomer of *trans*-2,3-dimethyloxirane

9.16B Reaction with Acids HZ


Acids **HZ** that contain a nucleophile **Z** also open epoxide rings by a two-step reaction sequence.



- **Step [1]:** Protonation of the epoxide oxygen with HZ makes the epoxide oxygen into a good leaving group (OH). It also provides a source of a good nucleophile (Z^-) to open the epoxide ring.
- **Step [2]:** The nucleophile Z^- then opens the protonated epoxide ring by **backside** attack.

These two steps—**protonation followed by nucleophilic attack**—are the exact reverse of the opening of epoxide rings with strong nucleophiles, where nucleophilic attack precedes protonation.

HCl, HBr, and **HI** all open an epoxide ring in this manner. **H_2O** and **ROH** can, too, but acid must also be added. Regardless of the reaction, the product has an OH group from the epoxide on one carbon and a new functional group Z from the nucleophile on the adjacent carbon. With epoxides fused to rings, ***trans*-1,2-disubstituted cycloalkanes** are formed.




enantiomers

Although backside attack of the nucleophile suggests that this reaction follows an S_N2 mechanism, the regioselectivity of the reaction with unsymmetrical epoxides does not.

- **With unsymmetrical epoxides, nucleophilic attack occurs at the *more* substituted carbon atom.**

For example, the treatment of 2,2-dimethyloxirane with HCl results in nucleophilic attack at the carbon with two methyl groups.



Backside attack of the nucleophile suggests an S_N2 mechanism, but attack at the more substituted carbon suggests an S_N1 mechanism. To explain these results, the **mechanism of nucleophilic attack is thought to be somewhere in between S_N1 and S_N2.**

Figure 9.10 illustrates two possible pathways for the reaction of 2,2-dimethyloxirane with HCl. Backside attack of Cl^- at the more substituted carbon proceeds via transition state **A,** whereas backside attack of Cl^- at the less substituted carbon proceeds via transition state **B. Transition state A has a partial positive charge on a more substituted carbon, making it more stable.** Thus, the preferred reaction path takes place by way of the lower energy transition state **A.**

Opening of an epoxide ring with either a strong nucleophile $:Nu^-$ or an acid HZ is **regioselective,** because one constitutional isomer is the major or exclusive product. The **site selectivity of these two reactions, however, is *exactly the opposite.***






Figure 9.10
Opening of an unsymmetrical epoxide ring with HCl



- Transition state **A** is lower in energy because the partial positive charge (δ+) is located on the more substituted carbon. In this case, therefore, nucleophilic attack occurs from the back side (an S_N2 characteristic) at the more substituted carbon (an S_N1 characteristic).

Figure 9.11
The synthesis of two bronchodilators uses the opening of an epoxide ring.




- A key step in each synthesis is the opening of an epoxide ring with a nitrogen nucleophile to form a new C – N bond, shown in red.

- **With a strong nucleophile, :Nu⁻ attacks at the less substituted carbon.**
- **With an acid HZ, the nucleophile attacks at the more substituted carbon.**

The reaction of epoxide rings with nucleophiles is important for the synthesis of many biologically active compounds, including **salmeterol** and **albuterol,** two bronchodilators used in the treatment of asthma (Figure 9.11).

Problem 9.36 Draw the product of each reaction.



9.17 Application: Epoxides, Leukotrienes, and Asthma


The opening of epoxide rings with nucleophiles is a key step in some important biological processes.

9.17A Asthma and Leukotrienes

Asthma is an obstructive lung disease that affects millions of Americans. Because it involves episodic constriction of small airways, bronchodilators such as albuterol (Figure 9.11) are used to treat symptoms by widening airways. Because asthma is also characterized by chronic inflammation, inhaled steroids that reduce inflammation are also commonly used.


Leukotrienes were first synthesized in 1980 in the laboratory of Professor E. J. Corey, the 1990 recipient of the Nobel Prize in Chemistry.

The **leukotrienes** are molecules that contribute to the asthmatic response. A typical example, **leukotriene C_4,** is shown. Although its biological activity was first observed in the 1930s, the chemical structure of leukotriene C_4 was not determined until 1979. Structure determination and chemical synthesis were difficult because leukotrienes are highly unstable and extremely potent, and are therefore present in tissues in exceedingly small amounts.




9.17B Leukotriene Synthesis and Asthma Drugs

Leukotrienes are synthesized in cells by the oxidation of **arachidonic acid** to 5-HPETE, which is then converted to an epoxide, **leukotriene A_4.** Opening of the epoxide ring with a sulfur nucleophile **RSH** yields leukotriene C_4.



New asthma drugs act by blocking the synthesis of leukotriene C_4 from arachidonic acid. For example, **zileuton** (trade name Zyflo CR) inhibits the enzyme (called a lipoxygenase) needed for the first step of this process. By blocking the synthesis of leukotriene C_4, a compound responsible for the disease, zileuton treats the **cause of asthma,** not just its symptoms.




Generic name zileuton
Trade name Zyflo CR
anti-asthma drug

9.18 Application: Benzo[*a*]pyrene, Epoxides, and Cancer



The sooty exhaust from trucks and buses contains PAHs such as benzo[*a*]pyrene.


Benzo[*a*]pyrene is a widespread environmental pollutant, produced during the combustion of all types of organic material—gasoline, fuel oil, wood, garbage, and cigarettes. It is a **polycyclic aromatic hydrocarbon (PAH),** a class of compounds that is discussed further in Chapter 17.



benzo[*a*]pyrene — **water insoluble**

a diol epoxide — **more water soluble**

After this nonpolar and water-insoluble hydrocarbon is inhaled or ingested, it is oxidized in the liver to a diol epoxide. Oxidation is a common fate of foreign substances that are not useful nutrients for the body. The oxidation product has three oxygen-containing functional groups, making it much more water soluble, and more readily excreted in urine. It is also a potent carcinogen. The strained three-membered ring of the epoxide reacts readily with biological nucleophiles $:Nu^-$ (such as DNA or an enzyme), leading to ring-opened products that often disrupt normal cell function, causing cancer or cell death.



carcinogen

These examples illustrate the central role of the nucleophilic opening of epoxide rings in two well-defined cellular processes.

KEY CONCEPTS


Alcohols, Ethers, and Related Compounds

General Facts about ROH, ROR, and Epoxides

- All three compounds contain an O atom that is sp^3 hybridized and tetrahedral (9.2).
- All three compounds have polar C – O bonds, but only alcohols have an O – H bond for intermolecular hydrogen bonding (9.4).
- Alcohols and ethers do not contain a good leaving group. Nucleophilic substitution can occur only after the OH (or OR) group is converted to a better leaving group (9.7A).
- Epoxides have a leaving group located in a strained three-membered ring, making them reactive to strong nucleophiles and acids HZ that contain a nucleophilic atom Z (9.16).

A New Reaction of Carbocations (9.9)

- Less stable carbocations rearrange to more stable carbocations by the shift of a hydrogen atom or an alkyl group.



- Besides rearranging, a carbocation can also react with a nucleophile (7.12) and a base (8.6).


Preparation of Alcohols, Ethers, and Epoxides (9.6)

[1] Preparation of alcohols



- The mechanism is S_N2.
- The reaction works best for CH_3X and 1° RX.

[2] Preparation of alkoxides—A Brønsted–Lowry acid–base reaction




[3] Preparation of ethers (Williamson ether synthesis)



- The mechanism is S_N2.
- The reaction works best for CH_3X and 1° RX.

[4] Preparation of epoxides—Intramolecular S_N2 reaction




- A two-step reaction sequence:
  [1] The removal of a proton with base forms an alkoxide.
  [2] An intramolecular S_N2 reaction forms the epoxide.

Reactions of Alcohols


[1] Dehydration to form alkenes

a. Using strong acid (9.8, 9.9)




- Order of reactivity: $R_3COH > R_2CHOH > RCH_2OH$.
- The mechanism for 2° and 3° ROH is E1—carbocations are intermediates and rearrangements occur.
- The mechanism for 1° ROH is E2.
- The Zaitsev rule is followed.

b. Using $POCl_3$ and pyridine (9.10)




- The mechanism is E2.
- No carbocation rearrangements occur.

[2] Reaction with HX to form RX (9.11)




- Order of reactivity: $R_3COH > R_2CHOH > RCH_2OH$.
- The mechanism for 2° and 3° ROH is S_N1—carbocations are intermediates and rearrangements occur.
- The mechanism for CH_3OH and 1° ROH is S_N2.

[3] Reaction with other reagents to form RX (9.12)



- Reactions occur with CH_3OH and 1° and 2° ROH.
- The reactions follow an S_N2 mechanism.


[4] Reaction with tosyl chloride to form alkyl tosylates (9.13A)



- The C – O bond is not broken, so the configuration at a stereogenic center is retained.

Reactions of Alkyl Tosylates


Alkyl tosylates undergo either substitution or elimination, depending on the reagent (9.13B).



- Substitution is carried out with a strong :Nu⁻, so the mechanism is S_N2.
- Elimination is carried out with a strong base, so the mechanism is E2.

Reactions of Ethers

Only one reaction is useful: cleavage with strong acids (9.14).



- With 2° and 3° R groups, the mechanism is S_N1.
- With CH_3 and 1° R groups, the mechanism is S_N2.

Reactions Involving Thiols and Sulfides (9.15)


[1] Preparation of thiols




- The mechanism is S_N2.
- The reaction works best for CH_3X and 1° RX.

[2] Oxidation and reduction involving thiols


a. Oxidation of thiols to disulfides



b. Reduction of disulfides to thiols



[3] Preparation of sulfides



- The mechanism is S_N2.
- The reaction works best for CH_3X and 1° RX.


[4] Reaction of sulfides to form sulfonium ions



- The mechanism is S_N2.
- The reaction works best for CH_3X and 1° RX.

Reactions of Epoxides

Epoxide rings are opened with nucleophiles :Nu⁻ and acids HZ (9.16).





- The reaction occurs with backside attack, resulting in trans or anti products.
- With :Nu⁻, the mechanism is S_N2, and nucleophilic attack occurs at the less substituted C.
- With HZ, the mechanism is between S_N1 and S_N2, and attack of Z^- occurs at the more substituted C.


PROBLEMS

Problems Using Three-Dimensional Models


9.37 Name each compound depicted in the ball-and-stick models.

a. 

b. 

c. 


9.38 Answer each question using the ball-and-stick model of compound **A.**



A


a. Give the IUPAC name for **A,** including *R,S* designations for stereogenic centers.
b. Classify **A** as a 1°, 2°, or 3° alcohol.
c. Draw a stereoisomer for **A** and give its IUPAC name.
d. Draw a constitutional isomer that contains an OH group and give its IUPAC name.
e. Draw a constitutional isomer that contains an ether and give its IUPAC name.
f. Draw the products formed (including stereochemistry) when **A** is treated with each reagent: [1] NaH; [2] H_2SO_4; [3] $POCl_3$, pyridine; [4] HCl; [5] $SOCl_2$, pyridine; [6] TsCl, pyridine.


9.39 Draw the product and indicate the stereochemistry when the given alcohol is treated with each reagent: (a) HBr; (b) PBr_3; (c) HCl; (d) $SOCl_2$ and pyridine.





Nomenclature


9.40 Give the IUPAC name for each alcohol.

a. 


c. 


e. 


b. 


d. 


9.41 Name each ether, epoxide, thiol, and sulfide.

a. 

c. 

e. 

b. 

d. 

9.42 Give the structure corresponding to each name.

a. *trans*-2-methylcyclohexanol
b. 2,3,3-trimethylbutan-2-ol
c. 6-*sec*-butyl-7,7-diethyldecan-4-ol
d. 3-chloropropane-1,2-diol
e. 1,2-epoxy-1,3,3-trimethylcyclohexane
f. 1-ethoxy-3-ethylheptane
g. (2*R*,3*S*)-3-isopropylhexan-2-ol
h. (*S*)-2-ethoxy-1,1-dimethylcyclopentane
i. 4-ethylheptane-3-thiol
j. 1-isopropylthio-2-methylcyclohexane

Physical Properties

9.43 Explain the observed trend in the melting points of the following three isomeric alcohols: $(CH_3)_2CHCH_2OH$ (–108 °C), $CH_3CH_2CH_2CH_2OH$ (–90 °C), $(CH_3)_3COH$ (26 °C).


9.44 Why is the boiling point of propane-1,3-diol ($HOCH_2CH_2CH_2OH$) higher than the boiling point of propane-1,2-diol [$HOCH_2CH(OH)CH_3$] (215 °C vs. 187 °C)? Why do both diols have a higher boiling point than butan-1-ol ($CH_3CH_2CH_2CH_2OH$, 118 °C)?

Alcohols

9.45 Draw the organic product(s) formed when $CH_3CH_2CH_2OH$ is treated with each reagent.

a. H_2SO_4
b. NaH
c. $HCl + ZnCl_2$
d. HBr
e. $SOCl_2$, pyridine
f. PBr_3
g. TsCl, pyridine
h. [1] NaH; [2] CH_3CH_2Br
i. [1] TsCl, pyridine; [2] NaSH
j. $POCl_3$, pyridine


9.46 What alkenes are formed when each alcohol is dehydrated with TsOH? Label the major product when a mixture results.




9.47 Draw the products formed when $CH_3CH_2CH_2CH_2OTs$ is treated with each reagent.

a. CH_3SH b. $NaOCH_2CH_3$ c. NaOH d. $KOC(CH_3)_3$

9.48 Draw the products of each reaction and indicate stereochemistry around stereogenic centers.




9.49 Draw the product of the following reaction, one step in the synthesis of the antiplatelet agent clopidogrel used to reduce the risk of strokes.




9.50 Draw the substitution product formed (including stereochemistry) when (*R*)-hexan-2-ol is treated with each series of reagents: (a) NaH, followed by CH_3I; (b) TsCl and pyridine, followed by $NaOCH_3$; (c) PBr_3, followed by $NaOCH_3$. Which two routes produce identical products?

9.51 (a) What is the major alkene formed when **A** is dehydrated with H_2SO_4? (b) What is the major alkene formed when **A** is treated with $POCl_3$ and pyridine? Explain why the major product is different in these reactions.




A


9.52 Draw a stepwise mechanism for the following reaction.



9.53 Although alcohol **V** gives a single alkene **W** when treated with $POCl_3$ and pyridine, three isomeric alkenes (**X–Z**) are formed on dehydration with H_2SO_4. Draw a stepwise mechanism for each reaction and explain why the difference occurs.




9.54 Sometimes carbocation rearrangements can change the size of a ring. Draw a stepwise, detailed mechanism for the following reaction.




9.55 Explain the following observation. When 3-methylbutan-2-ol is treated with HBr, a single alkyl bromide is isolated, resulting from a 1,2-shift. When 2-methylpropan-1-ol is treated with HBr, no rearrangement occurs to form an alkyl bromide.


9.56 An allylic alcohol contains an OH group on a carbon atom adjacent to a C – C double bond. Treatment of allylic alcohol **A** with HCl forms a mixture of two allylic chlorides, **B** and **C.** Draw a stepwise mechanism that illustrates how both products are formed.



9.57 Draw a stepwise, detailed mechanism for the following reaction.




9.58 Draw a stepwise, detailed mechanism for the following intramolecular reaction that forms a cyclic ether.




Ethers

9.59 Draw two different routes to each of the following ethers using a Williamson ether synthesis. Indicate the preferred route (if there is one).





9.60 Explain why it is not possible to prepare *tert*-butyl phenyl ether using a Williamson ether synthesis.

9.61 Draw the products formed when each ether is treated with two equivalents of HBr.




9.62 Draw a stepwise mechanism for each reaction.

a. 

b. 

9.63 Draw a stepwise, detailed mechanism for the following reaction.





Epoxides


9.64 Draw the products formed when ethylene oxide is treated with each reagent.


a. HBr
b. H_2O (H_2SO_4)
c. [1] $CH_3CH_2O^-$; [2] H_2O
d. [1] $HC\equiv C^-$; [2] H_2O
e. [1] ^-OH; [2] H_2O
f. [1] CH_3S^-; [2] H_2O

9.65 Draw the products of each reaction.


a. 


b. 


c. 

d. 

9.66 When each halohydrin is treated with NaH, a product of molecular formula C_4H_8O is formed. Draw the structure of the product and indicate its stereochemistry.


a. 

b. 

c. 

9.67 (a) What reaction conditions are needed to convert (*R*)-2-ethyl-2-methyloxirane to (*R*)-2-methylbutane-1,2-diol? (b) What reaction conditions are needed to convert (*R*)-2-ethyl-2-methyloxirane to (*S*)-2-methylbutane-1,2-diol?


9.68 Draw a stepwise mechanism for the following reaction and explain the stereochemistry of the product.





General Problems


9.69 Draw the products of each reaction, and indicate the stereochemistry where appropriate.


a. 


b. 

c. 


d. 

e. 


f. 




9.70 The following two-step procedure was used to prepare a sulfide from a diol. Draw the intermediate formed in Reaction [1] and draw a mechanism for Reaction [2].




9.71 Prepare each compound from cyclopentanol. More than one step may be needed.



9.72 Identify the reagents (a–h) needed to carry out each reaction.




9.73 **Propranolol,** an antihypertensive agent used in the treatment of high blood pressure, can be prepared from 1-naphthol, epichlorohydrin, and isopropylamine using two successive nucleophilic substitution reactions. Devise a stepwise synthesis of propranolol from these starting materials.




propranolol 1-naphthol epichlorohydrin isopropylamine

Challenge Problems


9.74 Treatment of *cis*-4-bromocyclohexanol with HO^- affords compound **A** and cyclohex-3-en-1-ol. Treatment of *trans*-4-bromocyclohexanol under the same conditions forms compound **B** and cyclohex-3-en-1-ol. **A** and **B** contain different functional groups and are not isomers of each other. Propose structures for **A** and **B** and offer an explanation for their formation.



cis-4-bromocyclohexanol




trans-4-bromocyclohexanol




cyclohex-3-en-1-ol

9.75 Epoxides are converted to allylic alcohols with nonnucleophilic bases such as lithium diethylamide [$LiN(CH_2CH_3)_2$]. Draw a stepwise mechanism for the conversion of 1,2-epoxycyclohexane to cyclohex-2-en-1-ol with this base. Explain why a strong bulky base must be used in this reaction.




cyclohex-2-en-1-ol

9.76 Rearrangements can occur during the dehydration of 1° alcohols even though no 1° carbocation is formed—that is, a 1,2-shift occurs as the $C-OH_2^+$ bond is broken, forming a more stable 2° or 3° carbocation, as shown. Using this information, draw a stepwise mechanism that shows how $CH_3CH_2CH_2CH_2OH$ is dehydrated with H_2SO_4 to form a mixture of $CH_3CH_2CH=CH_2$ and the cis and trans isomers of $CH_3CH=CHCH_3$. We will see another example of this type of rearrangement in Section 18.5C.




1° alcohol — no 1° carbocation at this step — 2° carbocation

9.77 Draw a stepwise, detailed mechanism for the following reaction.




9.78 Dehydration of 1,2,2-trimethylcyclohexanol with H_2SO_4 affords 1-*tert*-butylcyclopentene as a minor product. (a) Draw a stepwise mechanism that shows how this alkene is formed. (b) Draw other alkenes formed in this dehydration. At least one must contain a five-membered ring.




1,2,2-trimethylcyclohexanol — 1-*tert*-butylcyclopentene

9.79 1,2-Diols are converted to carbonyl compounds when treated with strong acids, in a reaction called the *pinacol rearrangement.* (a) Draw a stepwise mechanism for this reaction. (Hint: The reaction proceeds by way of carbocation intermediates.) (b) Assuming that the pinacol rearrangement occurs via the more stable carbocation, draw the rearrangement product formed from diol **D.**




pinacol — pinacolone — **D**


9.80 Draw a stepwise mechanism for the following reaction.




9.81 Aziridines are heterocycles that contain an N atom in a three-membered ring. Like epoxides, aziridines are strained and reactive because the 60° bond angles of the three-membered ring deviate greatly from the theoretical tetrahedral bond angle. One step in the synthesis of the drug oseltamivir (trade name Tamiflu, Section 3.2) involves the conversion of amine **X** to diamine **Y,** a reaction that occurs by way of an intermediate aziridine. Draw a stepwise mechanism for the conversion of **X** to **Y.** Indicate the structure of the aziridine intermediate, and explain the trans stereochemistry of the two amines in **Y.**



oseltamivir




X



Y

Alkenes

10



10.1 Introduction
10.2 Calculating degrees of unsaturation
10.3 Nomenclature
10.4 Physical properties
10.5 Interesting alkenes
10.6 Lipids—Part 2
10.7 Preparation of alkenes
10.8 Introduction to addition reactions
10.9 Hydrohalogenation—Electrophilic addition of HX
10.10 Markovnikov's rule
10.11 Stereochemistry of electrophilic addition of HX
10.12 Hydration—Electrophilic addition of water
10.13 Halogenation—Addition of halogen
10.14 Stereochemistry of halogenation
10.15 Halohydrin formation
10.16 Hydroboration–oxidation
10.17 Keeping track of reactions
10.18 Alkenes in organic synthesis

Candlenuts, known as kukui nuts in Hawai'i, are rich in **linoleic acid** and **linolenic acid,** two unsaturated fatty acids. **Unsaturated fatty acids** have a carboxy group (CO_2H) bonded to a carbon chain that contains one or more carbon–carbon double bonds. The presence of the double bonds greatly affects the chemical and physical properties of these compounds. In Chapter 10, we learn about alkenes, organic compounds that contain carbon–carbon double bonds.


In Chapters 10 and 11 we turn our attention to **alkenes** and **alkynes,** compounds that contain one and two π bonds, respectively. Because π bonds are easily broken, alkenes and alkynes undergo **addition,** the third general type of organic reaction. These multiple bonds also make carbon atoms electron rich, so alkenes and alkynes react with a wide variety of electrophilic reagents in addition reactions.

In Chapter 10 we review the properties and synthesis of alkenes first, and then concentrate on reactions. **Every new reaction in Chapter 10 is an *addition reaction*.** The most challenging part is learning the reagents, mechanism, and stereochemistry that characterize each individual reaction.


10.1 Introduction

Alkenes are also called **olefins.**

Alkenes are compounds that contain a carbon–carbon double bond. **Terminal alkenes** have the double bond at the end of the carbon chain, whereas **internal alkenes** have at least one carbon atom bonded to each end of the double bond. **Cycloalkenes** contain a double bond in a ring.



The double bond of an alkene consists of one σ bond and one π bond. Each carbon is sp^2 hybridized and trigonal planar, and all bond angles are approximately 120° (Section 8.2A).




Bond dissociation energies of the C−C bonds in ethane (a σ bond only) and ethylene (one σ and one π bond) can be used to estimate the strength of the π component of the double bond. If we assume that the σ bond in ethylene is similar in strength to the σ bond in ethane (368 kJ/mol), then the π bond is worth 267 kJ/mol.

$CH_2=CH_2$		CH_3-CH_3		
635 kJ/mol (σ + π bond)	−	368 kJ/mol (σ bond)	=	267 kJ/mol **π bond only**


- **The π bond is much weaker than the σ bond of a C−C double bond, making it much more easily broken. As a result, alkenes undergo many reactions that alkanes do not.**

Other features of the carbon–carbon double bond, which were presented in Chapter 8, are summarized in Table 10.1.

Table 10.1 Properties of the Carbon–Carbon Double Bond

Property	Result
Restricted rotation	• **The rotation around the C – C double bond is restricted.** Rotation can occur only if the π bond breaks and then re-forms, a process that is unfavorable (Section 8.2B).
Stereoisomerism	• Whenever the two groups on each end of a C=C are different from each other, two diastereomers are possible. *Cis*- and *trans*-but-2-ene (drawn at the bottom of Table 10.1) are diastereomers (Section 8.2B).
Stability	• **Trans** alkenes are generally more stable than **cis** alkenes. • **The stability of an alkene increases as the number of R groups on the C=C increases** (Section 8.2C).  but-1-ene *cis*-but-2-ene *trans*-but-2-ene **Increasing stability**

Cycloalkenes having fewer than eight carbon atoms have a cis geometry. A trans cycloalkene must have a carbon chain long enough to connect the ends of the double bond without introducing too much strain. *trans*-Cyclooctene is the smallest, isolable trans cycloalkene, but it is considerably less stable than *cis*-cyclooctene, making it one of the few alkenes having a higher energy trans isomer.



Problem 10.1 Draw the six alkenes of molecular formula C_5H_{10}. Label one pair of diastereomers.

10.2 Calculating Degrees of Unsaturation

An acyclic alkene has the general molecular formula C_nH_{2n}, giving it two fewer hydrogens than an acyclic alkane with the same number of carbons.

- **Alkenes are *unsaturated hydrocarbons* because they have fewer than the maximum number of hydrogen atoms per carbon.**

In Chapter 12 we will learn how to use the hydrogenation of π bonds to determine how many degrees of unsaturation result from π bonds and how many result from rings.

Cycloalkanes also have the general molecular formula C_nH_{2n}. Thus, **each π bond or ring removes two hydrogen atoms from a molecule, and this introduces one *degree of unsaturation*.** The number of degrees of unsaturation for a given molecular formula can be calculated by comparing the actual number of H atoms in a compound and the maximum number of H atoms possible. Remember that for n carbons, the maximum number of H atoms is $2n + 2$ (Section 4.1). This procedure gives the total number of rings and π bonds in a molecule.

Sample Problem 10.1

Calculate the number of degrees of unsaturation in a compound of molecular formula C_4H_6, and propose possible structures.

Solution

[1] Calculate the maximum number of H's possible.

- For *n* carbons, the maximum number of H's is $2n + 2$; in this example, $2n + 2 = 2(4) + 2 = 10$.


[2] Subtract the actual number of H's from the maximum number and divide by two.

- 10 H's (maximum) – 6 H's (actual) = 4 H's fewer than the maximum number.

$$\frac{\text{4 H's fewer than the maximum}}{\text{2 H's removed for each degree of unsaturation}} =$$

Answer: two degrees of unsaturation

A compound with two degrees of unsaturation has:

	two rings	*or*	**two π bonds**	*or*	**one ring and one π bond**
Possible structures for C_4H_6:					

This procedure can be extended to compounds that contain heteroatoms such as oxygen, nitrogen, and halogen, as illustrated in Sample Problem 10.2.

Sample Problem 10.2

Calculate the number of degrees of unsaturation for each molecular formula: (a) C_5H_8O; (b) $C_6H_{11}Cl$; (c) C_8H_9N. Propose one possible structure for each compound.

Solution

a. When a compound contains an oxygen atom, **use the given number of C's and H's and ignore the O atom** in the calculation; that is, C_5H_8O is equivalent to C_5H_8 when calculating degrees of unsaturation.

[1] For 5 C's, the maximum number of H's = $2n + 2 = 2(5) + 2 = 12$.

[2] Because the compound contains only 8 H's, it has 12 – 8 = 4 H's fewer than the maximum number.

[3] Each degree of unsaturation removes 2 H's, so the answer in Step [2] must be divided by 2.
Answer: two degrees of unsaturation

b. A compound with a halogen atom is equivalent to a hydrocarbon having one more H; that is, $C_6H_{11}Cl$ is equivalent to C_6H_{12} when calculating degrees of unsaturation.

[1] For 6 C's, the maximum number of H's = $2n + 2 = 2(6) + 2 = 14$.

[2] Because the compound contains only 12 H's, it has 14 – 12 = 2 H's fewer than the maximum number.

[3] Each degree of unsaturation removes 2 H's, so the answer in Step [2] must be divided by 2.
Answer: one degree of unsaturation


c. A compound with a nitrogen atom is equivalent to a hydrocarbon having one fewer H; that is, C_8H_9N is equivalent to C_8H_8 when calculating degrees of unsaturation.


[1] For 8 C's, the maximum number of H's = $2n + 2 = 2(8) + 2 = 18$.


[2] Because the compound contains only 8 H's, it has 18 – 8 = 10 H's fewer than the maximum number.

[3] Each degree of unsaturation removes 2 H's, so the answer in Step [2] must be divided by 2.
Answer: five degrees of unsaturation

Possible structures:

a. 

b. 

c. 

Problem 10.2 How many degrees of unsaturation are present in each compound?

a. C_6H_6 b. C_8H_{18} c. C_7H_8O d. $C_7H_{11}Br$ e. C_5H_9N

Problem 10.3 How many degrees of unsaturation does each of the following drugs contain?

a. zolpidem (sleep aid sold as Ambien), $C_{19}H_{21}N_3O$

b. mefloquine (antimalarial drug), $C_{17}H_{16}F_6N_2O$


10.3 Nomenclature

- In the IUPAC system, an alkene is identified by the suffix *-ene.*


10.3A General IUPAC Rules

How To Name an Alkene

Example Give the IUPAC name of the following alkene:



Step [1] Find the longest chain that contains *both* carbon atoms of the double bond.




6 C's in the longest chain

hex*ane* ----→ **hex*ene***

- Change the ***-ane*** ending of the parent alkane to ***-ene.***

Step [2] Number the carbon chain to give the double bond the lower number, and apply all other rules of nomenclature.


a. **Number** the chain, and name using the ***first* number** assigned to the C=C.



- Number the chain to put the C=C at C2, not C4.

hex-2-ene

b. **Name** and **number** the substituents.



three methyl groups at C2, C3, and C5

Answer: 2,3,5-trimethylhex-2-ene

Compounds with two double bonds are named as **dienes** by changing the ***-ane*** ending of the parent alkane to the suffix ***-adiene.*** Compounds with three double bonds are named as **trienes,** and so forth. Always choose the longest chain that contains *both* atoms of the double bond. In

Figure 10.1, the alkene is named as a derivative of heptene because the seven-carbon chain contains both atoms of the double bond, but the eight-carbon chain does not.

Figure 10.1
Naming an alkene in which the longest carbon chain does not contain both atoms of the double bond



7 C's ----→ heptene

Both C's of the C=C are contained in this long chain.

Correct: 2-ethylhept-1-ene




8 C's

Both C's of the C=C are NOT contained in this long chain.


Incorrect

In naming cycloalkenes, the double bond is located between C1 and C2, and the "1" is usually omitted in the name. The ring is numbered clockwise or counterclockwise to give the first substituent the lower number. Representative examples are given in Figure 10.2.

Figure 10.2
Examples of cycloalkene nomenclature




1-methylcyclopentene



3-methylcycloheptene

Number clockwise beginning at the C=C and place the CH_3 at C3.




1,6-dimethylcyclohexene


Number counterclockwise beginning at the C=C and place the first CH_3 at C1.

$CH_3CH_2CH=CH_2$ is named as 1-butene using the 1979 IUPAC recommendations and but-1-ene using the 1993 IUPAC recommendations.

Compounds that contain both a double bond and a hydroxy group are named as **alkenols,** and the chain (or ring) is numbered to give the OH group the lower number.




prop-2-en-1-ol




6-methylhept-6-en-2-ol

Problem 10.4 Give the IUPAC name for each alkene.



Problem 10.5 Give the IUPAC name for each polyfunctional compound.



10.3B Naming Stereoisomers

A prefix is needed to distinguish two alkenes when diastereomers are possible.

Using Cis and Trans as Prefixes

An alkene having one alkyl group bonded to each carbon atom can be named using the prefixes **cis** and **trans** to designate the relative location of the two alkyl groups. For example, *cis*-hex-3-ene

has two ethyl groups on the **same side** of the double bond, whereas *trans*-hex-3-ene has two ethyl groups on **opposite sides** of the double bond.



2 R's on the same side
cis-hex-3-ene

2 R's on opposite sides
trans-hex-3-ene

Using the Prefixes *E* and *Z*

E stands for the German word ***e****ntgegen* meaning "opposite." *Z* stands for the German word ***z****usammen,* meaning "together." Using *E,Z* nomenclature, a cis isomer has the *Z* configuration and a trans isomer has the *E* configuration.

Although the prefixes cis and trans can be used to distinguish diastereomers when two alkyl groups are bonded to the C=C, they cannot be used when there are three or four alkyl groups bonded to the C=C.



A
3-methylpent-2-ene


B
3-methylpent-2-ene

For example, alkenes **A** and **B** are two *different* compounds that are both called 3-methylpent-2-ene. In **A** the two CH_3 groups are cis, whereas in **B** the CH_3 and CH_2CH_3 groups are cis. The ***E,Z* system of nomenclature** has been devised to unambiguously name these kinds of alkenes.

How To Assign the Prefixes *E* and *Z* to an Alkene

Step [1] **Assign priorities to the two substituents on each end of the C=C by using the priority rules for *R,S* nomenclature (Section 5.6).**


- Divide the double bond in half, and assign the numbers **1** and **2** to indicate the relative priority of the two groups on each end—the higher priority group is labeled **1,** and the lower priority group is labeled **2.**



Assign priorities to each side separately.


Step [2] **Assign *E* or *Z* based on the location of the two higher priority groups (1).**

Two higher priority groups on **opposite sides**



***E* isomer**
(*E*)-3-methylpent-2-ene


Two higher priority groups on the **same side**





***Z* isomer**
(*Z*)-3-methylpent-2-ene

- The ***E*** isomer has the two higher priority groups on the **opposite sides.**
- The ***Z*** isomer has the two higher priority groups on the **same side.**

Problem 10.6 Label each C – C double bond as *E* or *Z*. Kavain is a naturally occurring relaxant isolated from kava root.

a. 

b. 


c. 

kavain

Problem 10.7 Label each carbon–carbon double bond in 11-*cis*-retinal as *E* or *Z*. As we will learn in Section 21.11, the isomerization of one double bond in this compound to a less crowded stereoisomer takes place when light strikes the retina of the eye.



11-*cis*-Retinal is the light-sensitive aldehyde that plays a key role in the chemistry of vision for all vertebrates, arthropods, and mollusks.



11-*cis*-retinal

Problem 10.8 Draw the structure corresponding to each IUPAC name.

a. (*Z*)-4-ethylhept-3-ene b. (*E*)-3,5,6-trimethyloct-2-ene c. (*Z*)-2-bromo-1-iodohex-1-ene

Problem 10.9 Draw the structure of (2*Z*,6*E*)-3-ethyl-7-methyldeca-2,6-dien-1-ol, the sex pheromone of the codling moth, a common agricultural pest that destroys apple crops. By spraying an apple orchard with this pheromone, the mating of male and female moths is disrupted, and moth populations can be controlled.

10.3C Common Names


The simplest alkene, $\mathbf{CH_2{=}CH_2}$, named in the IUPAC system as **ethene,** is often called **ethylene,** its common name. The common names for three **alkyl groups** derived from alkenes are also used. Three examples of naming organic molecules using these common names are shown in Figure 10.3.



methylene group




vinyl group




allyl group


Figure 10.3
Naming alkenes with common substituent names



methylenecyclohexane



1-vinylcyclopentene




2-allylcycloheptanol

10.4 Physical Properties

Most alkenes exhibit only weak van der Waals interactions, so their physical properties are similar to alkanes of comparable molecular weight.


- Alkenes have low melting points and boiling points.
- Melting points and boiling points increase as the number of carbons increases because of increased surface area.
- Alkenes are soluble in organic solvents and insoluble in water.

Cis and trans alkenes often have somewhat different physical properties. For example, *cis*-but-2-ene has a higher boiling point (4 °C) than *trans*-but-2-ene (1 °C). This difference arises because the C−C single bond between an alkyl group and one of the double-bond carbons of an alkene is slightly polar. **The sp^3 hybridized alkyl carbon donates electron density to the sp^2 hybridized alkenyl carbon.**




Related arguments involving $C_{sp^3} - C_{sp^2}$ bonds were used in Section 8.2C to explain why the stability of an alkene increases with increasing alkyl substitution.

The bond dipole places a partial negative charge on the alkenyl carbon (sp^2) relative to the alkyl carbon (sp^3) because an sp^2 hybridized orbital has greater percent *s*-character (33%) than an sp^3 hybridized orbital (25%). **In a cis isomer, the two $C_{sp^3} - C_{sp^2}$ bond dipoles reinforce each other, yielding a small net molecular dipole. In a trans isomer, the two bond dipoles cancel.**



- A cis alkene is more polar than a trans alkene, giving it a slightly higher boiling point and making it more soluble in polar solvents.


Problem 10.10 Rank the following isomers in order of increasing boiling point.



10.5 Interesting Alkenes


Ethylene is prepared from petroleum by a process called **cracking.** Ethylene is the most widely produced organic chemical, serving as the starting material not only for the polymer **polyethylene,** a widely used plastic, but also for many other useful organic compounds, as shown in Figure 10.4.

Figure 10.4
Ethylene, an industrial starting material for many useful products



Numerous organic compounds containing carbon–carbon double bonds have been isolated from natural sources (Figure 10.5).


Figure 10.5
Four naturally occurring alkenes




10.6 Lipids—Part 2

Lipids are water-insoluble biomolecules composed largely of nonpolar C – C and C – H bonds (Section 4.15).

General structure of an ester:



Understanding the geometry of C−C double bonds provides an insight into the properties of **triacylglycerols,** the most abundant lipids. Triacylglycerols contain three ester groups, each having a long carbon chain (abbreviated as R, R', and R") bonded to a carbonyl group (C=O).



triacylglycerol

R groups have 11–19 C's.


[Three ester groups are labeled in red.]

10.6A Fatty Acids



Linoleic and linolenic acids are essential fatty acids, meaning they cannot be synthesized in the body and must therefore be obtained in the diet. A common source of these essential fatty acids is whole milk. Babies fed a diet of nonfat milk in their early months do not thrive because they do not obtain enough of these essential fatty acids.

Triacylglycerols are hydrolyzed to glycerol (a triol), and three **fatty acids** of general structure RCO_2H. Naturally occurring fatty acids contain 12–20 carbon atoms, with a carboxy group (CO_2H) at one end.



triacylglycerol





- **Saturated fatty acids have no double bonds in their long hydrocarbon chains, and unsaturated fatty acids have one or more double bonds in their hydrocarbon chains.**
- **Double bonds in naturally occurring fatty acids have the *Z* configuration.**

Table 10.2 lists the structure and melting point of four fatty acids containing 18 carbon atoms. Stearic acid is one of the two most common saturated fatty acids, and oleic and linoleic acids are the most common unsaturated ones. The data show the effect of *Z* double bonds on the melting point of fatty acids.

- **As the number of double bonds in the fatty acid *increases,* the melting point *decreases.***


The three-dimensional structures of the fatty acids in Figure 10.6 illustrate how *Z* double bonds introduce kinks in the long hydrocarbon chain, decreasing the ability of the fatty acid to pack well in a crystalline lattice. **The larger the number of *Z* double bonds, the more kinks in the hydrocarbon chain, and the lower the melting point.**

Table 10.2 The Effect of Double Bonds on the Melting Point of Fatty Acids

Name	Structure	Mp (°C)
Stearic acid (0 C=C)		69
Oleic acid (1 C=C)		4
Linoleic acid (2 C=C)		–5
Linolenic acid (3 C=C)		–11


Increasing number of double bonds

Problem 10.11 Linolenic acid (Table 10.2) and stearidonic acid are omega-3 fatty acids, unsaturated fatty acids that contain the first double bond located at C3, when numbering begins at the methyl end of the chain. Predict how the melting point of stearidonic acid compares with the melting points of linolenic and stearic acids. A current avenue of research is examining the use of soybean oil enriched in stearidonic acid as a healthier alternative to vegetable oils that contain fewer degrees of unsaturation.




stearidonic acid


Figure 10.6 Three-dimensional structure of four C_{18} fatty acids




stearic acid



oleic acid



linoleic acid



linolenic acid

10.6B Fats and Oils



Canola, soybeans, and flaxseed are excellent dietary sources of linolenic acid, an essential fatty acid. Oils derived from omega-3 fatty acids (Problem 10.11) are currently thought to be especially beneficial for individuals at risk of developing coronary artery disease.

Fats and **oils** are triacylglycerols with different physical properties.

- Fats have higher melting points—they are *solids* at room temperature.
- Oils have lower melting points—they are *liquids* at room temperature.

The identity of the three fatty acids in the triacylglycerol determines whether it is a fat or an oil. **Increasing the number of double bonds in the fatty acid side chains decreases the melting point of the triacylglycerol.**

- Fats are derived from fatty acids having few double bonds.
- Oils are derived from fatty acids having a larger number of double bonds.


Saturated fats are typically obtained from animal sources, whereas unsaturated oils are common in vegetable sources. Thus, butter and lard are high in saturated triacylglycerols, and olive oil and safflower oil are high in unsaturated triacylglycerols. An exception to this generalization is coconut oil, which is composed largely of saturated alkyl side chains.

Considerable evidence suggests that an elevated cholesterol level is linked to an increased risk of heart disease. Saturated fats stimulate cholesterol synthesis in the liver, thus increasing the cholesterol concentration in the blood.


10.7 Preparation of Alkenes

Recall from Chapters 8 and 9 that alkenes can be prepared from alkyl halides and alcohols via elimination reactions. For example, **dehydrohalogenation of alkyl halides with strong base yields alkenes via an E2 mechanism** (Sections 8.4 and 8.5).


- Typical bases include $^-$OH and $^-$OR [especially $^-OC(CH_3)_3$], and nonnucleophilic bases such as DBU and DBN.
- Alkyl tosylates can also be used as starting materials under similar reaction conditions (Section 9.13).




The acid-catalyzed dehydration of alcohols with H_2SO_4 or TsOH yields alkenes, too (Sections 9.8 and 9.9). The reaction occurs via an E1 mechanism for 2° and 3° alcohols, and an E2 mechanism for 1° alcohols. E1 reactions involve carbocation intermediates, so rearrangements are possible. **Dehydration can also be carried out with $POCl_3$ and pyridine** by an E2 mechanism (Section 9.10).



These elimination reactions are stereoselective and regioselective, so the most stable alkene is usually formed as the major product.




major product



major product
trans disubstituted alkene

Problem 10.12 Draw the products of each elimination reaction.



10.8 Introduction to Addition Reactions

Because the C−C π bond of an alkene is much weaker than a C−C σ bond, the characteristic reaction of alkenes is **addition: the π bond is broken and two new σ bonds are formed.**



A π bond is broken. Two σ bonds are formed.


Alkenes are electron rich, as seen in the electrostatic potential plot in Figure 10.7. The electron density of the π bond is concentrated above and below the plane of the molecule, making the π bond more exposed than the σ bond.

What kinds of reagents add to the weak, electron-rich π bond of alkenes? There are many of them, and that can make alkene chemistry challenging. To help you organize this information, keep in mind the following:

The addition reactions of alkenes are discussed in Sections 10.9–10.16 and in Chapter 12 (Oxidation and Reduction).


- **Every reaction of alkenes involves addition: the π bond is always broken.**
- **Because alkenes are electron rich, simple alkenes do *not* react with nucleophiles or bases, reagents that are themselves electron rich. Alkenes react with *electrophiles*.**

Figure 10.7
Electrostatic potential plot of ethylene



- The red electron-rich region of the π bond is located above and below the plane of the molecule. Because the plane of the alkene depicted in this electrostatic potential plot is tipped, only the red region above the molecule is visible.


The stereochemistry of addition is often important in delineating a reaction's mechanism. Because the carbon atoms of a double bond are both trigonal planar, the elements of X and Y can be added to them from the **same side** or from **opposite sides.**



- *Syn addition* takes place when both X and Y are added from the *same* side.
- *Anti addition* takes place when X and Y are added from *opposite* sides.

Five reactions of alkenes are discussed in Chapter 10 and each is illustrated in Figure 10.8, using cyclohexene as the starting material.

Figure 10.8
Five addition reactions of cyclohexene




- In each reaction, the π bond is broken and two new σ bonds are formed.

10.9 Hydrohalogenation—Electrophilic Addition of HX

Hydrohalogenation of an alkene to form an alkyl halide is the reverse of the dehydrohalogenation of an alkyl halide to form an alkene, a reaction discussed in detail in Sections 8.4 and 8.5.


Hydrohalogenation results in the addition of hydrogen halides HX (X = Cl, Br, and I) to alkenes to form alkyl halides.



Two bonds are broken in this reaction—the weak π bond of the alkene and the HX bond—and two new σ bonds are formed—one to H and one to X. Because X is more electronegative than H, the H−X bond is polarized, with a partial positive charge on H. Because the electrophilic (H) end of HX is attracted to the electron-rich double bond, these reactions are called **electrophilic additions.**


To draw the products of an addition reaction:

- Locate the C−C double bond.
- Identify the σ bond of the reagent that breaks—namely, the H−X bond in hydrohalogenation.
- Break the π bond of the alkene and the σ bond of the reagent, and form two new σ bonds to the C atoms of the double bond.



Addition reactions are exothermic because the two σ bonds formed in the product are stronger than the σ and π bonds broken in the reactants. For example, $\Delta H°$ for the addition of HBr to ethylene is –60 kJ/mol, as illustrated in Figure 10.9.

Figure 10.9
The addition of HBr to $CH_2=CH_2$, an exothermic reaction



$\Delta H°$ calculation:

[1] Bonds broken

	$\Delta H°$ (kJ/mol)
$CH_2=CH_2$ π bond	+267
H–Br	+368
Total	+635 kJ/mol

Energy needed to break bonds.

[2] Bonds formed

	$\Delta H°$ (kJ/mol)
$BrCH_2CH_2$–H	−410
CH_3CH_2–Br	−285
Total	−695 kJ/mol

Energy released in forming bonds.

[3] Overall $\Delta H°$ =

sum in Step [1]
+
sum in Step [2]

+635 kJ/mol
−695 kJ/mol


$\Delta H°$ = –60 kJ/mol

The reaction is exothermic.

[Values taken from Appendix C.]

The mechanism of electrophilic addition of HX consists of **two steps:** addition of H^+ to form a carbocation, followed by nucleophilic attack of X^-. The mechanism is illustrated for the reaction of *cis*-but-2-ene with HBr in Mechanism 10.1.

Mechanism 10.1 Electrophilic Addition of HX to an Alkene




1. The π bond of the alkene attacks the H of HBr to form a new C–H bond and a carbocation in the rate-determining in step.
2. Nucleophilic attack of Br^- on the carbocation forms the new C–Br bond.


The mechanism of electrophilic addition consists of two successive Lewis acid–base reactions. In Step [1], the **alkene is the Lewis base** that donates an electron pair to **H–Br, the Lewis acid,** while in Step [2], **Br^- is the Lewis base** that donates an electron pair to the **carbocation, the Lewis acid.**

An energy diagram for the reaction of $CH_3CH{=}CHCH_3$ with HBr is given in Figure 10.10. Each step has its own energy barrier with a transition state at each energy maximum. Because Step [1] has a higher energy transition state, it is rate-determining. $\Delta H°$ for Step [1] is positive because more bonds are broken than formed, whereas $\Delta H°$ for Step [2] is negative because only bond making occurs.

Figure 10.10


Energy diagram for electrophilic addition: $CH_3CH{=}CHCH_3 + HBr \rightarrow CH_3CH_2CH(Br)CH_3$






- The mechanism has two steps, so there are two energy barriers.
- Step [1] is rate-determining.

Problem 10.13 What product is formed when each alkene is treated with HCl?


a.  b.  c. 

Problem 10.14 Draw a stepwise mechanism for the following reaction. Draw the transition state for each step.



10.10 Markovnikov's Rule

With an unsymmetrical alkene, HX can add to the double bond to give two constitutional isomers.




propene — 1-chloropropane — 2-chloropropane

only product

For example, HCl addition to propene could in theory form 1-chloropropane by addition of H and Cl to C2 and C1, respectively, and 2-chloropropane by addition of H and Cl to C1 and C2, respectively. In fact, **electrophilic addition forms *only* 2-chloropropane.** This is a specific example of a general trend called **Markovnikov's rule,** named for the Russian chemist who first determined the regioselectivity of electrophilic addition of HX.

- **Markovnikov's rule: In the addition of HX to an unsymmetrical alkene, the H atom bonds to the less substituted carbon atom—that is, the carbon that has more H atoms to begin with.**


The basis of Markovnikov's rule is the formation of a carbocation in the rate-determining step of the mechanism. With propene, there are two possible paths for this first step, depending on which carbon atom of the double bond forms the new bond to hydrogen.



The Hammond postulate was first introduced in Section 7.15 to explain the relative rate of S_N1 reactions with 1°, 2°, and 3° RX.

Path [1] forms a highly unstable 1° carbocation, whereas Path [2] forms a **more stable 2° carbocation.** According to the Hammond postulate, Path [2] is faster because formation of the carbocation is an endothermic process, so **the transition state to form the more stable 2° carbocation is lower in energy** (Figure 10.11).

Figure 10.11
Electrophilic addition and the Hammond postulate





- The E_a for formation of the more stable 2° carbocation is lower than the E_a for formation of the 1° carbocation. The 2° carbocation is formed faster.

- **In the addition of HX to an unsymmetrical alkene, the H atom is added to the less substituted carbon to form the more stable, more substituted carbocation.**

Similar results are seen in any electrophilic addition involving an intermediate carbocation: **the more stable, more substituted carbocation is formed by addition of the electrophile to the less substituted carbon.**


Problem 10.15 Draw the products formed when each alkene is treated with HCl.

a. 
b. 
c. 

Problem 10.16 Use the Hammond postulate to explain why $(CH_3)_2C{=}CH_2$ reacts faster than $CH_3CH{=}CH_2$ in electrophilic addition of HX.

Because carbocations are formed as intermediates in hydrohalogenation, carbocation rearrangements can occur, as illustrated in Sample Problem 10.3.


Sample Problem 10.3 Draw a stepwise mechanism for the following reaction.



Solution

Because the carbon skeletons of the starting material and product are different—the alkene reactant has a 4° carbon and the product alkyl halide does not—a carbocation rearrangement must have occurred.


Markovnikov addition of HBr adds H^+ to the less substituted end of the double bond, forming a 2° carbocation in Step [1].



2° carbocation
new bond shown in red




—Continued

Rearrangement of the 2° carbocation by a 1,2-methyl shift forms a more stable 3° carbocation in Step [2]. Nucleophilic attack of Br^- forms the product, a 3° alkyl halide, in Step [3].



Problem 10.17 Treatment of 3-methylcyclohexene with HCl yields two products, 1-chloro-3-methylcyclohexane and 1-chloro-1-methylcyclohexane. Draw a mechanism to explain this result.

Problem 10.18 Addition of HBr to which of the following alkenes will lead to a rearrangement?


a.  b.  c. 

10.11 Stereochemistry of Electrophilic Addition of HX

To understand the stereochemistry of electrophilic addition, recall two stereochemical principles learned in Chapters 7 and 9.

- **Trigonal planar atoms react with reagents from two directions with equal probability (Section 7.13C).**
- **Achiral starting materials yield achiral or racemic products (Section 9.15).**


Many hydrohalogenation reactions begin with an **achiral reactant** and form an **achiral product.** For example, the addition of HBr to cyclohexene, an achiral alkene, forms bromocyclohexane, an achiral alkyl halide.



cyclohexene
achiral starting material

bromocyclohexane
achiral product

Because addition converts sp^2 hybridized carbons to sp^3 hybridized carbons, sometimes new stereogenic centers are formed from hydrohalogenation. Markovnikov addition of HCl to 1,3,3-trimethylcyclohexene, an achiral alkene, forms one constitutional isomer, 1-chloro-1,3,3-trimethylcyclohexane. Because this product now has a stereogenic center at one of the newly formed sp^3 hybridized carbons (labeled in blue), **an equal amount of two enantiomers—a racemic mixture—**must form.




1,3,3-trimethylcyclohexene
achiral starting material

1-chloro-1,3,3-trimethyl-
cyclohexane


A **B**

The product has one new stereogenic center, so two enantiomers are formed.

The mechanism of hydrohalogenation illustrates why two enantiomers are formed. Initial addition of the electrophile H^+ (from HCl) occurs from **either side of the planar double bond** to form a carbocation. Both modes of addition (from above and below) generate the same **achiral carbocation.** Either representation of this carbocation can then be used to draw the second step of the mechanism.



Nucleophilic attack of Cl^- on the trigonal planar carbocation also occurs from two different directions, forming two products, **A** and **B,** having a new stereogenic center. **A** and **B** are not superimposable, so they are **enantiomers.** Because attack from either direction occurs with equal probability, a **racemic mixture** of **A** and **B** is formed.



The terms **cis** and **trans** refer to the arrangement of groups in a particular compound, usually an alkene or a disubstituted cycloalkane. The terms **syn** and **anti** describe the stereochemistry of a process—for example, how two groups are added to a double bond.

Because hydrohalogenation begins with a **planar** double bond and forms a **planar** carbocation, addition of H and Cl occurs in two different ways. The elements of H and Cl can both be added from the same side of the double bond—that is, **syn addition**—or they can be added from opposite sides—that is, **anti addition.** *Both* modes of addition occur in this two-step reaction mechanism.


- **Hydrohalogenation occurs with syn and anti addition of HX.**


Table 10.3 summarizes the characteristics of electrophilic addition of HX to alkenes.

Table 10.3 Summary: Electrophilic Addition of HX to Alkenes

	Observation
Mechanism	• The mechanism involves two steps. • The rate-determining step forms a carbocation. • Rearrangements can occur.
Regioselectivity	• Markovnikov's rule is followed. In unsymmetrical alkenes, H bonds to the less substituted C to form the more stable carbocation.
Stereochemistry	• Syn and anti addition occur.

Problem 10.19 Draw the products, including stereochemistry, of each reaction.


a. 

b. 

Problem 10.20 Draw all stereoisomers formed when 1,2-dimethylcyclohexene is treated with HCl. Label pairs of enantiomers.


10.12 Hydration—Electrophilic Addition of Water

Hydration results in the addition of water to an alkene to form an alcohol. H_2O itself is too weak an acid to protonate an alkene, but with added H_2SO_4, H_3O^+ is formed and addition readily occurs.



Hydration is simply another example of **electrophilic addition.** The first two steps of the mechanism are similar to those of electrophilic addition of HX—that is, addition of H^+ (from H_3O^+) to generate a carbocation, followed by nucleophilic attack of H_2O. Mechanism 10.2 illustrates the addition of H_2O to cyclohexene to form cyclohexanol.

Mechanism 10.2 Electrophilic Addition of H_2O to an Alkene—Hydration




1. The π bond of the alkene attacks the H of H_3O^+ to form a new C–H bond and a carbocation in the rate-determining step.
2. Nucleophilic attack of H_2O on the carbocation forms the new C–O bond.
3. Removal of a proton with H_2O forms a neutral alcohol. Because the acid used in Step [1] is regenerated in Step [3], the reaction is acid-catalyzed.

Hydration of an alkene to form an alcohol is the reverse of the dehydration of an alcohol to form an alkene, a reaction discussed in detail in Section 9.8.

There are three consequences to the formation of carbocation intermediates.


- **In unsymmetrical alkenes, H adds to the less substituted carbon to form the more stable carbocation; that is, Markovnikov's rule holds.**
- **Addition of H and OH occurs in both a syn and anti fashion.**
- **Carbocation rearrangements can occur.**

Alcohols add to alkenes, forming ethers, using the same mechanism. Addition of CH_3OH to 2-methylpropene, for example, forms *tert*-butyl methyl ether (**MTBE**), a high octane fuel additive described in Section 3.4C.



tert-butyl methyl ether
MTBE

Problem 10.21 What two alkenes give rise to each alcohol as the major product of acid-catalyzed hydration?




Problem 10.22 What stereoisomers are formed when pent-1-ene is treated with H_2O and H_2SO_4?

10.13 Halogenation—Addition of Halogen

Halogenation results in the addition of halogen X_2 (X = Cl or Br) to an alkene, forming a **vicinal dihalide.**




Bromination is a simple chemical test for the presence of π bonds. When an alkene such as hex-1-ene is treated with red liquid bromine, the Br_2 adds to the double bond to give a colorless product, as seen in the right test tube. With the alkane hexane (test tube on the left), no reaction occurs, and the red color of Br_2 remains.




A bond is broken. vicinal dihalide

Halogenation is synthetically useful only with Cl_2 and Br_2. The dichlorides and dibromides formed in this reaction serve as starting materials for the synthesis of alkynes, as we learned in Section 8.10.



Halogens add to π bonds because halogens are **polarizable.** The electron-rich double bond induces a dipole in an approaching halogen molecule, making one halogen atom electron deficient and the other electron rich ($X^{\delta+}-X^{\delta-}$). **The electrophilic halogen atom is then attracted to the nucleophilic double bond,** making addition possible.


Two facts demonstrate that halogenation follows a different mechanism from that of hydrohalogenation or hydration. First, no rearrangements occur, and second, only anti addition of X_2 is observed. For example, treatment of cyclohexene with Br_2 yields two **trans** enantiomers formed by **anti addition.**



enantiomers


These facts suggest that **carbocations are *not* intermediates in halogenation.** Unstable carbocations rearrange, and both syn and anti addition is possible with carbocation intermediates. The accepted mechanism for halogenation comprises **two steps,** but it does *not* proceed with formation of a carbocation, as shown in Mechanism 10.3.

Mechanism 10.3 Addition of X_2 to an Alkene—Halogenation




1. Four bonds are broken or formed to generate an unstable bridged halonium ion that contains a three-membered ring. The electron pair in the π bond and a lone pair on a halogen are used to form two new C – X bonds, and the X – X bond is cleaved.
2. Nucleophilic attack of X^- ring opens the bridged halonium ion and forms a new C – X bond.

Bridged halonium ions resemble carbocations in that they are short-lived intermediates that react readily with nucleophiles. Carbocations are inherently unstable because only six electrons surround carbon, whereas **halonium ions are unstable because they contain a strained three-membered ring** with a positively charged halogen atom.




Problem 10.23 Draw the products of each reaction, including stereochemistry.




10.14 Stereochemistry of Halogenation

How does the proposed mechanism invoking a bridged halonium ion intermediate explain the observed **trans products of halogenation?** For example, chlorination of cyclopentene affords both enantiomers of *trans*-1,2-dichlorocyclopentane, with *no* cis products.




trans enantiomers

Initial addition of the electrophile Cl^+ (from Cl_2) occurs from either side of the planar double bond to form the bridged chloronium ion. In this example, both modes of addition (from above and below) generate the same **achiral** intermediate, so either representation can be used to draw the second step.



The opening of bridged halonium ion intermediates resembles the opening of epoxide rings with nucleophiles discussed in Section 9.15.

In the second step, **nucleophilic attack of Cl^- must occur from the back side**—that is, from the side of the five-membered ring opposite to the side having the bridged chloronium ion. Because the nucleophile attacks from below in this example and the leaving group departs from above, the two Cl atoms in the product are oriented **trans** to each other. Backside attack occurs with equal probability at either carbon of the three-membered ring to yield an equal amount of two enantiomers—**a racemic mixture.**



In summary, the mechanism for halogenation of alkenes occurs in two steps:

- **Addition of X^+ forms an unstable bridged halonium ion in the rate-determining step.**
- **Nucleophilic attack of X^- occurs from the back side to form trans products. The overall result is anti addition of X_2 across the double bond.**

Because halogenation occurs exclusively in an anti fashion, cis and trans alkenes yield different stereoisomers. Halogenation of alkenes is a **stereospecific reaction.**

- **A reaction is stereospecific when each of two specific stereoisomers of a starting material yields a particular stereoisomer of a product.**

cis-But-2-ene yields two enantiomers, whereas *trans*-but-2-ene yields a single achiral meso compound, as shown in Figure 10.12.

Problem 10.24 Draw all stereoisomers formed in each reaction.





Figure 10.12
Halogenation of *cis*- and *trans*-but-2-ene





To draw the products of halogenation:

- Add Br_2 in an **anti** fashion across the double bond, leaving all other groups in their original orientations. With the alkene drawn in the plane of the page, one Br adds from the front (ending up on a wedge), and one Br adds from the back (ending up on a dashed wedge).
- Sometimes this reaction produces two stereoisomers, as in the case of *cis*-but-2-ene, which forms an equal amount of two enantiomers. Sometimes it produces a single compound, as in the case of *trans*-but-2-ene, where a meso compound is formed.

10.15 Halohydrin Formation


Treatment of an alkene with a halogen $\mathbf{X_2}$ and $\mathbf{H_2O}$ forms a **halohydrin** by addition of the elements of **X** and **OH** to the double bond.





The mechanism for halohydrin formation is similar to the mechanism for halogenation: addition of the electrophile X^+ (from X_2) to form a **bridged halonium ion,** followed by nucleophilic attack by H_2O from the back side on the three-membered ring (Mechanism 10.4). Even though X^- is formed in Step [1] of the mechanism, its concentration is small compared to H_2O (often the solvent), so H_2O and *not* X^- is the nucleophile.


Mechanism 10.4 Addition of X and OH—Halohydrin Formation



1. Four bonds are broken or formed to generate an unstable bridged halonium ion that contains a three-membered ring. The electron pair in the π bond and a lone pair on a halogen are used to form two new C – X bonds, and the X – X bond is cleaved.
2. Nucleophilic attack of H_2O ring opens the bridged halonium ion and forms a new C – O bond.
3. Loss of a proton forms the halohydrin.

Recall from Section 7.8C that DMSO (dimethyl sulfoxide) is a polar aprotic solvent.

Although the combination of Br_2 and H_2O effectively forms **bromohydrins** from alkenes, other reagents can also be used. Bromohydrins are also formed with ***N*-bromosuccinimide** (abbreviated as **NBS**) in **aqueous DMSO** [$(CH_3)_2S{=}O$]. NBS serves as a source of Br_2, which then goes on to form a bromohydrin by the same reaction mechanism.




N-bromosuccinimide
NBS

bromohydrin


10.15A Stereochemistry and Regioselectivity of Halohydrin Formation

Because the bridged halonium ion ring is opened by backside attack of H_2O, addition of X and OH occurs in an **anti** fashion and **trans** products are formed.



trans enantiomers


Sample Problem 10.4 Draw the products of the following reaction, including stereochemistry.



trans-but-2-ene

Solution


The reagent (Br_2 + H_2O) adds the elements of Br and OH to a double bond in an **anti** fashion—that is, from **opposite** sides. To draw two products of anti addition: add Br from above and OH from below in one product; then add Br from below and OH from above in the other product. In this example, the two products are nonsuperimposable mirror images—**enantiomers.**



trans-but-2-ene

enantiomers

With unsymmetrical alkenes, two constitutional isomers are possible from addition of X and OH, but only one is formed. **The preferred product has the electrophile X^+ bonded to the less substituted carbon atom**—that is, the carbon that has more H atoms to begin with in the reacting alkene. Thus, the **nucleophile (H_2O) bonds to the more substituted carbon.**



only product

Br ends up on the less substituted C.

not formed

This result is reminiscent of the opening of epoxide rings with acids HZ (Z = a nucleophile), which we encountered in Section 9.15B. As in the opening of an epoxide ring, **nucleophilic attack occurs at the more substituted carbon end of the bridged halonium ion** because that carbon is better able to accommodate a partial positive charge in the transition state.




Table 10.4 summarizes the characteristics of halohydrin formation.

Problem 10.25 Draw the products of each reaction and indicate their stereochemistry.




Table 10.4 Summary: Conversion of Alkenes to Halohydrins

	Observation
Mechanism	• The mechanism involves three steps. • The rate-determining step forms a bridged halonium ion. • No rearrangements can occur.
Regioselectivity	• The electrophile X^+ bonds to the less substituted carbon.
Stereochemistry	• Anti addition occurs.

10.15B Halohydrins: Useful Compounds in Organic Synthesis

Because halohydrins are easily converted to epoxides by intramolecular S_N2 reaction (Section 9.6), they have been used in the synthesis of many naturally occurring compounds. Key steps in the synthesis of estrone, a female sex hormone, are illustrated in Figure 10.13.


Figure 10.13 The synthesis of estrone from a chlorohydrin



- Chlorohydrin **B,** prepared from alkene **A** by addition of Cl and OH, is converted to epoxide **C** with base. **C** is converted to estrone in one step.


10.16 Hydroboration–Oxidation

Hydroboration–oxidation is a two-step reaction sequence that converts an alkene to an alcohol.




- *Hydroboration* is the addition of borane (BH_3) to an alkene, forming an alkylborane.
- *Oxidation* converts the C – B bond of the alkylborane to a C – O bond.

Hydroboration–oxidation results in **addition of H_2O** to an alkene.




Borane (BH_3) is a reactive gas that exists mostly as the dimer, diborane (B_2H_6). Borane is a strong **Lewis acid** that reacts readily with Lewis bases. For ease in handling in the laboratory, it is commonly used as a complex with tetrahydrofuran (THF).



10.16A Hydroboration


The first step in hydroboration–oxidation is **addition of the elements of H and BH_2** to the π bond of the alkene, forming an intermediate alkylborane.



Because syn addition to the double bond occurs and no carbocation rearrangements are observed, carbocations are *not* formed during hydroboration, as shown in Mechanism 10.5. The proposed

Mechanism 10.5 Addition of H and BH_2—Hydroboration


One step The π bond and H – BH_2 bonds break as the C – H and C – B bonds form.



mechanism involves a **concerted addition of H and BH_2 from the same side of the planar double bond:** the π bond and H−BH_2 bond are broken as two new σ bonds are formed. Because four atoms are involved, the transition state is said to be **four-centered.**


Because the alkylborane formed by reaction with one equivalent of alkene still has two B−H bonds, it can react with two more equivalents of alkene to form a trialkylborane. This is illustrated in Figure 10.14 for the reaction of $CH_2{=}CH_2$ with BH_3.

Figure 10.14
Conversion of BH_3 to a trialkylborane with three equivalents of $CH_2{=}CH_2$




- We often draw hydroboration as if addition stopped after one equivalent of alkene reacts with BH_3. Instead, all three B−H bonds actually react with three equivalents of an alkene to form a trialkylborane. The term **organoborane** is used for any compound with a carbon–boron bond.

Because only one B−H bond is needed for hydroboration, commercially available dialkylboranes having the general structure **R_2BH** are sometimes used instead of BH_3. A common example is 9-borabicyclo[3.3.1]nonane **(9-BBN).** 9-BBN undergoes hydroboration in the same manner as BH_3.



9-borabicyclo[3.3.1]nonane
9-BBN

Hydroboration is regioselective. **With unsymmetrical alkenes, the boron atom bonds to the less substituted carbon atom.** For example, addition of BH_3 to propene forms an alkylborane with the B bonded to the terminal carbon atom.



only product *not* formed

B bonds to the terminal C.


Because H is more electronegative than B, the B−H bond is polarized to give boron a partial positive charge ($H^{\delta-}-B^{\delta+}$), making BH_2 the electrophile in hydroboration.

Steric factors explain this regioselectivity. The larger boron atom bonds to the less sterically hindered, more accessible carbon atom.



Electronic factors are also used to explain this regioselectivity. If bond breaking and bond making are not completely symmetrical, boron bears a partial negative charge in the transition state and carbon bears a partial positive charge. Because alkyl groups stabilize a positive charge, the more stable transition state has the partial positive charge on the more substituted carbon, as illustrated in Figure 10.15.

- **In hydroboration, the boron atom bonds to the less substituted carbon.**

Figure 10.15
Hydroboration of an unsymmetrical alkene




Problem 10.26 What alkylborane is formed from hydroboration of each alkene?

a. 
b. 
c. 

10.16B Oxidation of the Alkylborane

Because alkylboranes react rapidly with water and spontaneously burn when exposed to the air, they are oxidized, without isolation, with basic hydrogen peroxide (H_2O_2, HO^-). **Oxidation replaces the C−B bond with a C−O bond, forming a new OH group with retention of configuration;** that is, the **OH group replaces the BH_2 group in the same position** relative to the other three groups on carbon.




Thus, to draw the product of a hydroboration–oxidation reaction, keep in mind two stereochemical facts:

- **Hydroboration occurs with syn addition.**
- **Oxidation occurs with retention of configuration.**


The overall result of this two-step sequence is **syn addition of the elements of H and OH** to a double bond, as illustrated in Sample Problem 10.5. The OH group bonds to the *less* substituted carbon.

Sample Problem 10.5 Draw the product of the following reaction sequence, including stereochemistry.






Solution

In Step [1], **syn addition of BH_3 to the unsymmetrical alkene adds the BH_2 group to the less substituted carbon from above and below the planar double bond.** Two enantiomeric alkylboranes are formed. In Step [2], oxidation replaces the BH_2 group with OH in each enantiomer with **retention of configuration** to yield two alcohols that are also enantiomers.





Hydroboration–oxidation results in the **addition of H and OH in a syn fashion** across the double bond. The achiral alkene is converted to an equal mixture of two enantiomers—that is, a **racemic mixture of alcohols.**

Problem 10.27 Draw the products formed when each alkene is treated with BH_3 followed by H_2O_2, HO^-. Include the stereochemistry at all stereogenic centers.

a.  b.  c. 

Problem 10.28 What alkene can be used to prepare each alcohol as the exclusive product of a two-step hydroboration–oxidation sequence?

a. 

b. 


c. 


Table 10.5 summarizes the features of hydroboration–oxidation.

Table 10.5 Summary: Hydroboration–Oxidation of Alkenes

	Observation
Mechanism	• The addition of H and BH_2 occurs in one step. • No rearrangements can occur.
Regioselectivity	• The OH group bonds to the less substituted carbon atom.
Stereochemistry	• Syn addition occurs. • OH replaces BH_2 with retention of configuration.

Hydroboration–oxidation is a very common method for adding H_2O across a double bond. One example is shown in the synthesis of **artemisinin** (or **qinghaosu**), the active component of **qinghao,** a Chinese herbal remedy used for the treatment of malaria (Figure 10.16). As mentioned in the Prologue, currently recommended antimalarial therapy consists of a combination of drugs

Figure 10.16 An example of hydroboration–oxidation in synthesis



• The carbon atoms of artemisinin that come from alcohol **A** are indicated in red.

that includes artemisinin or a related compound. A ball-and-stick model of artemisinin appears on the cover of this text.


10.16C A Comparison of Hydration Methods

Hydration (H_2O, H^+) and hydroboration–oxidation (BH_3 followed by H_2O_2, HO^-) both add the elements of H_2O across a double bond. Despite their similarities, these reactions often form different constitutional isomers, as shown in Sample Problem 10.6.


Sample Problem 10.6 Draw the product formed when $CH_3CH_2CH_2CH_2CH{=}CH_2$ is treated with (a) H_2O, H_2SO_4; and (b) BH_3 followed by H_2O_2, HO^-.

Solution

With H_2O + H_2SO_4, electrophilic addition of H and OH places the **H atom on the less substituted carbon** of the alkene to yield a **2° alcohol.** In contrast, addition of BH_3 gives an alkylborane with the **BH_2 group on the less substituted terminal carbon** of the alkene. Oxidation replaces BH_2 by OH to yield a **1° alcohol.**



Problem 10.29 Draw the constitutional isomer formed when the following alkenes are treated with each set of reagents: [1] H_2O, H_2SO_4; or [2] BH_3 followed by H_2O_2, ^-OH.



10.17 Keeping Track of Reactions

Chapters 7–10 have introduced three basic kinds of organic reactions: **nucleophilic substitution, β elimination,** and **addition.** In the process, many specific reagents have been discussed and the stereochemistry that results from many different mechanisms has been examined. **How can we keep track of all the reactions?**

To make the process easier, **remember that most organic molecules undergo only one or two different kinds of reactions.** For example:

- **Alkyl halides undergo substitution and elimination because they have good leaving groups.**
- **Alcohols also undergo substitution and elimination, but can do so only when OH is made into a good leaving group.**
- **Alkenes undergo addition because they have easily broken π bonds.**

You must still learn many reaction details, and in truth, there is no one method to learn them. ***You must practice these reactions over and over again, not by merely looking at them, but by writing them.*** Some students do this by making a list of specific reactions for each functional group, and then rewriting them with different starting materials. Others make flash cards: index cards that have the starting material and reagent on one side and the product on the other. Whatever method you choose, **the details must become second nature,** much like the answers to simple addition problems, such as, what is the sum of 2 + 2?


Learning reactions is really a two-step process.


- **First, learn the basic type of reaction for a functional group. This provides an overall organization to the reactions.**
- **Then, learn the specific reagents for each reaction. It helps to classify the reagent according to its properties. Is it an acid or a base? Is it a nucleophile or an electrophile? Is it an oxidizing agent or a reducing agent?**

Sample Problem 10.7 illustrates this process.

Sample Problem 10.7

Draw the product of each reaction.


a. 

b. 

Solution


In each problem, identify the functional group to determine the general reaction type—substitution, elimination, or addition. Then, determine if the reagent is an electrophile, nucleophile, acid, base, and so forth.

a. The reactant is a **1° alkyl halide,** which can undergo substitution and elimination. The reagent [$KOC(CH_3)_3$] is a **strong nonnucleophilic base,** favoring elimination by an E2 mechanism.




E2 product

b. The reactant is an **alkene,** which undergoes addition reactions to its π bond. The reagent (Br_2 + H_2O) serves as the source of the **electrophile Br^+,** resulting in **addition** of Br and OH to the double bond (Section 10.15).



addition product


Problem 10.30 Draw the products of each reaction using the two-part strategy from Sample Problem 10.7.



10.18 Alkenes in Organic Synthesis

Alkenes are a central functional group in organic chemistry. **Alkenes are easily prepared by elimination reactions** such as dehydrohalogenation and dehydration. **Because their π bond is easily broken, they undergo many addition reactions** to prepare a variety of useful compounds.

Suppose, for example, that we must synthesize 1,2-dibromocyclohexane from cyclohexanol, a cheap and readily available starting material. Because there is no way to accomplish this transformation in one step, this synthesis must have at least two steps.




cyclohexanol — **starting material**

1,2-dibromocyclohexane — **product**

To solve this problem we must:

- **Work backwards from the product by asking: What type of reactions introduce the functional groups in the product?**
- **Work forwards from the starting material by asking: What type of reactions does the starting material undergo?**




In Chapter 11 we will learn about retrosynthetic analysis in more detail.

Working backwards from the product to determine the starting material from which it is made is called *retrosynthetic analysis*.

We know reactions that answer each of these questions.

Working backwards:


[1] 1,2-Dibromocyclohexane, a vicinal dibromide, can be prepared by the addition of Br_2 to **cyclohexene.**



cyclohexene — 1,2-dibromocyclohexane

Working forwards:


[2] Cyclohexanol can undergo acid-catalyzed dehydration to form **cyclohexene.**



cyclohexanol — cyclohexene


A **reactive intermediate** is an unstable intermediate like a carbocation, which is formed during the conversion of a stable starting material to a stable product. A **synthetic intermediate** is a stable compound that is the product of one step and the starting material of another in a multistep synthesis.

Cyclohexene is called a **synthetic intermediate,** or simply an **intermediate,** because it is the **product of one step and the starting material of another.** We now have a two-step sequence to convert cyclohexanol to 1,2-dibromocyclohexane, and the synthesis is complete. Take note of the central role of the alkene in this synthesis.



a synthetic intermediate

Problem 10.31 Devise a synthesis of each compound from the indicated starting material.



KEY CONCEPTS

Alkenes


General Facts About Alkenes

- Alkenes contain a carbon–carbon double bond consisting of a stronger σ bond and a weaker π bond. Each carbon is sp^2 hybridized and trigonal planar (10.1).
- Alkenes are named using the suffix ***-ene*** (10.3).
- Alkenes with different groups on each end of the double bond exist as a pair of diastereomers, identified by the prefixes *E* and *Z* (10.3B).
- Alkenes have weak intermolecular forces, giving them low mp's and bp's, and making them water insoluble. A cis alkene is more polar than a trans alkene, giving it a slightly higher boiling point (10.4).
- Because a π bond is electron rich and much weaker than a σ bond, alkenes undergo addition reactions with electrophiles (10.8).

Stereochemistry of Alkene Addition Reactions (10.8)


A reagent XY adds to a double bond in one of three different ways:

- **Syn addition**—X and Y add from the same side.




- Syn addition occurs in **hydroboration.**

- **Anti addition**—X and Y add from opposite sides.



- Anti addition occurs in **halogenation** and **halohydrin formation.**


- **Both syn and anti addition** occur when carbocations are intermediates.



- Syn and anti addition occur in **hydrohalogenation** and **hydration.**


Addition Reactions of Alkenes

[1] Hydrohalogenation (Addition of HX; X = Cl, Br, I) (10.9–10.11)



- The mechanism has two steps.
- Carbocations are formed as intermediates.
- Carbocation rearrangements are possible.
- Markovnikov's rule is followed. H bonds to the less substituted C to form the more stable carbocation.
- Syn and anti addition occur.


[2] Hydration and related reactions (Addition of H_2O or ROH) (10.12)



For both reactions:


- The mechanism has three steps.
- Carbocations are formed as intermediates.
- Carbocation rearrangements are possible.
- Markovnikov's rule is followed. H bonds to the less substituted C to form the more stable carbocation.
- Syn and anti addition occur.

[3] Halogenation (Addition of X_2; X = Cl or Br) (10.13–10.14)




- The mechanism has two steps.
- Bridged halonium ions are formed as intermediates.
- No rearrangements can occur.
- Anti addition occurs.

[4] Halohydrin formation (Addition of OH and X; X = Cl, Br) (10.15)



- The mechanism has three steps.
- Bridged halonium ions are formed as intermediates.
- No rearrangements can occur.
- X bonds to the less substituted C.
- Anti addition occurs.
- NBS in DMSO and H_2O adds Br and OH in the same fashion.

[5] Hydroboration–oxidation (Addition of H_2O) (10.16)





- Hydroboration has a one-step mechanism.
- No rearrangements can occur.
- OH bonds to the less substituted C.
- Syn addition of H_2O results.

PROBLEMS


Problems Using Three-Dimensional Models

10.32 Give the IUPAC name for each compound.

a. 


b. 

10.33 (a) Label the carbon–carbon double bond in **A** as *E* or *Z*. (b) Draw the products (including stereoisomers) formed when **A** is treated with H_2O in the presence of H_2SO_4.



A

10.34 Name the alkene depicted in the ball-and-stick model and draw the constitutional isomers formed when the alkene is treated with each reagent: (a) Br_2; (b) Br_2 in H_2O; (c) Br_2 in CH_3OH.



Degrees of Unsaturation

10.35 Calculate the number of degrees of unsaturation for each molecular formula.


a. C_6H_8 c. $C_{10}H_{16}O_2$ e. C_8H_9ClO g. C_4H_8BrN
b. $C_{40}H_{56}$ d. C_8H_9Br f. $C_7H_{11}N$ h. $C_{10}H_{18}ClNO$

10.36 How many rings and π bonds does a compound with molecular formula $C_{10}H_{14}$ possess? List all possibilities.

Nomenclature and Stereochemistry


10.37 Label the alkene in each drug as *E* or *Z*. Enclomiphene is one component of the fertility drug Clomid. Tamoxifen is an anticancer drug. Clavulanic acid is sold in combination with the antibiotic amoxicillin under the trade name Augmentin.

a.




enclomiphene

b.



tamoxifen


c.




clavulanic acid

10.38 Give the IUPAC name for each compound.


a.




c.




e.




b.



d.



f.




10.39 Give the structure corresponding to each name.

a. (*E*)-4-ethylhept-3-ene
b. 3,3-dimethylcyclopentene
c. 4-vinylcyclopentene
d. (*Z*)-3-isopropylhept-2-ene
e. *cis*-3,4-dimethylcyclopentene
f. 1-isopropyl-4-propylcyclohexene
g. 3,4-dimethylcyclohex-2-enol
h. 3,5-diethylhex-5-en-3-ol


10.40 (a) Draw all possible stereoisomers of 4-methylnon-2-ene, and name each isomer, including its *E,Z* and *R,S* prefixes. (b) Label two pairs of enantiomers. (c) Label four pairs of diastereomers.

10.41 (a) Draw the structure of (1*E*,4*R*)-1,4-dimethylcyclodecene. (b) Draw the enantiomer and name it, including its *E,Z* and *R,S* prefixes. (c) Draw two diastereomers and name them, including the *E,Z* and *R,S* prefixes.

10.42 Now that you have learned how to name alkenes in Section 10.3, name each of the following epoxides as an alkene oxide, as described in Section 9.3.




10.43 Iejimalide B, an anticancer agent with a 24-membered ring, is isolated from a tunicate found off Ie Island in Okinawa. (a) Label each double bond in iejimalide B as *E* or *Z*. (b) Label each tetrahedral stereogenic center as *R* or *S*. (c) How many stereoisomers are possible for iejimalide B?



iejimalide B


Lipids

10.44 Although naturally occurring unsaturated fatty acids generally have the *Z* configuration, elaidic acid, a C_{18} fatty acid having an *E* double bond, is present in processed foods such as margarine and cooking oils. Predict how the melting point of elaidic acid compares with the melting points of stearic and oleic acids (Table 10.2).



elaidic acid

10.45 Eleostearic acid is an unsaturated fatty acid obtained from the seeds of the tung oil tree (*Aleurites fordii*), a deciduous tree native to China. (a) Draw the structure of a stereoisomer that has a higher melting point than eleostearic acid. (b) Draw the structure of a stereoisomer that has a lower melting point.




eleostearic acid

Reactions of Alkenes


10.46 Draw the products formed when $(CH_3)_2C{=}CH_2$ is treated with each reagent.


a. HBr
b. H_2O, H_2SO_4
c. CH_3CH_2OH, H_2SO_4
d. Cl_2
e. Br_2, H_2O
f. NBS (aqueous DMSO)
g. [1] BH_3; [2] H_2O_2, HO^-


10.47 What alkene can be used to prepare each alkyl halide or dihalide as the exclusive or major product of an addition reaction?




10.48 Which alcohols can be prepared as a single product by hydroboration–oxidation of an alkene? Which alcohols can be prepared as a single product by the acid-catalyzed addition of H_2O to an alkene?


a. 


b. 


c. 


d. 


10.49 Draw the constitutional isomer formed in each reaction.


a. 

b. 

c. 


d. 


e. 


f. 

10.50 What three alkenes (excluding stereoisomers) can be used to prepare 3-chloro-3-methylhexane by addition of HCl?


10.51 Draw all stereoisomers formed in each reaction.


a. 


b. 


c. 


10.52 Draw the products of each reaction, including stereoisomers.


a. 

b. 


c. 

d. 


e. 

f. 

10.53 Which alkene reacts faster with HBr? Explain your choice.



A




B

10.54 (a) Which diastereomer of oct-4-ene yields a mixture of two enantiomers, (4*R*,5*R*)- and (4*S*,5*S*)-4,5-dibromooctane on reaction with Br_2? (b) Which diastereomer of oct-4-ene yields a single meso compound, (4*R*,5*S*)-4,5-dibromooctane?


10.55 Using *cis*- and *trans*-hex-3-ene, demonstrate that the addition of HCl is not a stereospecific reaction. Draw the structure of the stereoisomers formed from each alkene.

Mechanisms


10.56 Draw a stepwise mechanism for the following reaction.



10.57 Draw a stepwise mechanism for the following reaction, which results in ring expansion of a six-membered ring to a seven-membered ring.




10.58 Draw a stepwise mechanism for the conversion of hex-5-en-1-ol to the cyclic ether **A**.



A


10.59 Draw a stepwise mechanism that shows how all three alcohols are formed from the bicyclic alkene.



10.60 Less stable alkenes can be isomerized to more stable alkenes by treatment with strong acid. For example, 2,3-dimethylbut-1-ene is converted to 2,3-dimethylbut-2-ene when treated with H_2SO_4. Draw a stepwise mechanism for this isomerization process.


10.61 When buta-1,3-diene ($CH_2=CH-CH=CH_2$) is treated with HBr, two constitutional isomers are formed, $CH_3CHBrCH=CH_2$ and $BrCH_2CH=CHCH_3$. Draw a stepwise mechanism that accounts for the formation of both products.

10.62 Explain why the addition of HBr to alkenes **A** and **C** is regioselective, forming addition products **B** and **D,** respectively.




A **B** **C** **D**

10.63 Bromoetherification, the addition of the elements of Br and OR to a double bond, is a common method for constructing rings containing oxygen atoms. This reaction has been used in the synthesis of the polyether antibiotic monensin (Problem 21.37). Draw a stepwise mechanism for the following intramolecular bromoetherification reaction.




Synthesis

10.64 Devise a synthesis of each product from the given starting material. More than one step is required.



10.65 Devise a synthesis of each compound from cyclohexene as the starting material. More than one step is needed.



+ enantiomer

Challenge Problems


10.66 Explain why **A** is a stable compound but **B** is not.



A **B**


10.67 (a) What product(s) are formed when the *E* isomer of $C_6H_5CH=CHC_6H_5$ is treated with Br_2, followed by one equivalent of KOH? Label the resulting alkene(s) as *E* or *Z*. (b) What product(s) are formed when the *Z* isomer of $C_6H_5CH=CHC_6H_5$ is subjected to the same reaction sequence? (c) How are the compounds in parts (a) and (b) related to each other?

10.68 Alkene **A** can be isomerized to isocomene, a natural product isolated from goldenrod, by treatment with TsOH. Draw a stepwise mechanism for this conversion. (Hint: Look for a carbocation rearrangement.)




A isocomene

10.69 Lactones, cyclic esters such as compound **A,** are prepared by **halolactonization,** an addition reaction to an alkene. For example, iodolactonization of **B** forms lactone **C,** a key intermediate in the synthesis of prostaglandin $PGF_{2\alpha}$ (Section 4.15). Draw a stepwise mechanism for this addition reaction.




A **B** **C** $PGF_{2\alpha}$


10.70 Draw a stepwise mechanism for the following reaction.



10.71 Like other electrophiles, carbocations add to alkenes to form new carbocations, which can then undergo substitution or elimination reactions depending on the reaction conditions. With this in mind, consider the following reactions of nerol, a natural product isolated from lemon grass and other plant sources. Treatment of nerol with TsOH forms α-terpineol as the major product, whereas treatment of nerol with chlorosulfonic acid, HSO_3Cl, forms a constitutional isomer, α-cyclogeraniol. Write stepwise mechanisms for both processes. Each mechanism involves the addition of an electrophile—a carbocation—to a double bond.




10.72 Draw a stepwise mechanism for the following reaction. This reaction combines two processes together: the opening of an epoxide ring with a nucleophile and the addition of an electrophile to a carbon–carbon double bond. (Hint: Begin the mechanism by protonating the epoxide ring.)



11 Alkynes

11.1 Introduction
11.2 Nomenclature
11.3 Physical properties
11.4 Interesting alkynes
11.5 Preparation of alkynes
11.6 Introduction to alkyne reactions
11.7 Addition of hydrogen halides
11.8 Addition of halogen
11.9 Addition of water
11.10 Hydroboration–oxidation
11.11 Reaction of acetylide anions
11.12 Synthesis




Ethynylestradiol is a synthetic compound whose structure closely resembles the carbon skeleton of female estrogen hormones. Because it is more potent than its naturally occurring analogues, it is a component of several widely used oral contraceptives. Ethynylestradiol and related compounds with similar biological activity contain a carbon–carbon triple bond. In Chapter 11 we learn about alkynes, hydrocarbons that contain triple bonds.

In Chapter 11 we continue our focus on organic molecules with electron-rich functional groups by examining ***alkynes,*** **compounds that contain a carbon–carbon triple bond.** Like alkenes, **alkynes are nucleophiles with easily broken π bonds,** and as such, they undergo **addition** reactions with electrophilic reagents.

Alkynes also undergo a reaction that has no analogy in alkene chemistry. Because a C−H bond of an alkyne is more acidic than a C−H bond in an alkene or an alkane, alkynes are readily deprotonated with strong base. The resulting nucleophiles react with electrophiles to form new carbon–carbon σ bonds, so that complex molecules can be prepared from simple starting materials. The study of alkynes thus affords an opportunity to learn more about organic synthesis.


11.1 Introduction

Alkynes contain a carbon–carbon triple bond. A **terminal alkyne** has the triple bond at the end of the carbon chain, so that a hydrogen atom is directly bonded to a carbon atom of the triple bond. An **internal alkyne** has a carbon atom bonded to each carbon atom of the triple bond.




An alkyne has the general molecular formula $\mathbf{C_nH_{2n-2}}$, giving it four fewer hydrogens than the maximum number possible. Because every degree of unsaturation removes two hydrogens, a **triple bond introduces two degrees of unsaturation.**

Each carbon of a triple bond is ***sp*** hybridized and **linear,** and all bond angles are **180°** (Section 1.10C).




The triple bond of an alkyne consists of one σ bond and two π bonds.




- The σ bond is formed by end-on overlap of the two *sp* hybrid orbitals.
- Each π bond is formed by side-by-side overlap of two 2*p* orbitals.

Bond dissociation energies of the C−C bonds in ethylene (one σ and one π bond) and acetylene (one σ and two π bonds) can be used to estimate the strength of the second π bond of the triple bond. If we assume that the σ bond and first π bond in acetylene are similar in strength to the σ and π bonds in ethylene (368 and 267 kJ/mol, respectively), then the second π bond is worth 202 kJ/mol.




- Both π bonds of a C – C triple bond are weaker than a C – C σ bond, making them much more easily broken. As a result, alkynes undergo many addition reactions.
- Alkynes are more polarizable than alkenes because the electrons in their π bonds are more loosely held.

Skeletal structures for alkynes may look somewhat unusual, but they follow the customary convention: a carbon atom is located at the intersection of any two lines and at the end of any line; thus,




Like trans cycloalkenes, cycloalkynes with small rings are unstable. The carbon chain must be long enough to connect the two ends of the triple bond without introducing too much strain. Cyclooctyne is the smallest isolated cycloalkyne, though it decomposes upon standing at room temperature after a short time. To accommodate the triple bond in the ring, bending occurs around the *sp* hybridized C's, destabilizing cyclooctyne.



cyclooctyne

Problem 11.1 Nepheliosyne B is a novel acetylenic fatty acid isolated from a New Caledonian marine sponge. (a) Label the most acidic H atom. (b) Which carbon–carbon σ bond is shortest? (c) How many degrees of unsaturation does nepheliosyne B contain? (d) How many bonds are formed from $C_{sp}—C_{sp^3}$? (e) Label each triple bond as internal or terminal.




nepheliosyne B

11.2 Nomenclature

Alkynes are named in the same way that alkenes were named in Section 10.3.


- In the IUPAC system, change the *-ane* ending of the parent alkane to the suffix *-yne.*
- Choose the longest carbon chain that contains both atoms of the triple bond and number the chain to give the triple bond the lower number.
- Compounds with two triple bonds are named as *diynes,* those with three are named as *triynes,* and so forth.
- Compounds with both a double and a triple bond are named as *enynes.* The chain is numbered to give the first site of unsaturation (either C=C or C≡C) the lower number.

Sample Problem 11.1 Give the IUPAC name for the following alkyne.



Solution


[1] Find the longest chain that contains both carbons of the triple bond.



8 C's in the longest chain

octane ----→ octyne

[2] Number the long chain; then name and number the substituents.





two methyl groups at C6


Answer: 6,6-dimethyloct-3-yne


The simplest alkyne, HC≡CH, named in the IUPAC system as **ethyne,** is more often called **acetylene,** its common name. The two-carbon alkyl group derived from acetylene is called an **ethynyl group** (HC≡C−). Examples of alkyne nomenclature are shown in Figure 11.1.

Problem 11.2 Give the IUPAC name for each compound.

a. 

c. 

b. 

d. 


Problem 11.3 Give the structure corresponding to each of the following names.

a. *trans*-2-ethynylcyclopentanol b. 4-*tert*-butyldec-5-yne c. 3,3,5-trimethylcyclononyne


Figure 11.1
Examples of alkyne nomenclature




2,5-dimethylhept-3-yne



1-ethynyl-2-isopropylcyclohexane



hexa-1,3-diyne



5-methylhex-4-en-1-yne

11.3 Physical Properties

The physical properties of alkynes resemble those of hydrocarbons having a similar shape and molecular weight.

- **Alkynes have low melting points and boiling points.**
- **Melting points and boiling points increase as the number of carbons increases.**
- **Alkynes are soluble in organic solvents and insoluble in water.**

Problem 11.4 Explain why an alkyne often has a slightly higher boiling point than an alkene of similar molecular weight. For example, the bp of pent-1-yne is 39°C, and the bp of pent-1-ene is 30°C.

11.4 Interesting Alkynes

Acetylene, HC≡CH, is a colorless gas with an ethereal odor that burns in oxygen to form CO_2 and H_2O. Because the combustion of acetylene releases more energy per mole of product formed than other hydrocarbons, it burns with a very hot flame, making it an excellent fuel for welding torches.

Ethynylestradiol, the molecule that opened Chapter 11, and **norethindrone** are two components of oral contraceptives that contain a carbon–carbon triple bond (Figure 11.2). Both molecules are synthetic analogues of the naturally occurring female hormones estradiol and progesterone, but are more potent so they can be administered in lower doses. Most oral contraceptives contain two of these synthetic hormones. They act by artificially elevating hormone levels in a woman, thereby preventing pregnancy.





Figure 11.2 How oral contraceptives work




Monthly cycles of hormones from the pituitary gland cause ovulation, the release of an egg from an ovary. To prevent pregnancy, the two synthetic hormones in many oral contraceptives have different effects on the female reproductive system.

A: The elevated level of **ethynylestradiol,** a synthetic estrogen, "fools" the pituitary gland into thinking a woman is pregnant, so ovulation does not occur.

B: The elevated level of **norethindrone,** a synthetic progesterone, stimulates the formation of a thick layer of mucus in the cervix, making it difficult for sperm to reach the uterus.

Two other synthetic hormones with alkynyl appendages are **RU 486** and **levonorgestrel.** RU 486 blocks the effects of progesterone, and because of this, prevents implantation of a fertilized egg. RU 486 is used to induce abortions within the first few weeks of pregnancy. Levonorgestrel interferes with ovulation, and so it prevents pregnancy if taken within a few days of unprotected sex.




RU 486
(Trade name Mifepristone)

levonorgestrel
(Trade name Plan B)

Histrionicotoxin is a diyne isolated in small quantities from the skin of *Dendrobates histrionicus,* a colorful South American frog (Figure 11.3). This toxin, secreted by the frog as a natural defense mechanism, was used as a poison on arrow tips by the Choco tribe of South America.


Figure 11.3
Histrionicotoxin




- Histrionicotoxin is a defensive toxin that protects *Dendrobates histrionicus* from potential predators. These small "poison dart" frogs inhabit the moist humid floor of tropical rainforests, and are commonly found in western Ecuador and Colombia. Histrionicotoxin acts by interfering with nerve transmission in mammals, resulting in prolonged muscle contraction.

11.5 Preparation of Alkynes

Alkynes are prepared by elimination reactions, as discussed in Section 8.10. **A strong base removes two equivalents of HX from a vicinal or geminal dihalide to yield an alkyne** by two successive E2 eliminations.




geminal dichloride



vicinal dibromide

Because vicinal dihalides are synthesized by adding halogens to alkenes, an alkene can be converted to an alkyne by the two-step process illustrated in Sample Problem 11.2.


Sample Problem 11.2 Convert alkene **A** into alkyne **B** by a stepwise method.



Solution


A two-step method is needed:

- **Addition of X_2** forms a vicinal dihalide.
- **Elimination** of two equivalents of HX forms two π bonds.



- **This two-step process introduces one degree of unsaturation:** an alkene with one π bond is converted to an alkyne with two π bonds.

Problem 11.5 Convert each compound to hex-1-yne, $HC{\equiv}CCH_2CH_2CH_2CH_3$.




11.6 Introduction to Alkyne Reactions

All reactions of alkynes occur because they contain **easily broken π bonds** or, in the case of terminal alkynes, an **acidic, *sp* hybridized C−H bond.**

11.6A Addition Reactions


Like alkenes, **alkynes undergo addition reactions because they contain weak π bonds.** Two sequential reactions take place: addition of one equivalent of reagent forms an alkene, which then adds a second equivalent of reagent to yield a product having **four new bonds.**



Alkynes are electron rich, as shown in the electrostatic potential map of acetylene in Figure 11.4. The two π bonds form a cylinder of electron density between the two *sp* hybridized carbon atoms, and this exposed electron density makes a triple bond nucleophilic. As a result, **alkynes react with electrophiles.** Four addition reactions are discussed in Chapter 11 and illustrated in Figure 11.5 with but-1-yne as the starting material.


The oxidation and reduction of alkynes, reactions that also involve addition, are discussed in Chapter 12.

Figure 11.4
Electrostatic potential map of acetylene



- The red electron-rich region is located between the two carbon atoms, forming a cylinder of electron density.


Figure 11.5
Four addition reactions of but-1-yne



- In each addition, both π bonds of the triple bond are broken, and four new bonds are formed.

11.6B Terminal Alkynes—Reaction as an Acid

Because ***sp* hybridized C−H bonds are more acidic than *sp*^{2} and *sp*^{3} hybridized C−H bonds,** terminal alkynes are readily deprotonated with strong base in a Brønsted–Lowry acid–base reaction. The resulting anion is called an **acetylide anion.**




Recall from Section 2.5D that the acidity of a C−H bond increases as the percent *s*-character of C increases. Thus, the following order of relative acidity results: $C_{sp^3}-H < C_{sp^2}-H < C_{sp}-H$.

What bases can be used for this reaction? Because an acid–base equilibrium favors the weaker acid and base, only **bases having conjugate acids with pK_a values *higher* than the terminal alkyne—that is, pK_a values > 25—are strong enough** to form a significant concentration of acetylide anion. As shown in Table 11.1, $^-NH_2$ and H^- are strong enough to deprotonate a terminal alkyne, but ^-OH and ^-OR are not.

Table 11.1 A Comparison of Bases for Alkyne Deprotonation

	Base	pK_a of the conjugate acid
These bases are **strong** enough to deprotonate an alkyne.	$^-NH_2$	38
	H^-	35
These bases are ***not*** strong enough to deprotonate an alkyne.	^-OH	15.7
	^-OR	15.5–18

Why is this reaction useful? The acetylide anions formed by deprotonating terminal alkynes are **strong nucleophiles** that can react with a variety of electrophiles, as shown in Section 11.11.




Problem 11.6 Which bases can deprotonate acetylene? The pK_a values of the conjugate acids are given in parentheses.

a. CH_3NH^- (pK_a = 40) b. CO_3^{2-} (pK_a = 10.2) c. $CH_2{=}CH^-$ (pK_a = 44) d. $(CH_3)_3CO^-$ (pK_a = 18)


11.7 Addition of Hydrogen Halides

Alkynes undergo **hydrohalogenation, the addition of hydrogen halides, HX** (X = Cl, Br, I). Two equivalents of HX are usually used: addition of one mole forms a **vinyl halide,** which then reacts with a second mole of HX to form a **geminal dihalide.**




Addition of HX to an alkyne is another example of **electrophilic addition,** because the electrophilic (H) end of the reagent is attracted to the electron-rich triple bond.

- **With two equivalents of HX, both H atoms bond to the *same* carbon.**
- **With a terminal alkyne, both H atoms bond to the *terminal* carbon; that is, the hydrohalogenation of alkynes follows Markovnikov's rule.**




- **With only one equivalent of HX, the reaction stops with formation of the vinyl halide.**



Because of the instability of a vinyl carbocation, other mechanisms for HX addition that avoid formation of a discrete carbocation have been proposed. It is likely that more than one mechanism occurs, depending in part on the identity of the alkyne substrate.


One proposed mechanism for the addition of two equivalents of HX to an alkyne involves **two steps for each addition of HX:** addition of H^+ (from HX) to form a carbocation, followed by nucleophilic attack of X^-. Mechanism 11.1 illustrates the addition of HBr to but-1-yne to yield 2,2-dibromobutane. Each two-step mechanism is similar to the two-step addition of HBr to *cis*-but-2-ene discussed in Section 10.9.

Mechanism 11.1 Electrophilic Addition of HX to an Alkyne




1. Addition of H^+ forms a vinyl carbocation and follows Markovinkov's rule. The H atom bonds to the terminal C to form the more substituted carbocation.
2. Nucleophilic attack of Br^- forms a vinyl bromide. One equivalent of HBr adds in two steps.
3. The addition of a second equivalent of HBr follows in the same two-step manner. Addition of H^+ to the vinyl bromide forms a carbocation.
4. Nucleophilic attack of Br^- forms a geminal dibromide, 2,2-dibromobutane.


The formation of both carbocations (in Steps [1] and [3]) deserves additional scrutiny. **The vinyl carbocation formed in Step [1] is *sp* hybridized and therefore less stable than a 2° sp^2 hybridized carbocation** (Section 7.18). This makes electrophilic addition of HX to an alkyne ***slower*** than electrophilic addition of HX to an alkene, even though alkynes are more polarizable and have more loosely held π electrons than alkenes.



In Step [3] two carbocations are possible but only one is formed. Markovnikov addition in Step [3] places the H on the terminal carbon (C1) to form the more substituted carbocation **A,** rather than the less substituted carbocation **B.** Because the more stable carbocation is formed faster—another example of the Hammond postulate—carbocation **A** must be more stable than carbocation **B.**






Why is carbocation **A,** having a positive charge on a carbon that also has a Br atom, more stable? Shouldn't the electronegative Br atom withdraw electron density from the positive charge, and thus destabilize it? It turns out that **A is stabilized by resonance** but **B** is not. Two resonance structures can be drawn for carbocation **A,** but only one Lewis structure can be drawn for carbocation **B.**






- Resonance stabilizes a molecule by delocalizing charge and electron density.
- Thus, halogens stabilize an adjacent positive charge by resonance.

Markovnikov's rule applies to the addition of HX to vinyl halides because **addition of H^+ forms a resonance-stabilized carbocation.** As a result, addition of each equivalent of HX to a triple bond forms the more stable carbocation, so that both H atoms bond to the less substituted C.

Problem 11.7 Draw the organic products formed when each alkyne is treated with two equivalents of HBr.


a.  b.  c. 

Problem 11.8 Draw additional resonance structures for each cation.

a.  b.  c. 


11.8 Addition of Halogen

Halogens, X_2 (X = Cl or Br), add to alkynes in much the same way they add to alkenes (Section 10.13). Addition of one mole of X_2 forms a **trans dihalide,** which can then react with a second mole of X_2 to yield a **tetrahalide.**



Each addition of X_2 involves a two-step process with a **bridged halonium ion** intermediate, reminiscent of the addition of X_2 to alkenes (Section 10.13). A trans dihalide is formed after addition of one equivalent of X_2 because the intermediate **halonium ion ring is opened upon backside attack of the nucleophile.** Mechanism 11.2 illustrates the addition of two equivalents of Cl_2 to $CH_3C{\equiv}CCH_3$ to form $CH_3CCl_2CCl_2CH_3$.

Mechanism 11.2 Addition of X_2 to an Alkyne—Halogenation




1. Four bonds are broken or formed to generate a bridged halonium ion that contains a three-membered ring.
2. Nucleophilic attack of Cl^- ring opens the bridged halonium ion from the back side and forms the trans dihalide.
3. Electrophilic addition of Cl^+ forms the bridged halonium ion.
4. Nucleophilic attack of Cl^- ring opens the bridged halonium ion and forms the tetrahalilde.

Problem 11.9 Draw the products formed when $CH_3CH_2C{\equiv}CCH_2CH_3$ is treated with each reagent: (a) Br_2 (2 equiv); (b) Cl_2 (1 equiv).

Problem 11.10 Explain the following result. Although alkenes are generally more reactive than alkynes towards electrophiles, the reaction of Cl_2 with but-2-yne can be stopped after one equivalent of Cl_2 has been added.

11.9 Addition of Water


Although the addition of H_2O to an alkyne resembles the acid-catalyzed addition of H_2O to an alkene in some ways, an important difference exists. In the presence of strong acid or Hg^{2+} catalyst, the **elements of H_2O add to the triple bond,** but the initial addition product, an **enol,** is unstable and rearranges to a product containing a **carbonyl group**—that is, a **C=O.** A carbonyl compound having two alkyl groups bonded to the C=O carbon is called a **ketone.**



Internal alkynes undergo hydration with concentrated acid, whereas terminal alkynes require the presence of an additional Hg^{2+} catalyst—usually $HgSO_4$—to yield methyl ketones by **Markovnikov addition of H_2O.**

Because an enol contains both a C=C and a hydroxy group, the name **enol** comes from alk**en**e + alcoh**ol.**


$HgSO_4$ is often used in the hydration of internal alkynes as well, because hydration can be carried out under milder reaction conditions.



Markovnikov addition of H_2O
H adds to the terminal C.

Let's first examine the conversion of a general enol **A** to the carbonyl compound **B. A** and **B** are called **tautomers: A is the *enol form* and B is the *keto form* of the tautomer.**

- ***Tautomers*** **are constitutional isomers that differ in the location of a double bond and a hydrogen atom. Two tautomers are in equilibrium with each other.**




Tautomers differ in the position of a double bond and a hydrogen atom. In Chapter 23 an in-depth discussion of keto–enol tautomers is presented.

- **An enol tautomer has an O–H group bonded to a C=C.**
- **A keto tautomer has a C=O and an additional C–H bond.**

Equilibrium favors the keto form largely because a C=O is much stronger than a C=C. Tautomerization, the process of converting one tautomer into another, is catalyzed by both acid and base. Under the strongly acidic conditions of hydration, tautomerization of the enol to the keto form occurs rapidly by a two-step process: **protonation,** followed by **deprotonation** as shown in Mechanism 11.3.

Mechanism 11.3 Tautomerization in Acid



new bond shown in red

two resonance structures for the carbocation


1 Protonation of the double bond forms a resonance-stabilized carbocation.

2 Loss of a proton, which can be drawn with either resonance structure, forms the carbonyl group. Because acid is re-formed in this step, tautomerizataion is acid-catalyzed.

Hydration of an internal alkyne with strong acid forms an enol by a mechanism similar to that of the acid-catalyzed hydration of an alkene (Section 10.12). Mechanism 11.4 illustrates the hydration of but-2-yne with H_2O and H_2SO_4. Once formed, the enol then tautomerizes to the more stable keto form by protonation followed by deprotonation.

Mechanism 11.4 Hydration of an Alkyne

Part [1] Addition of H_2O to form an enol



but-2-yne

vinyl carbocation
new bond shown in red


(*E* and *Z* isomers)

enol

1 Addition of H^+ forms a vinyl carbocation.

2 – 3 Nucleophilic attack followed by loss of a proton forms the enol.

Part [2] Tautomerization



enol

new bond shown in red


resonance-stabilized carbocation

ketone

4 Tautomerization of the enol to the keto form begins with protonation of the double bond to form a carbocation.


5 Loss of a proton, which can be drawn with either resonance structure, forms the ketone.

Sample Problem 11.3 Draw the enol intermediate and the ketone product formed in the following reaction.




Solution

First, form the enol by adding H_2O to the triple bond with the **H bonded to the less substituted terminal carbon**, according to Markovnikov's rule.




To convert the enol to the keto tautomer, add a proton to the C = C and remove a proton from the OH group. In tautomerization, the C – OH bond is converted to a C = O, and a new C – H bond is formed on the other enol carbon.




- **The overall result is the addition of H_2O to a triple bond to form a ketone.**

Problem 11.11 Draw the keto tautomer of each enol.



Problem 11.12 What two enols are formed when pent-2-yne is treated with H_2O, H_2SO_4, and $HgSO_4$? Draw the ketones formed from these enols after tautomerization.


Problem 11.13 (a) Draw two different enol tautomers of 2-methylcyclohexanone. (b) Draw two constitutional isomers that are not tautomers, but contain a C = C and an OH group.



2-methylcyclohexanone


11.10 Hydroboration–Oxidation

Hydroboration–oxidation is a two-step reaction sequence that converts an alkyne to a carbonyl compound.



- Addition of borane forms an organoborane.
- Oxidation with basic H_2O_2 forms an enol.
- Tautomerization of the enol forms a carbonyl compound.
- The overall result is addition of H_2O to a triple bond.

Hydroboration–oxidation of an *internal* alkyne forms a **ketone. Hydroboration of a *terminal* alkyne adds boron to the less substituted, terminal carbon.** After oxidation to the enol, tautomerization yields an **aldehyde,** a carbonyl compound having a hydrogen atom bonded to the carbonyl carbon. Hydroboration of a terminal alkyne is generally carried out with a dialkylborane (R_2BH), which has been prepared from BH_3 (Section 10.16).




Hydration (H_2O, H_2SO_4, and $HgSO_4$) and **hydroboration–oxidation** (BH_3 or R_2BH followed by H_2O_2, HO^-) both **add the elements of H_2O across a triple bond.** Sample Problem 11.4 shows that different constitutional isomers are formed from terminal alkynes in these two reactions.

Sample Problem 11.4 Draw the product formed when $CH_3CH_2C{\equiv}CH$ is treated with each of the following sets of reagents: (a) H_2O, H_2SO_4, $HgSO_4$; and (b) R_2BH, followed by H_2O_2, HO^-.



Solution

(a) With H_2O + H_2SO_4 + $HgSO_4$, electrophilic addition of H and OH places the **H atom on the less substituted carbon** of the alkyne to form a **ketone** after tautomerization. (b) In contrast, addition of R_2BH places the **R_2B group on the less substituted terminal carbon** of the alkyne. Oxidation and tautomerization yield an **aldehyde.** The ketone and aldehyde formed in these reactions are constitutional isomers.




- Addition of H_2O using H_2O, H_2SO_4, and $HgSO_4$ forms methyl ketones from terminal alkynes.
- Addition of H_2O using an organoborane, then H_2O_2, HO^- forms aldehydes from terminal alkynes.

Problem 11.14 Draw the products formed when the following alkynes are treated with each set of reagents: [1] H_2O, H_2SO_4, $HgSO_4$; or [2] R_2BH followed by H_2O_2, ^-OH.

a.  b. 


11.11 Reaction of Acetylide Anions

Terminal alkynes are readily converted to acetylide anions with strong bases such as $NaNH_2$ and NaH. These anions are strong nucleophiles, capable of reacting with electrophiles such as alkyl halides and epoxides.




11.11A Reaction of Acetylide Anions with Alkyl Halides

Acetylide anions react with unhindered alkyl halides to yield products of nucleophilic substitution.



Because acetylide anions are strong nucleophiles, the mechanism of nucleophilic substitution is **S_N2,** and thus the **reaction is fastest with CH_3X and 1° alkyl halides.** Terminal alkynes (Reaction [1]) or internal alkynes (Reaction [2]) can be prepared depending on the identity of the acetylide anion.




- Nucleophilic substitution with acetylide anions forms new carbon–carbon bonds.

Because organic compounds consist of a carbon framework, reactions that form carbon–carbon bonds are especially useful. In Reaction [2], for example, nucleophilic attack of a seven-carbon acetylide anion on a seven-carbon alkyl halide yields a 14-carbon alkyne as product.

Although nucleophilic substitution with acetylide anions is a very valuable carbon–carbon bond-forming reaction, it has the same limitations as any S_N2 reaction. **Steric hindrance around the leaving group causes 2° and 3° alkyl halides to undergo elimination by an E2 mechanism,**

as shown with 2-bromo-2-methylpropane. Thus, nucleophilic substitution with acetylide anions forms new carbon–carbon bonds in high yield only with unhindered CH_3X and 1° alkyl halides.




2-bromo-2-methylpropane
3° alkyl halide

E2 product


Steric hindrance prevents an S_N2 reaction.

Sample Problem 11.5 Draw the organic products formed in each reaction.




Solution

a. Because the alkyl halide is **1°** and the acetylide anion is a strong nucleophile, substitution occurs by an **S_N2** mechanism, resulting in a new C – C bond.




b. Because the alkyl halide is **2°,** the major product is formed from elimination by an **E2** mechanism.




major product

Problem 11.15 Draw the organic products formed in each reaction.




Problem 11.16 What acetylide anion and alkyl halide can be used to prepare each alkyne? Indicate all possibilities when more than one route will work.




Because acetylene has two *sp* hybridized C – H bonds, two sequential reactions can occur to form **two new carbon–carbon bonds,** as shown in Sample Problem 11.6.

Sample Problem 11.6 Identify the terminal alkyne **A** and the internal alkyne **B** in the following reaction sequence.




Solution

In each step, the base $^{-}NH_2$ removes a proton on an *sp* hybridized carbon, and the resulting acetylide anion reacts as a nucleophile with an alkyl halide to yield an S_N2 product. The first two-step reaction sequence forms the **terminal alkyne A** by nucleophilic attack of the acetylide anion on $CH_3CH_2CH_2Br$.




The second two-step reaction sequence forms the **internal alkyne B** by nucleophilic attack of the acetylide anion on CH_3CH_2Cl.






The soft coral *Capnella imbricata* is the source of the natural product capnellene.

Sample Problem 11.6 illustrates how a seven-carbon product can be prepared from three smaller molecules by forming two new carbon–carbon bonds.



Carbon–carbon bond formation with acetylide anions is a valuable reaction used in the synthesis of numerous natural products. Two examples include **capnellene,** isolated from the soft coral *Capnella imbricata,* and **niphatoxin B,** isolated from a red sea sponge, as shown in Figure 11.6.

Figure 11.6
Use of acetylide anion reactions in the synthesis of two marine natural products




- New carbon–carbon bonds formed from acetylide anions are shown in red.

Problem 11.17 Show how HC≡CH, CH_3CH_2Br, and $(CH_3)_2CHCH_2CH_2Br$ can be used to prepare $CH_3CH_2C{\equiv}CCH_2CH_2CH(CH_3)_2$. Show all reagents, and use curved arrows to show movement of electron pairs.


Problem 11.18 Explain why 2,2,5,5-tetramethylhex-3-yne can't be made using acetylide anions.

11.11B Reaction of Acetylide Anions with Epoxides

Acetylide anions are strong nucleophiles that open epoxide rings by an S_N2 mechanism. This reaction also results in the formation of a **new carbon–carbon bond.** Backside attack occurs at the less substituted end of the epoxide.




Opening of epoxide rings with strong nucleophiles was first discussed in Section 9.15A.



enantiomers
new C–C bonds shown in red

Problem 11.19 Draw the products of each reaction.



Problem 11.20 Draw the products formed when $CH_3CH_2C{\equiv}C^-Na^+$ reacts with each compound.

a. $CH_3CH_2CH_2Br$
b. $(CH_3)_2CHCH_2CH_2Cl$
c. $(CH_3CH_2)_3CCl$
d. $BrCH_2CH_2CH_2CH_2OH$
e. ethylene oxide followed by H_2O
f. propene oxide followed by H_2O

11.12 Synthesis

The reactions of acetylide anions give us an opportunity to examine organic synthesis more systematically. Performing a multistep synthesis can be difficult. Not only must you know the reactions for a particular functional group, but you must also put these reactions in a logical order, a process that takes much practice to master.

11.12A General Terminology and Conventions

To plan a synthesis of more than one step, we use the process of **retrosynthetic analysis**—that is, working backwards from the desired product to determine the starting materials from which it is made (Section 10.18). To write a synthesis working backwards from the product to the starting material, an **open arrow** (⇒) is used to indicate that the product is drawn on the left and the starting material on the right.

Carefully read the directions for each synthesis problem. Sometimes a starting material is specified, whereas at other times you must begin with a compound that meets a particular criterion; for example, you may be asked to synthesize a compound from alcohols having five or fewer carbon atoms. These limitations are meant to give you some direction in planning a multistep synthesis.

The product of a synthesis is often called the **target compound.** Using retrosynthetic analysis, we must determine what compound can be converted to the target compound by a single reaction. That is, **what is the immediate precursor of the target compound?** After an appropriate precursor is identified, this process is continued until we reach a specified starting material. Sometimes multiple retrosynthetic pathways are examined before a particular route is decided upon.



In designing a synthesis, reactions are often divided into two categories:

- **Those that form new carbon–carbon bonds.**
- **Those that convert one functional group into another—that is, functional group interconversions.**

Appendix D lists the carbon–carbon bond-forming reactions encountered in this text.

Carbon–carbon bond-forming reactions are central to organic synthesis because simpler and less valuable starting materials can be converted to more complex products. Keep in mind that whenever the product of a synthesis has more carbon–carbon bonds than the starting material, the synthesis must contain at least one of these reactions.

How To Develop a Retrosynthetic Analysis

Step [1] Compare the carbon skeletons of the starting material and product.

- If the product has more carbon–carbon σ bonds than the starting material, the synthesis must form one or more C – C bonds. If not, only functional group interconversion occurs.
- Match the carbons in the starting material with those in the product, to see where new C – C bonds must be added or where functional groups must be changed.

Step [2] Concentrate on the functional groups in the starting material and product and ask:

- What methods introduce the functional groups in the product?
- What kind of reactions does the starting material undergo?

Step [3] Work backwards from the product and forwards from the starting material.

- Ask: What is the immediate precursor of the product?
- Compare each precursor to the starting material to determine if there is a one-step reaction that converts one to the other. Continue this process until the starting material is reached.
- Always generate simpler precursors when working backwards.
- Use fewer steps when multiple routes are possible.
- Keep in mind that you may need to evaluate several different precursors for a given compound.

Step [4] Check the synthesis by writing it in the synthetic direction.

- To check a retrosynthetic analysis, write out the steps beginning with the starting material, indicating all necessary reagents.


11.12B Examples of Multistep Synthesis

Retrosynthetic analysis with acetylide anions is illustrated in Sample Problems 11.7 and 11.8.

Sample Problem 11.7 Devise a synthesis of $HC{\equiv}CCH_2CH_2CH_3$ from $HC{\equiv}CH$ and any other organic or inorganic reagents.

Retrosynthetic Analysis

The two C's in the starting material match up with the two *sp* hybridized C's in the product, so a three-carbon unit must be added.




Thinking backwards . . .

[1] Form a new C – C bond using an acetylide anion and a 1° alkyl halide.
[2] Prepare the acetylide anion from acetylene by treatment with base.

Synthesis

Deprotonation of $HC{\equiv}CH$ with NaH forms the acetylide anion, which undergoes S_N2 reaction with an alkyl halide to form the target compound, a five-carbon alkyne.

A two-step process:




Sample Problem 11.8 Devise a synthesis of the following compound from starting materials having two carbons or fewer.



Retrosynthetic Analysis

A carbon–carbon bond-forming reaction must be used to convert the two-carbon starting materials to the four-carbon product.




Thinking backwards . . .

[1] Form the carbonyl group by hydration of a triple bond.
[2] Form a new C – C bond using an acetylide anion and a 1° alkyl halide.
[3] Prepare the acetylide anion from acetylene by treatment with base.

Synthesis

Three steps are needed to complete the synthesis. Treatment of HC≡CH with NaH forms the acetylide anion, which undergoes an S_N2 reaction with an alkyl halide to form a four-carbon terminal alkyne. Hydration of the alkyne with H_2O, H_2SO_4, and $HgSO_4$ yields the target compound.



These examples illustrate the synthesis of organic compounds by multistep routes. In Chapter 12 we will learn other useful reactions that expand our capability to do synthesis.

Problem 11.21 Use retrosynthetic analysis to show how hex-3-yne can be prepared from acetylene and any other organic and inorganic compounds. Then draw the synthesis in the synthetic direction, showing all needed reagents.

Problem 11.22 Devise a synthesis of $CH_3CH_2CH_2CHO$ from two-carbon starting materials.

KEY CONCEPTS


Alkynes

General Facts About Alkynes

- Alkynes contain a carbon–carbon triple bond consisting of a strong σ bond and two weak π bonds. Each carbon is *sp* hybridized and linear (11.1).
- Alkynes are named using the suffix ***-yne*** (11.2).
- Alkynes have weak intermolecular forces, giving them low mp's and low bp's, and making them water insoluble (11.3).
- Because its weaker π bonds make an alkyne electron rich, alkynes undergo addition reactions with electrophiles (11.6).


Addition Reactions of Alkynes

[1] Hydrohalogenation—Addition of HX (X = Cl, Br, I) (11.7)




- Markovnikov's rule is followed. H bonds to the less substituted C to form the more stable carbocation.

[2] Halogenation—Addition of X_2 (X = Cl or Br) (11.8)




- Bridged halonium ions are formed as intermediates.
- Anti addition of X_2 occurs.

[3] Hydration—Addition of H_2O (11.9)



- Markovnikov's rule is followed. H bonds to the less substituted C to form the more stable carbocation.
- An unstable enol is first formed, which rearranges to a carbonyl group.


[4] Hydroboration–oxidation—Addition of H_2O (11.10)



- The unstable enol, first formed after oxidation, rearranges to a carbonyl group.


Reactions Involving Acetylide Anions

[1] Formation of acetylide anions from terminal alkynes (11.6B)




- Typical bases used for the reaction are $NaNH_2$ and NaH.

[2] Reaction of acetylide anions with alkyl halides (11.11A)



- The reaction follows an S_N2 mechanism.
- The reaction works best with CH_3X and RCH_2X.

[3] Reaction of acetylide anions with epoxides (11.11B)





- The reaction follows an S_N2 mechanism.
- Opening of the ring occurs from the back side at the less substituted end of the epoxide.

PROBLEMS


Problems Using Three-Dimensional Models


11.23 Give the IUPAC name for each compound.

a. 

b. 


11.24 Draw the enol tautomer of (a) and the keto tautomer of (b).

a. 


b. 

Structure and Nomenclature

11.25 Answer the following questions about erlotinib and phomallenic acid C. Erlotinib, sold under the trade name Tarceva, was introduced in 2004 for the treatment of lung cancer. Phomallenic acid C is an inhibitor of bacterial fatty acid synthesis.









erlotinib



phomallenic acid C

a. Which C – H bond in erlotinib is most acidic?
b. What orbitals are used to form the shortest C – C single bond in erlotinib?
c. Which H atom in phomallenic acid C is most acidic?
d. How many *sp* hybridized carbons are contained in phomallenic acid C?
e. Rank the labeled bonds in phomallenic acid C in order of increasing bond strength.

11.26 Give the IUPAC name for each alkyne.









a. 
c. 
e. 
b. 
d. 
f. 

11.27 Give the structure corresponding to each name.





a. 5,6-dimethylhept-2-yne
b. 5-*tert*-butyl-6,6-dimethylnon-3-yne
c. (*S*)-4-chloropent-2-yne
d. *cis*-1-ethynyl-2-methylcyclopentane
e. 3,4-dimethylocta-1,5-diyne
f. (*Z*)-6-methyloct-6-en-1-yne

Tautomers


11.28 Which of the following pairs of compounds represent keto–enol tautomers?





a.  and 
c.  and 
b.  and 
d.  and 

11.29 Draw the enol form of each keto tautomer in parts (a) and (b), and the keto form of each enol tautomer in parts (c) and (d).

a. 
b. 
(two different enols)
c. 
d. 


11.30 How is each compound related to **A?** Choose from tautomers, constitutional isomers but not tautomers, or neither.


A

a. 
b. 
c. 
d. 

11.31 Ignoring stereoisomers, draw the two possible enols for butan-2-one ($CH_3COCH_2CH_3$), and predict which one is more stable.

11.32 Enamines and imines are tautomers that contain N atoms. Draw a stepwise mechanism for the acid-catalyzed conversion of enamine **X** to imine **Y.**




X enamine **Y** imine


Reactions


11.33 Draw the products formed when hex-1-yne is treated with each reagent.


a. HCl (2 equiv)
b. HBr (2 equiv)
c. Cl_2 (2 equiv)
d. $H_2O + H_2SO_4 + HgSO_4$
e. [1] R_2BH; [2] H_2O_2, HO^-
f. NaH
g. [1] $^-NH_2$; [2] CH_3CH_2Br
h. [1] $^-NH_2$; [2]  ; [3] H_2O

11.34 What reagents are needed to convert $(CH_3CH_2)_3CC{\equiv}CH$ to each compound?

a. 


b. 


c. 


d. 


11.35 Explain the apparent paradox. Although the addition of one equivalent of HX to an alkyne is more exothermic than the addition of HX to an alkene, an alkene reacts faster with HX.

11.36 What alkynes give each of the following ketones as the only product after hydration with H_2O, H_2SO_4, and $HgSO_4$?


a. 


b. 

c. 


d. 


11.37 What alkyne gives each compound as the only product after hydroboration–oxidation?


a. 


b. 


11.38 Draw the organic products formed in each reaction.


a. 


b. 


c. 


d. 

e. $HC{\equiv}C^-$ + D_2O ⟶
f. 


g. 

h. 


i. 

j. 


11.39 Draw the structure of compounds **A–E** in the following reaction scheme.




11.40 When alkyne **A** is treated with $NaNH_2$ followed by CH_3I, a product having molecular formula $C_6H_{10}O$ is formed, but it is *not* compound **B.** What is the structure of the product and why is it formed?




11.41 Draw the products formed in each reaction and indicate stereochemistry.



11.42 What reactions are needed to convert alcohol **A** into either alkyne **B** or alkyne **C?**




11.43 Identify the lettered compounds in the following reaction schemes. Each reaction sequence was used in the synthesis of a natural product.




Mechanisms

11.44 Treatment of 2,2-dibromobutane with two equivalents of strong base affords but-1-yne and but-2-yne, as well as a small amount of buta-1,2-diene. Draw a mechanism showing how each compound is formed. Which alkyne should be the major product?




11.45 Explain the following statement. Although $HC{\equiv}C^-$ is more stable than $CH_2{=}CH^-$, $HC{\equiv}C^+$ is less stable than $CH_2{=}CH^+$.

11.46 Tautomerization in base resembles tautomerization in acid, but deprotonation precedes protonation in the two-step mechanism. (a) Draw a stepwise mechanism for the following tautomerization. (b) Then draw a stepwise mechanism for the reverse reaction, the conversion of the keto form to the enol.



enol form keto form

11.47 Draw a stepwise mechanism for each reaction.




11.48 From what you have learned about enols and the hydration of alkynes, predict what product is formed by the acid-catalyzed hydration of $CH_3CH_2CH_2C{\equiv}COCH_3$. Draw a stepwise mechanism that illustrates how it is formed.


Synthesis

11.49 What steps are needed to prepare phenylacetylene, $C_6H_5C{\equiv}CH$, from each compound: (a) $C_6H_5CH_2CHBr_2$; (b) $C_6H_5CHBrCH_3$; (c) $C_6H_5CH_2CH_2OH$?


11.50 What acetylide anion and alkyl halide are needed to synthesize each alkyne?




11.51 Synthesize each compound from acetylene. You may use any other organic or inorganic reagents.



11.52 Devise a synthesis of each compound using $CH_3CH_2CH{=}CH_2$ as the starting material. You may use any other organic compounds or inorganic reagents.





11.53 Devise a synthesis of the following compound from cyclohexene and acetylene. You may use any other inorganic reagents.



+ enantiomer

11.54 Devise a synthesis of each compound. You may use HC≡CH, ethylene oxide, and alkyl halides as organic starting materials and any inorganic reagents.

a. 

b. 

11.55 Devise a synthesis of the ketone hexan-3-one, $CH_3CH_2COCH_2CH_2CH_3$, from CH_3CH_2Br as the only organic starting material; that is, all the carbon atoms in hexan-3-one must come from CH_3CH_2Br. You may use any other needed reagents.


11.56 Devise a synthesis of each compound using $CH_3CH_2CH_2OH$ as the only organic starting material: (a) $CH_3C{\equiv}CCH_2CH_2CH_3$; (b) $CH_3C{\equiv}CCH_2CH(OH)CH_3$. You may use any other needed inorganic reagents.

11.57 Devise a synthesis of $CH_3CH_2C{\equiv}CCH_2CH_2OH$ from CH_3CH_2OH as the only organic starting material. You may use any other needed reagents.

Challenge Problems


11.58 Explain why the C=C of an enol is more nucleophilic than the C=C of an alkene, despite the fact that the electronegative oxygen atom of the enol inductively withdraws electron density from the carbon–carbon double bond.

11.59 *N*-Chlorosuccinimide (NCS) serves as a source of Cl^+ in electrophilic addition reactions to alkenes and alkynes. Keeping this in mind, draw a stepwise mechanism for the following addition to but-2-yne.




N-chlorosuccinimide


11.60 Draw a stepwise mechanism for the following reaction.



11.61 Draw a stepwise mechanism for the following reaction.




11.62 Why is compound **X** formed in the following reaction, instead of its constitutional isomer **Y?**




X **Y**

11.63 Write a stepwise mechanism for each of the following reactions. Explain why a more stable alkyne (but-2-yne) is isomerized to a less stable alkyne (but-1-yne), but under similar conditions, 2,5-dimethylhex-3-yne forms 2,5-dimethylhexa-2,3-diene.



but-2-yne


but-1-yne




2,5-dimethylhex-3-yne

2,5-dimethylhexa-2,3-diene

11.64 Draw a stepwise mechanism for the following intramolecular reaction.



11.65 Explain why an optically active solution of (*R*)-α-methylbutyrophenone loses its optical activity when dilute acid is added to the solution.



(*R*)-α-methylbutyrophenone

Oxidation and Reduction

12



12.1 Introduction
12.2 Reducing agents
12.3 Reduction of alkenes
12.4 Application: Hydrogenation of oils
12.5 Reduction of alkynes
12.6 The reduction of polar C–X σ bonds
12.7 Oxidizing agents
12.8 Epoxidation
12.9 Dihydroxylation
12.10 Oxidative cleavage of alkenes
12.11 Oxidative cleavage of alkynes
12.12 Oxidation of alcohols
12.13 Green chemistry
12.14 Biological oxidation
12.15 Sharpless epoxidation

Throughout history, humans have ingested alcoholic beverages for their pleasant taste and the feeling of euphoria they impart. Wine, beer, and similar products contain **ethanol** (CH_3CH_2OH), a 1° alcohol that is quickly absorbed in the stomach and small intestines and rapidly transported in the bloodstream to other organs. Like other 1° alcohols, ethanol is easily oxidized, and as a result, ethanol is metabolized in the body by a series of enzyme-catalyzed oxidation reactions that take place in the liver. In Chapter 12, we learn about oxidation and reduction reactions of organic molecules like ethanol.

In Chapter 12, we discuss the oxidation and reduction of **alkenes** and **alkynes,** as well as compounds with **polar C–X σ bonds**—alcohols, alkyl halides, and epoxides. Although there will be many different reagents and mechanisms, discussing these reactions as a group allows us to more easily compare and contrast them.

The word *mechanism* will often be used loosely here. In contrast to the S_N1 reaction of alkyl halides or the electrophilic addition reactions of alkenes, the details of some of the mechanisms presented in Chapter 12 are known with less certainty. For example, although the identity of a particular intermediate might be confirmed by experiment, other details of the mechanism are suggested by the structure or stereochemistry of the final product.

Oxidation and reduction reactions are very versatile, and knowing them allows us to design many more complex organic syntheses.

12.1 Introduction


Recall from Section 4.14 that the way to determine whether an organic compound has been oxidized or reduced is to compare the **relative number of C–H and C–Z bonds** (Z = an element *more electronegative* than carbon) in the starting material and product.

- ***Oxidation*** **results in an *increase* in the number of C–Z bonds (usually C–O bonds) *or* a *decrease* in the number of C–H bonds.**
- ***Reduction*** **results in a *decrease* in the number of C–Z bonds (usually C–O bonds) *or* an *increase* in the number of C–H bonds.**

Two components are always present in an oxidation or reduction reaction—**one component is oxidized and one is reduced.** When an organic compound is *oxidized* by a reagent, the reagent itself must be *reduced.* Similarly, when an organic compound is *reduced* by a reagent, the reagent becomes *oxidized.*


Thus, an organic compound such as CH_4 can be oxidized by replacing C–H bonds with C–O bonds, as shown in Figure 12.1. Reduction is the opposite of oxidation, so Figure 12.1 also shows how a compound can be reduced by replacing C–O bonds with C–H bonds. The symbols **[O]** and **[H]** indicate oxidation and reduction, respectively.

Figure 12.1
A general scheme for the oxidation and reduction of a carbon compound




Sometimes two carbon atoms are involved in a single oxidation or reduction reaction, and the net change in the number of C–H or C–Z bonds at *both* atoms must be taken into account. The conversion of an **alkyne to an alkene** and an **alkene to an alkane** are examples of reduction, because each process adds two new C–H bonds to the starting material, as shown in Figure 12.2.

Figure 12.2
Oxidation and reduction of hydrocarbons




Problem 12.1 Classify each reaction as oxidation, reduction, or neither.



12.2 Reducing Agents

Reducing agents provide the equivalent of two hydrogen atoms, but **there are three types of reductions,** differing in how H_2 is added. The simplest reducing agent is molecular **H_2.** Reductions of this sort are carried out in the presence of a metal catalyst that acts as a surface on which the reaction occurs.

The second way to deliver H_2 in a reduction is to add two protons and two electrons to a substrate—that is, **$H_2 = 2\ H^+ + 2\ e^-$.** Reducing agents of this sort use alkali metals as a source of electrons and liquid ammonia (NH_3) as a source of protons. Reductions with **Na in NH_3** are called **dissolving metal reductions.**




The third way to deliver the equivalent of two hydrogen atoms is to add **hydride (H^-)** and a **proton (H^+).** The most common hydride reducing agents contain a hydrogen atom bonded to boron or aluminum. Simple examples include **sodium borohydride ($NaBH_4$)** and **lithium aluminum hydride ($LiAlH_4$).** These reagents deliver H^- to a substrate, and then a proton is added from H_2O or an alcohol.



sodium borohydride
$NaBH_4$

lithium aluminum hydride
$LiAlH_4$

- **Metal hydride reagents act as a source of H^- because they contain polar metal–hydrogen bonds that place a partial negative charge on hydrogen.**




a polar metal–hydrogen bond

M = B or Al


12.3 Reduction of Alkenes

Reduction of an alkene forms an alkane by addition of H_2. Two bonds are broken—the **weak π bond** of the alkene and the H_2 σ bond—and two new C−H σ bonds are formed.




Hydrogenation catalysts are insoluble in common solvents, thus creating a **heterogeneous** reaction mixture. This insolubility has a practical advantage. These catalysts contain expensive metals, but they can be filtered away from the other reactants after the reaction is complete, and then reused.


The addition of H_2 occurs only in the presence of a **metal catalyst,** and thus, the reaction is called **catalytic hydrogenation.** The catalyst consists of a metal—usually Pd, Pt, or Ni—adsorbed onto a finely divided inert solid, such as charcoal. For example, the catalyst 10% Pd on carbon is composed of 10% Pd and 90% carbon, by weight. H_2 adds in a **syn** fashion, as shown in Equation [2].



Problem 12.2 What alkane is formed when each alkene is treated with H_2 and a Pd catalyst?



Problem 12.3 Draw all alkenes that react with one equivalent of H_2 in the presence of a palladium catalyst to form each alkane. Consider constitutional isomers only.



12.3A Hydrogenation and Alkene Stability

Hydrogenation reactions are **exothermic** because the bonds in the product are stronger than the bonds in the starting materials, making them similar to other alkene addition reactions. The $\Delta H°$ for hydrogenation, called the **heat of hydrogenation,** can be used as a measure of the relative stability of two different alkenes that are hydrogenated to the same alkane.

Recall from Chapter 8 that **trans alkenes are generally more stable than cis alkenes.**

For example, both *cis*- and *trans*-but-2-ene are hydrogenated to butane, and the heat of hydrogenation for the trans isomer is less than that for the cis isomer. **Because less energy is released in converting the trans alkene to butane, it must be lower in energy (more stable) to begin with.** The relative energies of the butene isomers are illustrated in Figure 12.3.







Figure 12.3
Relative energies of *cis*- and *trans*-but-2-ene



- **When hydrogenation of two alkenes gives the same alkane, the more stable alkene has the *smaller* heat of hydrogenation.**

Problem 12.4 Which alkene in each pair has the larger heat of hydrogenation?

a.  or 


b.  or 

Problem 12.5 Explain why heats of hydrogenation cannot be used to determine the relative stability of 2-methylpent-2-ene and 3-methylpent-1-ene.

12.3B The Mechanism of Catalytic Hydrogenation


In the generally accepted mechanism for catalytic hydrogenation, the surface of the metal catalyst binds both H_2 and the alkene, and H_2 is transferred to the π bond in a rapid but stepwise process (Mechanism 12.1).

Mechanism 12.1 Addition of H_2 to an Alkene—Hydrogenation



1 **H_2 adsorbs to the catalyst surface** with partial or complete cleavage of the H – H bond.


2 The π bond of the alkene complexes with the metal.




3 – 4 **Two H atoms are transferred sequentially** to the π bond in Steps [3] and [4], forming the alkane. Because the product alkane no longer has a π bond with which to complex to the metal, it is released from the catalyst surface.


The mechanism explains two facts about hydrogenation:


- **Rapid, sequential addition of H_2 occurs from the side of the alkene complexed to the metal surface, resulting in syn addition.**
- **Less crowded double bonds complex more readily to the catalyst surface, resulting in faster reaction.**



Problem 12.6 Given that syn addition of H_2 occurs from both sides of a trigonal planar double bond, draw all stereoisomers formed when each alkene is treated with H_2.

a. 

b. 

c. 

12.3C Hydrogenation Data and Degrees of Unsaturation

Recall from Section 10.2 that the **number of degrees of unsaturation gives the *total* number of rings and π bonds in a molecule.** Because H_2 adds to π bonds but does *not* add to the C−C σ bonds of rings, hydrogenation allows us to determine how many degrees of unsaturation are due to π bonds and how many are due to rings. This is done by comparing the number of degrees of unsaturation before and after a molecule is treated with H_2, as illustrated in Sample Problem 12.1.

Sample Problem 12.1 How many rings and π bonds are contained in a compound of molecular formula C_8H_{12} that is hydrogenated to a compound of molecular formula C_8H_{14}?

Solution

[1] Determine the number of degrees of unsaturation in the compounds before and after hydrogenation.

Before H_2 addition—C_8H_{12}

- The maximum number of H's possible for *n* C's is $2n + 2$; in this example, $2n + 2 = 2(8) + 2 = 18$.
- 18 H's (maximum) – 12 H's (actual) = 6 H's fewer than the maximum number.

$$\frac{\text{6 H's fewer than the maximum}}{\text{2 H's removed for each degree of unsaturation}} = \textbf{three degrees of unsaturation}$$

After H_2 addition—C_8H_{14}

- The maximum number of H's possible for *n* C's is $2n + 2$; in this example, $2n + 2 = 2(8) + 2 = 18$.
- 18 H's (maximum) – 14 H's (actual) = 4 H's fewer than the maximum number.

$$\frac{\text{4 H's fewer than the maximum}}{\text{2 H's removed for each degree of unsaturation}} = \textbf{two degrees of unsaturation}$$

[2] Assign the number of degrees of unsaturation to rings or π bonds as follows:

- The number of degrees of unsaturation that remain in the product after H_2 addition = the **number of rings** in the starting material.
- The number of degrees of unsaturation that react with H_2 = the **number of π bonds.**

In this example, **two** degrees of unsaturation remain after hydrogenation, so the starting material has **two** rings. Thus:


Before H_2 addition:		After H_2 addition:		
three degrees of unsaturation	−	**two** degrees of unsaturation	=	**one** degree of unsaturation that reacted with H_2
three rings or π bonds in C_8H_{12}	=	**two rings**	+	**one π bond** **ANSWER**

Problem 12.7 Complete the missing information for compounds **A, B,** and **C,** each subjected to hydrogenation. The number of rings and π bonds refers to the reactant (**A, B,** or **C**) prior to hydrogenation.

Compound	Molecular formula before hydrogenation	Molecular formula after hydrogenation	Number of rings	Number of π bonds
A	$C_{10}H_{12}$	$C_{10}H_{16}$	?	?
B	?	C_4H_{10}	0	1
C	C_6H_8	?	1	?

12.3D Hydrogenation of Other Double Bonds

Compounds that contain a carbonyl group also react with H_2 and a metal catalyst. For example, **aldehydes and ketones are reduced to 1° and 2° alcohols,** respectively. We return to this reaction in Chapter 20.




aldehyde 1° alcohol ketone 2° alcohol

12.4 Application: Hydrogenation of Oils

Many processed foods, such as peanut butter, margarine, and some brands of crackers, contain *partially hydrogenated* vegetable oils. These oils are produced by hydrogenating the long hydrocarbon chains of triacylglycerols.

In Section 10.6 we learned that **fats and oils are triacylglycerols that differ in the number of degrees of unsaturation** in their long alkyl side chains.

- **Fats—usually animal in origin—are solids with triacylglycerols having few degrees of unsaturation.**
- **Oils—usually vegetable in origin—are liquids with triacylglycerols having a larger number of degrees of unsaturation.**



triacylglycerol

The number of double bonds in the R groups of the triacylglycerol determines whether it is a fat or an oil.



Peanut butter is a common consumer product that contains partially hydrogenated vegetable oil.

When an unsaturated vegetable oil is treated with hydrogen, some (or all) of the π bonds add H_2, decreasing the number of degrees of unsaturation (Figure 12.4). This increases the melting point of the oil. For example, margarine is prepared by partially hydrogenating vegetable oil to give a product having a semi-solid consistency that more closely resembles butter. This process is sometimes called ***hardening.***


If unsaturated oils are healthier than saturated fats, why does the food industry hydrogenate oils? There are two reasons—aesthetics and shelf life. Consumers prefer the semi-solid consistency of margarine to a liquid oil. Imagine pouring vegetable oil on a piece of toast or pancakes.

Furthermore, unsaturated oils are more susceptible than saturated fats to oxidation at the **allylic carbon atoms**—the carbons adjacent to the double bond carbons—a process discussed in Chapter 15. Oxidation makes the oil rancid and inedible. Hydrogenating the double bonds reduces the number of allylic carbons (also illustrated in Figure 12.4), thus reducing the likelihood of oxidation and increasing the shelf life of the food product. This process reflects a delicate balance between providing consumers with healthier food products, while maximizing shelf life to prevent spoilage.

One other fact is worthy of note. Because the steps in hydrogenation are reversible and H atoms are added in a sequential rather than concerted fashion, a cis double bond can be isomerized to a trans double bond. After addition of one H atom (Step [3] in Mechanism 12.1), an intermediate can lose a hydrogen atom to re-form a double bond with either the cis or trans configuration.


As a result, some of the cis double bonds in vegetable oils are converted to trans double bonds during hydrogenation, forming so-called "trans fats." The shape of the resulting fatty acid chain is very different, closely resembling the shape of a *saturated* fatty acid chain. Consequently, trans

Figure 12.4 Partial hydrogenation of the double bonds in a vegetable oil




- **Decreasing** the number of degrees of unsaturation **increases** the melting point. Only one long chain of the triacylglycerol is drawn.
- When an oil is *partially* hydrogenated, some double bonds react with H_2, whereas some double bonds remain in the product.
- Partial hydrogenation **decreases** the number of allylic sites (shown in blue), making a triacylglycerol **less** susceptible to oxidation, thereby increasing its shelf life.

fats are thought to have the same negative effects on blood cholesterol levels as saturated fats; that is, trans fats stimulate cholesterol synthesis in the liver, thus increasing blood cholesterol levels, a factor linked to increased risk of heart disease.



Problem 12.8 Draw the products formed when triacylglycerol **A** is treated with each reagent, forming compounds **B** and **C.** Rank **A, B,** and **C** in order of increasing melting point.




a. H_2 (excess), Pd-C (Compound **B**)
b. H_2 (1 equiv), Pd-C (Compound **C**)


12.5 Reduction of Alkynes

Reduction of an alkyne adds H_2 to one or both of the π bonds. There are three different ways by which the elements of H_2 can be added to a triple bond.


- **Adding two equivalents of H_2 forms an alkane.**



- **Adding one equivalent of H_2 in a syn fashion forms a cis alkene.**




- **Adding one equivalent of H_2 in an anti fashion forms a trans alkene.**



12.5A Reduction of an Alkyne to an Alkane


When an alkyne is treated with two or more equivalents of H_2 and a Pd catalyst, reduction of *both* π bonds occurs. **Syn addition** of one equivalent of H_2 forms a cis alkene, which adds a second equivalent of H_2 to form an **alkane. Four new C−H bonds are formed.** By using a Pd-C catalyst, it is not possible to stop the reaction after addition of only one equivalent of H_2.



Problem 12.9 Which alkyne has the smaller heat of hydrogenation, $HC{\equiv}CCH_2CH_2CH_3$ or $CH_3C{\equiv}CCH_2CH_3$? Explain your choice.


12.5B Reduction of an Alkyne to a Cis Alkene

Palladium metal is too active a catalyst to allow the hydrogenation of an alkyne to stop after one equivalent of H_2. To prepare a cis alkene from an alkyne and H_2, a less active Pd catalyst is used—Pd adsorbed onto $CaCO_3$ with added lead(II) acetate and quinoline. This catalyst is called the **Lindlar catalyst** after the chemist who first prepared it. Compared to Pd metal, the **Lindlar catalyst is deactivated or "poisoned."**



Reduction of an alkyne to a cis alkene is a **stereoselective reaction,** because only one stereoisomer is formed.


With the Lindlar catalyst, one equivalent of H_2 adds to an alkyne, and the cis alkene product is unreactive to further reduction.






Jasmine flowers are the source of *cis*-jasmone, a perfume component.

Problem 12.10 What is the structure of *cis*-jasmone, a natural product isolated from jasmine flowers, formed by treatment of alkyne **A** with H_2 in the presence of the Lindlar catalyst?




Problem 12.11 (a) Draw the structure of a compound of molecular formula C_6H_{10} that reacts with H_2 in the presence of Pd-C but does not react with H_2 in the presence of Lindlar catalyst. (b) Draw the structure of a compound of molecular formula C_6H_{10} that reacts with H_2 when either catalyst is present.

12.5C Reduction of an Alkyne to a Trans Alkene




NH_3 has a boiling point of –33 °C, making it a gas at room temperature. To carry out a Na, NH_3 reduction, NH_3 gas is condensed into a flask kept at –78 °C by a cooling bath of solid CO_2 in acetone. When Na is added to the liquid NH_3, a brilliant blue solution is formed.

Although catalytic hydrogenation is a convenient method for preparing cis alkenes from alkynes, it cannot be used to prepare trans alkenes. With a dissolving metal reduction (such as Na in NH_3), however, the elements of H_2 are added in an **anti** fashion to the triple bond, thus forming a **trans alkene.** For example, but-2-yne reacts with Na in NH_3 to form *trans*-but-2-ene.




The **mechanism** for the dissolving metal reduction using Na in NH_3 features sequential addition of electrons and protons to the triple bond. Half-headed arrows denoting the movement of a single electron must be used in two steps when Na donates *one* electron. The mechanism can be divided conceptually into two parts, each of which consists of two steps: **addition of an electron followed by protonation of the resulting negative charge,** as shown in Mechanism 12.2.

Mechanism 12.2 Dissolving Metal Reduction of an Alkyne to a Trans Alkene




1. Addition of an electron to the triple bond forms a **radical anion,** a species that contains *both* a negative charge *and* an unpaired electron.
2. Protonation of the anion with the solvent NH_3 yields a radical. The net result of the first two steps is the addition of a H atom.
3. Addition of a second electron forms a carbanion.
4. Protonation of the carbanion forms the trans alkene. Steps [3] and [4] add the second H atom to the triple bond.

Although the vinyl carbanion formed in Step [3] could have two different arrangements of its R groups, only the trans alkene is formed from the more stable vinyl carbanion; this carbanion has the larger R groups farther away from each other to avoid steric interactions. Protonation of this anion leads to the more stable trans product.



The larger R groups are farther away from each other.

This more stable vinyl carbanion forms the trans alkene.




Steric interactions between closer R groups destabilize this carbanion.

Dissolving metal reduction of a triple bond with Na in NH_3 is a **stereoselective reaction** because it forms a trans product exclusively.

- **Dissolving metal reductions always form the more stable trans product preferentially.**

The three methods to reduce a triple bond are summarized in Figure 12.5 using hex-3-yne as starting material.

Figure 12.5
Summary: Three methods to reduce a triple bond




Problem 12.12 What product is formed when $CH_3OCH_2CH_2C{\equiv}CCH_2CH(CH_3)_2$ is treated with each reagent: (a) H_2 (excess), Pd-C; (b) H_2 (1 equiv), Lindlar catalyst; (c) H_2 (excess), Lindlar catalyst; (d) Na, NH_3?

Problem 12.13 A chiral alkyne **A** with molecular formula C_6H_{10} is reduced with H_2 and Lindlar catalyst to **B** having the *R* configuration at its stereogenic center. What are the structures of **A** and **B?**


12.6 The Reduction of Polar C–X σ Bonds

Compounds containing polar C–X σ bonds that react with strong nucleophiles are reduced with metal hydride reagents, most commonly lithium aluminum hydride. Two functional groups possessing both of these characteristics are **alkyl halides** and **epoxides.** Alkyl halides are reduced to alkanes with loss of X^- as the leaving group. Epoxide rings are opened to form alcohols.



Reduction of these C−X σ bonds is another example of nucleophilic substitution, in which $LiAlH_4$ serves as a source of a hydride nucleophile (H^-). Because H^- is a strong nucleophile, the reaction follows an **S_N2 mechanism,** illustrated for the one-step reduction of an alkyl halide in Mechanism 12.3.

Mechanism 12.3 Reduction of RX with $LiAlH_4$




- The nucleophile H^- replaces the leaving group X^- in a single step.

Because the reaction follows an S_N2 mechanism:


- **Unhindered CH_3X and 1° alkyl halides are more easily reduced than more substituted 2° and 3° halides.**
- **In unsymmetrical epoxides, nucleophilic attack of H^- (from $LiAlH_4$) occurs at the less substituted carbon atom.**

Examples are shown in Figure 12.6.

Figure 12.6
Examples of reduction of C−X σ bonds with $LiAlH_4$



Problem 12.14 Draw the products of each reaction.




12.7 Oxidizing Agents

Oxidizing agents fall into two main categories:


- **Reagents that contain an oxygen–oxygen bond**
- **Reagents that contain metal–oxygen bonds**

Oxidizing agents containing an O−O bond include **O_2, O_3** (ozone), **H_2O_2** (hydrogen peroxide), **$(CH_3)_3COOH$** (*tert*-butyl hydroperoxide), and peroxyacids. **Peroxyacids,** a group of reagents with the general structure **RCO_3H,** have one more O atom than carboxylic acids (RCO_2H). Some peroxyacids are commercially available whereas others are prepared and used without isolation.

Two common peroxyacids are peroxyacetic acid and *meta*-chloroperoxybenzoic acid, abbreviated as **mCPBA.**



The most common oxidizing agents with metal–oxygen bonds contain either chromium in the +6 oxidation state (six Cr−O bonds) or manganese in the +7 oxidation state (seven Mn−O bonds). Common Cr^{6+} reagents include chromium(VI) oxide **(CrO_3)** and sodium or potassium dichromate **($Na_2Cr_2O_7$** and **$K_2Cr_2O_7$). These reagents are strong oxidants** used in the presence of a strong aqueous acid such as H_2SO_4. **Pyridinium chlorochromate (PCC),** a Cr^{6+} reagent that is soluble in halogenated organic solvents, can be used without strong acid present. This makes it a **more selective Cr^{6+} oxidant,** as described in Section 12.12.




CrO_3 **PCC**

The most common Mn^{7+} reagent is **$KMnO_4$** (potassium permanganate), a strong, water-soluble oxidant. Other oxidizing agents that contain metals include **OsO_4** (osmium tetroxide) and **Ag_2O** [silver(I) oxide].

In the remainder of Chapter 12, the oxidation of alkenes, alkynes, and alcohols—three functional groups already introduced in this text—is presented (Figure 12.7). Addition reactions to alkenes and alkynes that increase the number of C−O bonds are described in Sections 12.8–12.11. Oxidation of alcohols to carbonyl compounds appears in Sections 12.12–12.14.

Figure 12.7
Oxidation reactions of alkenes, alkynes, and alcohols



epoxidation (Sections 12.8, 12.15)

dihydroxylation (Section 12.9)


oxidative cleavage (Section 12.10)

oxidative cleavage (Section 12.11)


(Sections 12.12–12.14)

12.8 Epoxidation

Epoxidation is the addition of a single oxygen atom to an alkene to form an **epoxide.**




The weak π bond of the alkene is broken and two new C−O σ bonds are formed. Epoxidation is typically carried out with a peroxyacid, resulting in cleavage of the weak O−O bond of the reagent.





Epoxidation occurs via the concerted addition of one oxygen atom of the peroxyacid to the π bond as shown in Mechanism 12.4. Epoxidation resembles the formation of the bridged halonium ion in Section 10.13, in that two bonds in a three-membered ring are formed in one step.

Mechanism 12.4 Epoxidation of an Alkene with a Peroxyacid



- All bonds are broken and formed in a single step. The two epoxide C−O bonds are formed from one electron pair of the π bond and one lone pair of the peroxyacid. The weak O−O bond is broken.

Problem 12.15 What epoxide is formed when each alkene is treated with mCPBA?

a. 
b. 
c. 

12.8A The Stereochemistry of Epoxidation

Epoxidation occurs via **syn addition** of an O atom from either side of the planar double bond, so that both C−O bonds are formed on the same side. The relative position of substituents in the alkene reactant is **retained** in the epoxide product.


- **A cis alkene gives an epoxide with cis substituents. A trans alkene gives an epoxide with trans substituents.**

Epoxidation is a **stereospecific** reaction because cis and trans alkenes yield different stereoisomers as products, as illustrated in Sample Problem 12.2.

Sample Problem 12.2 Draw the stereoisomers formed when *cis*- and *trans*-but-2-ene are epoxidized with mCPBA.


Solution

To draw each product of epoxidation, add an O atom from either side of the alkene, and keep all substituents in their *original* orientations. The **cis** methyl groups in *cis*-but-2-ene become **cis** substituents in the epoxide. Addition of an O atom from either side of the trigonal planar alkene leads to the same compound—an **achiral meso compound that contains two stereogenic centers,** labeled in blue.




Epoxidation of *cis*- and *trans*-but-2-ene illustrates the general rule about the stereochemistry of reactions: **an achiral starting material gives achiral or racemic products.**

The **trans** methyl groups in *trans*-but-2-ene, become **trans** substituents in the epoxide. Addition of an O atom from either side of the trigonal planar alkene yields an equal mixture of two enantiomers—a **racemic mixture**—with two stereogenic centers labeled in blue**.**



Problem 12.16 Draw all stereoisomers formed when each alkene is treated with mCPBA.

a. b. c.




12.8B The Synthesis of Disparlure



In 1869, the gypsy moth was introduced into New England in an attempt to develop a silk industry. Some moths escaped into the wild and the population flourished. Mature gypsy moth caterpillars eat an average of one square foot of leaf surface per day, defoliating shade trees and entire forests. Many trees die after a single defoliation.

Disparlure, the sex pheromone of the female gypsy moth, is synthesized by a stepwise reaction sequence that uses an epoxidation reaction as the final step. Retrosynthetic analysis of disparlure illustrates three key operations:




- **Step [1]** The cis epoxide in disparlure is prepared from a cis alkene **A** by epoxidation.
- **Step [2]** **A** is prepared from an internal alkyne **B** by reduction.
- **Step [3]** **B** is prepared from acetylene and two 1° alkyl halides (**C** and **D**) by using S_N2 reactions with acetylide anions.

Figure 12.8 illustrates the synthesis of disparlure beginning with acetylene. The synthesis is conceptually divided into three parts:


- **Part [1]** Acetylene is converted to an internal alkyne **B** by forming two C−C bonds. Each bond is formed by treating an alkyne with base ($NaNH_2$) to form an acetylide anion, which reacts with an alkyl halide (**C** or **D**) in an S_N2 reaction (Section 11.11).
- **Part [2]** The internal alkyne **B** is reduced to a cis alkene **A** by syn addition of H_2 using the Lindlar catalyst (Section 12.5B).
- **Part [3]** The cis alkene **A** is epoxidized to disparlure using a peroxyacid such as mCPBA.

Figure 12.8
The synthesis of disparlure


Part [1] Formation of two C−C bonds using acetylide anions



Part [2] Reduction of alkyne **B** to form cis alkene **A**



Part [3] Epoxidation of **A** to form disparlure




- Disparlure has been used to control the spread of the gypsy moth caterpillar, a pest that has periodically devastated forests in the northeastern United States by defoliating many shade and fruit-bearing trees. The active pheromone is placed in a trap containing a poison or sticky substance, and the male moth is lured to the trap by the pheromone. Alternatively, thousands of disparlure-baited traps are placed along the edges of infestation. When the pheromone permeates the air, males are confused and can't locate individual females, so that mating is disrupted. Such a species-specific method presents a new way of controlling an insect population that avoids the widespread use of harmful, nonspecific pesticides.

How to separate a racemic mixture into its component enantiomers is discussed in Section 29.3.

Epoxidation of the cis alkene **A** from two different sides of the double bond affords two cis epoxides in the last step—a racemic mixture of two enantiomers. Thus, half of the product is the desired pheromone disparlure, but the other half is its biologically inactive enantiomer. Separating the desired from the undesired enantiomer is difficult and expensive, because both compounds have identical physical properties. A reaction that affords a chiral epoxide from an achiral precursor without forming a racemic mixture is discussed in Section 12.15.


12.9 Dihydroxylation

Dihydroxylation is the addition of two hydroxy groups to a double bond, forming a **1,2-diol** or **glycol.** Depending on the reagent, the two new OH groups can be added to the opposite sides (**anti** addition) or the same side (**syn** addition) of the double bond.




12.9A Anti Dihydroxylation

Anti dihydroxylation is achieved in two steps—epoxidation followed by opening of the ring with $^-$OH or H_2O. Cyclohexene, for example, is converted to a racemic mixture of two *trans*-cyclohexane-1,2-diols by anti addition of two OH groups.



The stereochemistry of the products can be understood by examining the stereochemistry of each step.




Epoxidation of cyclohexene adds an O atom from either above or below the plane of the double bond to form a single **achiral epoxide,** so only one representation is shown. Opening of the epoxide ring then occurs with **backside attack at either C−O bond.** Because the epoxide is drawn above the plane of the six-membered ring, nucleophilic attack occurs from **below** the plane. This reaction is a specific example of the opening of epoxide rings with strong nucleophiles, first presented in Section 9.15.

Because one OH group of the 1,2-diol comes from the epoxide and one OH group comes from the nucleophile ($^-$OH), the overall result is **anti addition of two OH groups** to an alkene.

Problem 12.17 Draw the products formed when both *cis*- and *trans*-but-2-ene are treated with a peroxyacid followed by $^{-}$OH (in H_2O). Explain how these reactions illustrate that anti dihydroxylation is stereospecific.

12.9B Syn Dihydroxylation


Syn dihydroxylation results when an alkene is treated with either **$KMnO_4$** or **OsO_4.**




cis-cyclohexane-1,2-diol *cis*-cyclopentane-1,2-diol

Each reagent adds two oxygen atoms to the same side of the double bond—that is, in a **syn** fashion—to yield a cyclic intermediate. Hydrolysis of the cyclic intermediate cleaves the metal–oxygen bonds, forming the *cis*-1,2-diol. With OsO_4, sodium bisulfite ($NaHSO_3$) is also added in the hydrolysis step.

Two O atoms are added to the **same** side of the C=C.




cis-1,2-diol




Although $KMnO_4$ is inexpensive and readily available, its use is limited by its insolubility in organic solvents. To prevent further oxidation of the product 1,2-diol, the reaction mixture must be kept basic with added $^{-}$OH.

Although OsO_4 is a more selective oxidant than $KMnO_4$ and is soluble in organic solvents, it is toxic and expensive. To overcome these limitations, dihydroxylation can be carried out by using a *catalytic* amount of OsO_4, if the oxidant ***N*-methylmorpholine *N*-oxide** (**NMO**) is also added.

NMO is an **amine oxide.** It is not possible to draw a Lewis structure of an amine oxide having only neutral atoms.




amine oxide



N-methylmorpholine *N*-oxide

NMO

In the catalytic process, dihydroxylation of the double bond converts the Os^{8+} oxidant into an Os^{6+} product, which is then re-oxidized by NMO to Os^{8+}. This Os^{8+} reagent can then be used for dihydroxylation once again, and the catalytic cycle continues.



NMO oxidizes the **Os^{6+} product** back to **Os^{8+}** to begin the cycle again.


Problem 12.18 Draw the products formed when both *cis*- and *trans*-but-2-ene are treated with OsO_4, followed by hydrolysis with $NaHSO_3 + H_2O$. Explain how these reactions illustrate that syn dihydroxylation is stereospecific.

12.10 Oxidative Cleavage of Alkenes




Lightning produces O_3 from O_2 during an electrical storm. Moreover, the pungent odor around a heavily used photocopy machine is O_3 produced from O_2 during the process. O_3 at ground level is an unwanted atmospheric pollutant. In the stratosphere, however, it protects us from harmful ultraviolet radiation, as discussed in Chapter 15.


Oxidative cleavage of an alkene breaks both the σ and π bonds of the double bond to form two carbonyl groups. Depending on the number of R groups bonded to the double bond, oxidative cleavage yields either **ketones** or **aldehydes.**



One method of oxidative cleavage relies on a two-step procedure using **ozone (O_3) as the oxidant** in the first step. Cleavage with ozone is called **ozonolysis.**




Addition of ozone to the π bond of the alkene forms an unstable intermediate called a **molozonide,** which then rearranges to an **ozonide** by a stepwise process. The unstable ozonide is then reduced without isolation to afford carbonyl compounds. **Zn (in H_2O) or dimethyl sulfide (CH_3SCH_3)** are two common reagents used to convert the ozonide to carbonyl compounds.



To draw the product of any oxidative cleavage:


- **Locate all π bonds in the molecule.**
- **Replace each C=C by two C=O bonds.**

Sample Problem 12.3 Draw the products when each alkene is treated with O_3 followed by CH_3SCH_3.




Solution

a. Cleave the double bond and replace it with two carbonyl groups.




b. For a cycloalkene, oxidative cleavage results in a **single molecule with two carbonyl groups—a dicarbonyl compound.**




Problem 12.19 Draw the products formed when each alkene is treated with O_3 followed by Zn, H_2O.

a. b. c.




Ozonolysis of dienes (and other polyenes) results in oxidative cleavage of all C=C bonds. The number of carbonyl groups formed in the products is *twice* the number of double bonds in the starting material. The *two* double bonds in limonene are converted to products containing *four* carbonyl groups.




Oxidative cleavage is a valuable tool for structure determination of unknown compounds. The ability to determine what alkene gives rise to a particular set of oxidative cleavage products is thus a useful skill, illustrated in Sample Problem 12.4.

Sample Problem 12.4 What alkene forms the following products after reaction with O_3 followed by CH_3SCH_3?




Solution

To draw the starting material, ignore the O atoms in the carbonyl groups and join the carbonyl carbons together by a C=C.






Problem 12.20 What alkene yields each set of oxidative cleavage products?

a. b. c. only




Problem 12.21 Draw the products formed when each diene is treated with O_3 followed by CH_3SCH_3.


a.  b.  c. 

12.11 Oxidative Cleavage of Alkynes




Alkynes also undergo oxidative cleavage of the σ bond and both π bonds of the triple bond. Internal alkynes are oxidized to **carboxylic acids (RCOOH),** whereas terminal alkynes afford carboxylic acids and **CO_2** from the *sp* hybridized C−H bond.




Oxidative cleavage is commonly carried out with O_3, followed by cleavage of the intermediate ozonide with H_2O.





Problem 12.22 Draw the products formed when each alkyne is treated with O_3 followed by H_2O.


a.  b.  c. 

Problem 12.23 What alkyne (or diyne) yields each set of oxidative cleavage products?

a. CO_2 + 

b.  only


c. 

d. 


12.12 Oxidation of Alcohols

Alcohols are oxidized to a variety of carbonyl compounds, depending on the type of alcohol and reagent. Oxidation occurs by replacing the C−H bonds *on the carbon bearing the OH group* by C−O bonds.


- **1° Alcohols** are oxidized to either **aldehydes** or **carboxylic acids** by replacing either one or two C–H bonds by C–O bonds.




- **2° Alcohols** are oxidized to **ketones** by replacing the one C–H bond by a C–O bond.



- **3° Alcohols have no H atoms on the carbon with the OH group,** so they are not easily oxidized.



Alcohol oxidations often occur by a pathway that involves bonding a leaving group Z to the oxygen, where Z is typically a metal in a high oxidation state. Elimination with a base then forms a C=O and a metal in a lower oxidation state.




The oxidation of alcohols to carbonyl compounds is typically carried out with Cr^{6+} oxidants, which are reduced to Cr^{3+} products.

- **CrO_3, $Na_2Cr_2O_7$, and $K_2Cr_2O_7$** are **strong, nonselective oxidants** used in aqueous acid ($H_2SO_4 + H_2O$).
- **PCC** (Section 12.7) is soluble in CH_2Cl_2 (dichloromethane), and can be used without strong acid present, making it a **more selective, milder oxidant.**


12.12A Oxidation of 2° Alcohols

Any of the Cr^{6+} oxidants effectively oxidizes 2° alcohols to ketones.



The mechanism for alcohol oxidation has two key parts: **formation of a chromate ester and loss of a proton.** Mechanism 12.5 is drawn for the oxidation of a general 2° alcohol with CrO_3.

Mechanism 12.5 Oxidation of an Alcohol with CrO_3



1—2 Nucleophilic attack of the alcohol on the electrophilic metal (Cr^{6+} oxidation state) followed by proton transfer forms a chromate ester.


3 A base removes a proton and the electron pair in the C–H bond forms the new π bond of the C=O. Carbon is oxidized because the number of C–O bonds increases, and Cr^{6+} is reduced to Cr^{4+}.

These three steps convert the Cr^{6+} oxidant to a Cr^{4+} product, which is then further reduced to a Cr^{3+} product by a series of steps.

12.12B Oxidation of 1° Alcohols


1° Alcohols are oxidized to either aldehydes or carboxylic acids, depending on the reagent.

- **1° Alcohols are oxidized to aldehydes (RCHO) under mild reaction conditions—using PCC in CH_2Cl_2.**
- **1° Alcohols are oxidized to carboxylic acids (RCOOH) under harsher reaction conditions: $Na_2Cr_2O_7$, $K_2Cr_2O_7$, or CrO_3 in the presence of H_2O and H_2SO_4.**



The mechanism for the oxidation of 1° alcohols to aldehydes parallels the oxidation of 2° alcohols to ketones detailed in Section 12.12A. Oxidation of a 1° alcohol to a carboxylic acid requires three operations: **oxidation first to the aldehyde, reaction with water,** and then further **oxidation to the carboxylic acid,** as shown in Mechanism 12.6.

Mechanism 12.6 Oxidation of a 1° Alcohol to a Carboxylic Acid




Part 1 The 1° alcohol is oxidized to an aldehyde by the three-step sequence in Mechanism 12.5.

Part 2 Water adds to the C=O to form a hydrate, a compound with two OH groups bonded to the same carbon, by a mechanism discussed in Section 21.13.

Part 3 Oxidation of the C–H bond of the hydrate follows Mechanism 12.5—formation of a chromate ester and loss of a proton.


Cr^{6+} oxidations are characterized by a color change, as the **red-orange Cr^{6+} reagent** is reduced to **green Cr^{3+}.** The first devices used to measure blood alcohol content in individuals suspected of "driving under the influence" made use of this color change. Oxidation of CH_3CH_2OH, the 1° alcohol in alcoholic beverages, with orange $K_2Cr_2O_7$ forms CH_3COOH and green Cr^{3+}.



Blood alcohol level can be determined by having an individual blow into a tube containing $K_2Cr_2O_7$, H_2SO_4, and an inert solid. The alcohol in the exhaled breath is oxidized by the Cr^{6+} reagent, which turns green in the tube (Figure 12.9). The higher the concentration of CH_3CH_2OH in the breath, the more Cr^{6+} is reduced, and the farther the green Cr^{3+} color extends down the length of the sample tube. This value is then correlated with blood alcohol content to determine if an individual has surpassed the legal blood alcohol limit.

Figure 12.9 Blood alcohol screening

a. Schematic of an alcohol testing device




b. Consumer product



- The oxidation of CH_3CH_2OH with $K_2Cr_2O_7$ to form CH_3COOH and Cr^{3+} was the first available method for the routine testing of alcohol concentration in exhaled air. Some consumer products for alcohol screening are still based on this technology.

Problem 12.24 Draw the organic products in each of the following reactions.




12.13 Green Chemistry

Several new methods of oxidation are based on green chemistry. ***Green chemistry* is the use of environmentally benign methods to synthesize compounds.** Its purpose is to use safer reagents and less solvent, and develop reactions that form fewer by-products and generate less waste.

Green polymer synthesis using starting materials derived from renewable resources (rather than petroleum) is discussed in Chapter 30.


Since many oxidation methods use toxic reagents (such as OsO_4 and O_3) and corrosive acids (such as H_2SO_4), or they generate carcinogenic by-products (such as Cr^{3+}), alternative reactions have been developed. One method uses a polymer-supported Cr^{6+} reagent—$HCrO_4^-$–Amberlyst A-26 resin—that avoids the use of strong acid, and forms a Cr^{3+} by-product that can be easily removed from the product by filtration.

The Amberlyst A-26 resin consists of a complex hydrocarbon network with cationic ammonium ion appendages that serve as counterions to the anionic chromium oxidant, $HCrO_4^-$. Heating the insoluble polymeric reagent with an alcohol results in oxidation to a carbonyl compound, with formation of an insoluble Cr^{3+} by-product. Not only can the metal by-product be removed by filtration without added solvent, it can also be regenerated and reused in a subsequent reaction.




Amberlyst A-26 resin Cr^{6+} oxidant


With $HCrO_4^-$–Amberlyst A-26 resin, 1° alcohols are oxidized to aldehydes and 2° alcohols are oxidized to ketones.



1° alcohol




2° alcohol




potassium peroxymonosulfate

Many other green approaches to oxidation that avoid the generation of metal by-products entirely are also under active investigation. For example, potassium peroxymonosulfate, $KHSO_5$, is a sulfate derivative of hydrogen peroxide, sold as a triple salt ($2\ KHSO_5 \bullet KHSO_4 \bullet K_2SO_4$) under the trade name of Oxone. Oxone oxidizes a variety of substrates without the presence of a heavy metal like chromium or manganese, and in some cases, oxidation reactions can be carried out in water or aqueous solutions. The weak oxygen–oxygen bond of the reagent is cleaved during oxidation, and a sulfate salt (K_2SO_4) is formed as by-product. Two examples of oxidations of alcohols are shown.



Problem 12.25 What carbonyl compound is formed when each alcohol is treated with $HCrO_4^-$–Amberlyst A-26 resin?




Problem 12.26 Sodium hypochlorite (NaOCl, the oxidant in household bleach) in aqueous CH_3COOH is also touted as a "green" oxidizing agent. For example, oxidation of $(CH_3)_2CHOH$ with NaOCl forms $(CH_3)_2C{=}O$ along with NaCl and H_2O. (a) What advantages and/or disadvantages does this method have over oxidation with $HCrO_4^-$–Amberlyst A-26 resin? (b) What advantages and/or disadvantages does this method have over oxidation with CrO_3, H_2SO_4, H_2O?

12.14 Biological Oxidation


Many reactions in biological systems involve oxidation or reduction. Instead of using Cr^{6+} reagents for oxidation, cells use two organic compounds—a high molecular weight **enzyme** and a simpler **coenzyme** that serves as the oxidizing agent.

The coenzyme often used to oxidize alcohols in biological systems is **nicotinamide adenine dinucleotide,** abbreviated as **NAD^+.** Although the structure is complex, only a portion of the molecule, labeled in red, participates in redox reactions.




nicotinamide adenine dinucleotide
NAD^+

Biological oxidation of an alcohol occurs by transferring a hydride, a hydrogen atom with two electrons, from the alcohol to NAD^+ to form a carbonyl group. In the process, NAD^+ is reduced to nicotinamide adenine dinucleotide (reduced form), abbreviated as **NADH.** NADH is a biological reducing agent that converts carbonyl compounds to alcohols, as discussed in Section 20.6.




nicotinamide adenine dinucleotide
(reduced form)
NADH

For example, when CH_3CH_2OH (ethanol) is ingested, it is oxidized in the liver by NAD^+ to CH_3CHO (acetaldehyde), and then to CH_3COO^- (acetate anion, the conjugate base of acetic acid). Acetate is the starting material for the synthesis of fatty acids and cholesterol. Both oxidations are catalyzed by a dehydrogenase enzyme.




If more ethanol is ingested than can be metabolized in a given time, the concentration of acetaldehyde builds up. This toxic compound is responsible for the feelings associated with a hangover.


antabuse

Antabuse, a drug given to alcoholics to prevent them from consuming alcoholic beverages, acts by interfering with the normal oxidation of ethanol. Antabuse inhibits the oxidation of acetaldehyde to the acetate anion. Because the first step in ethanol metabolism occurs but the second does not, the concentration of acetaldehyde rises, causing an individual to become violently ill.


Like ethanol, methanol is oxidized by the same enzymes to give an aldehyde and an acid: formaldehyde and formic acid. These oxidation products are extremely toxic because they cannot be used by the body. As a result, the pH of the blood decreases, and blindness and death can follow.



Because the enzymes have a higher affinity for ethanol than methanol, methanol poisoning is treated by giving ethanol to the afflicted individual. With both methanol and ethanol in the patient's system, the enzymes react more readily with ethanol, allowing the methanol to be excreted unchanged without the formation of methanol's toxic oxidation products.

12.15 Sharpless Epoxidation

In all of the reactions discussed so far, an **achiral starting material has reacted with an achiral reagent to give either an achiral product or a racemic mixture of two enantiomers.** If you are trying to make a chiral product, this means that only half of the product mixture is the desired enantiomer and the other half is the undesired one. The synthesis of disparlure, outlined in Figure 12.8, exemplifies this dilemma.




K. Barry Sharpless shared the 2001 Nobel Prize in Chemistry for his work on chiral oxidation reactions.

K. Barry Sharpless, currently at The Scripps Research Institute, reasoned that using a chiral reagent might make it possible to favor the formation of one enantiomer over the other.

- An *enantioselective* reaction affords predominantly or exclusively one enantiomer.
- A reaction that converts an achiral starting material into predominantly one enantiomer is also called an *asymmetric reaction.*

The Sharpless asymmetric epoxidation is an enantioselective reaction that oxidizes alkenes to epoxides. Only the double bonds of **allylic alcohols**—that is, alcohols having a hydroxy group on the carbon adjacent to a C=C—are oxidized in this reaction.




allylic alcohol — O added from above — O added from below

Sharpless reagent
$(CH_3)_3C-OOH$
$Ti[OCH(CH_3)_2]_4$
(+)- or (–)-diethyl tartrate


- With Sharpless reagent, one enantiomer is favored.
- The new stereogenic center is labeled in blue.

The **Sharpless reagent** consists of three components: *tert*-butyl hydroperoxide, **$(CH_3)_3COOH$;** a titanium catalyst—usually titanium(IV) isopropoxide, **$Ti[OCH(CH_3)_2]_4$;** and **diethyl tartrate (DET).** There are two different chiral diethyl tartrate isomers, labeled as (+)-DET or (–)-DET to indicate the direction in which they rotate polarized light.



(+)-(*R*,*R*)-diethyl tartrate

(+)-DET



(–)-(*S*,*S*)-diethyl tartrate


(–)-DET

The identity of the DET isomer determines which enantiomer is the major product obtained in the epoxidation of an allylic alcohol with the Sharpless reagent.



(+)-DET is prepared from (+)-(*R*,*R*)-tartaric acid [$HO_2CCH(OH)CH(OH)CO_2H$], a naturally occurring carboxylic acid found in grapes and sold as a by-product of the wine industry.

Enantiomeric excess = ***ee*** = % of one enantiomer – % of the other enantiomer.




Stereogenic centers are labeled in blue.

The degree of enantioselectivity of a reaction is measured by its enantiomeric excess (***ee***) (Section 5.12D). Reactions [1] and [2] are highly enantioselective because each has an enantiomeric excess of 95% (97.5% of the major enantiomer – 2.5% of the minor enantiomer).


To determine which enantiomer is formed for a given isomer of DET, draw the allylic alcohol in a plane, with the **C=C horizontal and the OH group in the upper right corner;** then:

- **Epoxidation with (–)-DET adds an oxygen atom from above the plane.**
- **Epoxidation with (+)-DET adds an oxygen atom from below the plane.**



Sample Problem 12.5

Predict the major product in each epoxidation.




Solution


To draw an epoxidation product:

- Draw the allylic alcohol with the **C = C horizontal and the OH group in the upper right corner of the alkene.** Re-draw the alkene if necessary.
- **(+)-DET** adds the O atom from **below,** and **(–)-DET** adds the O atom from **above.**


a. Since the C = C is drawn horizontal with the OH group in the upper right corner, it is not necessary to re-draw the alkene. With **(+)-DET,** the O atom is added from **below.**



b. The allylic alcohol must be re-drawn with the C = C horizontal and the OH group in the upper right corner. Because **(–)-DET** is used, the O atom is then added from **above.**




Problem 12.27 Draw the products of each Sharpless epoxidation.




The Sharpless epoxidation has been used to synthesize many chiral natural products, including two insect pheromones—(+)-α-multistriatin and (–)-frontalin, as shown in Figure 12.10.

Figure 12.10
The synthesis of chiral insect pheromones using asymmetric epoxidation



- The bonds in the products that originate from the epoxide intermediate are indicated in red.

Problem 12.28 Explain why only one C = C of geraniol is epoxidized with the Sharpless reagent.



geraniol

KEY CONCEPTS

Oxidation and Reduction

Summary: Terms that Describe Reaction Selectivity

- A **regioselective reaction** forms predominantly or exclusively one constitutional isomer (8.5).
- A **stereoselective reaction** forms predominantly or exclusively one stereoisomer (8.5).
- An **enantioselective reaction** forms predominantly or exclusively one enantiomer (12.15).

Definitions of Oxidation and Reduction (12.1)

Oxidation reactions result in:


- an increase in the number of C–Z bonds, *or*
- a decrease in the number of C–H bonds

Reduction reactions result in:

- a decrease in the number of C–Z bonds, *or*
- an increase in the number of C–H bonds

Reduction Reactions

[1] Reduction of alkenes—Catalytic hydrogenation (12.3)




alkane

- **Syn addition** of H_2 occurs.
- Increasing alkyl substitution on the C=C decreases the rate of reaction.

[2] Reduction of alkynes


a. R—≡—R $\xrightarrow[\text{Pd-C}]{2\ H_2}$



alkane

- Two equivalents of H_2 are added and four new C–H bonds are formed (12.5A).


b. R—≡—R $\xrightarrow[\text{Lindlar catalyst}]{H_2}$



cis alkene

- **Syn addition** of H_2 occurs, forming a **cis** alkene (12.5B).
- The Lindlar catalyst is deactivated; reaction stops after one equivalent of H_2 has added.

c. R—≡—R $\xrightarrow[NH_3]{Na}$



trans alkene

- **Anti addition** of H_2 occurs, forming a **trans** alkene (12.5C).


[3] Reduction of alkyl halides (12.6)

$$R{-}X \xrightarrow[[2]\ H_2O]{[1]\ LiAlH_4} R{-}H$$

alkane

- The reaction follows an $\mathbf{S_N2}$ mechanism.
- CH_3X and RCH_2X react faster than a more substituted RX.

[4] Reduction of epoxides (12.6)




alcohol

- The reaction follows an $\mathbf{S_N2}$ mechanism.
- In unsymmetrical epoxides, H^- (from $LiAlH_4$) attacks at the less substituted carbon.

Oxidation Reactions


[1] Oxidation of alkenes

a. Epoxidation (12.8)




- The mechanism has **one step.**
- **Syn addition** of an O atom occurs.
- The reaction is stereospecific.

b. Anti dihydroxylation (12.9A)




- Opening of an epoxide ring intermediate with ^-OH or H_2O forms a 1,2-diol with two OH groups added in an **anti** fashion.

c. Syn dihydroxylation (12.9B)




- Each reagent adds two new C – O bonds to the C = C in a **syn** fashion.


d. Oxidative cleavage (12.10)



- Both the σ and π bonds of the alkene are cleaved to form two carbonyl groups.

[2] Oxidative cleavage of alkynes (12.11)






- The σ bond and both π bonds of the alkyne are cleaved.


[3] Oxidation of alcohols (12.12, 12.13)

a.




- Oxidation of a 1° alcohol with PCC or $HCrO_4^-$–Amberlyst A-26 resin stops at the aldehyde stage. Only one C – H bond is replaced by a C–O bond.

b.




- Oxidation of a 1° alcohol under harsher reaction conditions—CrO_3 (or $Na_2Cr_2O_7$ or $K_2Cr_2O_7$) + H_2O + H_2SO_4—leads to a RCOOH. Two C–H bonds are replaced by two C – O bonds.

c.



- Because a 2° alcohol has only one C–H bond on the carbon bearing the OH group, all Cr^{6+} reagents—PCC, CrO_3, $Na_2Cr_2O_7$, $K_2Cr_2O_7$, or $HCrO_4^-$ (Amberlyst A-26 resin)—oxidize a 2° alcohol to a ketone.


[4] Asymmetric epoxidation of allylic alcohols (12.15)



PROBLEMS


Problems Using Three-Dimensional Models

12.29 Draw the products formed when **A** is treated with each reagent: (a) H_2 + Pd-C; (b) mCPBA; (c) PCC; (d) CrO_3, H_2SO_4, H_2O; (e) Sharpless reagent with (+)-DET.




A

12.30 Draw the products formed when the following diene is treated with O_3 followed by CH_3SCH_3.




12.31 Devise a synthesis of the following compound from acetylene and organic compounds containing two carbons or fewer. You may use any other required reagents.




Classifying Reactions as Oxidation or Reduction

12.32 Label each reaction as oxidation, reduction, or neither.



Hydrogenation

12.33 Draw the organic products formed when each alkene is treated with H_2, Pd-C. Indicate the three-dimensional structure of all stereoisomers formed.




12.34 Match each alkene to its heat of hydrogenation.
Alkenes: 3-methylbut-1-ene, 2-methylbut-1-ene, 2-methylbut-2-ene
$\Delta H°$ (hydrogenation) kJ/mol: –119, –127, –112


12.35 How many rings and π bonds are contained in compounds **A–C?** Draw one possible structure for each compound.

a. Compound **A** has molecular formula C_5H_8 and is hydrogenated to a compound having molecular formula C_5H_{10}.
b. Compound **B** has molecular formula $C_{10}H_{16}$ and is hydrogenated to a compound having molecular formula $C_{10}H_{18}$.
c. Compound **C** has molecular formula C_8H_8 and is hydrogenated to a compound having molecular formula C_8H_{16}.

12.36 For alkenes **A, B,** and **C:** (a) Rank **A, B,** and **C** in order of increasing heat of hydrogenation; (b) rank **A, B,** and **C** in order of increasing rate of reaction with H_2, Pd-C; (c) draw the products formed when each alkene is treated with ozone, followed by Zn, H_2O.



12.37 Stearidonic acid ($C_{18}H_{28}O_2$) is an unsaturated fatty acid obtained from oils isolated from hemp and blackcurrant (see also Problem 10.11).



stearidonic acid


a. What fatty acid is formed when stearidonic acid is hydrogenated with excess H_2 and a Pd catalyst?
b. What fatty acids are formed when stearidonic acid is hydrogenated with one equivalent of H_2 and a Pd catalyst?
c. Draw the structure of a possible product formed when stearidonic acid is hydrogenated with one equivalent of H_2 and a Pd catalyst, and one double bond is isomerized to a trans isomer.
d. How do the melting points of the following fatty acids compare: stearidonic acid; one of the products formed in part (b); the product drawn in part (c)?

Reactions—General

12.38 Draw the organic products formed when cyclopentene is treated with each reagent. With some reagents, no reaction occurs.

a. H_2 + Pd-C
b. H_2 + Lindlar catalyst
c. Na, NH_3
d. CH_3CO_3H
e. [1] CH_3CO_3H; [2] H_2O, HO^-
f. [1] OsO_4 + NMO; [2] $NaHSO_3$, H_2O
g. $KMnO_4$, H_2O, HO^-
h. [1] $LiAlH_4$; [2] H_2O
i. [1] O_3; [2] CH_3SCH_3
j. $(CH_3)_3COOH$, $Ti[OCH(CH_3)_2]_4$, (–)-DET
k. mCPBA
l. Product in (k); then [1] $LiAlH_4$; [2] H_2O


12.39 Draw the organic products formed when allylic alcohol **A** is treated with each reagent.




A

a. H_2 + Pd-C
b. mCPBA
c. PCC
d. CrO_3, H_2SO_4, H_2O
e. $(CH_3)_3COOH$, $Ti[OCH(CH_3)_2]_4$, (+)-DET
f. $(CH_3)_3COOH$, $Ti[OCH(CH_3)_2]_4$, (–)-DET
g. [1] PBr_3; [2] $LiAlH_4$; [3] H_2O
h. $HCrO_4^-$–Amberlyst A-26 resin

12.40 Draw the organic products formed in each reaction.




12.41 One step in the degradation of fats involves the reaction of (*R*)-glycerol phosphate with NAD^+ in the presence of the enzyme glycerol phosphate dehydrogenase. What products are formed if reaction occurs only at the 2° alcohol?



(*R*)-glycerol phosphate


12.42 Draw the structure of two different epoxides that would yield 2-methylpentan-2-ol [$(CH_3)_2C(OH)CH_2CH_2CH_3$] when reduced with $LiAlH_4$.

12.43 Hydrogenation of alkene **A** with D_2 in the presence of Pd-C affords a single product **B.** Keeping this result in mind, what compound is formed when **A** is treated with each reagent: (a) mCPBA; (b) Br_2, H_2O followed by base? Explain these results.




A **B**

12.44 What alkene is needed to synthesize each 1,2-diol using [1] OsO_4 followed by $NaHSO_3$ in H_2O; or [2] CH_3CO_3H followed by ^-OH in H_2O?




12.45 Draw a stepwise mechanism for the reduction of epoxide **A** to alcohol **B** using $LiAlH_4$. What product would be formed if $LiAlD_4$ were used as reagent? Indicate the stereochemistry of all stereogenic centers in the product using wedges and dashed wedges.




A **B**

12.46 (a) What product is formed in Step [1] of the following reaction sequence? (b) Draw a mechanism for Step [2] that accounts for the observed stereochemistry. (c) What reaction conditions are necessary to form chiral **A** from prop-2-en-1-ol ($CH_2=CHCH_2OH$)?




A **B**

12.47 Draw the products formed after Steps [1] and [2] in the following three-step sequence. Then draw stepwise mechanisms for each step.




Oxidative Cleavage


12.48 Draw the products formed in each oxidative cleavage.




12.49 What alkene or alkyne yields each set of products after oxidative cleavage with ozone?



12.50 Identify the starting material in each reaction.



12.51 Draw the products formed when each naturally occurring compound is treated with O_3 followed by Zn, H_2O.



Identifying Compounds from Reactions


12.52 Identify compounds **A, B,** and **C.**

a. Compound **A** has molecular formula C_8H_{12} and reacts with two equivalents of H_2. **A** gives $HCOCH_2CH_2CHO$ as the only product of oxidative cleavage with O_3 followed by CH_3SCH_3.


b. Compound **B** has molecular formula C_6H_{10} and gives $(CH_3)_2CHCH_2CH_2CH_3$ when treated with excess H_2 in the presence of Pd. **B** reacts with $NaNH_2$ and CH_3I to form compound **C** (molecular formula C_7H_{12}).

12.53 Oximene and myrcene, two hydrocarbons isolated from alfalfa that have the molecular formula $C_{10}H_{16}$, both yield 2,6-dimethyloctane when treated with H_2 and a Pd catalyst. Ozonolysis of oximene forms $(CH_3)_2C=O$, $CH_2=O$, $CH_2(CHO)_2$, and CH_3COCHO. Ozonolysis of myrcene yields $(CH_3)_2C=O$, $CH_2=O$ (two equiv), and $HCOCH_2CH_2COCHO$. Identify the structures of oximene and myrcene.

12.54 An unknown compound **A** of molecular formula $C_{10}H_{18}O$ reacts with H_2SO_4 to form two compounds (**B** and **C**) of molecular formula $C_{10}H_{16}$. **B** and **C** both react with H_2 in the presence of Pd-C to form decalin. Ozonolysis of **B** forms **D,** and ozonolysis of **C** forms a diketone **E** of molecular formula $C_{10}H_{16}O_2$. Identify the structures of compounds **A, B, C,** and **E.**



decalin




D


12.55 DHA is a fatty acid derived from fish oil and an abundant fatty acid in vertebrate brains. Hydrogenation of DHA forms docosanoic acid $[CH_3(CH_2)_{20}CO_2H]$ and ozonolysis forms CH_3CH_2CHO, $CH_2(CHO)_2$ (five equivalents), and $HCOCH_2CH_2CO_2H$. What is the structure of DHA if all double bonds have the *Z* configuration?

12.56 One compound that contributes to the "seashore smell" at beaches in Hawai'i is dictyopterene D', a component of a brown edible seaweed called limu lipoa. Hydrogenation of dictyopterene D' with excess H_2 in the presence of a Pd catalyst forms butylcycloheptane. Ozonolysis with O_3 followed by $(CH_3)_2S$ forms $CH_2(CHO)_2$, $HCOCH_2CH(CHO)_2$, and CH_3CH_2CHO. What are possible structures of dictyopterene D'?


Sharpless Asymmetric Epoxidation

12.57 Draw the product of each asymmetric epoxidation reaction.

a. 

b. 


12.58 Epoxidation of the following allylic alcohol using the Sharpless reagent with (–)-DET gives two epoxy alcohols in a ratio of 87:13.




a. Assign structures to the major and minor product.

b. What is the enantiomeric excess in this reaction?


12.59 What allylic alcohol and DET isomer are needed to make each chiral epoxide using a Sharpless asymmetric epoxidation reaction?

a. 

b. 

c. 


12.60 Identify **A** in the following reaction sequence and draw a mechanism for the conversion of **A** to **B. B** has been converted to (*S,S*)-reboxetine, an antidepressant marketed outside the United States.



(*S*,*S*)-reboxetine

Synthesis


12.61 Devise a synthesis of muscalure, the sex pheromone of the common housefly, from acetylene and any other required reagents.




muscalure


12.62 It is sometimes necessary to isomerize a cis alkene to a trans alkene in a synthesis, a process that cannot be accomplished in a single step. Using the reactions you have learned in Chapters 8–12, devise a stepwise method to convert *cis*-but-2-ene to *trans*-but-2-ene.

12.63 Devise a synthesis of each compound from acetylene and any other required reagents.



12.64 Devise a synthesis of compound **A** from the given starting materials. You may use any other inorganic reagents or organic alcohols. **A** was used to prepare aliskiren, a drug used to treat hypertension (see also Problem 5.7).






aliskiren

12.65 Devise a synthesis of (*E*)-hex-2-ene from pent-1-ene and any needed organic compounds or inorganic reagents.


12.66 Devise a synthesis of each compound from the indicated starting material, organic compounds containing one or two carbons, and any other required reagents.



12.67 Devise a synthesis of 1-phenyl-5-methylhexane [$C_6H_5(CH_2)_4CH(CH_3)_2$] from acetylene, alkyl halides, and any required inorganic reagents.


12.68 Devise a synthesis of (3*R*,4*S*)-3,4-dichlorohexane from acetylene and any needed organic compounds or inorganic reagents.

12.69 Devise a synthesis of each compound from CH_3CH_2OH as the only organic starting material; that is, every carbon in the product must come from a molecule of ethanol. You may use any other needed inorganic reagents.



Challenge Problems


12.70 The Birch reduction is a dissolving metal reaction that converts substituted benzenes to cyclohexa-1,4-dienes using Li and liquid ammonia in the presence of an alcohol. Draw a stepwise mechanism for the following Birch reduction.



12.71 In the Cr^{6+} oxidation of cyclohexanols, it is generally true that sterically hindered alcohols react faster than unhindered alcohols. Which of the following alcohols should be oxidized more rapidly?




12.72 Draw a stepwise mechanism for the following reaction.




R = alkyl group R = alkyl group

12.73 Dihydroxylation of an alkene can be carried out with H_2O_2 in HCO_2H. In this reaction, *trans*-but-2-ene affords (2*R*,3*S*)-butane-2,3-diol, whereas *cis*-but-2-ene affords a mixture of (2*R*,3*R*)-butane-2,3-diol and (2*S*,3*S*)-butane-2,3-diol. Does dihydroxylation by this method occur with syn or anti addition?

12.74 Draw a stepwise mechanism for the following reaction.




12.75 Sharpless epoxidation of allylic alcohol **X** forms compound **Y.** Treatment of **Y** with NaOH and C_6H_5SH in an alcohol–water mixture forms **Z.** Identify the structure of **Y** and draw a mechanism for the conversion of **Y** to **Z.** Account for the stereochemistry of the stereogenic centers in **Z. Z** has been used as an intermediate in the synthesis of chiral carbohydrates.



Mass Spectrometry and Infrared Spectroscopy

13



13.1 Mass spectrometry
13.2 Alkyl halides and the M + 2 peak
13.3 Fragmentation
13.4 Other types of mass spectrometry
13.5 Electromagnetic radiation
13.6 Infrared spectroscopy
13.7 IR absorptions
13.8 IR and structure determination

The serendipitous discovery of **penicillin** by Scottish bacteriologist Sir Alexander Fleming in 1928 is considered one of the single most important events in the history of medicine. Penicillin G and related compounds are members of the β-lactam family of antibiotics, all of which contain a strained four-membered amide ring that is responsible for their biological activity. Penicillin was first used to cure a streptococcal infection in 1942, and by 1944 penicillin production was given high priority by the United States government, because it was needed to treat the many injured soldiers in World War II. The unusual structure of penicillin was elucidated by modern instrumental methods in the 1940s. In Chapter 13, we learn about mass spectrometry and infrared spectroscopy, two techniques for characterizing organic compounds like penicillin.

Whether a compound is prepared in the laboratory or isolated from a natural source, a chemist must determine its identity. Seventy years ago, determining the structure of an organic compound involved a series of time-consuming operations: measuring physical properties (melting point, boiling point, solubility, and density), identifying the functional groups using a series of chemical tests, and converting an unknown compound into another compound whose physical and chemical properties were then characterized as well.

Although still a challenging task, structure determination has been greatly simplified by modern instrumental methods. These techniques have both decreased the time needed for compound characterization, and increased the complexity of compounds whose structures can be completely determined.

In Chapter 13 we examine **mass spectrometry (MS),** which is used to determine the molecular weight and molecular formula of a compound, and **infrared (IR) spectroscopy,** a tool used to identify a compound's functional groups. Chapter 14 is devoted to **nuclear magnetic resonance (NMR) spectroscopy,** which is used to identify the carbon–hydrogen framework in a compound, making it the most powerful spectroscopic tool for organic structure analysis. **Each of these methods relies on the interaction of an energy source with a molecule to produce a change that is recorded in a spectrum.**


13.1 Mass Spectrometry

***Mass spectrometry* is a technique used for measuring the molecular weight and determining the molecular formula of an organic molecule.**

13.1A General Features


In the most common type of **mass spectrometer,** a molecule is vaporized and ionized, usually by bombardment with a beam of high-energy electrons, as shown in Figure 13.1. The energy of these electrons is typically about 6400 kJ, or 70 electron volts (eV). Because it takes ~400 kJ of energy to cleave a typical σ bond, 6400 kJ is an enormous amount of energy to come into contact with a molecule. This electron beam ionizes a molecule by causing it to eject an electron.

Figure 13.1
Schematic of a mass spectrometer



In a mass spectrometer, a sample is vaporized and bombarded by a beam of electrons to form an unstable radical cation, which then decomposes to smaller fragments. The positively charged ions are accelerated toward a negatively charged plate, and then passed through a curved analyzer tube in a magnetic field, where they are deflected by different amounts depending on their ratio of mass to charge (*m*/*z*). A mass spectrum plots the intensity of each ion versus its *m*/*z* ratio.


The term **spectro*scopy*** is usually used for techniques that use electromagnetic radiation as an energy source. Because the energy source in MS is a beam of electrons, the term **mass spectro*metry*** is used instead.



The species formed is a **radical cation,** symbolized $\mathbf{M^{+\bullet}}$**.** It is a radical because it has an unpaired electron, and it is a cation because it has one fewer electron than it started with.

- **The radical cation $M^{+\bullet}$ is called the *molecular ion* or the *parent ion.***


A single electron has a negligible mass, so the **mass of $M^{+\bullet}$ represents the molecular weight** of M. Because the molecular ion $\mathbf{M^{+\bullet}}$ is inherently unstable, it decomposes. Single bonds break to form ***fragments,* radicals and cations having a lower molecular weight than the molecular ion.** A mass spectrometer analyzes the masses of cations only. The cations are accelerated in an electric field and deflected in a curved path in a magnetic field, thus sorting the molecular ion and its fragments by their **mass-to-charge (*m*/*z*) ratio.** Because z is almost always +1, m/z actually measures the mass (m) of the individual ions.



- **A *mass spectrum* plots the amount of each cation (its relative abundance) versus its mass.**

The whole-number mass of CH_4 is (1 C × 12 amu) + (4 H × 1 amu) = 16 amu; amu = atomic mass unit.


A mass spectrometer analyzes the masses of *individual* molecules, not the weighted average mass of a group of molecules, so the whole-number masses of the most common individual isotopes must be used to calculate the mass of the molecular ion. Thus, the mass of the molecular ion for CH_4 should be 16. As a result, the mass spectrum of CH_4 shows a line for the molecular ion—the parent peak or **M** peak—at $m/z = 16$.



m/*z*	relative abundance
17	1.2
16	100.0 ← base peak
15	85.9
14	16.1
13	8.1
12	2.8

The tallest peak in a mass spectrum is called the **base peak.** For CH_4, the base peak is also the M peak, although this may *not* always be the case for all organic compounds.


The mass spectrum of CH_4 consists of more peaks than just the M peak. What is responsible for the peaks at $m/z < 16$? Because the molecular ion is unstable, it fragments into other cations and radical cations containing one, two, three, or four fewer hydrogen atoms than methane itself. Thus, the peaks at m/z = 15, 14, 13, and 12, are due to these lower molecular weight fragments. The decomposition of a molecular ion into lower molecular weight fragments is called **fragmentation.**



What is responsible for the small peak at $m/z = 17$ in the mass spectrum of CH_4? Although most carbon atoms have an atomic mass of 12, 1.1% of them have an additional neutron in the nucleus, giving them an atomic mass of 13. When one of these carbon-13 isotopes forms methane, it gives a molecular ion peak at $m/z = 17$ in the mass spectrum. This peak is called the **M + 1** peak.

These key features—the molecular ion, the base peak, and the M + 1 peak—are illustrated in the mass spectrum of hexane in Figure 13.2.

Figure 13.2
Mass spectrum of hexane ($CH_3CH_2CH_2CH_2CH_2CH_3$)





- The molecular ion for hexane (molecular formula C_6H_{14}) is at $m/z = 86$.
- The base peak (relative abundance = 100) occurs at $m/z = 57$.
- A small M + 1 peak occurs at $m/z = 87$.


13.1B Analyzing Unknowns Using the Molecular Ion

Because the **mass of the molecular ion equals the molecular weight of a compound,** a mass spectrum can be used to distinguish between compounds that have similar physical properties but different molecular weights, as illustrated in Sample Problem 13.1.

Sample Problem 13.1 Pentane, pent-1-ene, and pent-1-yne are low-boiling hydrocarbons that have different molecular ions in their mass spectra. Match each hydrocarbon to its mass spectrum.







Solution

To solve this problem, first determine the molecular formula and molecular weight of each compound. Then, because the molecular weight of the compound equals the mass of the molecular ion, match the molecular weight to *m/z* for the molecular ion:

Compound	Molecular formula	Molecular weight = *m/z* of molecular ion	Spectrum
 pentane	C_5H_{12}	72	[3]
 pent-1-ene	C_5H_{10}	70	[2]
 pent-1-yne	C_5H_8	68	[1]

Problem 13.1 What is the mass of the molecular ion formed from compounds having each molecular formula: (a) C_3H_6O; (b) $C_{10}H_{20}$; (c) $C_8H_8O_2$; (d) methamphetamine ($C_{10}H_{15}N$)?

How to use the mass of the molecular ion to propose molecular formulas for an unknown is shown in Sample Problem 13.2. In this process, keep in mind the following useful fact. Hydrocarbons like methane (CH_4) and hexane (C_6H_{14}), as well as compounds that contain only C, H, and O atoms, always have a molecular ion with an *even* mass. An odd molecular ion generally indicates that a compound contains nitrogen.

Sample Problem 13.2 Propose possible molecular formulas for a compound with a molecular ion at *m/z* = 86.

Solution

Because the molecular ion has an **even** mass, the compound likely contains C, H, and possibly O atoms. Begin by determining the molecular formula for a hydrocarbon having a molecular ion at 86. Then, because the mass of an O atom is 16 (the mass of CH_4), replace CH_4 by O to give a molecular formula containing one O atom. Repeat this last step to give possible molecular formulas for compounds with two or more O atoms.

For a molecular ion at *m/z* = 86:

Possible hydrocarbons:

- Divide 86 by 12 (mass of 1 C atom). This gives the maximum number of C's possible.

$$\frac{86}{12} = 7 \text{ C's maximum (remainder = 2)} \longrightarrow C_7H_2$$

- Replace 1 C by 12 H's for another possible molecular formula.

$$C_7H_2 \xrightarrow[+\ 12\ H\text{'s}]{-\ 1\ C} C_6H_{14}$$

Possible compounds with C, H, and O:


- Substitute 1 O for CH_4. (This can't be done for C_7H_2.)

$$C_6H_{14} \xrightarrow[+\ 1\ O]{-\ CH_4} C_5H_{10}O$$


- Repeat the process.

$$C_5H_{10}O \xrightarrow[+\ 1\ O]{-\ CH_4} C_4H_6O_2$$

The effect of N atoms on the mass of the molecular ion in a mass spectrum is called the **nitrogen rule: A compound that contains an *odd* number of N atoms gives an odd molecular ion.** Conversely, a compound that contains an *even* number of N atoms (including *zero*) gives an *even* molecular ion. Two "street" drugs that mimic the effects of heroin illustrate this principle: 3-methylfentanyl (two N atoms, even molecular weight) and MPPP (one N atom, odd molecular weight).



3-methylfentanyl
$C_{23}H_{30}N_2O$
molecular weight = 350



MPPP
(1-methyl-4-phenyl-4-propionoxypiperidine)
$C_{15}H_{21}NO_2$
molecular weight = 247

Problem 13.2 Propose two molecular formulas for each of the following molecular ions: (a) 72; (b) 100; (c) 73.

Problem 13.3 Use the following information to propose a molecular formula for nootkatone, a compound partly responsible for the characteristic odor of grapefruit. Nootkatone contains the elements C, H, and O, has five degrees of unsaturation, and a molecular ion in its mass spectrum at *m/z* = 218.



Nootkatone (Problem 13.3) occurs naturally in grapefruits, and has been used for many years as a flavoring in foods and beverages.

13.2 Alkyl Halides and the M + 2 Peak

Most of the elements found in organic compounds, such as carbon, hydrogen, oxygen, nitrogen, sulfur, phosphorus, fluorine, and iodine, have one major isotope. **Chlorine** and **bromine,** on the other hand, have two, giving characteristic patterns to the mass spectra of their compounds.

Chlorine has two common isotopes, ^{35}**Cl** and ^{37}**Cl,** which occur naturally in a 3:1 ratio. Thus, **there are two peaks in a 3:1 ratio for the molecular ion of an alkyl chloride.** The larger peak—the **M** peak—corresponds to the compound containing ^{35}Cl, and the smaller peak—the **M + 2** peak—corresponds to the compound containing ^{37}Cl.

- **When the molecular ion consists of two peaks (M and M + 2) in a 3:1 ratio, a Cl atom is present.**

Sample Problem 13.3 What molecular ions will be present in a mass spectrum of 2-chloropropane, $(CH_3)_2CHCl$?

Solution

Calculate the molecular weight using each of the common isotopes of Cl.

Molecular formula	Mass of molecular ion (*m/z*)
$C_3H_7{}^{35}Cl$	78 (M peak)
$C_3H_7{}^{37}Cl$	80 (M + 2 peak)

There should be two peaks in a ratio of 3:1, at *m/z* = 78 and 80, as illustrated in the mass spectrum of 2-chloropropane in Figure 13.3.

Bromine has two common isotopes, ^{79}**Br** and ^{81}**Br,** which occur naturally in a 1:1 ratio. Thus, **there are two peaks in a 1:1 ratio for the molecular ion of an alkyl bromide.** In the mass spectrum of 2-bromopropane (Figure 13.4), for example, there is an M peak at *m/z* = 122 and an M + 2 peak at *m/z* = 124.

- **When the molecular ion consists of two peaks (M and M + 2) in a 1:1 ratio, a Br atom is present in the molecule.**

Figure 13.3
Mass spectrum of 2-chloropropane [$(CH_3)_2CHCl$]






Figure 13.4
Mass spectrum of 2-bromopropane [$(CH_3)_2CHBr$]



Problem 13.4 What molecular ions would you expect for compounds having each of the following molecular formulas: (a) C_4H_9Cl; (b) C_3H_7F; (c) $C_4H_{11}N$; (d) $C_4H_4N_2$?

Problem 13.5 What molecular ions would you expect for the compound depicted in the ball-and-stick model?




13.3 Fragmentation

While many chemists use a mass spectrum to determine only a compound's molecular weight and molecular formula, additional useful structural information can be obtained from fragmentation patterns. Although each organic compound fragments in a unique way, a particular functional group exhibits common fragmentation patterns.

13.3A General Features of Fragmentation

As an example, consider hexane, whose mass spectrum was shown in Figure 13.2. When hexane is bombarded by an electron beam, it forms a highly unstable radical cation (m/z = 86) that can decompose by cleavage of any of the C−C bonds. Thus, cleavage of the terminal C−C bond


forms $CH_3CH_2CH_2CH_2CH_2^+$ and CH_3•. Fragmentation generates a cation and a radical, and cleavage generally yields the more stable, more substituted carbocation.



- **Loss of a CH_3 group always forms a fragment with a mass 15 units less than the molecular ion.**

As a result, the mass spectrum of hexane shows a peak at $m/z = 71$ due to $CH_3CH_2CH_2CH_2CH_2^+$. Figure 13.5 illustrates how cleavage of other C–C bonds in hexane gives rise to other fragments that correspond to peaks in its mass spectrum.

Figure 13.5
Identifying fragments in the mass spectrum of hexane




- Cleavage of C–C bonds (labeled [1]–[4]) in hexane forms lower molecular weight fragments that correspond to lines in the mass spectrum. Although the mass spectrum is complex, possible structures can be assigned to some of the fragments, as shown.

Sample Problem 13.4 The mass spectrum of 2,3-dimethylpentane $[(CH_3)_2CHCH(CH_3)CH_2CH_3]$ shows fragments at m/z = 85 and 71. Propose possible structures for the ions that give rise to these peaks.

Solution

To solve a problem of this sort, first calculate the mass of the molecular ion. Draw out the structure of the compound, break a C–C bond, and calculate the mass of the resulting fragments. Repeat this process on different C–C bonds until fragments of the desired mass-to-charge ratio are formed.




In this example, 2,3-dimethylpentane has a molecular ion at *m/z* = 100. Cleavage of bond [1] forms a 2° carbocation with *m/z* = 85 and CH_3•. Cleavage of bond [2] forms another 2° carbocation with *m/z* = 71 and CH_3CH_2•. Thus, the fragments at *m/z* = 85 and 71 are possibly due to the two carbocations drawn.

Problem 13.6 The mass spectrum of 2,3-dimethylpentane also shows peaks at *m/z* = 57 and 43. Propose possible structures for the ions that give rise to these peaks.


Problem 13.7 The base peak in the mass spectrum of 2,2,4-trimethylpentane [$(CH_3)_3CCH_2CH(CH_3)_2$] occurs at *m/z* = 57. What ion is responsible for this peak and why is this ion the most abundant fragment?

13.3B Fragmentation Patterns of Some Common Functional Groups


Each functional group exhibits characteristic fragmentation patterns that help to analyze a mass spectrum. For example, aldehydes and ketones often undergo the process of **α cleavage, breaking the bond between the carbonyl carbon and the carbon adjacent to it.** Cleavage yields a neutral radical and a resonance-stabilized acylium ion.



Alcohols undergo fragmentation in two different ways—α cleavage and dehydration. Alpha (α) cleavage occurs by breaking a bond between an alkyl group and the carbon that bears the OH group, resulting in an alkyl radical and a resonance-stabilized carbocation.



Likewise, alcohols undergo dehydration, the elimination of H_2O, from two adjacent atoms. Unlike fragmentations discussed thus far, dehydration results in the cleavage of two bonds and forms H_2O and the radical cation derived from an alkene.




- **Loss of H_2O from an alcohol always forms a fragment with a mass 18 units less than the molecular ion.**

Sample Problem 13.5 What mass spectral fragments are formed from α cleavage of pentan-2-one, $CH_3COCH_2CH_2CH_3$?

Solution


Alpha (α) cleavage breaks the bond between the carbonyl carbon and the carbon adjacent to it, yielding a neutral radical and a resonance-stabilized acylium ion. A ketone like pentan-2-one with two different alkyl groups bonded to the carbonyl carbon has two different pathways for α cleavage.



As a result, two fragments are formed by α cleavage of pentan-2-one, giving peaks at m/z = 71 and 43.

Problem 13.8 (a) What mass spectral fragments are formed by α cleavage of butan-2-ol, $CH_3CH(OH)CH_2CH_3$?
(b) What fragments are formed by dehydration of butan-2-ol?

Problem 13.9 What cations are formed in the mass spectrometer by α cleavage of each of the following compounds?



13.4 Other Types of Mass Spectrometry

Recent advances have greatly expanded the information obtained from mass spectrometry.

13.4A High-Resolution Mass Spectrometry

The mass spectra described thus far have been low-resolution spectra; that is, they report m/z values to the nearest whole number. As a result, the mass of a given molecular ion can correspond to many different molecular formulas, as shown in Sample Problem 13.2.

Table 13.1
Exact Masses of Some Common Isotopes

Isotope	Mass
^{12}C	12.0000
^{1}H	1.00783
^{16}O	15.9949
^{14}N	14.0031

High-resolution mass spectrometers measure m/z ratios to four (or more) decimal places. This is valuable because except for carbon-12, whose mass is defined as 12.0000, the masses of all other nuclei are very close to—but not exactly—whole numbers. Table 13.1 lists the exact mass values of a few common nuclei. Using these values it is possible to determine the single molecular formula that gives rise to a molecular ion.


For example, a compound having a molecular ion at m/z = 60 using a low-resolution mass spectrometer could have the following molecular formulas:

Formula	Exact mass
C_3H_8O	60.0575
$C_2H_4O_2$	60.0211
$C_2H_8N_2$	60.0688

If the molecular ion had an exact mass of 60.0578, the compound's molecular formula is C_3H_8O, because its mass is closest to the observed value.

Problem 13.10 The low-resolution mass spectrum of an unknown analgesic **X** had a molecular ion of 151. Possible molecular formulas include $C_7H_5NO_3$, $C_8H_9NO_2$, and $C_{10}H_{17}N$. High-resolution mass spectrometry gave an exact mass of 151.0640. What is the molecular formula of **X?**

13.4B Gas Chromatography–Mass Spectrometry (GC–MS)



Like other controlled substances, the tetrahydrocannabinol from marijuana leaves can be detected in minute amounts by GC–MS.


Two analytical tools—**gas chromatography (GC)** and **mass spectrometry (MS)**—can be combined into a single instrument **(GC–MS)** to analyze mixtures of compounds (Figure 13.6a). The gas chromatograph separates the mixture, and then the mass spectrometer records a spectrum of the individual components.

A gas chromatograph consists of a thin capillary column containing a viscous, high-boiling liquid, all housed in an oven. When a sample is injected into the GC, it is vaporized and swept by an inert gas through the column. The components of the mixture travel through the column at different rates, often separated by boiling point, with lower boiling compounds exiting the column before higher boiling compounds. Each compound then enters the mass spectrometer, where it is ionized to form its molecular ion and lower molecular weight fragments. The GC–MS records a gas chromatogram for the mixture, which plots the amount of each component versus its **retention time**—that is, the time required to travel through the column. Each component of a mixture is characterized by its retention time in the gas chromatogram and its molecular ion in the mass spectrum (Figure 13.6b).


GC–MS is widely used for characterizing mixtures containing environmental pollutants. It is also used to analyze urine and hair samples for the presence of illegal drugs or banned substances thought to improve athletic performance.

Figure 13.6 Compound analysis using GC–MS

a. Schematic of a GC–MS instrument




b. GC trace of a three-component mixture. The mass spectrometer gives a spectrum for each component.




To analyze a urine sample for THC (tetrahydrocannabinol), the principal psychoactive component of marijuana, the organic compounds are extracted from urine, purified, concentrated, and injected into the GC–MS. THC appears as a GC peak with a characteristic retention time (for a given set of experimental parameters), and gives a molecular ion at 314, its molecular weight, as shown in Figure 13.7.

Figure 13.7
Mass spectrum of tetrahydrocannabinol (THC)



Problem 13.11 Benzene, toluene, and *p*-xylene (BTX) are often added to gasoline to boost octane ratings. What would be observed if a mixture of these three compounds were subjected to GC–MS analysis? How many peaks would be present in the gas chromatogram? What would be the relative order of the peaks? What molecular ions would be observed in the mass spectra?



benzene toluene *p*-xylene

13.4C Mass Spectra of High Molecular Weight Biomolecules

Dr. John Fenn shared the 2002 Nobel Prize in Chemistry for his development of ESI mass spectrometry.

Until the 1980s mass spectra were limited to molecules that could be readily vaporized with heat under vacuum, and thus had molecular weights of < 800. In the last 25 years, new methods have been developed to generate gas phase ions of large molecules, allowing mass spectra to be recorded for large biomolecules such as proteins and carbohydrates. **Electrospray ionization (ESI),** for example, forms ions by creating a fine spray of charged droplets in an electric field. Evaporation of the charged droplets forms gaseous ions that are then analyzed by their *m*/*z* ratio. ESI and related techniques have extended mass spectrometry into the analysis of nonvolatile compounds with molecular weights greater than 100,000 daltons (atomic mass units).

13.5 Electromagnetic Radiation

Infrared (IR) spectroscopy and **nuclear magnetic resonance (NMR)** spectroscopy (Chapter 14) both use a form of electromagnetic radiation as their energy source. To understand IR and NMR, therefore, you need to understand some of the properties of **electromagnetic radiation**—radiant energy having dual properties of both waves and particles.

The particles of electromagnetic radiation are called **photons,** each having a discrete amount of energy called a **quantum.** Because electromagnetic radiation also has wave properties, it can be characterized by its **wavelength** and **frequency.**

Length units used to report wavelength include:


Unit	Length
meter (m)	1 m
centimeter (cm)	10^{-2} m
micrometer (μm)	10^{-6} m
nanometer (nm)	10^{-9} m
Angstrom (Å)	10^{-10} m

- **Wavelength (λ) is the distance from one point on a wave (e.g., the peak or trough) to the same point on the adjacent wave. A variety of different length units are used for λ, depending on the type of radiation.**
- **Frequency (ν) is the number of waves passing a point per unit time. Frequency is reported in cycles per second (s^{-1}), which is also called hertz (Hz).**

You come into contact with many different kinds of electromagnetic radiation in your daily life. You use visible light to see the words on this page, you may cook with microwaves, and you should use sunscreen to protect your skin from the harmful effects of ultraviolet radiation.

The different forms of electromagnetic radiation make up the **electromagnetic spectrum.** The spectrum is arbitrarily divided into different regions, as shown in Figure 13.8. All electromagnetic radiation travels at the speed of light (c), 3.0×10^8 m/s.

Figure 13.8
The electromagnetic spectrum



- Visible light occupies only a small region of the electromagnetic spectrum.

The speed of electromagnetic radiation (c) is directly proportional to its wavelength and frequency:

$$c = \lambda\nu$$

The speed of light (c) is a constant, so wavelength and frequency are *inversely* related:

- **$\lambda = c/\nu$: Wavelength increases as frequency decreases.**
- **$\nu = c/\lambda$: Frequency increases as wavelength decreases.**

The energy (E) of a photon is directly proportional to its frequency where h = Planck's constant (6.63×10^{-34} J·s).

$$E = h\nu$$


Frequency and wavelength are *inversely* proportional ($\nu = c/\lambda$), however, so energy and wavelength are *inversely* proportional:

$$E = h\nu = \frac{hc}{\lambda}$$

- **The energy of electromagnetic radiation increases as frequency increases and wavelength decreases.**

When electromagnetic radiation strikes a molecule, some wavelengths—but not all—are absorbed. Only some wavelengths are absorbed because molecules have discrete energy levels. The energies of their electronic, vibrational, and nuclear spin states are *quantized,* not *continuous.*

- **For absorption to occur, the energy of the photon must match the difference between two energy states in a molecule.**



- **The *larger* the energy difference between two states, the *higher* the energy of radiation needed for absorption, the *higher* the frequency, and the *shorter* the wavelength.**

Problem 13.12 Which of the following has the higher frequency: (a) light having a wavelength of 10^2 or 10^4 nm; (b) light having a wavelength of 100 nm or 100 μm; (c) red light or blue light?

Problem 13.13 Which of the following has the higher energy: (a) light having a ν of 10^4 Hz or 10^8 Hz; (b) light having a λ of 10 nm or 1000 nm; (c) red light or blue light?

13.6 Infrared Spectroscopy

Organic chemists use infrared (IR) spectroscopy to identify the functional groups in a compound.

13.6A Background

Using the wavenumber scale results in IR values in a numerical range that is easier to report than the corresponding frequencies given in hertz (4000–400 cm^{-1} compared to 1.2×10^{14}–1.2×10^{15} Hz).

Infrared radiation (λ = 2.5–25 μm) is the energy source in infrared spectroscopy. These are somewhat longer wavelengths than visible light, so they are lower in frequency and lower in energy than visible light. Frequencies in IR spectroscopy are reported using a unit called the **wavenumber** ($\tilde{\nu}$):

$$\tilde{\nu} = \frac{1}{\lambda}$$

Wavenumber is *inversely* proportional to wavelength and reported in reciprocal centimeters **(cm^{-1}).** Wavenumber ($\tilde{\nu}$) is *proportional* to frequency (ν). **Frequency (and therefore energy) increases as the wavenumber increases.** Using the wavenumber scale, IR absorptions occur from **4000 cm^{-1}–400 cm^{-1}.**

- **Absorption of IR light causes changes in the vibrational motions of a molecule.**


Covalent bonds are not static. They are more like springs with weights on each end. When two atoms are bonded to each other, the bond stretches back and forth. When three or more atoms are

joined together, bonds can also bend. These bond stretching and bending vibrations represent the different vibrational modes available to a molecule.



A bond can stretch. Two bonds can bend.

These vibrations are quantized, so they occur only at specific frequencies, which correspond to the frequency of IR light. When the frequency of IR light matches the frequency of a particular vibrational mode, the IR light is absorbed, causing the amplitude of the particular bond stretch or bond bend to increase.




- **Different kinds of bonds vibrate at different frequencies, so they absorb different frequencies of IR light.**
- **IR spectroscopy distinguishes between the different kinds of bonds in a molecule, so it is possible to determine the functional groups present.**

Problem 13.14 Which of the following has higher energy: (a) IR light of 3000 cm^{-1} or 1500 cm^{-1} in wavenumber; (b) IR light having a wavelength of 10 μm or 20 μm?

13.6B Characteristics of an IR Spectrum

In an IR spectrometer, light passes through a sample. Frequencies that match vibrational frequencies are absorbed, and the remaining light is transmitted to a detector. A spectrum plots the amount of transmitted light versus its wavenumber. The IR spectrum of propan-1-ol, $CH_3CH_2CH_2OH$, illustrates several important features of IR spectroscopy.



- An IR spectrum has broad lines.
- The absorption peaks go *down* on a page. The *y* axis measures **percent transmittance:** 100% transmittance means that all the light shone on a sample is transmitted and none is absorbed; 0% transmittance means that none of the light shone on a sample is transmitted and all is absorbed. Most absorptions lie between these two extremes.
- **Each peak corresponds to a particular kind of bond, and each bond type (such as O–H and C–H) occurs at a characteristic frequency.**
- IR spectra have both a wavelength and a wavenumber scale on the *x* axis. Wavelengths are recorded in μm (2.5–25). Wavenumber, frequency, and energy *decrease* from left to right. Where a peak occurs is reported in reciprocal centimeters (cm^{-1}).

Conceptually, the IR spectrum is divided into two regions:

- **The functional group region occurs at ≥ 1500 cm^{-1}. Common functional groups give one or two peaks in this region, at a characteristic frequency.**
- **The fingerprint region occurs at < 1500 cm^{-1}. This region often contains a complex set of peaks and is unique for every compound.**

Compare, for example, the IR spectra of 5-methylhexan-2-one (**A**) and ethyl propanoate (**B**) in Figure 13.9. The IR spectra look similar in their functional group regions because both compounds contain a carbonyl group (C=O) and several sp^3 hybridized C–H bonds. Because **A** and **B** are different compounds, however, their fingerprint regions look very different.




Figure 13.9

Comparing the functional group region and fingerprint region of two compounds

- **A** and **B** show peaks in the same regions for their C=O group and sp^3 hybridized C–H bonds.
- **A** and **B** are different compounds, so their fingerprint regions are quite different.

13.7 IR Absorptions

13.7A Where Particular Bonds Absorb in the IR

Where a particular bond absorbs in the IR depends on **bond strength** and **atom mass.**


- **Bond strength: stronger bonds vibrate at higher frequency, so they absorb at higher $\tilde{\nu}$.**
- **Atom mass: bonds with lighter atoms vibrate at higher frequency, so they absorb at higher $\tilde{\nu}$.**

Thinking of bonds as springs with weights on each end illustrates these trends. The strength of the spring is analogous to bond strength, and the mass of the weights is analogous to atomic mass. For two springs with the same weights on each end, the **stronger spring vibrates at a higher**

frequency. For two springs of the same strength, **springs with lighter weights vibrate at higher frequency** than those with heavier weights. Hooke's law, as shown in Figure 13.10, describes the relationship of frequency to mass and bond strength.

Figure 13.10
Hooke's law: How the frequency of bond vibration depends on atom mass and bond strength

The frequency of bond vibration can be derived from Hooke's law, which describes the motion of a vibrating spring:




- The force constant (f) is the strength of the bond (or spring). The larger the value of f, the stronger the bond, and the higher the $\tilde{\nu}$ of vibration.
- The mass (m) is the mass of atoms (or weights). The smaller the value of m, the higher the $\tilde{\nu}$ of vibration.

As a result, **bonds absorb in four predictable regions in an IR spectrum.** These four regions, and the bonds that absorb there, are summarized in Figure 13.11. Remembering the information in this figure will help you analyze the spectra of unknown compounds. To help you remember it, keep in mind the following two points:

- **Absorptions for bonds to hydrogen always occur on the left side of the spectrum (the high wavenumber region). H has so little mass that H – Z bonds (where Z = C, O, and N) vibrate at high frequencies.**
- **Bond strength decreases in going from C≡C → C=C → C–C, so the frequency of vibration decreases—that is, the absorptions for these bonds move farther to the right side of the spectrum.**

The functional group region consists of absorptions for single bonds to hydrogen (all H–Z bonds), as well as absorptions for all multiple bonds. Most absorptions in the functional group region are due to bond stretching (rather than bond bending). The fingerprint region consists of absorptions due to all other single bonds (except H–Z bonds), often making it a complex region that is very difficult to analyze.

Figure 13.11
Summary: The four regions of the IR spectrum




Besides learning the general regions of the IR spectrum, it is also useful to learn the specific absorption values for common bonds. Table 13.2 lists the most important IR absorptions in the functional group region. Other details of IR absorptions will be presented in later chapters when new functional groups are introduced. Appendix E contains a detailed list of the characteristic IR absorption frequencies for common bonds.

Table 13.2 Important IR Absorptions


Bond type	Approximate $\tilde{\nu}$ (cm^{-1})	Intensity
O–H	3600–3200	strong, broad
N–H	3500–3200	medium
C–H	~3000	
• C_{sp^3}–H	3000–2850	strong
• C_{sp^2}–H	3150–3000	medium
• C_{sp}–H	3300	medium
C≡C	2250	medium
C≡N	2250	medium
C=O	1800–1650 (often ~1700)	strong
C=C	1650	medium
	1600, 1500	medium

Even subtle differences that affect bond strength affect the frequency of an IR absorption. Recall from Section 1.11 that the strength of a C–H bond increases as the percent *s*-character of the hybrid orbital on the carbon increases; thus:



- **The higher the percent *s*-character, the stronger the bond and the higher the wavenumber of absorption.**

Problem 13.15 Which highlighted bond in each pair absorbs at higher wavenumber?



Finally, almost all bonds in a molecule give rise to an absorption peak in an IR spectrum, but a few do not. **For a bond to absorb in the IR, there must be a change in dipole moment during the vibration.** Thus, symmetrical, nonpolar bonds do not absorb in the IR. The carbon–carbon

triple bond of but-2-yne, for example, does not have an IR stretching absorption at 2250 cm^{-1} because the C≡C bond is nonpolar and there is no change in dipole moment when the bond stretches along its axis. This type of vibration is said to be **IR inactive.**

Stretching along the bond axis
does not change the dipole moment.


$CH_3-C\equiv C-CH_3$

nonpolar bond
IR inactive


13.7B IR Absorptions in Hydrocarbons

The IR spectra of hexane, hex-1-ene, and hex-1-yne illustrate the important differences that characterize the IR spectra of hydrocarbons above 1500 cm^{-1}. Although all three compounds contain C–C bonds and sp^3 hybridized C–H bonds, the absorption peaks due to C=C and C≡C readily distinguish the alkene and alkyne.


Note, too, that the C–H absorptions in alkanes, alkenes, and alkynes have a characteristic appearance and position. The sp^3 hybridized C–H bonds are often seen as a broad, strong absorption at < 3000 cm^{-1}, whereas sp^2 and sp hybridized C–H bonds absorb at somewhat higher frequency.



- The **alkane** hexane has only C–C single bonds and sp^3 hybridized C atoms. Therefore, it has only one major absorption above 1500 cm^{-1}, its C_{sp^3}–H absorption at 3000–2850 cm^{-1}.



- The **alkene** hex-1-ene has a C=C and C_{sp^2}–H, in addition to its sp^3 hybridized C atoms. Therefore, there are three major absorptions above 1500 cm^{-1}:
  - C_{sp^2}–H at 3150–3000 cm^{-1}
  - C_{sp^3}–H at 3000–2850 cm^{-1}
  - C=C at 1650 cm^{-1}




- The **alkyne** hex-1-yne has a C≡C and C_{sp}–H, in addition to its sp^3 hybridized C atoms. Therefore, there are three major absorptions:
  - C_{sp}–H at 3300 cm^{-1}
  - C_{sp^3}–H at 3000–2850 cm^{-1}
  - C≡C at ~2250 cm^{-1}


Problem 13.16 How do the IR spectra of the isomers cyclopentane and pent-1-ene differ?

13.7C IR Absorptions in Oxygen-Containing Compounds


The most important IR absorptions for oxygen-containing compounds occur at **3600–3200 cm^{-1} for an OH group,** and at approximately **1700 cm^{-1} for a C=O,** as illustrated in the IR spectra of an **alcohol** (butan-2-ol), a **ketone** (butan-2-one), and an **ether** (diethyl ether). The peak at ~3000 cm^{-1} in each spectrum is due to C_{sp^3}–H bonds.



- The **OH** group in the alcohol butan-2-ol shows a strong absorption at 3600–3200 cm^{-1}.




- The **C=O** group in the ketone butan-2-one shows a strong absorption at ~1700 cm^{-1}.
- The exact location of the C=O absorption depends on the particular type of carbonyl group, whether the carbonyl carbon is part of a ring, and whether there are nearby double bonds. These details are discussed in Chapters 21 and 22.




- Diethyl ether has neither an OH group nor a C=O, so its only absorption above 1500 cm^{-1} occurs at ~3000 cm^{-1}, due to sp^3 hybridized C–H bonds. Compounds that contain an oxygen atom but do not show an OH or C=O absorption are **ethers.**

13.7D IR Absorptions in Nitrogen-Containing Compounds


Common functional groups that contain nitrogen atoms are also distinguishable by their IR absorptions above 1500 cm^{-1}, as illustrated by the IR spectra of an **amine** (octylamine), an **amide** (propanamide), and a **nitrile** (octanenitrile). Additional details on the IR spectra of these compounds are given in Chapters 22 and 25.



- The **N–H** bonds in the amine octylamine give rise to two weak absorptions at 3300 and 3400 cm^{-1}.



- The **amide** propanamide exhibits absorptions above 1500 cm^{-1} for both its **N–H** and **C=O** groups:
  - N–H (two peaks) at 3200 and 3400 cm^{-1}
  - C=O at 1660 cm^{-1}




- The **C≡N** group of the **nitrile** octanenitrile absorbs in the triple bond region at ~2250 cm^{-1}.

Sample Problem 13.6 How can the two isomers having molecular formula C_2H_6O be distinguished by IR spectroscopy?

Solution


First, draw the structures of the compounds and then locate the functional groups. One compound is an alcohol and one is an ether.



hydroxy group

ethanol

- C–H absorption at ~3000 cm^{-1}
- O–H absorption at 3600–3200 cm^{-1}




no OH group

dimethyl ether

- C–H absorption at ~3000 cm^{-1} **only**

Although both compounds have sp^3 hybridized C–H bonds, ethanol has an OH group that gives a strong absorption at 3600–3200 cm^{-1}, and dimethyl ether does not. This feature distinguishes the two isomers.


Problem 13.17 How do the three isomers of molecular formula C_3H_6O (**A, B,** and **C**) differ in their IR spectra?



A **B** **C**

Sample Problem 13.7 shows how the region above 1500 cm^{-1} in an IR spectrum can be used for functional group identification.

Sample Problem 13.7 What functional groups are responsible for the absorptions above 1500 cm^{-1} in compounds **A** and **B?**




Solution

a. Compound **A** has two major absorptions above 1500 cm^{-1}: The absorption at ~3000 cm^{-1} is due to C–H bonds and the absorption at ~1700 cm^{-1} is due to a C=O group.

b. Compound **B** has two major absorptions above 1500 cm^{-1}: The absorption at ~3000 cm^{-1} is due to C–H bonds and the absorption at ~2250 cm^{-1} is due to a triple bond, either a C≡C or a C≡N. Because there is no absorption due to an *sp* hybridized C–H bond at 3300 cm^{-1}, this IR spectrum can*not* be due to a terminal alkyne (HC≡CR) but may still be due to an internal alkyne.


Problem 13.18 What functional groups are responsible for the absorptions above 1500 cm^{-1} in the IR spectra for compounds **A** and **B?**




Problem 13.19 What are the major IR absorptions in the functional group region for each compound?

a.


b.




c.



d.




e.



capsaicin
(spicy component of hot peppers)

Problem 13.20 What are the major IR absorptions in the functional group region for oleic acid, a common unsaturated fatty acid (Section 10.6A)?



oleic acid




The antibiotic properties of penicillin were discovered when Sir Alexander Fleming noticed that a mold of the genus *Penicillium* inhibited the growth of certain bacteria. Penicillin was used as an antibiotic before its structure was determined conclusively.

13.8 IR and Structure Determination

Since its introduction, IR spectroscopy has proven to be a valuable tool for determining the functional groups in organic molecules.


In the 1940s, IR spectroscopy played a key role in elucidating the structure of the antibiotic penicillin G, the chapter-opening molecule. **β-Lactams,** four-membered rings that contain an amide, have a carbonyl group that absorbs at ~1760 cm^{-1}, a much higher frequency than that observed for most amides and many other carbonyl groups. Because penicillin G had an IR absorption at

this frequency, **A** became the leading candidate for the structure of penicillin rather than **B,** a possibility originally considered more likely. Structure **A** was later confirmed by X-ray analysis.



New instruments for determining blood alcohol concentration use IR spectroscopy for analyzing the C–H absorption of CH_3CH_2OH in exhaled air. Figure 12.10 illustrated an earlier method based on oxidation chemistry.


IR spectroscopy is often used to determine the outcome of a chemical reaction. For example, oxidation of the hydroxy group in **C** to form the carbonyl group in periplanone B is accompanied by the disappearance of the OH absorption (3600–3200 cm^{-1}) and the appearance of a carbonyl absorption near 1700 cm^{-1} in the IR spectrum of the product. Periplanone B is the sex pheromone of the female American cockroach.




The combination of IR and mass spectral data provides key information on the structure of an unknown compound. The mass spectrum reveals the molecular weight of the unknown (and the molecular formula if an exact mass is available), and the IR spectrum helps to identify the important functional groups.

How To Use MS and IR for Structure Determination

Example **What information is obtained from the mass spectrum and IR spectrum of an unknown compound X? Assume X contains the elements C, H, and O.**






How To, continued . . .

Step [1] Use the molecular ion to determine possible molecular formulas. Use an exact mass (when available) to determine a molecular formula.

- Use the procedure outlined in Sample Problem 13.2 to calculate possible molecular formulas. For a molecular ion at *m/z* = 88:



- Discounting C_7H_4 (a hydrocarbon) and C_6O (because it contains no H's) gives three possible formulas for **X.**
- If high-resolution mass spectral data are available, the molecular formula can be determined directly. If the molecular ion had an exact mass of 88.0580, the molecular formula of **X** is **$C_4H_8O_2$** (exact mass = 88.0524) rather than $C_5H_{12}O$ (exact mass = 88.0888) or $C_3H_4O_3$ (exact mass = 88.0160).

Step [2] Calculate the number of degrees of unsaturation (Section 10.2).


- For a compound of molecular formula $C_4H_8O_2$, the maximum number of H's = $2n + 2 = 2(4) + 2 = 10$.
- Because the compound contains only 8 H's, it has 10 – 8 = 2 H's fewer than the maximum number.
- Because each degree of unsaturation removes 2 H's, **X** has one degree of unsaturation. **X has one ring or one π bond.**

Step [3] Determine what functional group is present from the IR spectrum.

- The two major absorptions in the IR spectrum above 1500 cm^{-1} are due to sp^3 hybridized C–H bonds (~3000–2850 cm^{-1}) and a C=O group (1740 cm^{-1}). Thus, the one degree of unsaturation in **X** is due to the presence of the **C=O.**

Mass spectrometry and IR spectroscopy give valuable but limited information on the identity of an unknown. Although the mass spectral and IR data reveal that **X** has a molecular formula of $C_4H_8O_2$ and contains a carbonyl group, more data are needed to determine its complete structure. In Chapter 14, we will learn how other spectroscopic data can be used for that purpose.

Problem 13.21 Which of the following possible structures for **X** can be excluded on the basis of its IR spectrum?



Problem 13.22 Propose structures consistent with each set of data: (a) a hydrocarbon with a molecular ion at *m/z* = 68 and IR absorptions at 3310, 3000–2850, and 2120 cm^{-1}; (b) a compound containing C, H, and O with a molecular ion at *m/z* = 60 and IR absorptions at 3600–3200 and 3000–2850 cm^{-1}.

KEY CONCEPTS

Mass Spectrometry and Infrared Spectroscopy

Mass Spectrometry (MS; 13.1–13.4)

- Mass spectrometry measures the molecular weight of a compound (13.1A).
- The mass of the molecular ion (**M**) = the molecular weight of a compound. Except for isotope peaks at M + 1 and M + 2, the molecular ion has the highest mass in a mass spectrum (13.1A).
- The base peak is the tallest peak in a mass spectrum (13.1A).
- A compound with an odd number of N atoms gives an odd molecular ion. A compound with an even number of N atoms (including zero) gives an even molecular ion (13.1B).
- Organic monochlorides show two peaks for the molecular ion (M and M + 2) in a 3:1 ratio (13.2).


- Organic monobromides show two peaks for the molecular ion (M and M + 2) in a 1:1 ratio (13.2).
- The fragmentation of radical cations formed in a mass spectrometer gives lower molecular weight fragments, often characteristic of a functional group (13.3).
- High-resolution mass spectrometry gives the molecular formula of a compound (13.4A).

Electromagnetic Radiation (13.5)


- The wavelength and frequency of electromagnetic radiation are *inversely* related by the following equations: $\lambda = c/\nu$ or $\nu = c/\lambda$ (13.5).
- The energy of a photon is proportional to its frequency; the higher the frequency, the higher the energy: $E = h\nu$ (13.5).

Infrared Spectroscopy (IR; 13.6 and 13.7)

- Infrared spectroscopy identifies functional groups.
- IR absorptions are reported in wavenumbers, $\tilde{\nu} = 1/\lambda$.
- The functional group region from **4000–1500 cm^{-1}** is the most useful region of an IR spectrum.
- C−H, O−H, and N−H bonds absorb at high frequency, ≥ 2500 cm^{-1}.
- As bond strength increases, the $\tilde{\nu}$ of absorption increases; thus, triple bonds absorb at higher $\tilde{\nu}$ than double bonds.




- The higher the percent *s*-character, the stronger the bond, and the higher the $\tilde{\nu}$ of an IR absorption.




PROBLEMS

Problems Using Three-Dimensional Models

13.23 What major IR absorptions are present above 1500 cm^{-1} for each compound?

a. 



b. 


13.24 The mass spectrum of the following compound shows fragments at *m/z* = 127, 113, and 85. Propose structures for the ions that give rise to these peaks.




Mass Spectrometry

13.25 What molecular ion is expected for each compound?

a. 
b. 
c. 

d. 

e. 

13.26 Which compound gives a molecular ion at *m/z* = 122: $C_6H_5CH_2CH_2CH_3$, $C_6H_5COCH_2CH_3$, or $C_6H_5OCH_2CH_3$?


13.27 Propose two molecular formulas for each molecular ion: (a) 102; (b) 98; (c) 119; (d) 74.


13.28 Propose four possible structures for a hydrocarbon with a molecular ion at *m/z* = 112.


13.29 What is the molecular formula for α-himachalene, a hydrocarbon obtained from cedar wood, which has four degrees of unsaturation and a molecular ion in its mass spectrum at *m/z* = 204?


13.30 Propose a molecular formula for rose oxide, a rose-scented compound isolated from roses and geraniums, which contains the elements of C, H, and O, has two degrees of unsaturation, and a molecular ion in its mass spectrum at *m/z* = 154.

13.31 Match each structure to its mass spectrum.












13.32 Propose two possible structures for a hydrocarbon having an exact mass of 96.0939 that forms ethylcyclopentane upon hydrogenation with H_2 and Pd-C.

13.33 What cations are formed in the mass spectrometer by α cleavage of each of the following compounds?

a. 

b. 

c. 

13.34 2,3-Dimethylbutane and 2,2-dimethylbutane have the same molecular ion in the mass spectrum, but only one of these isomers gives a significant fragment at *m/z* = 57. (a) Which isomer shows an intense peak at *m/z* = 57? (b) Propose a structure for the ion that gives rise to this peak. (c) The base peak in the mass spectrum of the other isomer occurs at *m/z* = 43. What ion gives rise to this peak?

13.35 For each compound, assign likely structures to the fragments at each *m/z* value, and explain how each fragment is formed.

a. $C_6H_5CH_2CH_2OH$: peaks at *m/z* = 104, 91

b. $CH_2=C(CH_3)CH_2CH_2OH$: peaks at *m/z* = 71, 68, 41, 31

13.36 Suppose you have two bottles, labeled ketone **A** and ketone **B.** You know that one bottle contains $CH_3CO(CH_2)_5CH_3$ and one contains $CH_3CH_2CO(CH_2)_4CH_3$, but you do not know which ketone is in which bottle. Ketone **A** gives a fragment at *m/z* = 99 and ketone **B** gives a fragment at *m/z* = 113. What are the likely structures of ketones **A** and **B** from these fragmentation data?

13.37 Propose a structure consistent with each set of data.

a. a compound that contains a benzene ring and has a molecular ion at *m/z* = 107

b. a hydrocarbon that contains only sp^3 hybridized carbons and a molecular ion at *m/z* = 84

c. a compound that contains a carbonyl group and gives a molecular ion at *m/z* = 114

d. a compound that contains C, H, N, and O and has an exact mass for the molecular ion at 101.0841

13.38 A low-resolution mass spectrum of the neurotransmitter dopamine gave a molecular ion at *m/z* = 153. Two possible molecular formulas for this molecular ion are $C_8H_{11}NO_2$ and $C_7H_{11}N_3O$. A high-resolution mass spectrum provided an exact mass at 153.0680. Which of the possible molecular formulas is the correct one?

13.39 Primary (1°) alcohols often show a peak in their mass spectra at *m/z* = 31. Suggest a structure for this fragment.

13.40 Like alcohols, ethers undergo α cleavage by breaking a carbon–carbon bond between an alkyl group and the carbon bonded to the ether oxygen atom; that is, the red C−C bond in $R{-}CH_2OR'$ is broken. With this in mind, propose structures for the fragments formed by α cleavage of $(CH_3)_2CHCH_2OCH_2CH_3$. Suggest a reason why an ether fragments by α cleavage.

Infrared Spectroscopy


13.41 Which of the highlighted bonds absorbs at higher $\tilde{\nu}$ in an IR spectrum?


a. 

or 


b. 


or 


c. 


or 

13.42 What major IR absorptions are present above 1500 cm^{-1} for each compound?


a. 



b. 



c. 



d. 

13.43 How would each of the following pairs of compounds differ in their IR spectra?


a.  and


c.  and 


b.  and 

d.  and 


13.44 Morphine, heroin, and oxycodone are three addicting analgesic narcotics. How could IR spectroscopy be used to distinguish these three compounds from each other?



morphine



heroin



oxycodone

13.45 Reduction of cyclohex-2-enone can yield cyclohexanone, cyclohex-2-enol, or cyclohexanol, depending on the reagent and reaction conditions. How could you use IR spectroscopy to distinguish the three possible products?



cyclohex-2-enone



cyclohexanone



cyclohex-2-enol



cyclohexanol

13.46 Tell how IR spectroscopy could be used to determine when each reaction is complete.


a. 

b. 


c. 

d. 


13.47 Match each compound to its IR spectrum.


A **B** **C** **D** **E** **F**


Spectrum [1]




Spectrum [2]




Spectrum [3]




Spectrum [4]



Spectrum [5]



Spectrum [6]



Combined Spectroscopy Problems

13.48 Propose possible structures consistent with each set of data. Assume each compound has an sp^3 hybridized C – H absorption in its IR spectrum, and that other major IR absorptions above 1500 cm^{-1} are listed.


a. a compound having a molecular ion at 72 and an absorption in its IR spectrum at 1725 cm^{-1}

b. a compound having a molecular ion at 55 and an absorption in its IR spectrum at ~2250 cm^{-1}

c. a compound having a molecular ion of 74 and an absorption in its IR spectrum at 3600–3200 cm^{-1}

13.49 A chiral hydrocarbon **X** exhibits a molecular ion at 82 in its mass spectrum. The IR spectrum of **X** shows peaks at 3300, 3000–2850, and 2250 cm^{-1}. Propose a structure for **X.**

13.50 A chiral compound **Y** has a strong absorption at 2970–2840 cm^{-1} in its IR spectrum and gives the following mass spectrum. Propose a structure for **Y.**



13.51 Treatment of benzoic acid ($C_6H_5CO_2H$) with NaOH followed by 1-iodo-3-methylbutane forms **H. H** has a molecular ion at 192 and IR absorptions at 3064, 3035, 2960–2872, and 1721 cm^{-1}. Propose a structure for **H.**

13.52 Treatment of benzaldehyde (C_6H_5CHO) with Zn(Hg) in aqueous HCl forms a compound **Z** that has a molecular ion at 92 in its mass spectrum. **Z** shows absorptions at 3150–2950, 1605, and 1496 cm^{-1} in its IR spectrum. Give a possible structure for **Z.**

13.53 Reaction of *tert*-butyl pentyl ether [$CH_3CH_2CH_2CH_2CH_2OC(CH_3)_3$] with HBr forms 1-bromopentane ($CH_3CH_2CH_2CH_2CH_2Br$) and compound **B. B** has a molecular ion in its mass spectrum at 56 and gives peaks in its IR spectrum at 3150–3000, 3000–2850, and 1650 cm^{-1}. Propose a structure for **B** and draw a stepwise mechanism that accounts for its formation.


13.54 Reaction of 2-methylpropanoic acid [$(CH_3)_2CHCO_2H$] with $SOCl_2$ followed by 2-methylpropan-1-ol forms **X. X** has a molecular ion at 144 and IR absorptions at 2965, 2940, and 1739 cm^{-1}. Propose a structure for **X.**


13.55 Reaction of pentanoyl chloride ($CH_3CH_2CH_2CH_2COCl$) with lithium dimethyl cuprate [$LiCu(CH_3)_2$] forms a compound **J** that has a molecular ion in its mass spectrum at 100, as well as fragments at *m/z* = 85, 57, and 43 (base). The IR spectrum of **J** has strong peaks at 2962 and 1718 cm^{-1}. Propose a structure for **J.**

13.56 Benzonitrile (C_6H_5CN) is reduced to two different products depending on the reducing agent used. Treatment with lithium aluminum hydride followed by water forms **K,** which has a molecular ion in its mass spectrum at 107 and the following IR absorptions: 3373, 3290, 3062, 2920, and 1600 cm^{-1}. Treatment with a milder reducing agent forms **L,** which has a molecular ion in its mass spectrum at 106 and the following IR absorptions: 3086, 2820, 2736, 1703, and 1600 cm^{-1}. **L** shows fragments in its mass spectrum at *m/z* = 105 and 77. Propose structures for **K** and **L** and explain how you arrived at your conclusions.

13.57 Treatment of anisole ($CH_3OC_6H_5$) with Cl_2 and $FeCl_3$ forms **P,** which has peaks in its mass spectrum at *m/z* = 142 (M), 144 (M + 2), 129, and 127. **P** has absorptions in its IR spectrum at 3096–2837 (several peaks), 1582, and 1494 cm^{-1}. Propose possible structures for **P.**

13.58 Reaction of $BrCH_2CH_2CH_2CH_2NH_2$ with NaH forms compound **W,** which gives the IR and mass spectra shown below. Propose a structure for **W** and draw a stepwise mechanism that accounts for its formation.







Challenge Problems

13.59 The carbonyl absorption of an amide typically occurs in the 1630–1680 cm^{-1} range, while the carbonyl absorption of an ester occurs at much higher wavenumber, typically 1735–1745 cm^{-1}. Account for this difference.

13.60 Explain why a carbonyl absorption shifts to lower frequency in an α,β-unsaturated carbonyl compound—a compound having a carbonyl group bonded directly to a carbon–carbon double bond. For example, the carbonyl absorption occurs at 1720 cm^{-1} for cyclohexanone, and at 1685 cm^{-1} for cyclohex-2-enone.




cyclohexanone



cyclohex-2-enone
(an α,β-unsaturated
carbonyl compound)

13.61 Explain why a ketone carbonyl typically absorbs at a lower wavenumber than an aldehyde carbonyl (1715 vs. 1730 cm^{-1}).

13.62 Oxidation of citronellol, a constituent of rose and geranium oils, with PCC in the presence of added $NaOCOCH_3$ forms compound **A. A** has a molecular ion in its mass spectrum at 154 and a strong peak in its IR spectrum at 1730 cm^{-1}, in addition to C–H stretching absorptions. Without added $NaOCOCH_3$, oxidation of citronellol with PCC yields isopulegone, which is then converted to **B** with aqueous base. **B** has a molecular ion at 152, and a peak in its IR spectrum at 1680 cm^{-1} in addition to C–H stretching absorptions.




citronellol isopulegone

a. Identify the structures of **A** and **B.**
b. Draw a mechanism for the conversion of citronellol to isopulegone.
c. Draw a mechanism for the conversion of isopulegone to **B.**

Nuclear Magnetic Resonance Spectroscopy

14



Palau'amine is a complex natural product isolated from the sea sponge *Stylotella agminata* collected in the Pacific Ocean near the Republic of Palau. The initial structure proposed for palau'amine in 1993 was revised in 2007 using a variety of modern spectroscopic techniques, including nuclear magnetic resonance spectroscopy. The dense array of functional groups in palau'amine and its antitumor and immunosuppressive properties attracted the attention of dozens of organic chemists, leading to its total synthesis in the laboratory in early 2010. In Chapter 14, we learn how nuclear magnetic resonance spectroscopy plays a key role in structure determination.

14.1 An introduction to NMR spectroscopy
14.2 ^{1}H NMR: Number of signals
14.3 ^{1}H NMR: Position of signals
14.4 The chemical shift of protons on sp^2 and sp hybridized carbons
14.5 ^{1}H NMR: Intensity of signals
14.6 ^{1}H NMR: Spin–spin splitting
14.7 More complex examples of splitting
14.8 Spin–spin splitting in alkenes
14.9 Other facts about ^{1}H NMR spectroscopy
14.10 Using ^{1}H NMR to identify an unknown
14.11 ^{13}C NMR spectroscopy
14.12 Magnetic resonance imaging (MRI)

In Chapter 14 we continue our study of organic structure determination by learning about **nuclear magnetic resonance (NMR)** spectroscopy. NMR spectroscopy is the most powerful tool for characterizing organic molecules, because it can be used to identify the carbon–hydrogen framework in a compound.

14.1 An Introduction to NMR Spectroscopy

Two common types of NMR spectroscopy are used to characterize organic structure:

- **^{1}H NMR (proton NMR)** is used to determine the number and type of hydrogen atoms in a molecule; and
- **^{13}C NMR (carbon NMR)** is used to determine the type of carbon atoms in a molecule.

Before you can learn how to use NMR spectroscopy to determine the structure of a compound, you need to understand a bit about the physics behind it. Keep in mind, though, that NMR stems from the same basic principle as all other forms of spectroscopy. Energy interacts with a molecule, and absorptions occur only when the incident energy matches the energy difference between two states.

14.1A The Basis of NMR Spectroscopy

The source of energy in NMR is radio waves. Radiation in the radiofrequency region of the electromagnetic spectrum (so-called **RF** radiation) has very long wavelengths, so its corresponding frequency and energy are both low. **When these low-energy radio waves interact with a molecule, they can change the nuclear spins of some elements, including ^{1}H and ^{13}C.**



A spinning proton creates a magnetic field.

When a charged particle such as a proton spins on its axis, it creates a magnetic field. For the purpose of this discussion, therefore, a nucleus is a tiny bar magnet, symbolized by . Normally these nuclear magnets are randomly oriented in space, but in the presence of an external magnetic field, B_0, they are oriented with or against this applied field. More nuclei are oriented *with* the applied field because this arrangement is lower in energy, but the **energy difference between these two states is very small** (< 0.4 J/mol).




In a magnetic field, there are now two different energy states for a proton:

- **In the *lower energy state* the nucleus is aligned in the *same direction* as B_0.**
- **In the *higher energy state* the nucleus is aligned *opposed* to B_0.**

When an external energy source ($h\nu$) that matches the energy difference (ΔE) between these two states is applied, energy is absorbed, causing the **nucleus to "spin flip" from one orientation to another.** The energy difference between these two nuclear spin states corresponds to the low-frequency radiation in the RF region of the electromagnetic spectrum.

Absorbing RF radiation causes the nucleus to spin flip.



- **A nucleus is in *resonance* when it absorbs RF radiation and "spin flips" to a higher energy state.**

Thus, two variables characterize NMR:

- **An applied magnetic field, B_0.** Magnetic field strength is measured in tesla (T).
- **The frequency ν of radiation used for resonance,** measured in hertz (Hz) or megahertz (MHz); (1 MHz = 10^6 Hz).

The frequency needed for resonance and the applied magnetic field strength are proportionally related:

$$\nu \propto B_0$$

frequency — applied magnetic field strength


- **The stronger the magnetic field, the larger the energy difference between the two nuclear spin states, and the higher the ν needed for resonance.**

NMR spectrometers are referred to as 300 MHz instruments, 500 MHz instruments, and so forth, depending on the frequency of RF radiation used for resonance.

Early NMR spectrometers used a magnetic field strength of ~1.4 T, which required RF radiation of 60 MHz for resonance. Modern NMR spectrometers use stronger magnets, thus requiring higher frequencies of RF radiation for resonance. For example, a magnetic field strength of 7.05 T requires a frequency of 300 MHz for a proton to be in resonance. These spectrometers use very powerful magnetic fields to create a small, but measurable energy difference between the two possible spin states. A schematic of an NMR spectrometer is shown in Figure 14.1.

If all protons absorbed at the same frequency in a given magnetic field, the spectra of all compounds would consist of a single absorption, rendering NMR useless for structure determination. Fortunately, however, this is not the case.

Figure 14.1 Schematic of an NMR spectrometer



An NMR spectrometer. The sample is dissolved in a solvent, usually $CDCl_3$ (deuterochloroform), and placed in a magnetic field. A radiofrequency generator then irradiates the sample with a short pulse of radiation, causing resonance. When the nuclei fall back to their lower energy state, the detector measures the energy released, and a spectrum is recorded. The superconducting magnets in modern NMR spectrometers have coils that are cooled in liquid helium and conduct electricity with essentially no resistance.


- All protons do *not* absorb at the same frequency. Protons in different environments absorb at slightly different frequencies, and so they are distinguishable by NMR.

The frequency at which a particular proton absorbs is determined by its electronic environment, as discussed in Section 14.3. Because electrons are moving charged particles, they create a magnetic field opposed to the applied field B_0, and the size of the magnetic field generated by the electrons around a proton determines where it absorbs. Modern NMR spectrometers use a constant magnetic field strength B_0, and then a narrow range of frequencies is applied to achieve the resonance of all protons.

Only nuclei that contain odd mass numbers (such as ^{1}H, ^{13}C, ^{19}F, and ^{31}P) or odd atomic numbers (such as ^{2}H and ^{14}N) give rise to NMR signals. Because both ^{1}H and ^{13}C, the less abundant isotope of carbon, are NMR active, NMR allows us to map the carbon and hydrogen framework of an organic molecule.

14.1B A ^{1}H NMR Spectrum

An NMR spectrum plots the **intensity of a signal** against its **chemical shift** measured in **parts per million (ppm).** The common scale of chemical shifts is called the **δ (delta) scale.** The proton NMR spectrum of *tert*-butyl methyl ether [$CH_3OC(CH_3)_3$] illustrates several important features:



tert-Butyl methyl ether (MTBE) is the high-octane gasoline additive that has contaminated the water supply in some areas (Section 3.4).

- NMR absorptions generally appear as sharp signals. The ^{1}H NMR spectrum of $CH_3OC(CH_3)_3$ consists of two signals: a tall peak at 1.2 ppm due to the $(CH_3)_3C-$ group, and a smaller peak at 3.2 ppm due to the CH_3O- group.
- **Increasing chemical shift is plotted from *right to left.*** Most protons absorb somewhere from 0–12 ppm.
- The terms **upfield** and **downfield** describe the relative location of signals. Upfield means to the *right.* The $(CH_3)_3C-$ peak is upfield from the CH_3O- peak. Downfield means to the *left.* The CH_3O- peak is downfield from the $(CH_3)_3C-$ peak.

$(CH_3)_4Si$
tetramethylsilane
TMS

NMR absorptions are measured relative to the position of a reference signal at 0 ppm on the δ scale due to **tetramethylsilane (TMS). TMS** is a volatile and inert compound that gives a single peak upfield from other typical NMR absorptions.

Although chemical shifts are measured relative to the TMS signal at 0 ppm, this reference is often not plotted on a spectrum.

The *positive* direction of the δ scale is *downfield* from TMS. A very small number of absorptions occur upfield from the TMS signal, which is defined as the negative direction of the δ scale. (See Problem 14.74.)

The **chemical shift** on the *x* axis gives the position of an NMR signal, measured in ppm, according to the following equation:

$$\text{chemical shift (in ppm on the } \delta \text{ scale)} = \frac{\text{observed chemical shift (in Hz) downfield from TMS}}{\nu \text{ of the NMR spectrometer (in MHz)}}$$

A chemical shift gives absorptions as a fraction of the NMR operating frequency, making it independent of the spectrometer used to record a spectrum. Because the frequency of the radiation required for resonance is proportional to the strength of the applied magnetic field, B_0, reporting NMR absorptions in frequency is meaningless unless the value of B_0 is also reported. By reporting the absorption as a fraction of the NMR operating frequency, though, we get units—ppm—that are independent of the spectrometer.

Sample Problem 14.1 Calculate the chemical shift of an absorption that occurs at 1500 Hz downfield from TMS using a 300 MHz NMR spectrometer.

Solution

Use the equation that defines the chemical shift in ppm:

$$\text{chemical shift} = \frac{\text{1500 Hz downfield from TMS}}{\text{300 MHz operating frequency}} = 5 \text{ ppm}$$

Problem 14.1 The ^{1}H NMR spectrum of CH_3OH recorded on a 500 MHz NMR spectrometer consists of two signals, one due to the CH_3 protons at 1715 Hz and one due to the OH proton at 1830 Hz, both measured downfield from TMS. (a) Calculate the chemical shift of each absorption. (b) Do the CH_3 protons absorb upfield or downfield from the OH proton?

Problem 14.2 The ^{1}H NMR spectrum of 1,2-dimethoxyethane ($CH_3OCH_2CH_2OCH_3$) recorded on a 300 MHz NMR spectrometer consists of signals at 1017 Hz and 1065 Hz downfield from TMS. (a) Calculate the chemical shift of each absorption. (b) At what frequency would each absorption occur if the spectrum were recorded on a 500 MHz NMR spectrometer?

Four different features of a ^{1}H NMR spectrum provide information about a compound's structure:

[1] Number of signals (Section 14.2)
[2] Position of signals (Sections 14.3 and 14.4)
[3] Intensity of signals (Section 14.5)
[4] Spin–spin splitting of signals (Sections 14.6–14.8)

14.2 ^{1}H NMR: Number of Signals

How many ^{1}H NMR signals does a compound exhibit? The number of NMR signals *equals* the number of different types of protons in a compound.


14.2A General Principles

- **Protons in different environments give different NMR signals. Equivalent protons give the same NMR signal.**

In many compounds, deciding whether two protons are in identical or different environments is intuitive.

Any CH_3 group is different from any CH_2 group, which is different from any CH group in a molecule. Two CH_3 groups may be identical (as in CH_3OCH_3) or different (as in $CH_3OCH_2CH_3$), depending on what each CH_3 group is bonded to.

tert-Butyl methyl ether [$CH_3OC(CH_3)_3$] (Section 14.1) exhibits two NMR signals because it contains two different kinds of protons: one CH_3 group is bonded to $-OC(CH_3)_3$, whereas the other three CH_3 groups are each bonded to the same group, $[-C(CH_3)_2]OCH_3$.



- **CH_3OCH_3:** Each CH_3 group is bonded to the same group ($-OCH_3$), making both CH_3 groups equivalent.
- **CH_3CH_2Cl:** The protons of the CH_3 group are different from those of the CH_2 group.
- **$CH_3OCH_2CH_3$:** The protons of the CH_2 group are different from those in each CH_3 group. The two CH_3 groups are also different from each other; one CH_3 group is bonded to $-OCH_2CH_3$ and the other is bonded to $-CH_2OCH_3$.


In some cases, it is less obvious by inspection if two protons are equivalent or different. To rigorously determine whether two protons are in identical environments (and therefore give rise to one NMR signal), replace each H atom in question by another atom Z (for example, Z = Cl). **If substitution by Z yields the same compound or enantiomers, the two protons are equivalent,** as shown in Sample Problem 14.2.

Sample Problem 14.2


How many different kinds of H atoms does $CH_3CH_2CH_2CH_2CH_3$ contain?

Solution

In comparing two H atoms, replace each H by Z (for example, Z = Cl), and examine the substitution products that result. The two CH_3 groups are identical because substitution of one H by Cl on each carbon gives the same product, 1-chloropentane.



There are two different types of CH_2 groups. Substitution of Cl for H on C2 or C4 gives the same product, 2-chloropentane, so these H's are identical. Substitution of Cl for H on C3 gives a different product, 3-chloropentane, so this CH_2 group is different from the other two CH_2 groups.



Thus, $CH_3CH_2CH_2CH_2CH_3$ has three different types of protons and gives three different NMR signals.






Figure 14.2 gives the number of NMR signals exhibited by four additional molecules. All protons—not just protons bonded to carbon atoms—give rise to NMR signals. Ethanol (CH_3CH_2OH), for example, gives three NMR signals, one of which is due to its OH proton.

Figure 14.2
The number of ^1H NMR signals of some representative organic compounds




Problem 14.3 How many ^{1}H NMR signals does each compound show?




14.2B Determining Equivalent Protons in Alkenes and Cycloalkanes

To determine equivalent protons in cycloalkanes and alkenes that have restricted bond rotation, always draw in all bonds to hydrogen.




Then, in comparing two H atoms on a ring or double bond, **two protons are equivalent only if they are cis (or trans) to the same groups,** as illustrated with 1,1-dichloroethylene, 1-bromo-1-chloroethylene, and chloroethylene.




- **1,1-Dichloroethylene:** The two H atoms on the C=C are both cis to a Cl atom. Thus, both H atoms are equivalent.
- **1-Bromo-1-chloroethylene:** H_a is cis to a Cl atom and H_b is cis to a Br atom. Thus, H_a and H_b are different, giving rise to two NMR signals.
- **Chloroethylene:** H_a is bonded to the carbon with the Cl atom, making it different from H_b and H_c. Of the remaining two H atoms, H_b is cis to a Cl atom and H_c is cis to a H atom, making them different. All three H atoms in this compound are different.

Proton equivalency in cycloalkanes can be determined similarly.




- **Cyclopropane:** All H atoms are equivalent, so there is only one NMR signal.
- **Chlorocyclopropane:** There are now three kinds of H atoms: H_a is bonded to a carbon bonded to a Cl; both H_b protons are cis to the Cl whereas both H_c protons are cis to another H.


Problem 14.4 How many ^{1}H NMR signals does each dimethylcyclopropane show?




Problem 14.5 How many ^{1}H NMR signals does each compound give?

a. 

b. 


c. 

d. 

14.2C Homotopic, Enantiotopic, and Diastereotopic Protons

Let's look more closely at the protons of a single sp^3 hybridized CH_2 group to determine whether these two protons are always equivalent to *each other.* Three examples illustrate different outcomes.


$CH_3CH_2CH_3$ has two different types of protons—those of the CH_3 groups and those of the CH_2 group—meaning that the two H atoms of the CH_2 group are *equivalent to each other*. Replacement of each H by Z forms the same product, so they give one NMR signal.



H_a and H_b are homotopic. identical products

- **When substitution of two H atoms by Z forms the *same* product, these equivalent hydrogens are called *homotopic* protons.**


CH_3CH_2Br has two different types of protons—those of the CH_3 group and those of the CH_2 group—meaning that the two H atoms of the CH_2 group are *equivalent to each other.* Replacement of each H of the CH_2 group by an atom Z creates a new stereogenic center, forming two products that are **enantiomers.**



H_a and H_b are enantiotopic. enantiomers

- **When substitution of two H atoms by Z forms *enantiomers*, the two H atoms are equivalent and give a single NMR signal. These two H atoms are called *enantiotopic* protons.**

In contrast, the two H atoms of the CH_2 group in (*R*)-2-chlorobutane, which contains one stereogenic center, are *not* equivalent to each other. Substitution of each H by Z forms two **diastereomers,** and thus, these two H atoms give *different* NMR signals.






(*R*)-2-chlorobutane

H_a and H_b are diastereotopic. diastereomers

- **When substitution of two H atoms by Z forms *diastereomers*, the two H atoms are *not* equivalent, and give two NMR signals. These two H atoms are called *diastereotopic* protons.**


Sample Problem 14.3 Classify the protons in each labeled CH_2 group as homotopic, enantiotopic, or diastereotopic.

a.  b.  c. 


Solution

To determine equivalency in these cases, look for whether the compound has a stereogenic center to begin with and whether a new stereogenic center is formed when H is replaced by Z.


a. The compound is achiral and has no stereogenic center. Replacement of each H on the labeled CH_2 group by Z forms the same product, making them **homotopic.** The H's within the CH_2 group are *equivalent* to each other and give *one* NMR signal.





b. The compound is achiral and has no stereogenic center. Because a new stereogenic center is formed on substitution of H by Z, the protons of the CH_2 group are **enantiotopic.** These H's are *equivalent* to each other and give *one* NMR signal.




c. The compound has one stereogenic center to begin with. Because a new stereogenic center is formed on substitution of H by Z, the protons are **diastereotopic.** The H's within the CH_2 group are *different* from each other and give *different* NMR signals.





Problem 14.6 Label the protons in each highlighted CH_2 group as enantiotopic, diastereotopic, or homotopic.

a.  b.  c. 

Problem 14.7 How many ^{1}H NMR signals would you expect for each compound?

a. 

b. 

c. 


14.3 ^{1}H NMR: Position of Signals

In the NMR spectrum of *tert*-butyl methyl ether in Section 14.1B, why does the CH_3O- group absorb downfield from the $-C(CH_3)_3$ group?

- **Where a particular proton absorbs depends on its electronic environment.**

14.3A Shielding and Deshielding Effects

To understand how the electronic environment around a nucleus affects its chemical shift, recall that in a magnetic field, an electron creates a small magnetic field that opposes the applied magnetic field, B_0. **Electrons are said to *shield* the nucleus from B_0.**




In the vicinity of the nucleus, therefore, the magnetic field generated by the circulating electron *decreases* the external magnetic field that the proton "feels." Because the proton experiences a lower magnetic field strength, it needs a lower frequency to achieve resonance. Lower frequency is to the right in an NMR spectrum, toward lower chemical shift, so **shielding shifts an absorption *upfield,*** as shown in Figure 14.3a.

What happens if the electron density around a nucleus is *decreased,* instead? For example, how do the chemical shifts of the protons in CH_4 and CH_3Cl compare?

Figure 14.3 How chemical shift is affected by electron density around a nucleus

a. **Shielding effects**
- An electron shields the nucleus.
- The absorption shifts *upfield.*

b. **Deshielding effects**
- Decreased electron density deshields a nucleus.
- The absorption shifts *downfield.*




The less shielded the nucleus becomes, the more of the applied magnetic field (B_0) it feels. This *deshielded* nucleus experiences a higher magnetic field strength, so it needs a higher frequency to achieve resonance. Higher frequency is to the *left* in an NMR spectrum, toward higher chemical shift, so **deshielding shifts an absorption downfield,** as shown in Figure 14.3b for CH_3Cl versus CH_4. The electronegative Cl atom withdraws electron density from the carbon and hydrogen atoms in CH_3Cl, thus deshielding them relative to those in CH_4.

Remember the trend:
***D*ecreased electron density *d*eshields a nucleus and an absorption moves *d*ownfield.**


- **Protons near electronegative atoms are deshielded, so they absorb downfield.**

Figure 14.4 summarizes the effects of shielding and deshielding.


These electron density arguments explain the relative position of NMR signals in many compounds.



- The H_b protons are **deshielded** because they are closer to the electronegative Cl atom, so they absorb **downfield** from H_a.




- Because F is more electronegative than Br, the H_b protons are more **deshielded** than the H_a protons and absorb farther **downfield.**



- The larger number of electronegative Cl atoms (two vs. one) **deshields** H_b more than H_a, so it absorbs **downfield** from H_a.

Figure 14.4
Shielding and deshielding effects



- As the electron density around the nucleus increases, the nucleus feels a smaller resultant magnetic field, so a lower frequency is needed to achieve resonance.
- **The absorption shifts upfield.**

- As the electron density around the nucleus decreases, the nucleus feels a larger resultant magnetic field, so a higher frequency is needed to achieve resonance.
- **The absorption shifts downfield.**

Sample Problem 14.4 Which of the labeled protons in each pair absorbs farther downfield: (a) $CH_3CH_2CH_3$ or CH_3OCH_3; (b) CH_3OCH_3 or CH_3SCH_3?


Solution

a. The CH_3 group in CH_3OCH_3 is deshielded by the electronegative O atom. **Deshielding shifts the absorption downfield.**

b. Because oxygen is more electronegative than sulfur, the CH_3 group in CH_3OCH_3 is more **deshielded** and absorbs **downfield.**

Problem 14.8 For each compound, which of the protons on the highlighted carbons absorbs farther downfield?

a. 

b. 

c. 

14.3B Chemical Shift Values

Not only is the *relative* position of NMR absorptions predictable, but it is also possible to predict the approximate chemical shift value for a given type of proton.








- **Protons in a given environment absorb in a predictable region in an NMR spectrum.**

A more detailed list of characteristic chemical shift values is found in Appendix F.


Table 14.1 lists the typical chemical shift values for the most common bonds encountered in organic molecules.

Table 14.1 also illustrates that absorptions for a given type of C–H bond occur in a narrow range of chemical shift values, usually 1–2 ppm. For example, all sp^3 hybridized C–H bonds in alkanes and cycloalkanes absorb between 0.9 and 2.0 ppm. By contrast, absorptions due to N–H and O–H protons can occur over a broader range. For example, the OH proton of an alcohol is found anywhere in the 1–5 ppm range. The position of these absorptions is affected by the extent of hydrogen bonding, making it more variable.

Table 14.1 Characteristic Chemical Shifts of Common Types of Protons


Type of proton	Chemical shift (ppm)	Type of proton	Chemical shift (ppm)
	0.9–2		4.5–6
• RCH_3	~0.9		
• R_2CH_2	~1.3		6.5–8
• R_3CH	~1.7		
 Z = C, O, N	1.5–2.5		9–10
	~2.5		10–12
 Z = N, O, X	2.5–4		1–5


The chemical shift of a particular type of C–H bond is also affected by the number of R groups bonded to the carbon atom.




- **The chemical shift of a C–H bond increases with increasing alkyl substitution.**

Problem 14.9 For each compound, first label each different type of proton and then rank the protons in order of increasing chemical shift.

a. 

b. 


c. 

Problem 14.10 Label each statement as True or False.

a. When a nucleus is strongly shielded, the effective field is larger than the applied field and the absorption shifts downfield.
b. When a nucleus is strongly shielded, the effective field is smaller than the applied field and the absorption is shifted upfield.
c. A nucleus that is strongly deshielded requires a lower field strength for resonance.
d. A nucleus that is strongly shielded absorbs at a larger δ value.

14.4 The Chemical Shift of Protons on sp^2 and sp Hybridized Carbons

The chemical shift of protons bonded to benzene rings, C−C double bonds, and C−C triple bonds merits additional comment.




7.3 ppm 4.5–6 ppm 2.5 ppm

Each of these functional groups contains π bonds with **loosely held π electrons.** When placed in a magnetic field, these π electrons move in a circular path, inducing a new magnetic field. How this induced magnetic field affects the chemical shift of a proton depends on the direction of the induced field *in the vicinity of the absorbing proton.*


Protons on Benzene Rings

In a magnetic field, the six π electrons in **benzene** circulate around the ring, creating a ring current. The magnetic field induced by these moving electrons *reinforces* the applied magnetic field in the vicinity of the protons. The protons thus feel a stronger magnetic field and a higher frequency is needed for resonance, so the **protons are deshielded and the absorption is *downfield.***



Protons on Carbon–Carbon Double Bonds

A similar phenomenon occurs with protons on carbon–carbon double bonds. In a magnetic field, the loosely held π electrons create a magnetic field that *reinforces* the applied field in the vicinity of the protons. Because the protons now feel a stronger magnetic field, they require a higher frequency for resonance. **The protons are deshielded and the absorption is *downfield.***



Protons on Carbon–Carbon Triple Bonds

In a magnetic field, the π electrons of a carbon–carbon triple bond are induced to circulate, but in this case the induced magnetic field *opposes* the applied magnetic field (B_0). The proton thus feels a weaker magnetic field, so a lower frequency is needed for resonance. **The nucleus is shielded and the absorption is *upfield*.**






Table 14.2 summarizes the shielding and deshielding effects due to circulating π electrons.

Table 14.2 Effect of π Electrons on Chemical Shift Values

Proton type	Effect	Chemical shift (ppm)
	highly deshielded	6.5–8
	deshielded	4.5–6
	shielded	~2.5


To remember the chemical shifts of some common bond types, it is helpful to think of a ^{1}H NMR spectrum as being divided into six different regions (Figure 14.5).

Figure 14.5
Regions in the ^{1}H NMR spectrum



- Shielded protons absorb at lower chemical shift (to the right).
- Deshielded protons absorb at higher chemical shift (to the left).
- Note: The drawn chemical shift scale is not linear.


Sample Problem 14.5 Rank H_a, H_b, and H_c in order of increasing chemical shift.



Solution

The H_a protons are bonded to an sp^3 hybridized carbon, so they are shielded and absorb upfield compared to H_b and H_c. Because the H_b protons are deshielded by the electronegative oxygen atom on the C to which they are bonded, they absorb downfield from H_a. The H_c proton is deshielded by two factors. The electronegative O atom withdraws electron density from H_c. Moreover, because H_c is bonded directly to a C=C, the magnetic field induced by the π electrons causes further deshielding. Thus, in order of increasing chemical shift, $H_a < H_b < H_c$.

Problem 14.11 Rank each group of protons in order of increasing chemical shift.




14.5 ^{1}H NMR: Intensity of Signals

The relative intensity of ^{1}H NMR signals also provides information about a compound's structure.

- **The area under an NMR signal is proportional to the number of absorbing protons.**


For example, in the ^{1}H NMR spectrum of $CH_3OC(CH_3)_3$, the ratio of the area under the downfield peak (due to the CH_3O- group) to the upfield peak [due to the $-C(CH_3)_3$ group] is 1:3. An NMR spectrometer automatically integrates the area under the peaks, and prints out a digital display of the *relative* areas of the NMR signals. Older NMR spectrometers print out a stepped curve (an **integral**) on the spectrum. The height of each step is proportional to the area under the peak, which is in turn proportional to the number of absorbing protons.




Integrals can be manually measured, but modern NMR spectrometers automatically calculate and plot the value of each integral in arbitrary units. If the heights of two integrals are in a 1:3 ratio, then the ratio of absorbing protons is 1:3, or 2:6, or 3:9, and so forth. This tells the *ratio,* not the absolute number of protons.


Problem 14.12 Which compounds give a ^{1}H NMR spectrum with two signals in a ratio of 2:3?

a. 


b. 
c. 

d. 

Problem 14.13 Compound **A** exhibits two signals in its ^{1}H NMR spectrum at 2.64 and 3.69 ppm and the ratio of the absorbing signals is 2:3. Compound **B** exhibits two signals in its ^{1}H NMR spectrum at 2.09 and 4.27 ppm and the ratio of the absorbing signals is 3:2. Which compound corresponds to dimethyl succinate and which compound corresponds to ethylene diacetate?




dimethyl succinate



ethylene diacetate

14.6 ^{1}H NMR: Spin–Spin Splitting

The ^{1}H NMR spectra you have seen up to this point have been limited to one or more single absorptions called **singlets.** In the ^{1}H NMR spectrum of $BrCH_2CHBr_2$, however, the two signals for the two different kinds of protons are each split into more than one peak. The splitting patterns, the result of **spin–spin splitting,** can be used to determine how many protons reside on the carbon atoms near the absorbing proton.




To understand spin–spin splitting, we must distinguish between the **absorbing protons** that give rise to an NMR signal, and the **adjacent protons** that cause the signal to split. **The number of adjacent protons determines the observed splitting pattern.**

- The CH_2 signal appears as **two peaks,** called a ***doublet.*** The relative area under the peaks of a doublet is 1:1.
- The CH signal appears as **three peaks,** called a ***triplet.*** The relative area under the peaks of a triplet is 1:2:1.


Spin–spin splitting occurs between nonequivalent protons on the same carbon or adjacent carbons. To illustrate how spin–spin splitting arises, we'll examine nonequivalent protons on adjacent carbons, the more common example. Spin–spin splitting arises because protons are little magnets that can be aligned with or against an applied magnetic field, and this affects the magnetic field that a nearby proton feels.

14.6A Splitting: How a Doublet Arises

First, let's examine how the doublet due to the CH_2 group in $BrCH_2CHBr_2$ arises. The CH_2 group contains the absorbing protons and the CH group contains the adjacent proton that causes the splitting.



When placed in an applied magnetic field (B_0), the adjacent proton ($CHBr_2$) can be aligned with (↑) or against (↓) B_0. As a result, the absorbing protons (CH_2Br) feel two slightly different magnetic fields—one slightly larger than B_0 and one slightly smaller than B_0. Because the absorbing protons feel two different magnetic fields, they absorb at two different frequencies in the NMR spectrum, thus splitting a single absorption into a doublet.




Keep in mind the difference between an **NMR signal** and an **NMR peak.** An NMR signal is the entire absorption due to a particular kind of proton. NMR peaks are contained within a signal. A doublet constitutes one signal that is split into two peaks.

- One adjacent proton splits an NMR signal into a doublet.

The two peaks of a doublet are approximately equal in area. The area under both peaks—the entire NMR signal—is due to both protons of the CH_2 group of $BrCH_2CHBr_2$.


coupling constant, ***J,*** in Hz



The frequency difference (measured in Hz) between the two peaks of the doublet is called the **coupling constant,** denoted by ***J.*** Coupling constants are usually in the range of 0–18 Hz, and are **independent of the strength of the applied magnetic field, B_0.**

14.6B Splitting: How a Triplet Arises

Now let's examine how the triplet due to the CH group in $BrCH_2CHBr_2$ arises. The CH group contains the absorbing proton and the CH_2 group contains the adjacent protons (H_a and H_b) that cause the splitting.



When placed in an applied magnetic field (B_0), the adjacent protons H_a and H_b can each be aligned with (↑) or against (↓) B_0. As a result, the absorbing proton feels three slightly different magnetic fields—one slightly larger than B_0, one slightly smaller than B_0, and one the same strength as B_0.




Table 14.3
Names for a Given Number of Peaks in an NMR Signal

Number of peaks	Name
1	singlet
2	doublet
3	triplet
4	quartet
5	quintet
6	sextet
7	septet
> 7	multiplet

Because the absorbing proton feels three different magnetic fields, it absorbs at three different frequencies in the NMR spectrum, thus splitting a single absorption into a triplet. Because there are two different ways to align one proton with B_0 and one proton against B_0—that is, $\uparrow_a\downarrow_b$ and $\downarrow_a\uparrow_b$—the middle peak of the triplet is twice as intense as the two outer peaks, making the ratio of the areas under the three peaks 1:2:1.

- **Two adjacent protons split an NMR signal into a triplet.**

When two protons split each other's NMR signals, they are said to be ***coupled.*** In $BrCH_2CHBr_2$, the CH proton is coupled to the CH_2 protons. The spacing between peaks in a split NMR signal, measured by the J value, is *equal* for coupled protons.

14.6C Splitting: The Rules and Examples

Three general rules describe the splitting patterns commonly seen in the ^{1}H NMR spectra of organic compounds.


Rule [1] **Equivalent protons don't split each other's signals.**

Rule [2] **A set of *n* nonequivalent protons splits the signal of a nearby proton into *n* + 1 peaks.**

- In $BrCH_2CHBr_2$, for example, *one* adjacent CH proton splits an NMR signal into *two* peaks (a doublet), and *two* adjacent CH_2 protons split an NMR signal into *three* peaks (a triplet). Names for split NMR signals containing two to seven peaks are given in Table 14.3. An NMR signal having more than seven peaks is called a **multiplet.**
- The inside peaks of a split NMR signal are always most intense, with the area under the peaks decreasing from the inner to the outer peaks in a given splitting pattern.


Rule [3] **Splitting is observed for nonequivalent protons on the same carbon or adjacent carbons.**

If H_a and H_b are not equivalent, splitting is observed in each of the following cases.




The splitting of an NMR signal reveals the number of nearby nonequivalent protons. It tells nothing about the absorbing proton itself.

Splitting is not generally observed between protons separated by more than three σ bonds. Although H_a and H_b are not equivalent to each other in butan-2-one and ethyl methyl ether, H_a and H_b are separated by four σ bonds, so they are too far away to split each other's NMR signals.



butan-2-one
H_a and H_b are separated by four σ bonds.

no splitting between H_a and H_b











ethyl methyl ether
H_a and H_b are separated by four σ bonds.

no splitting between H_a and H_b

Table 14.4 illustrates common splitting patterns observed for adjacent nonequivalent protons.

Table 14.4 Common Splitting Patterns Observed in ^{1}H NMR


Example	Pattern	Analysis
[1] 		• H: one adjacent H proton ⟶ two peaks ⟶ a **doublet** • H: one adjacent H proton ⟶ two peaks ⟶ a **doublet**
[2] 		• H: two adjacent H protons ⟶ three peaks ⟶ a **triplet** • H: one adjacent H proton ⟶ two peaks ⟶ a **doublet**
[3] 		• H: two adjacent H protons ⟶ three peaks ⟶ a **triplet** • H: two adjacent H protons ⟶ three peaks ⟶ a **triplet**
[4] 		• H: three adjacent H protons ⟶ four peaks ⟶ a **quartet*** • H: two adjacent H protons ⟶ three peaks ⟶ a **triplet**
[5] 		• H: three adjacent H protons ⟶ four peaks ⟶ a **quartet*** • H: one adjacent H proton ⟶ two peaks ⟶ a **doublet**

*The relative area under the peaks of a quartet is 1:3:3:1.


Predicting splitting is always a two-step process:

- **Determine if two protons are equivalent or different.** Only nonequivalent protons split each other.
- **Determine if two nonequivalent protons are close enough to split each other's signals.** Splitting is observed only for nonequivalent protons on the *same* carbon or *adjacent* carbons.


Several examples of spin–spin splitting in specific compounds illustrate the result of this two-step strategy.




- All protons are equivalent, so there is no splitting and the NMR signal is one singlet.



- There are two NMR signals. H_a and H_b are nonequivalent protons bonded to adjacent C atoms, so they are close enough to split each other's NMR signals. The H_a signal is split into a triplet by the two H_b protons. The H_b signal is split into a triplet by the two H_a protons.





- There are three NMR signals. H_a has no adjacent nonequivalent protons, so its signal is a singlet. The H_b signal is split into a quartet by the three H_c protons. The H_c signal is split into a triplet by the two H_b protons.





- There are two NMR signals. H_a and H_b are nonequivalent protons on the same carbon, so they are close enough to split each other's NMR signals. The H_a signal is split into a doublet by H_b. The H_b signal is split into a doublet by H_a.


Problem 14.14 Into how many peaks will each proton shown in red be split?


a. 

b. 


c. 


d. 

e. 


f. 

Problem 14.15 For each compound give the number of ^{1}H NMR signals, and then determine how many peaks are present for each NMR signal.

a. 

b. 

c. 

d. 


Problem 14.16 Sketch the NMR spectrum of CH_3CH_2Cl, giving the approximate location of each NMR signal.

14.7 More Complex Examples of Splitting

Up to now you have studied examples of spin–spin splitting where the absorbing proton has nearby protons on *one* adjacent carbon only. What happens when the absorbing proton has nonequivalent protons on *two* adjacent carbons? Different outcomes are possible, depending on whether the adjacent nonequivalent protons are *equivalent to* or *different from* each other.

For example, 2-bromopropane [$(CH_3)_2CHBr$] has two types of protons—H_a and H_b—so it exhibits two NMR signals, as shown in Figure 14.6.

Figure 14.6
The ^{1}H NMR spectrum of 2-bromopropane, $(CH_3)_2CHBr$



- The H_a protons have only one adjacent nonequivalent proton (H_b), so they are split into two peaks, a **doublet.**
- H_b has three H_a protons on each side. Because the six H_a protons are *equivalent to each other,* the ***n* + 1** rule can be used to determine splitting: 6 + 1 = 7 peaks, a **septet.**

This is a specific example of a general rule:

- **Whenever two (or three) sets of adjacent protons are *equivalent to each other*, use the *n* + 1 rule to determine the splitting pattern.**

A different outcome results when an absorbing proton is flanked by adjacent protons that are *not equivalent to each other.* Consider the splitting pattern expected for the H_b protons in the ^{1}H NMR spectrum of $CH_3CH_2CH_2Z$. H_b has protons on both adjacent carbons, but since H_a and H_c are *not equivalent to each other,* we cannot merely add them together and use the $n + 1$ rule.


$$\underset{H_a\ \ H_b\ \ H_c}{CH_3CH_2CH_2{-}Z}$$

Instead, to determine the splitting of H_b, we must consider the effect of the H_a protons and the H_c protons *separately.* The three H_a protons split the H_b signal into four peaks, and the two H_c protons split each of these four peaks into three peaks—that is, the NMR signal due to H_b consists of $4 \times 3 =$ **12 peaks.** Figure 14.7 shows a splitting diagram illustrating how these 12 peaks arise.

- **When two sets of adjacent protons are *different from each other* (*n* protons on one adjacent carbon and *m* protons on the other), the number of peaks in an NMR signal = (*n* + 1)(*m* + 1).**


It is only possible to see 12 peaks in an NMR spectrum when the coupling constants between each set of nonequivalent protons—that is, J_{ab} and J_{bc} in this example—are different; in other words, $J_{ab} \neq J_{bc}$. Such is the case with the nonequivalent protons on carbon–carbon double bonds, which is discussed in Section 14.8. **In practice, with flexible alkyl chains it is more common for J_{ab} and J_{bc} to be very similar or identical.** In this case, peaks overlap and many fewer than 12 peaks are observed.

Figure 14.7
A splitting diagram for the H_b protons in $CH_3CH_2CH_2Z$



- The H_b signal is split into 12 peaks, a quartet of triplets. The number of peaks actually seen for the signal depends on the relative size of the coupling constants, J_{ab} and J_{bc}. When $J_{ab} >> J_{bc}$, as drawn in this diagram, all 12 lines of the pattern are visible. When J_{ab} and J_{bc} are similar in magnitude, peaks overlap and fewer lines are observed.

Figure 14.8
The ^{1}H NMR spectrum of 1-bromopropane, $CH_3CH_2CH_2Br$



- H_a and H_c are both triplets.
- The signal for H_b appears as a multiplet of six peaks (a sextet), due to peak overlap; the number of peaks = $n + m + 1 = 3 + 2 + 1 = 6$ peaks.


The ^{1}H NMR spectrum of 1-bromopropane ($CH_3CH_2CH_2Br$) illustrates the result of peak overlap (Figure 14.8).


$$\underset{H_a\ \ H_b\ \ H_c}{CH_3CH_2CH_2—Br}$$

$CH_3CH_2CH_2Br$ has three different types of protons—H_a, H_b, and H_c—so it exhibits three NMR signals. H_a and H_c are each triplets because they are adjacent to two H_b protons. H_b has protons on both adjacent carbons, and H_a and H_c are *not equivalent to each other.* The three H_a protons should split the H_b signal into four peaks, and the two H_c protons should split each of these four peaks into three peaks—that is, the NMR signal due to H_b should once again consist of $4 \times 3 =$ **12 peaks.** However, since $J_{ab} = J_{bc}$ in this case, peak overlap occurs and a multiplet of only six peaks is observed.


In $CH_3CH_2CH_2Br$, **the *n* protons on one adjacent carbon and the *m* protons on the other adjacent carbon split the observed signal into *n* + *m* + 1 peaks.** In other words, the 3 H_a protons and 2 H_c protons split the NMR signal into 3 + 2 + 1 = 6 peaks, as shown in the sextet in Figure 14.8.

Sample Problem 14.6 How many peaks are present in the NMR signal of the labeled protons of each compound?


a. 

b. 

Solution


a. 


- H_b has two H_a protons on each adjacent C. Because the four H_a protons are equivalent to each other, the $n + 1$ rule can be used to determine splitting: 4 + 1 = **5 peaks,** a quintet.


b. 


- H_b has two H_a protons on one adjacent C and two H_c protons on the other. Because H_a and H_c are *not equivalent to each other,* the maximum number of peaks for $H_b = (n + 1)(m + 1) = (2 + 1)(2 + 1) =$ **9 peaks.** However, since this molecule has a *flexible alkyl chain,* it is likely that J_{ab} and J_{bc} are very similar, so that peak overlap occurs, and the number of peaks for $H_b = n + m + 1 = 2 + 2 + 1 =$ **5 peaks.**

Problem 14.17 How many peaks are present in the NMR signal of each labeled proton?


a. 


b. 


c. 

d. 

Problem 14.18 Describe the ^{1}H NMR spectrum of each compound. State how many NMR signals are present, the splitting pattern for each signal, and the approximate chemical shift.

a. 


b. 

c. 

d. 


14.8 Spin–Spin Splitting in Alkenes

Protons on carbon–carbon double bonds often give characteristic splitting patterns. A disubstituted double bond can have two **geminal protons** (on the same carbon atom), two **cis protons,** or two **trans protons.** When these protons are different, each proton splits the NMR signal of the other, so that each proton appears as a doublet. **The magnitude of the coupling constant *J* for these doublets depends on the arrangement of hydrogen atoms.**



Thus, the *E* and *Z* isomers of 3-chloropropenoic acid both exhibit two doublets for the two alkenyl protons, but the coupling constant is larger when the protons are trans compared to when the protons are cis, as shown in Figure 14.9.

Figure 14.9
^{1}H NMR spectra for the alkenyl protons of (*E*)- and (*Z*)-3-chloropropenoic acid




- Although both (*E*)- and (*Z*)-3-chloropropenoic acid show two doublets in their ^{1}H NMR spectra for their alkenyl protons, $J_{trans} > J_{cis}$.

When a double bond is monosubstituted, there are three nonequivalent protons, and the pattern is more complicated because all three protons are coupled to each other. For example, vinyl acetate ($CH_2{=}CHOCOCH_3$) has four different types of protons, three of which are bonded to the double bond. Besides the singlet for the CH_3 group, each proton on the double bond is coupled to two other different protons on the double bond, giving the spectrum in Figure 14.10.

- H_b has two nearby nonequivalent protons that split its signal, the geminal proton H_c and the trans proton H_d. H_d splits the H_b signal into a doublet, and the H_c proton splits the doublet into two doublets. This pattern of four peaks is called a **doublet of doublets.**
- H_c has two nearby nonequivalent protons that split its signal, the geminal proton H_b and the cis proton H_d. H_d splits the H_c signal into a doublet, and the H_b proton splits the doublet into two doublets, forming another **doublet of doublets.**
- H_d has two nearby nonequivalent protons that split its signal, the trans proton H_b and the cis proton H_c. H_b splits the H_d signal into a doublet, and the H_c proton splits the doublet into two doublets, forming another **doublet of doublets.**


Figure 14.10
The ^{1}H NMR spectrum of vinyl acetate ($CH_2{=}CHOCOCH_3$)




Vinyl acetate is polymerized to poly(vinyl acetate) (Problem 15.26), a polymer used in paints, glues, and adhesives.

Splitting diagrams for the three alkenyl protons in vinyl acetate are drawn in Figure 14.11. Note that each pattern is different in appearance because the magnitude of the coupling constants forming them is different.

Figure 14.11
Splitting diagram for the alkenyl protons in vinyl acetate ($CH_2{=}CHOCOCH_3$)




J_{bc} = 1.2 Hz (geminal)
J_{cd} = 6.5 Hz (cis)
J_{bd} = 14 Hz (trans)



One nearby H splits the signal into a doublet.

The second nearby H splits the doublet into a doublet of doublets.

Problem 14.19 Draw a splitting diagram for H_b in *trans*-1,3-dichloropropene, given that J_{ab} = 13.1 Hz and J_{bc} = 7.2 Hz.



trans-1,3-dichloropropene

Problem 14.20 Identify **A** and **B,** isomers of molecular formula $C_3H_4Cl_2$, from the given ^{1}H NMR data: Compound **A** exhibits signals at 1.75 (doublet, 3 H, J = 6.9 Hz) and 5.89 (quartet, 1 H, J = 6.9 Hz) ppm. Compound **B** exhibits signals at 4.16 (singlet, 2 H), 5.42 (doublet, 1 H, J = 1.9 Hz), and 5.59 (doublet, 1 H, J = 1.9 Hz) ppm.

14.9 Other Facts About ^{1}H NMR Spectroscopy

14.9A OH Protons

- **Under usual conditions, an OH proton does not split the NMR signal of adjacent protons.**
- **The signal due to an OH proton is not split by adjacent protons.**

Ethanol (CH_3CH_2OH), for example, has three different types of protons, so there are three signals in its ^{1}H NMR spectrum, as shown in Figure 14.12.

- The H_a signal is split by the two H_b protons into three peaks, a **triplet.**
- The H_b signal is split by only the three H_a protons into four peaks, a **quartet.** The adjacent OH proton does *not* split the signal due to H_b.
- H_c is a **singlet** because OH protons are *not* split by adjacent protons.

Why is a proton bonded to an oxygen atom a singlet in a ^{1}H NMR spectrum? Protons on electronegative elements rapidly **exchange** between molecules in the presence of trace amounts of acid or base. It is as if the CH_2 group in ethanol never "feels" the presence of the OH proton, because the OH proton is rapidly moving from one molecule to another. We therefore see a peak due to the OH proton, but it is a single peak with no splitting. This phenomenon usually occurs with NH and OH protons.







Figure 14.12
The ^{1}H NMR spectrum of ethanol (CH_3CH_2OH)

Problem 14.21 How many signals are present in the ^{1}H NMR spectrum for each molecule? What splitting is observed in each signal?

a. 


b. 

c. 

14.9B Cyclohexane Conformations

How does the rotation around carbon–carbon σ bonds and the ring flip of cyclohexane rings affect an NMR spectrum? Because these processes are rapid at room temperature, an NMR spectrum records an **average** of all conformations that interconvert.

Thus, even though each cyclohexane carbon has two different types of hydrogens—one axial and one equatorial—the two chair forms of cyclohexane rapidly interconvert them, and an **NMR spectrum shows a single signal for the average environment** that it "sees."




Axial and equatorial H's rapidly interconvert. NMR sees an average environment and shows one signal.

14.9C Protons on Benzene Rings


We will learn more about the spectroscopic absorptions of benzene derivatives in Chapter 17.

Benzene has six equivalent, deshielded protons and exhibits a single peak in its ^{1}H NMR spectrum at 7.27 ppm. Monosubstituted benzene derivatives—that is, benzene rings with one H atom replaced by another substituent Z—contain five deshielded protons that are no longer all equivalent to each other. The identity of Z determines the appearance of this region of a ^{1}H NMR spectrum (6.5–8 ppm), as shown in Figure 14.13. We will not analyze the splitting patterns observed for the ring protons of monosubstituted benzenes.

Figure 14.13
The 6.5–8 ppm region of the ^{1}H NMR spectrum of three benzene derivatives




A monosubstituted benzene ring has three different types of H atoms: H_a, H_b, and H_c.




- The appearance of the signals in the 6.5–8 ppm region of the ^{1}H NMR spectrum depends on the identity of Z in C_6H_5Z.


Problem 14.22 What protons in alcohol **A** give rise to each signal in its ^{1}H NMR spectrum? Explain all splitting patterns observed for absorptions between 0–7 ppm.



A



Problem 14.23 How many peaks are observed in the ^{1}H NMR signal for each proton shown in red in palau'amine, the complex chapter-opening molecule?




palau'amine

14.10 Using ^{1}H NMR to Identify an Unknown

Once we know a compound's molecular formula from its mass spectral data and the identity of its functional group from its IR spectrum, we can then use its ^{1}H NMR spectrum to determine its structure. A suggested procedure is illustrated for compound **X,** whose molecular formula ($C_4H_8O_2$) and functional group (C=O) were determined in Section 13.8.

How To Use ^{1}H NMR Data to Determine a Structure

Example **Using its ^{1}H NMR spectrum, determine the structure of an unknown compound X that has molecular formula $C_4H_8O_2$ and contains a C=O absorption in its IR spectrum.**



Step [1] **Determine the number of different kinds of protons.**

- The number of NMR signals equals the number of different types of protons.
- This molecule has three NMR signals ([A], [B], and [C]) and therefore **three** types of protons (H_a, H_b, and H_c).


Step [2] **Use the relative area to determine the number of H atoms giving rise to each signal.**

- The relative area (printed on top of each signal) gives the *ratio* of absorbing protons responsible for each signal. In this case, the ratio is 3:2:3 for the signals from left-to-right.
- **When the sum of the relative areas *equals* the number of H's in the molecular formula, the relative area gives the number of absorbing H's responsible for the NMR signal.** In this example, the sum of the relative areas is 3 + 2 + 3 = 8, and the unknown has 8 H's, so the signals are due to 3 H's, 2 H's, and 3 H's from left-to-right in the spectrum.


3 H_a protons signal [A]	2 H_b protons signal [B]	3 H_c protons signal [C]
Three equivalent H's usually means a **CH_3** group.	Two equivalent H's usually means a **CH_2** group.	Three equivalent H's usually means a **CH_3** group.

Step [3] **Use individual splitting patterns to determine what carbon atoms are bonded to each other.**

- Start with the singlets. Signal [C] is due to a CH_3 group with no adjacent nonequivalent H atoms. Possible structures include:



- Because signal [A] is a **triplet,** there must be **2 H's** (CH_2 group) on the adjacent carbon.
- Because signal [B] is a **quartet,** there must be **3 H's** (CH_3 group) on the adjacent carbon.
- This information suggests that **X** has an **ethyl** group ⟶ CH_3CH_2-.




How To, continued . . .

To summarize, **X** contains CH_3-, CH_3CH_2-, and C=O (from the IR). Comparing these atoms with the molecular formula shows that one O atom is missing. Because O atoms do not absorb in a ^{1}H NMR spectrum, their presence can only be inferred by examining the chemical shift of protons near them. O atoms are more electronegative than C, thus deshielding nearby protons, and shifting their absorption downfield.

Step [4] Use chemical shift data to complete the structure.

- Put the structure together in a manner that preserves the splitting data and is consistent with the reported chemical shifts.
- In this example, two isomeric structures **(A** and **B)** are possible for **X** considering the splitting data only:




- Chemical shift information distinguishes the two possibilities. The electronegative O atom deshields adjacent H's, shifting them downfield between 3 and 4 ppm. If **A** is the correct structure, the singlet due to the CH_3 group (H_c) should occur downfield, whereas if **B** is the correct structure, the quartet due to the CH_2 group (H_b) should occur downfield.
- Because the NMR of **X** has a singlet (not a quartet) at 3.7, **A is the correct structure.**

Problem 14.24 Propose a structure for a compound of molecular formula $C_7H_{14}O_2$ with an IR absorption at 1740 cm^{-1} and the following ^{1}H NMR data:

Absorption	ppm	Relative area
singlet	1.2	9
triplet	1.3	3
quartet	4.1	2

Problem 14.25 Propose a structure for a compound of molecular formula C_3H_8O with an IR absorption at 3600–3200 cm^{-1} and the following NMR spectrum:



Problem 14.26 Identify products **A** and **B** from the given ^{1}H NMR data.


a. Treatment of $CH_2=CHCOCH_3$ with one equivalent of HCl forms compound **A.** **A** exhibits the following absorptions in its ^{1}H NMR spectrum: 2.2 (singlet, 3 H), 3.05 (triplet, 2 H), and 3.6 (triplet, 2 H) ppm. What is the structure of **A?**

b. Treatment of acetone [$(CH_3)_2C=O$] with dilute aqueous base forms **B.** Compound **B** exhibits four singlets in its ^{1}H NMR spectrum at 1.3 (6 H), 2.2 (3 H), 2.5 (2 H), and 3.8 (1 H) ppm. What is the structure of **B?**

14.11 ^{13}C NMR Spectroscopy

^{13}C NMR spectroscopy is also an important tool for organic structure analysis. The physical basis for ^{13}C NMR is the same as for ^{1}H NMR. When placed in a magnetic field, B_0, ^{13}C nuclei can align themselves with or against B_0. More nuclei are aligned with B_0 because this arrangement is lower in energy, but these nuclei can be made to spin flip against the applied field by applying RF radiation of the appropriate frequency.

^{13}C NMR spectra, like ^{1}H NMR spectra, plot peak intensity versus chemical shift, using TMS as the reference signal at 0 ppm. ^{13}C occurs in only 1.1% natural abundance, however, so ^{13}C NMR signals are much weaker than ^{1}H NMR signals. To overcome this limitation, modern spectrometers irradiate samples with many pulses of RF radiation and use mathematical tools to increase signal sensitivity and decrease background noise. The spectrum of acetic acid (CH_3COOH) illustrates the general features of a ^{13}C NMR spectrum.



^{13}C NMR spectra are easier to analyze than ^{1}H spectra because signals are not split. **Each type of carbon atom appears as a single peak.**

Why aren't ^{13}C signals split by nearby carbon atoms? Recall from Section 14.6 that splitting occurs when two NMR active nuclei—like two protons—are close to each other. Because of the low natural abundance of ^{13}C nuclei (1.1%), the chance of two ^{13}C nuclei being bonded to each other is very small (0.01%), and so no carbon–carbon splitting is observed.

A ^{13}C NMR signal can also be split by nearby protons. This ^{1}H – ^{13}C splitting is usually eliminated from a spectrum, however, by using an instrumental technique that decouples the proton–carbon interactions, so that every signal in a ^{13}C NMR spectrum is a singlet.

Two features of ^{13}C NMR spectra provide the most structural information: the **number of signals** observed and the **chemical shifts** of those signals.


14.11A ^{13}C NMR: Number of Signals

- **The number of signals in a ^{13}C spectrum gives the number of different types of carbon atoms in a molecule.**

Carbon atoms in the same environment give the same NMR signal, whereas carbons in different environments give different NMR signals. The ^{13}C NMR spectrum of CH_3COOH has two signals because there are two different types of carbon atoms—the C of the CH_3 group and the C of the carbonyl (C=O).

- **Because ^{13}C NMR signals are not split, the number of signals equals the number of lines in the ^{13}C NMR spectrum.**

Thus, the ^{13}C NMR spectra of dimethyl ether, chloroethane, and methyl acetate exhibit one, two, and three lines, respectively, because these compounds contain one, two, and three different types of carbon atoms.




dimethyl ether

1 ^{13}C NMR signal
Both C's are equivalent.



chloroethane

2 ^{13}C NMR signals





methyl acetate


3 ^{13}C NMR signals

In contrast to what occurs in proton NMR, peak intensity is not proportional to the number of absorbing carbons, so ^{13}C NMR signals are not integrated.

Sample Problem 14.7 How many lines are observed in the ^{13}C NMR spectrum of each compound?

a. 

b. 


c. 

Solution

The number of different types of carbons equals the number of lines in a ^{13}C NMR spectrum.

a. 

3 types of C's
3 ^{13}C NMR signals


b. 


4 types of C's
4 ^{13}C NMR signals


c. 


2 types of C's
2 ^{13}C NMR signals

Problem 14.27 How many lines are observed in the ^{13}C NMR spectrum of each compound?

a. 

b. 


c. 

d. 

Problem 14.28 Draw all constitutional isomers of molecular formula $C_3H_6Cl_2$.

a. How many signals does each isomer exhibit in its ^{1}H NMR spectrum?
b. How many lines does each isomer exhibit in its ^{13}C NMR spectrum?
c. When only the number of signals in both ^{1}H and ^{13}C NMR spectroscopy is considered, is it possible to distinguish all of these constitutional isomers?

Problem 14.29 Esters of chrysanthemic acid are naturally occurring insecticides. How many lines are present in the ^{13}C NMR spectrum of chrysanthemic acid?



chrysanthemic acid


Esters of chrysanthemic acid are obtained from the flowers of *Chrysanthemum cinerariifolium.* Because they are biodegradable and active against numerous insect species, these esters are widely used insecticides (see also Section 26.4).



14.11B ^{13}C NMR: Position of Signals

In contrast to the small range of chemical shifts in ^{1}H NMR (0–12 ppm usually), ^{13}C NMR absorptions occur over a much broader range, 0–220 ppm. The chemical shifts of carbon atoms in ^{13}C NMR depend on the same effects as the chemical shifts of protons in ^{1}H NMR:


- **The sp^3 hybridized C atoms of alkyl groups are shielded and absorb upfield.**
- **Electronegative elements like halogen, nitrogen, and oxygen shift absorptions downfield.**
- **The sp^2 hybridized C atoms of alkenes and benzene rings absorb downfield.**
- **Carbonyl carbons are highly deshielded, and absorb farther downfield than other carbon types.**


Table 14.5 lists common ^{13}C chemical shift values. The ^{13}C NMR spectra of propan-1-ol ($CH_3CH_2CH_2OH$) and methyl acetate ($CH_3CO_2CH_3$) in Figure 14.14 illustrate these principles.


Table 14.5 Common ^{13}C Chemical Shift Values

Type of carbon	Chemical shift (ppm)	Type of carbon	Chemical shift (ppm)
	5–45		100–140
 Z = N, O, X	30–80		120–150
	65–100		160–210

Problem 14.30 Which of the highlighted carbon atoms in each molecule absorbs farther downfield?

a. 


b. 

c. 

d. 


Figure 14.14 Representative ^{13}C NMR spectra

a. Propan-1-ol



- The three types of C's in propan-1-ol—identified as C_a, C_b, and C_c—give rise to three ^{13}C NMR signals.
- Deshielding increases with increasing proximity to the electronegative O atom, and the absorption shifts downfield; thus, in order of increasing chemical shift: $C_a < C_b < C_c$.


b. Methyl acetate




- The three types of C's in methyl acetate—identified as C_a, C_b, and C_c—give rise to three ^{13}C NMR signals.
- The carbonyl carbon (C_b) is highly deshielded, so it absorbs farthest downfield.
- C_a, an sp^3 hybridized C that is not bonded to an O atom, is the most shielded, and so it absorbs farthest upfield.
- Thus, in order of increasing chemical shift: $C_a < C_c < C_b$.

Problem 14.31 Identify the carbon atoms that give rise to each NMR signal.

a.



b.



Problem 14.32 A compound of molecular formula $C_4H_8O_2$ shows no IR peaks at 3600–3200 or 1700 cm^{-1}. It exhibits one singlet in its ^{1}H NMR spectrum at 3.69 ppm, and one line in its ^{13}C NMR spectrum at 67 ppm. What is the structure of this unknown?

Problem 14.33 Draw the structure of a compound of molecular formula C_4H_8O that has a signal in its ^{13}C NMR spectrum at > 160 ppm. Then draw the structure of an isomer of molecular formula C_4H_8O that has all of its ^{13}C NMR signals at < 160 ppm.

14.12 Magnetic Resonance Imaging (MRI)

Magnetic resonance imaging (MRI)—NMR spectroscopy in medicine—is a powerful diagnostic technique (Figure 14.15a). The "sample" is the patient, who is placed in a large cavity in a magnetic field, and then irradiated with RF energy. Because RF energy has very low frequency and low energy, the method is safer than X-rays or computed tomography (CT) scans that employ high-frequency, high-energy radiation that is known to damage living cells.


Living tissue contains protons (especially the H atoms in H_2O) in different concentrations and environments. When irradiated with RF energy, these protons are excited to a higher energy spin state, and then fall back to the lower energy spin state. These data are analyzed by a computer that generates a plot that delineates tissues of different proton density (Figure 14.15b). MRIs can be recorded in any plane. Moreover, because the calcium present in bones is not NMR active, an MRI instrument can "see through" bones such as the skull and visualize the soft tissue underneath.

Figure 14.15
Magnetic resonance imaging

a.



b.



a. An MRI instrument: An MRI instrument is especially useful for visualizing soft tissue. In 2002, 60 million MRI procedures were performed. The 2003 Nobel Prize in Physiology or Medicine was awarded to chemist Paul C. Lauterbur and physicist Sir Peter Mansfield for their contributions in developing magnetic resonance imaging.
b. An MRI image of the lower back: **A** labels spinal cord compression from a herniated disc. **B** labels the spinal cord, which would not be visualized with conventional X-rays.

KEY CONCEPTS

Nuclear Magnetic Resonance Spectroscopy

^{1}H NMR Spectroscopy

[1] The **number of signals** equals the number of different types of protons (14.2).

[2] The **position of a signal** (its chemical shift) is determined by shielding and deshielding effects.

- Shielding shifts an absorption upfield; deshielding shifts an absorption downfield.
- Electronegative atoms withdraw electron density, deshield a nucleus, and shift an absorption downfield (14.3).




This proton is shielded. Its absorption is upfield, 0.9–2 ppm.



This proton is deshielded. Its absorption is farther downfield, 2.5–4 ppm.

- Loosely held π electrons can either shield or deshield a nucleus. Protons on benzene rings and double bonds are deshielded and absorb downfield, whereas protons on triple bonds are shielded and absorb upfield (14.4).



deshielded H
downfield absorption



deshielded H
downfield absorption



shielded H
upfield absorption

[3] The **area under an NMR signal** is proportional to the number of absorbing protons (14.5).

[4] **Spin–spin splitting** tells about nearby nonequivalent protons (14.6–14.8).

- Equivalent protons do not split each other's signals.
- A set of *n* nonequivalent protons on the same carbon or adjacent carbons splits an NMR signal into *n* + 1 peaks.
- OH and NH protons do not cause splitting (14.9).
- When an absorbing proton has two sets of nearby nonequivalent protons that are equivalent to each other, use the *n* + 1 rule to determine splitting.
- When an absorbing proton has two sets of nearby nonequivalent protons that are not equivalent to each other, the number of peaks in the NMR signal = (*n* + 1)(*m* + 1). In flexible alkyl chains, peak overlap often occurs, resulting in *n* + *m* + 1 peaks in an NMR signal.

^{13}C NMR Spectroscopy (14.11)

[1] **The number of signals equals the number of different types of carbon atoms.** All signals are **single peaks.**


[2] The relative position of ^{13}C signals is determined by shielding and deshielding effects.

- Carbons that are sp^3 hybridized are shielded and absorb upfield.
- Electronegative elements (N, O, and halogen) shift absorptions downfield.
- The carbons of alkenes and benzene rings absorb downfield.
- Carbonyl carbons are highly deshielded, and absorb farther downfield than other carbon types.


PROBLEMS

Problems Using Three-Dimensional Models

14.34 (a) How many ^{1}H NMR signals does each of the following compounds exhibit? (b) How many ^{13}C NMR signals does each compound exhibit?




A




B

14.35 (a) How many ^{1}H NMR signals does each compound show? (b) Into how many peaks is each signal split?



C





D


^{1}H NMR Spectroscopy—Determining Equivalent Protons


14.36 How many different types of protons are present in each compound?


a. 


b. 

c. 

d. 


e. 

f. 


g. 

h. 


i. 

j. 


14.37 How many ^{1}H NMR signals does each natural product exhibit?

a. 


caffeine
(from coffee and tea leaves)

b. 

vanillin
(from the vanilla bean)

c. 

thymol
(from thyme)

d. 

capsaicin
(from hot peppers)


^{1}H NMR—Chemical Shift and Integration


14.38 Using a 300 MHz NMR instrument:

a. How many Hz downfield from TMS is a signal at 2.5 ppm?
b. If a signal comes at 1200 Hz downfield from TMS, at what ppm does it occur?
c. If two signals are separated by 2 ppm, how many Hz does this correspond to?

14.39 What effect does increasing the operating frequency of a ^{1}H NMR spectrum have on each value: (a) the chemical shift in δ; (b) the frequency of an absorption in Hz; (c) the magnitude of a coupling constant J in Hz?

14.40 Which of the labeled protons in each pair absorbs farther downfield?


a. 


b. 


14.41 How could you use chemical shift and integration data in ^{1}H NMR spectroscopy to distinguish between $CH_3OCH_2CH_2OCH_3$ and $CH_3OCH_2OCH_3$? The ^{1}H NMR spectrum of each compound contains only singlets.


^{1}H NMR—Splitting


14.42 Into how many peaks will the signal for each of the labeled protons be split?


a. 


c. 


e. 


g. 


i. 

b. 

d. 


f. 


h. 

j. 


14.43 How can you use ^{1}H NMR spectroscopy to distinguish between $CH_2=C(Br)CO_2CH_3$ and methyl (*E*)-3-bromopropenoate, $BrCH=CHCO_2CH_3$?

14.44 What splitting pattern is observed for each proton in the following compounds?


a. 

b. 

14.45 Label the signals due to H_a, H_b, and H_c in the ^{1}H NMR spectrum of acrylonitrile ($CH_2=CHCN$). Draw a splitting diagram for the absorption due to the H_a proton.



J_{ab} = 11.8 Hz
J_{bc} = 0.9 Hz
J_{ac} = 18 Hz





^{13}C NMR


14.46 Draw the four constitutional isomers having molecular formula C_4H_9Br and indicate how many different kinds of carbon atoms each has.


14.47 Explain why the carbonyl carbon of an aldehyde or ketone absorbs farther downfield than the carbonyl carbon of an ester in a ^{13}C NMR spectrum.


14.48 How many ^{13}C NMR signals does each compound exhibit?

a. 


c. 

e. 

g. 


b. 


d. 

f. 

h. 

14.49 Rank the highlighted carbon atoms in each compound in order of increasing chemical shift.

a. 

b. 

14.50 Identify the carbon atoms that give rise to the signals in the ^{13}C NMR spectrum of each compound.

a. $CH_3CH_2CH_2CH_2OH$; ^{13}C NMR: 14, 19, 35, and 62 ppm
b. $(CH_3)_2CHCHO$; ^{13}C NMR: 16, 41, and 205 ppm
c. $CH_2=CHCH(OH)CH_3$; ^{13}C NMR: 23, 69, 113, and 143 ppm


14.51 a. How many signals does dimethyl fumarate ($CH_3O_2CCH=CHCO_2CH_3$, with a trans C=C) exhibit in its ^{13}C NMR spectrum?

b. Draw the structure of an isomer of dimethyl fumarate that has each of the following number of signals in its ^{13}C NMR spectrum: [1] three; [2] four; [5] five.

Combined Spectroscopy Problems


Additional spectroscopy problems are located at the end of Chapters 15–23 and 25.

14.52 Answer the following questions about each of the hydroxy ketones: 1-hydroxybutan-2-one (**A**) and 4-hydroxybutan-2-one (**B**).



1-hydroxybutan-2-one

A



4-hydroxybutan-2-one

B

a. What is the molecular ion in the mass spectrum?

b. What IR absorptions are present in the functional group region?

c. How many lines are observed in the ^{13}C NMR spectrum?

d. How many signals are observed in the ^{1}H NMR spectrum?

e. Give the splitting observed for each type of proton as well as its approximate chemical shift.

14.53 Propose a structure consistent with each set of spectral data:

a. $C_4H_8Br_2$: IR peak at 3000–2850 cm^{-1}; NMR (ppm):
1.87 (singlet, 6 H)
3.86 (singlet, 2 H)

b. $C_3H_6Br_2$: IR peak at 3000–2850 cm^{-1}; NMR (ppm):
2.4 (quintet)
3.5 (triplet)

c. $C_5H_{10}O_2$: IR peak at 1740 cm^{-1}; NMR (ppm):
1.15 (triplet, 3 H) 2.30 (quartet, 2 H)
1.25 (triplet, 3 H) 4.72 (quartet, 2 H)

d. $C_6H_{14}O$: IR peak at 3600–3200 cm^{-1}; NMR (ppm):
0.8 (triplet, 6 H) 1.5 (quartet, 4 H)
1.0 (singlet, 3 H) 1.6 (singlet, 1 H)

e. $C_6H_{14}O$: IR peak at 3000–2850 cm^{-1}; NMR (ppm):
1.10 (doublet, relative area = 6)
3.60 (septet, relative area = 1)

f. C_3H_6O: IR peak at 1730 cm^{-1}; NMR (ppm):
1.11 (triplet)
2.46 (multiplet)
9.79 (triplet)

14.54 Identify the structures of isomers **A** and **B** (molecular formula $C_9H_{10}O$).

Compound **A:** IR peak at 1742 cm^{-1}; ^{1}H NMR data (ppm) at 2.15 (singlet, 3 H), 3.70 (singlet, 2 H), and 7.20 (broad singlet, 5 H).

Compound **B:** IR peak at 1688 cm^{-1}; ^{1}H NMR data (ppm) at 1.22 (triplet, 3 H), 2.98 (quartet, 2 H), and 7.28–7.95 (multiplet, 5 H).

14.55 Reaction of $C_6H_5CH_2CH_2OH$ with CH_3COCl affords compound **W,** which has molecular formula $C_{10}H_{12}O_2$. **W** shows prominent IR absorptions at 3088–2897, 1740, and 1606 cm^{-1}. **W** exhibits the following signals in its ^{1}H NMR spectrum: 2.02 (singlet), 2.91 (triplet), 4.25 (triplet), and 7.20–7.35 (multiplet) ppm. What is the structure of **W?** We will learn about this reaction in Chapter 22.


14.56 Treatment of 2-methylpropanenitrile [$(CH_3)_2CHCN$] with $CH_3CH_2CH_2MgBr$, followed by aqueous acid, affords compound **V,** which has molecular formula $C_7H_{14}O$. **V** has a strong absorption in its IR spectrum at 1713 cm^{-1}, and gives the following ^{1}H NMR data: 0.91 (triplet, 3 H), 1.09 (doublet, 6 H), 1.6 (multiplet, 2 H), 2.43 (triplet, 2 H), and 2.60 (septet, 1 H) ppm. What is the structure of **V?** We will learn about this reaction in Chapter 22.

14.57 Compound **C** has a molecular ion in its mass spectrum at 146 and a prominent absorption in its IR spectrum at 1762 cm^{-1}. **C** shows the following ^{1}H NMR spectral data: 1.47 (doublet, 3 H), 2.07 (singlet, 6 H), and 6.84 (quartet, 1 H) ppm. What is the structure of **C?**


14.58 As we will learn in Chapter 20, reaction of $(CH_3)_2CO$ with LiC≡CH followed by H_2O affords compound **D,** which has a molecular ion in its mass spectrum at 84 and prominent absorptions in its IR spectrum at 3600–3200, 3303, 2938, and 2120 cm^{-1}. **D** shows the following ^1H NMR spectral data: 1.53 (singlet, 6 H), 2.37 (singlet, 1 H), and 2.43 (singlet, 1 H) ppm. What is the structure of **D?**

14.59 Identify the structures of isomers **E** and **F** (molecular formula $C_4H_8O_2$). Relative areas are given above each signal.

a. **Compound E:** IR absorption at 1743 cm^{-1}




b. **Compound F:** IR absorption at 1730 cm^{-1}




14.60 Identify the structures of isomers **H** and **I** (molecular formula $C_8H_{11}N$).

a. **Compound H:** IR absorptions at 3365, 3284, 3026, 2932, 1603, and 1497 cm^{-1}




b. **Compound I:** IR absorptions at 3367, 3286, 3027, 2962, 1604, and 1492 cm^{-1}




14.61 Propose a structure consistent with each set of data.

a. $C_9H_{10}O_2$: IR absorption at 1718 cm^{-1}




b. C_9H_{12}: IR absorption at 2850–3150 cm^{-1}



14.62 Reaction of $(CH_3)_3CCHO$ with $(C_6H_5)_3P{=}C(CH_3)OCH_3$, followed by treatment with aqueous acid, affords **R** ($C_7H_{14}O$). **R** has a strong absorption in its IR spectrum at 1717 cm^{-1} and three singlets in its ^1H NMR spectrum at 1.02 (9 H), 2.13 (3 H), and 2.33 (2 H) ppm. What is the structure of **R?** We will learn about this reaction in Chapter 21.

14.63 Reaction of aldehyde **D** with amino alcohol **E** in the presence of NaH forms **F** (molecular formula $C_{11}H_{15}NO_2$). **F** absorbs at 1730 cm^{-1} in its IR spectrum. **F** also shows eight lines in its ^{13}C NMR spectrum, and gives the following ^1H NMR spectrum: 2.32 (singlet, 6 H), 3.05 (triplet, 2 H), 4.20 (triplet, 2 H), 6.97 (doublet, 2 H), 7.82 (doublet, 2 H), and 9.97 (singlet, 1 H) ppm. Propose a structure for **F.** We will learn about this reaction in Chapter 18.




D **E**

14.64 Propose a structure consistent with each set of data.


a. Compound **J:** molecular ion at 72; IR peak at 1710 cm^{-1}; ^1H NMR data (ppm) at 1.0 (triplet, 3 H), 2.1 (singlet, 3 H), and 2.4 (quartet, 2 H)


b. Compound **K:** molecular ion at 88; IR peak at 3600–3200 cm^{-1}; ^1H NMR data (ppm) at 0.9 (triplet, 3 H), 1.2 (singlet, 6 H), 1.5 (quartet, 2 H), and 1.6 (singlet, 1 H)

14.65 In the presence of a small amount of acid, a solution of acetaldehyde (CH_3CHO) in methanol (CH_3OH) was allowed to stand and a new compound **L** was formed. **L** has a molecular ion in its mass spectrum at 90 and IR absorptions at 2992 and 2941 cm^{-1}. **L** shows three signals in its ^{13}C NMR at 19, 52, and 101 ppm. The ^1H NMR spectrum of **L** is given below. What is the structure of **L?**




14.66 Treatment of $(CH_3)_2CHCH(OH)CH_2CH_3$ with TsOH affords two products (**M** and **N**) with molecular formula C_6H_{12}. The ^1H NMR spectra of **M** and **N** are given below. Propose structures for **M** and **N** and draw a mechanism to explain their formation.







14.67 Compound **O** has molecular formula $C_{10}H_{12}O$ and shows an IR absorption at 1687 cm^{-1}. The ^1H NMR spectrum of **O** is given below. What is the structure of **O?**





14.68 Compound **P** has molecular formula $C_5H_9ClO_2$. Deduce the structure of **P** from its ^1H and ^{13}C NMR spectra.





14.69 Treatment of butan-2-one ($CH_3COCH_2CH_3$) with strong base followed by CH_3I forms a compound **Q,** which gives a molecular ion in its mass spectrum at 86. The IR (> 1500 cm^{-1} only) and ^1H NMR spectrum of **Q** are given below. What is the structure of **Q?**






14.70 When 2-bromo-3,3-dimethylbutane is treated with $K^+ {}^-OC(CH_3)_3$, a single product **T** having molecular formula C_6H_{12} is formed. When 3,3-dimethylbutan-2-ol is treated with H_2SO_4, the major product **U** has the same molecular formula. Given the following ^1H NMR data, what are the structures of **T** and **U?** Explain in detail the splitting patterns observed for the three split signals in **T.**

^1H NMR of **T:** 1.01 (singlet, 9 H), 4.82 (doublet of doublets, 1 H, J = 10, 1.7 Hz), 4.93 (doublet of doublets, 1 H, J = 18, 1.7 Hz), and 5.83 (doublet of doublets, 1 H, J = 18, 10 Hz) ppm


^1H NMR of **U:** 1.60 (singlet) ppm

14.71 Propose a structure consistent with each set of data.

a. A compound **X** (molecular formula $C_6H_{12}O_2$) gives a strong peak in its IR spectrum at 1740 cm^{-1}. The ^1H NMR spectrum of **X** shows only two singlets, including one at 3.5 ppm. The ^{13}C NMR spectrum is given below. Propose a structure for **X.**




b. A compound **Y** (molecular formula C_6H_{10}) gives four lines in its ^{13}C NMR spectrum (27, 30, 67, and 93 ppm), and the IR spectrum given here. Propose a structure for **Y.**




Challenge Problems

14.72 Reaction of unknown **A** with HCl forms chlorohydrin **B** as the major product. **A** shows no absorptions in its IR spectrum at 1700 cm^{-1} or 3600–3200 cm^{-1}, and gives the following ^1H NMR data: 1.4 (doublet, 3 H), 3.0 (quartet of doublets, 1 H), 3.5 (doublet, 1 H), 3.8 (singlet, 3 H), 6.9 (doublet, 2 H), and 7.2 (doublet, 2 H) ppm. (a) Propose a structure for **A,** including stereochemistry. (b) Explain why **B** is the major product in this reaction.




B

14.73 The ^{1}H NMR spectrum of *N,N*-dimethylformamide shows three singlets at 2.9, 3.0, and 8.0 ppm. Explain why the two CH_3 groups are not equivalent to each other, thus giving rise to two NMR signals.



N,N-dimethylformamide

14.74 18-Annulene shows two signals in its ^{1}H NMR spectrum, one at 8.9 (12 H) and one at –1.8 (6 H) ppm. Using a similar argument to that offered for the chemical shift of benzene protons, explain why both shielded and deshielded values are observed for 18-annulene.




18-annulene

14.75 Explain why the ^{13}C NMR spectrum of 3-methylbutan-2-ol shows five signals.

14.76 Because ^{31}P has an odd mass number, ^{31}P nuclei absorb in the NMR and, in many ways, these nuclei behave similarly to protons in NMR spectroscopy. With this in mind, explain why the ^{1}H NMR spectrum of methyl dimethylphosphonate, $CH_3PO(OCH_3)_2$, consists of two doublets at 1.5 and 3.7 ppm.


14.77 Cyclohex-2-enone has two protons on its carbon–carbon double bond (labeled H_a and H_b) and two protons on the carbon adjacent to the double bond (labeled H_c). (a) If J_{ab} = 11 Hz and J_{bc} = 4 Hz, sketch the splitting pattern observed for each proton on the sp^2 hybridized carbons. (b) Despite the fact that H_a is located adjacent to an electron-withdrawing C=O, its absorption occurs upfield from the signal due to H_b (6.0 vs. 7.0 ppm). Offer an explanation.



cyclohex-2-enone

15 Radical Reactions

15.1 Introduction
15.2 General features of radical reactions
15.3 Halogenation of alkanes
15.4 The mechanism of halogenation
15.5 Chlorination of other alkanes
15.6 Chlorination versus bromination
15.7 Halogenation as a tool in organic synthesis
15.8 The stereochemistry of halogenation reactions
15.9 Application: The ozone layer and CFCs
15.10 Radical halogenation at an allylic carbon
15.11 Application: Oxidation of unsaturated lipids
15.12 Application: Antioxidants
15.13 Radical addition reactions to double bonds
15.14 Polymers and polymerization



Polystyrene, an inexpensive polymer synthesized from the monomer styrene, $C_6H_5CH{=}CH_2$, is one of the six compounds that account for three-quarters of the synthetic polymers produced in the United States. The polystyrene foam used in packaging materials and drinking cups for hot beverages is called Styrofoam, a trademark of the Dow Chemical Company. Polystyrene is also used to form the housings of small kitchen appliances, televisions, computers, and CD cases. Although recycled polystyrene can be molded into trays and trash cans, the polystyrene used in food packaging and beverage cups is contaminated with food, making it difficult to clean and recycle. In Chapter 15, we learn about the synthesis of polymers like polystyrene.


A small but significant group of reactions involves the homolysis of nonpolar bonds to form highly reactive **radical intermediates.** Although they are unlike other organic reactions, radical transformations are important in many biological and industrial processes. The gases O_2 and NO (nitric oxide) are both radicals. Many oxidation reactions with O_2 involve radical intermediates, and biological processes mediated by NO such as blood clotting and neurotransmission may involve radicals. Many useful industrial products such as Styrofoam and polyethylene are prepared by radical processes.

In Chapter 15 we examine the cleavage of nonpolar bonds by radical reactions.

15.1 Introduction


Radicals were first discussed in Section 6.3.

- A *radical* is a reactive intermediate with a single unpaired electron, formed by homolysis of a covalent bond.




A radical contains an atom that does not have an octet of electrons, making it reactive and unstable. Radical processes involve single electrons, so half-headed arrows are used to show the movement of electrons. **One half-headed arrow is used for each electron.**

Carbon radicals are classified as **primary (1°), secondary (2°),** or **tertiary (3°)** by the number of R groups bonded to the carbon with the unpaired electron. A carbon radical is ***sp*^{2}** hybridized and **trigonal planar,** like sp^2 hybridized carbocations. The unhybridized *p* orbital contains the unpaired electron and extends above and below the trigonal planar carbon.




Bond dissociation energies for the cleavage of C−H bonds are used as a measure of radical stability. For example, two different radicals can be formed by cleavage of the C−H bonds in $CH_3CH_2CH_3$.




Cleavage of the stronger 1° C−H bond to form the 1° radical ($CH_3CH_2CH_2$·) requires more energy than cleavage of the weaker 2° C−H bond to form the 2° radical [$(CH_3)_2CH$·]—410 versus 397 kJ/mol. This makes the 2° radical more stable, because less energy is required for its

Figure 15.1
The relative stability of 1° and 2° carbon radicals



formation, as illustrated in Figure 15.1. Thus, **cleavage of the weaker bond forms the more stable radical,** a specific example of a general trend.

- **The stability of a radical increases as the number of alkyl groups bonded to the radical carbon increases.**




The **lower** the bond dissociation energy for a C–H bond, the **more stable** the resulting carbon radical.


Thus, a 3° radical is more stable than a 2° radical, and a 2° radical is more stable than a 1° radical. Increasing alkyl substitution increases radical stability in the same way it increases carbocation stability. Alkyl groups are more polarizable than hydrogen atoms, so they can more easily donate electron density to the electron-deficient carbon radical, thus increasing stability.

Unlike carbocations, however, **less stable radicals generally do *not* rearrange to more stable radicals.** This difference can be used to distinguish between reactions involving radical intermediates and those involving carbocations.

Problem 15.1 Classify each radical as 1°, 2°, or 3°.



Problem 15.2 Draw the most stable radical that can result from cleavage of a C–H bond in each molecule.



15.2 General Features of Radical Reactions

Radicals are formed from covalent bonds by adding energy in the form of **heat (Δ)** or **light (*hν*).** Some radical reactions are carried out in the presence of a **radical initiator,** a compound that contains an especially weak bond that serves as a source of radicals. **Peroxides,** compounds with the general structure **RO–OR,** are the most commonly used radical initiators. Heating a peroxide readily causes homolysis of the weak O–O bond, forming two RO· radicals.

15.2A Two Common Reactions of Radicals

Radicals undergo two main types of reactions: **they react with σ bonds,** and **they add to π bonds,** in both cases achieving an octet of electrons.

[1] Reaction of a Radical X· with a C–H Bond

A radical X· abstracts a hydrogen atom from a C–H σ bond to form H–X and a carbon radical. One electron from the C–H bond is used to form the new H–X bond, and the other electron in the C–H bond remains on carbon. The result is that the original radical X· is now surrounded by an octet of electrons, and a new radical is formed.



new radical

- One electron in H–X comes from the radical.
- One electron in H–X comes from the C–H bond.

This radical reaction is typically seen with the nonpolar C–H bonds of **alkanes,** which cannot react with polar or ionic electrophiles and nucleophiles.

[2] Reaction of a Radical X· with a C=C

A radical X· also adds to the π bond of a carbon–carbon double bond. One electron from the double bond is used to form a new C–X bond, and the other electron remains on the other carbon originally part of the double bond.



new radical


- One electron in C–X comes from the radical.
- One electron in C–X comes from the π bond.

Whenever a radical reacts with a stable single or double bond, a new radical is formed in the products.

Although the electron-rich double bond of an **alkene** reacts with electrophiles by ionic addition mechanisms, it also reacts with radicals because these reactive intermediates are also electron deficient.


15.2B Two Radicals Reacting with Each Other

A radical, once formed, rapidly reacts with whatever is available. Usually that means a stable σ or π bond. Occasionally, however, two radicals come into contact with each other, and they react to form a σ bond.



- One electron in X–X comes from each radical.

The reaction of a radical with oxygen, a diradical in its ground state electronic configuration, is another example of two radicals reacting with each other. In this case, the reaction of O_2 with X· forms a new radical, thus preventing X· from reacting with an organic substrate.



a diradical

Compounds that prevent radical reactions from occurring are called *radical inhibitors* or *radical scavengers.* Besides O_2, vitamin E and related compounds, discussed in Section 15.12, are radical scavengers, too. The fact that these compounds inhibit a reaction often suggests that the reaction occurs via radical intermediates.


Problem 15.3 Draw the products formed when a chlorine atom (Cl·) reacts with each species.

a. 

b. $CH_2=CH_2$ c. Cl· d. O_2


15.3 Halogenation of Alkanes

In the presence of light or heat, alkanes react with halogens to form alkyl halides. Halogenation is a **radical substitution reaction,** because a halogen atom X replaces a hydrogen via a mechanism that involves radical intermediates.




Halogenation of alkanes is useful only with Cl_2 and Br_2. Reaction with F_2 is too violent and reaction with I_2 is too slow to be useful. With an alkane that has more than one type of hydrogen atom, a mixture of alkyl halides may result (Reaction [2]).

When asked to draw the products of halogenation of an alkane, **draw the products of monohalogenation only,** unless specifically directed to do otherwise.



In these examples of halogenation, a halogen has replaced a single hydrogen atom on the alkane. Can the other hydrogen atoms be replaced, too? Figure 15.2 shows that when CH_4 is treated with excess Cl_2, all four hydrogen atoms can be successively replaced by Cl to form CCl_4. **Monohalogenation**—the substitution of a single H by X—can be achieved experimentally by adding halogen X_2 to an excess of alkane.


Figure 15.2
Complete halogenation of CH_4 using excess Cl_2



Sample Problem 15.1 Draw all the constitutional isomers formed by monohalogenation of $(CH_3)_2CHCH_2CH_3$ with Cl_2 and *hν*.

Solution

Substitute Cl for H on every carbon, and then check to see if any products are identical. The starting material has five C's, but replacement of one H atom on two C's gives the same product. Thus, **$(CH_3)_2CHCH_2CH_3$ affords four monochloro substitution products.**



Problem 15.4 Draw all constitutional isomers formed by monochlorination of each alkane.

a.  b.  c. 

Problem 15.5 Compounds **A** and **B** are isomers having molecular formula C_5H_{12}. Heating **A** with Cl_2 gives a single product of monohalogenation, whereas heating **B** under the same conditions forms three constitutional isomers. What are the structures of **A** and **B?**


15.4 The Mechanism of Halogenation

Unlike nucleophilic substitution, which proceeds by two different mechanisms depending on the starting material and reagent, all halogenation reactions of alkanes—regardless of the halogen and alkane used—proceed by the *same* mechanism. Three facts about halogenation suggest that the mechanism involves radical, not ionic, intermediates.

Fact	Explanation
[1] Light, heat, or added peroxide is necessary for the reaction.	• Light or heat provides the energy needed for homolytic bond cleavage to form radicals. Breaking the weak O–O bond of peroxides initiates radical reactions as well.
[2] O_2 inhibits the reaction.	• The diradical O_2 removes radicals from a reaction mixture, thus preventing reaction.
[3] No rearrangements are observed.	• Radicals do not rearrange.

15.4A The Steps of Radical Halogenation

The chlorination of cyclopentane illustrates the **three distinct parts of radical halogenation** (Mechanism 15.1):



cyclopentane chlorocyclopentane


- ***Initiation:*** **Two radicals are formed by homolysis of a σ bond and this begins the reaction.**
- ***Propagation:*** **A radical reacts with another reactant to form a new σ bond and another radical.**
- ***Termination:*** **Two radicals combine to form a stable bond. Removing radicals from the reaction mixture without generating any new radicals stops the reaction.**

Although initiation generates the Cl· radicals needed to begin the reaction, the propagation steps ([2] and [3]) form the two reaction products—chlorocyclopentane and HCl. Once the process has begun, propagation occurs over and over without the need for Step [1] to occur. **A mechanism such as radical halogenation that involves two or more repeating steps is called a *chain mechanism*.** Each propagation step involves a reactive radical abstracting an atom from a stable bond to form a new bond and another radical that continues the chain.

Usually a radical reacts with a stable bond to propagate the chain, but occasionally two radicals combine, and this reaction terminates the chain. Depending on the reaction and the reaction conditions, some radical chain mechanisms can repeat thousands of times before termination occurs.


Mechanism 15.1 Radical Halogenation of Alkanes

Part [1] Initiation



1 Bond cleavage forms two radicals. Homolysis of the weakest bond (Cl–Cl) requires light or heat and forms two chlorine radicals.


Part [2] Propagation



2 The Cl· radical abstracts a hydrogen from cyclopentane to form HCl (a reaction product) and a new carbon radical.

3 The carbon radical abstracts a chlorine atom from Cl_2 to form chlorocyclopentane (a reaction product) and Cl·. Because Cl· is a reactant in Step [2], Steps [2] and [3] can occur repeatedly without additional initiation (Step [1]).

Part [3] Termination



4 Termination of the chain occurs when any two radicals combine to form a bond.

Termination Step [4a] forms Cl_2, a reactant, whereas Step [4c] forms chlorocyclopentane, one of the reaction products. Termination Step [4b] forms **A,** which is neither a reactant nor a desired product. The formation of a small quantity of **A,** however, is evidence that radicals are formed in the reaction.

The most important steps of radical halogenation are those that lead to product formation—the propagation steps—so subsequent discussion of this reaction concentrates on these steps only.

Problem 15.6 Using Mechanism 15.1 as a guide, write the mechanism for the reaction of CH_4 with Br_2 to form CH_3Br and HBr. Classify each step as initiation, propagation, or termination.

15.4B Energy Changes During the Chlorination of Ethane

The chlorination of ethane illustrates how bond dissociation energies (Section 6.4) can be used to calculate $\Delta H°$ in chain propagation.






Figure 15.3
Energy changes in the propagation steps during the chlorination of ethane



As shown in Figure 15.3, chain propagation consists of the same two steps drawn in Mechanism 15.1: abstraction of a hydrogen atom to form $CH_3CH_2\cdot$ and HCl, followed by abstraction of a chlorine atom by $CH_3CH_2\cdot$ to form CH_3CH_2Cl and a chlorine radical ($Cl\cdot$). The $\Delta H°$ for each step is negative, making the overall $\Delta H°$ negative and the reaction exothermic. Because the transition state for the first propagation step is higher in energy than the transition state for the second propagation step, the **first step is rate-determining.** Both of these facts are illustrated in the energy diagram in Figure 15.4.

Problem 15.7 Calculate $\Delta H°$ for the rate-determining step of the reaction of CH_4 with I_2. Explain why this result illustrates that this reaction is extremely slow.


Figure 15.4
Energy diagram for the propagation steps in the chlorination of ethane



- Because radical halogenation consists of two propagation steps, the energy diagram has two energy barriers.
- The first step is rate-determining because its transition state is at higher energy.
- The reaction is exothermic because $\Delta H°_{overall}$ is negative.


15.5 Chlorination of Other Alkanes

Recall from Section 15.3 that the chlorination of $CH_3CH_2CH_3$ affords a 1:1 mixture of $CH_3CH_2CH_2Cl$ (formed by removal of a 1° hydrogen) and $(CH_3)_2CHCl$ (formed by removal of a 2° hydrogen).





$CH_3CH_2CH_3$ has six 1° hydrogen atoms and only two 2° hydrogens, so the expected product ratio of $CH_3CH_2CH_2Cl$ to $(CH_3)_2CHCl$ (assuming all hydrogens are *equally* reactive) is 3:1. Because the observed ratio is 1:1, however, the 2° C−H bonds must be more reactive; that is, **it must be easier to homolytically cleave a 2° C−H bond than a 1° C−H bond.** Recall from Section 15.2 that 2° C−H bonds are weaker than 1° C−H bonds. Thus,

- **The weaker the C−H bond, the more readily the hydrogen atom is removed in radical halogenation.**



When alkanes react with Cl_2, a mixture of products results, with more product formed by cleavage of the weaker C−H bond than you would expect on statistical grounds.

Problem 15.8 Which C−H bond in each compound is most readily broken during radical halogenation?


a.  b.  c. 

15.6 Chlorination Versus Bromination

Although alkanes undergo radical substitution reactions with both Cl_2 and Br_2, chlorination and bromination exhibit two important differences:

- **Chlorination is *faster* than bromination.**
- **Although chlorination is *unselective,* yielding a mixture of products, bromination is often *selective,* yielding one major product.**

For example, propane reacts rapidly with Cl_2 to form a 1:1 mixture of 1° and 2° alkyl chlorides. On the other hand, propane reacts with Br_2 much more slowly and forms 99% $(CH_3)_2CHBr$.




- **In bromination, the major (and sometimes exclusive) product results from cleavage of the *weakest* C–H bond.**


Sample Problem 15.2 Draw the major product formed when 3-ethylpentane is heated with Br_2.

Solution

Keep in mind: **the more substituted the carbon atom, the weaker the C–H bond.** The major bromination product in 3-ethylpentane is formed by cleavage of the sole **3° C–H bond,** its weakest C–H bond.



Problem 15.9 Draw the major product formed when each cycloalkane is heated with Br_2.




To explain the difference between chlorination and bromination, we return to the Hammond postulate (Section 7.15). The **rate-determining step in halogenation is the abstraction of a hydrogen atom by the halogen radical,** so we must compare these steps for bromination and chlorination. Keep in mind:

- **Transition states in endothermic reactions resemble the products. The more stable product is formed faster.**
- **Transition states in exothermic reactions resemble the starting materials. The relative stability of the products does not greatly affect the relative energy of the transition states, so a mixture of products often results.**

Bromination: $CH_3CH_2CH_3 + Br_2$


A bromine radical can abstract either a 1° or a 2° hydrogen from propane, generating either a 1° radical or a 2° radical. Calculating $\Delta H°$ using bond dissociation energies reveals that both reactions are **endothermic,** but **it takes less energy to form the more stable 2° radical.**



According to the Hammond postulate, the transition state of an endothermic reaction resembles the products, so the energy of activation to form the more stable 2° radical is lower and it is formed faster, as shown in the energy diagram in Figure 15.5. Because the 2° radical [$(CH_3)_2CH\cdot$] is converted to 2-bromopropane [$(CH_3)_2CHBr$] in the second propagation step, this **2° alkyl halide is the major product of bromination.**

- **Conclusion: Because the rate-determining step in bromination is endothermic, the more stable radical is formed faster, and often a single radical halogenation product predominates.**

Figure 15.5
Energy diagram for a selective endothermic reaction



- The transition state to form the less stable 1° radical ($CH_3CH_2CH_2\cdot$) is higher in energy than the transition state to form the more stable 2° radical [$(CH_3)_2CH\cdot$]. Thus, **the 2° radical is formed faster.**

Chlorination: $CH_3CH_2CH_3 + Cl_2$

A chlorine radical can also abstract either a 1° or a 2° hydrogen from propane, generating either a 1° radical or a 2° radical. Calculating $\Delta H°$ using bond dissociation energies reveals that both reactions are **exothermic.**





Figure 15.6
Energy diagram for a nonselective exothermic reaction



Because chlorination has an *exothermic* rate-determining step, the transition state to form both radicals resembles the same starting material, $CH_3CH_2CH_3$. As a result, the relative stability of the two radicals is much less important and both radicals are formed. An energy diagram for these processes is drawn in Figure 15.6. Because the 1° and 2° radicals are converted to 1-chloropropane ($CH_3CH_2CH_2Cl$) and 2-chloropropane [$(CH_3)_2CHCl$], respectively, in the second propagation step, **both alkyl halides are formed in chlorination.**

- **Conclusion: Because the rate-determining step in chlorination is exothermic, the transition state resembles the starting material, both radicals are formed, and a mixture of products results.**

Problem 15.10 Reaction of $(CH_3)_3CH$ with Cl_2 forms two products: $(CH_3)_2CHCH_2Cl$ (63%) and $(CH_3)_3CCl$ (37%). Why is the major product formed by cleavage of the stronger 1° C–H bond?

15.7 Halogenation as a Tool in Organic Synthesis

Halogenation is a useful tool because it adds a functional group to a previously unfunctionalized molecule, making an **alkyl halide.** These alkyl halides can then be converted to alkenes by elimination, and to alcohols and ethers by nucleophilic substitution.


Sample Problem 15.3 Show how cyclohexane can be converted to cyclohexene by a stepwise sequence.




Solution

There is no one-step method to convert an alkane to an alkene. A two-step method is needed:


[1] **Radical halogenation** produces an alkyl halide.




[2] **Elimination of HCl** with a strong base produces cyclohexene.



Problem 15.11 Synthesize each compound from $(CH_3)_3CH$.

a. 

b. 

c. 

Problem 15.12 Show all steps and reagents needed to convert cyclohexane into each compound: (a) the two enantiomers of *trans*-1,2-dibromocyclohexane; and (b) 1,2-epoxycyclohexane.

15.8 The Stereochemistry of Halogenation Reactions


The stereochemistry of a reaction product depends on whether the reaction occurs at a stereogenic center or at another atom, and whether a new stereogenic center is formed. The rules predicting the stereochemistry of reaction products are summarized in Table 15.1.

Table 15.1 Rules for Predicting the Stereochemistry of Reaction Products

Starting material	Result
Achiral	• An achiral starting material always gives either an achiral or a racemic product.
Chiral	• If a reaction does not occur at a stereogenic center, the configuration at a stereogenic center is retained in the product. • If a reaction occurs at a stereogenic center, we must know the mechanism to predict the stereochemistry of the product.


15.8A Halogenation of an Achiral Starting Material

Halogenation of the **achiral starting material $CH_3CH_2CH_2CH_3$** forms two constitutional isomers by replacement of either a 1° or 2° hydrogen.



- 1-Chlorobutane ($CH_3CH_2CH_2CH_2Cl$) has no stereogenic center, so it is an **achiral** compound.
- 2-Chlorobutane [$CH_3CH(Cl)CH_2CH_3$] has a new stereogenic center, so an **equal amount of two enantiomers** must form—**a racemic mixture.**


A racemic mixture results when a new stereogenic center is formed because the first propagation step generates a **planar, sp^2 hybridized radical.** Cl_2 then reacts with the planar radical from either the front or back side to form an equal amount of two enantiomers.



Thus, the achiral starting material butane forms an achiral product (1-chlorobutane) and a racemic mixture of two enantiomers [(*R*)- and (*S*)-2-chlorobutane].


15.8B Halogenation of a Chiral Starting Material

Let's now examine chlorination of the chiral starting material (*R*)-2-bromobutane at C2 and C3.



(*R*)-2-bromobutane

Chlorination at C2 occurs at the stereogenic center. Abstraction of a hydrogen atom at C2 forms a trigonal planar sp^2 hybridized radical that is now achiral. This achiral radical then reacts with Cl_2 from either side to form a new stereogenic center, resulting in an **equal amount of two enantiomers—a racemic mixture.**



planar achiral radical + HCl


attack from the front

attack from the back

enantiomers

- Radical halogenation reactions occur with racemization at a stereogenic center.

Chlorination at C3 does *not* occur at the stereogenic center, but it forms a new stereogenic center. Because no bond is broken to the stereogenic center at C2, **its configuration is retained** during the reaction. Abstraction of a hydrogen atom at C3 forms a trigonal planar sp^2 hybridized radical that still contains this stereogenic center. Reaction of the radical with Cl_2 from either side forms a new stereogenic center, so the products have two stereogenic centers: the configuration at C2 is the same in both compounds, but the configuration at C3 is different, making them **diastereomers.**



The configuration at C2 is **retained.** + HCl

attack from the front


attack from the back

diastereomers


Thus, four isomers are formed by chlorination of (*R*)-2-bromobutane at C2 and C3. Attack at the stereogenic center (C2) gives a product with one stereogenic center, resulting in a mixture of enantiomers. Attack at C3 forms a new stereogenic center, giving a mixture of diastereomers.


Problem 15.13 What products are formed from monochlorination of (*R*)-2-bromobutane at C1 and C4? Assign *R* and *S* designations to each stereogenic center.

Problem 15.14 Draw the monochlorination products formed when each compound is heated with Cl_2. Include the stereochemistry at any stereogenic center.

a. 

b. 

c. 


d. 

(Consider attack at C2 and C3 only.)

15.9 Application: The Ozone Layer and CFCs

The 1995 Nobel Prize in Chemistry was awarded to Mario Molina, Paul Crutzen, and F. Sherwood Rowland for their work in elucidating the interaction of ozone with CFCs.

Ozone is formed in the upper atmosphere by reaction of oxygen molecules with oxygen atoms. Ozone is also decomposed with sunlight back to these same two species. The overall result of these reactions is to convert high-energy ultraviolet light into heat.



Ozone is vital to life; it acts like a shield, protecting the earth's surface from destructive ultraviolet radiation. A decrease in ozone concentration in this protective layer would have some immediate consequences, including an increase in the incidence of skin cancer and eye cataracts. Other long-term effects include a reduced immune response, interference with photosynthesis in plants, and harmful effects on the growth of plankton, the mainstay of the ocean food chain.




Propane and butane are now used as propellants in spray cans in place of CFCs.

Current research suggests that **chlorofluorocarbons (CFCs)** are responsible for destroying ozone in the upper atmosphere. **CFCs** are simple halogen-containing organic compounds manufactured under the trade name Freons.




$CFCl_3$

trichlorofluoromethane
CFC 11
Freon 11



CF_2Cl_2

dichlorodifluoromethane
CFC 12
Freon 12




O_3 destruction is most severe in the region of the South Pole, where a large ozone hole is visible with satellite imaging.

CFCs are inert, odorless, and nontoxic, and they have been used as refrigerants, solvents, and aerosol propellants. Because CFCs are volatile and water insoluble, they readily escape into the upper atmosphere, where they are decomposed by high-energy sunlight to form radicals that destroy ozone by the radical chain mechanism shown in Figure 15.7.


The overall result is that O_3 is consumed as a reactant and O_2 molecules are formed. In this way, a small amount of CFC can destroy a large amount of O_3. These findings led to a ban on the use of CFCs in aerosol propellants in the United States in 1978 and to the phasing out of their use in refrigeration systems.

Newer alternatives to CFCs are **hydrochlorofluorocarbons (HCFCs)** and **hydrofluorocarbons (HFCs)** such as CH_2FCF_3. These compounds have many properties in common with CFCs, but they are largely decomposed by HO· before they reach the stratosphere and therefore they have little impact on stratospheric O_3.



CH_2FCF_3


HFC-134a



HFC-134a is decomposed before it reaches the stratosphere.

Problem 15.15 Nitric oxide, NO·, is another radical also thought to cause ozone destruction by a similar mechanism. One source of NO· in the stratosphere is supersonic aircraft whose jet engines convert small amounts of N_2 and O_2 to NO·. Write the propagation steps for the reaction of O_3 with NO·.

Figure 15.7
CFCs and the destruction of the ozone layer



Initiation: CFCs are decomposed by sunlight to form chlorine radicals.

$$CFCl_3 \xrightarrow{h\nu} \cdot CFCl_2 + \cdot\ddot{\underset{\cdot\cdot}{Cl}}:$$

Propagation: Ozone is destroyed by a chain reaction with radical intermediates.


$$\cdot\ddot{\underset{\cdot\cdot}{Cl}}: + O_3 \longrightarrow :\ddot{\underset{\cdot\cdot}{Cl}}-\ddot{\underset{\cdot\cdot}{O}}\cdot + O_2$$

$$:\ddot{\underset{\cdot\cdot}{Cl}}-\ddot{\underset{\cdot\cdot}{O}}\cdot + \cdot\ddot{\underset{\cdot\cdot}{O}}\cdot \longrightarrow \cdot\ddot{\underset{\cdot\cdot}{Cl}}: + O_2$$

- The chain reaction is initiated by homolysis of a C – Cl bond in $CFCl_3$.
- Propagation consists of two steps. Reaction of Cl· with O_3 forms chlorine monoxide (ClO·), which reacts with oxygen atoms to form O_2 and Cl·.

15.10 Radical Halogenation at an Allylic Carbon


Now let's examine radical halogenation at an ***allylic carbon*—the carbon adjacent to a double bond.** Homolysis of the allylic C – H bond of propene generates the **allyl radical,** which has an unpaired electron on the carbon adjacent to the double bond.



propene
allylic C–H (in red)


allyl radical

The bond dissociation energy for this process (364 kJ/mol) is even less than that for a 3° C – H bond (381 kJ/mol). Because the weaker the C – H bond, the more stable the resulting radical, an **allyl radical is more stable than a 3° radical,** and the following order of radical stability results:



The allyl radical is more stable than other radicals because two resonance structures can be drawn for it.




The position of the atoms and the σ bonds stays the same in drawing resonance structures. Resonance structures differ in the location of only π bonds and nonbonded electrons.



two resonance structures for the allyl radical — hybrid


- **The "true" structure of the allyl radical is a hybrid of the two resonance structures. In the hybrid, the π bond and the unpaired electron are delocalized.**
- **Delocalizing electron density lowers the energy of the hybrid, thus stabilizing the allyl radical.**

Problem 15.16 Draw a second resonance structure for each radical. Then draw the hybrid.


a. 
b. 
c. 
d. 

15.10A Selective Bromination at Allylic C–H Bonds

Because allylic C–H bonds are weaker than other sp^3 hybridized C–H bonds, the **allylic carbon can be selectively halogenated** by using *N*-bromosuccinimide (**NBS,** Section 10.15) in the presence of light or peroxides. Under these conditions only the allylic C–H bond in cyclohexene reacts to form an allylic halide.



N-bromosuccinimide
NBS




allylic C — allylic halide

NBS contains a weak N–Br bond that is homolytically cleaved with light to generate a bromine radical, initiating an allylic halogenation reaction. Propagation then consists of the usual two steps of radical halogenation as shown in Mechanism 15.2.

Mechanism 15.2 Allylic Bromination with NBS


Part [1] Initiation



NBS

1 Homolysis of the weak N–Br bond with light energy forms a Br· radical that initiates radical halogenation.

Part [2] Propagation




allylic radical — (from NBS)

2 The Br· radical abstracts an allylic H to afford an allylic radical. (Only one resonance structure is drawn.)


3 The allylic radical reacts with Br_2 to form the allylic halide. The radical Br· formed in Step [3] can now react in Step [2], so Steps [2] and [3] can repeatedly occur without additional initiation.

Besides acting as a source of Br· to initiate the reaction, NBS generates a low concentration of Br_2 needed in the second chain propagation step (Step [3] of the mechanism). The HBr formed in Step [2] reacts with NBS to form Br_2, which is then used for halogenation in Step [3] of the mechanism.




A **low concentration of Br_2** (from NBS) **favors allylic substitution** (over addition) in part because bromine is needed for only *one* step of the mechanism. When Br_2 adds to a double bond, a low Br_2 concentration would first form a low concentration of bridged bromonium ion (Section 10.13), which must then react with more bromine (in the form of Br^-) in a second step to form a dibromide. **If concentrations of both intermediates—bromonium ion and Br^-—are low, the overall rate of addition is very slow.**

Thus, an alkene with allylic C−H bonds undergoes two different reactions depending on the reaction conditions.




- Treatment of cyclohexene with Br_2 (in an organic solvent like CCl_4) leads to **addition** via **ionic intermediates** (Section 10.13).
- Treatment of cyclohexene with NBS (+ *h*ν or ROOR) leads to **allylic substitution,** via **radical intermediates.**

Problem 15.17 Draw the products of each reaction.





15.10B Product Mixtures in Allylic Halogenation

Halogenation at an allylic carbon often results in a mixture of products. For example, bromination of 3-methylbut-1-ene under radical conditions forms a mixture of 3-bromo-3-methylbut-1-ene and 1-bromo-3-methylbut-2-ene.



A mixture is obtained because the reaction proceeds by way of a **resonance-stabilized radical.** Abstraction of an allylic hydrogen from the alkene with a Br· radical (from NBS) forms an allylic radical for which **two different Lewis structures** can be drawn.






As a result, two different C atoms have partial radical character (indicated by δ·), so that Br_2 reacts at two different sites and two allylic halides are formed.


- **Whenever two different resonance structures can be drawn for an allylic radical, two different allylic halides are formed by radical substitution.**

Sample Problem 15.4 Draw the products formed when **A** is treated with NBS + *hν*.






Solution

Hydrogen abstraction at the allylic C forms a resonance-stabilized radical (with two different resonance structures) that reacts with Br_2 to form two constitutional isomers as products.




two constitutional isomers


Problem 15.18 Draw all constitutional isomers formed when each alkene is treated with NBS + *hν*.


a.  b.  c. 

Problem 15.19 Draw the structure of the four allylic halides formed when 3-methylcyclohexene undergoes allylic halogenation with NBS + *hν*.

Problem 15.20 Which compounds can be prepared in good yield by allylic halogenation of an alkene?

a. 

b. 

c. 


15.11 Application: Oxidation of Unsaturated Lipids

Oils—triacylglycerols having one or more sites of unsaturation in their long carbon chains—are susceptible to oxidation at their allylic carbon atoms. Oxidation occurs by way of a radical chain mechanism, as shown in Figure 15.8.

- **Step [1]** Oxygen in the air abstracts an allylic hydrogen atom to form an allylic radical because the allylic C−H bond is weaker than the other C−H bonds.
- **Step [2]** The allylic radical reacts with another molecule of O_2 to form a peroxy radical.
- **Step [3]** The peroxy radical abstracts an allylic hydrogen from another lipid molecule to form a hydroperoxide and another allylic radical that continues the chain. Steps [2] and [3] can repeat again and again until some other radical terminates the chain.

The hydroperoxides formed by this process are unstable and decompose to other oxidation products, many of which have a disagreeable odor and taste. **This process turns an oil rancid. Unsaturated lipids are more easily oxidized than saturated ones** because they contain weak allylic C−H bonds that are readily cleaved in Step [1] of this reaction, forming resonance-stabilized allylic radicals.


Figure 15.8
The oxidation of unsaturated lipids with O_2



- Oxidation is shown at one allylic carbon only. Reaction at the other labeled allylic carbon is also possible.

Because saturated fats have no double bonds and thus no weak allylic C–H bonds, they are much less susceptible to air oxidation, resulting in increased shelf life of products containing them.

Problem 15.21 Which C–H bond is most readily cleaved in linoleic acid? Draw all possible resonance structures for the resulting radical. Draw all the hydroperoxides formed by reaction of this resonance-stabilized radical with O_2.






The purported health benefits of antioxidants have made them a popular component in anti-aging formulations.


15.12 Application: Antioxidants

An *antioxidant* is a compound that stops an oxidation reaction from occurring.

- Naturally occurring antioxidants such as **vitamin E** prevent radical reactions that can cause cell damage.
- Synthetic antioxidants such as **BHT**—**b**utylated **h**ydroxy **t**oluene—are added to packaged and prepared foods to prevent oxidation and spoilage.



vitamin E




BHT
(butylated hydroxy toluene)



Hazelnuts, almonds, and many other types of nuts are an excellent source of the natural antioxidant vitamin E.

Vitamin E and BHT are radical inhibitors, so they terminate radical chain mechanisms by reacting with radicals. How do they trap radicals? Both vitamin E and BHT use a hydroxy group bonded to a benzene ring—a general structure called a **phenol.**


Radicals (R·) abstract a hydrogen atom from the OH group of an antioxidant, forming a new resonance-stabilized radical. **This new radical does not participate in chain propagation,** but rather terminates the chain and halts the oxidation process. All phenols (including vitamin E and BHT) inhibit oxidation by this radical process.



R· = a general organic radical

The many nonpolar C−C and C−H bonds of vitamin E make it fat soluble, and thus it dissolves in the nonpolar interior of the cell membrane, where it is thought to inhibit the oxidation of the unsaturated fatty acid residues in the phospholipids. Oxidative damage to lipids in cells via radical mechanisms is thought to play an important role in the aging process. For this reason, many anti-aging formulas with antioxidants like vitamin E are now popular consumer products.

Problem 15.22 Rosmarinic acid is an antioxidant isolated from rosemary. Draw resonance structures for the radical that results from removal of the labeled H atom in rosmarinic acid.






Rosemary extracts contain rosmarinic acid, an antioxidant that helps prevent the oxidation of unsaturated vegetable oils.

15.13 Radical Addition Reactions to Double Bonds


We now turn our attention to the second common reaction of radicals, addition to double bonds. Because an alkene contains an electron-rich, easily broken π bond, it reacts with an electron-deficient radical.




Radicals react with alkenes via a radical chain mechanism that consists of initiation, propagation, and termination steps analogous to those discussed previously for radical substitution.

15.13A Addition of HBr

HBr adds to alkenes to form alkyl bromides in the presence of light, heat, or peroxides.






The regioselectivity of addition to an unsymmetrical alkene is *different* from the addition of HBr without added light, heat, or peroxides.



- HBr addition to propene *without* added light, heat, or peroxides gives 2-bromopropane: the **H atom is added to the less substituted carbon.** This reaction occurs via **carbocation** intermediates (Section 10.10).
- HBr addition to propene *with* added light, heat, or peroxides gives 1-bromopropane: the **Br atom is added to the less substituted carbon.** This reaction occurs via **radical** intermediates.

Problem 15.23 Draw the product(s) formed when each alkene is treated with either [1] HBr alone; or [2] HBr in the presence of peroxides.


a.  b.  c. 

15.13B The Mechanism of the Radical Addition of HBr to an Alkene

In the presence of added light, heat, or peroxides, HBr addition to an alkene forms radical intermediates, and like other radical reactions, proceeds by a mechanism with three distinct parts: initiation, propagation, and termination. Mechanism 15.3 is written for the reaction of $CH_3CH{=}CH_2$ with HBr and ROOR to form $CH_3CH_2CH_2Br$.


Mechanism 15.3 Radical Addition of HBr to an Alkene

Part [1] Initiation



1 – 2 Initiation with ROOR occurs in two steps—homolysis of the weak O–O bond and abstraction of H to form a bromine radical.

Part [2] Propagation



3 Addition of Br· to the terminal carbon forms a 2° radical.


4 Abstraction of H from HBr forms a new C–H bond and a bromine radical, so Steps [3] and [4] can occur repeatedly.

Part [3] Termination




5 Termination of the chain occurs when any two radicals combine to form a bond.

The first propagation step (Step [3] of the mechanism, the addition of Br· to the double bond) is worthy of note. With propene there are two possible paths for this step, depending on which carbon atom of the double bond forms the new bond to bromine. Path [A] forms a less stable 1° radical whereas Path [B] forms a more stable 2° radical. **The more stable 2° radical forms faster,** so Path [B] is preferred.



The mechanism also illustrates why the regioselectivity of HBr addition is different depending on the reaction conditions. In both reactions, H and Br add to the double bond, but the *order* of addition depends on the mechanism.



- In radical addition (HBr with added light, heat, or ROOR), *Br· adds first* to generate the more stable radical.
- In ionic addition (HBr alone), *H^+ adds first* to generate the more stable carbocation.

Problem 15.24 When HBr adds to $(CH_3)_2C{=}CH_2$ under radical conditions, two radicals are possible products in the first step of chain propagation. Draw the structure of both radicals and indicate which one is formed. Then draw the preferred product from HBr addition under radical conditions.

Problem 15.25 What reagents are needed to convert 1-ethylcyclohexene into (a) 1-bromo-2-ethylcyclohexane; (b) 1-bromo-1-ethylcyclohexane; (c) 1,2-dibromo-1-ethylcyclohexane?

15.13C Energy Changes in the Radical Addition of HBr

The energy changes during propagation in the radical addition of HBr to $CH_2{=}CH_2$ can be calculated from bond dissociation energies, as shown in Figure 15.9.




Figure 15.9
Energy changes during the propagation steps: $CH_2{=}CH_2 + HBr \rightarrow CH_3CH_2Br$

Both propagation steps for the addition of HBr are exothermic, so propagation is exothermic (energetically favorable) overall. For the addition of HCl or HI, however, one of the chain-propagating steps is quite endothermic, and thus too difficult to be part of a repeating chain mechanism. Thus, **HBr adds to alkenes under radical conditions, but HCl and HI do not.**

15.14 Polymers and Polymerization

Polymers*—large molecules made up of repeating units of smaller molecules called *monomers—include such biologically important compounds as proteins and carbohydrates. They also include such industrially important plastics as polyethylene, poly(vinyl chloride) (PVC), and polystyrene.




HDPE (high-density polyethylene) and **LDPE** (low-density polyethylene) are two common types of polyethylene prepared under different reaction conditions and having different physical properties. HDPE is opaque and rigid, and is used in milk containers and water jugs. LDPE is less opaque and more flexible, and is used in plastic bags and electrical insulation. Products containing HDPE and LDPE (and other plastics) are often labeled with a symbol indicating recycling ease: the lower the number, the easier to recycle.

15.14A Synthetic Polymers

Many synthetic polymers—that is, those synthesized in the lab—are among the most widely used organic compounds in modern society. Although some synthetic polymers resemble natural substances, many have different and unusual properties that make them more useful than naturally occurring materials. Soft drink bottles, plastic bags, food wrap, compact discs, Teflon, and Styrofoam are all made of synthetic polymers. In this section we examine polymers derived from alkene monomers. Chapter 30 is devoted to a detailed discussion of the synthesis and properties of several different types of synthetic polymers.

- *Polymerization* is the joining together of monomers to make polymers.


For example, joining **ethylene monomers** together forms the polymer **polyethylene,** a plastic used in milk containers and sandwich bags.



ethylene monomers






polyethylene polymer
new bonds shown in red

Many ethylene derivatives having the general structure $\mathbf{CH_2{=}CHZ}$ are also used as monomers for polymerization. The identity of Z affects the physical properties of the resulting polymer, making some polymers more suitable for one consumer product (e.g., plastic bags or food wrap) than another (e.g., soft drink bottles or compact discs). Polymerization of $CH_2{=}CHZ$ usually affords polymers with the Z groups on every other carbon atom in the chain. Table 15.2 lists some common monomers and polymers used in medicine or dentistry.



new bonds shown in red


Table 15.2 Common Monomers and Polymers Used in Medicine and Dentistry

Monomer	Polymer	Consumer product
 vinyl chloride	 **poly(vinyl chloride) PVC**	 PVC blood bags and tubing
 propene	 **polypropylene**	 polypropylene syringes
 tetrafluoroethylene	 **polytetrafluoroethylene Teflon**	 dental floss

Sample Problem 15.5 What polymer is formed when $CH_2=CHCO_2H$ (acrylic acid) is polymerized? The resulting polymer, poly(acrylic acid), is used in disposable diapers because it absorbs 30 times its weight in water.

Solution


Draw three or more alkene monomers, break one bond of each double bond, and join the alkenes together with single bonds. With unsymmetrical alkenes, substituents are bonded to every other carbon.



Join a C labeled in blue with a C labeled in red.

poly(acrylic acid)

Problem 15.26 (a) Draw the structure of polystyrene, the chapter-opening molecule, which is formed by polymerizing the monomer styrene, $C_6H_5CH=CH_2$. (b) What monomer is used to form poly(vinyl acetate), a polymer used in paints and adhesives?



poly(vinyl acetate)


15.14B Radical Polymerization

The alkene monomers used in polymerization are prepared from petroleum.

The polymers described in Section 15.14A are prepared by polymerization of alkene monomers by **adding a radical to a π bond.** The mechanism resembles the radical addition of HBr to an alkene, except that a **carbon radical rather than a bromine atom is added to the double bond.** Mechanism 15.4 is written with the general monomer $CH_2{=}CHZ$, and again has three parts: initiation, propagation, and termination.


Mechanism 15.4 Radical Polymerization of $CH_2{=}CHZ$

Part [1] Initiation



1 – 2 Initiation with ROOR occurs in two steps—homolysis of the weak O–O bond and addition of RO· to the alkene to form a carbon radical.

Part [2] Propagation



3 Chain propagation consists of a single step. The carbon radical adds to another alkene to form a new C–C bond and another carbon radical. Addition forms the radical with the unpaired electron on the atom with the Z substituent.


Part [3] Termination



4 Termination of the chain occurs when any two radicals combine to form a bond.

In radical polymerization, the more substituted radical always adds to the less substituted end of the monomer, a process called **head-to-tail polymerization.**

The **more substituted radical** adds to the **less substituted end** of the double bond.



The new radical is always located on the C bonded to Z.

Problem 15.27 Draw the steps of the mechanism that converts vinyl chloride ($CH_2{=}CHCl$) into poly(vinyl chloride).

KEY CONCEPTS


Radical Reactions

General Features of Radicals

- A radical is a reactive intermediate with a single unpaired electron (15.1).
- A carbon radical is sp^2 hybridized and trigonal planar (15.1).
- The stability of a radical increases as the number of C atoms bonded to the radical carbon increases (15.1).
- Allylic radicals are stabilized by resonance, making them more stable than 3° radicals (15.10).


Radical Reactions

[1] Halogenation of alkanes (15.4)




- The reaction follows a radical chain mechanism.
- The weaker the C – H bond, the more readily the hydrogen is replaced by X.
- Chlorination is faster and less selective than bromination (15.6).
- Radical substitution at a stereogenic center results in racemization (15.8).

[2] Allylic halogenation (15.10)




- The reaction follows a radical chain mechanism.

[3] Radical addition of HBr to an alkene (15.13)



- A radical addition mechanism is followed.
- Br bonds to the less substituted carbon atom to form the more substituted, more stable radical.

[4] Radical polymerization of alkenes (15.14)




- A radical addition mechanism is followed.

PROBLEMS


Problems Using Three-Dimensional Models

15.28 (a) Draw all constitutional isomers formed by monochlorination of each alkane with Cl_2 and *hν*. (b) Draw the major monobromination product formed by heating each alkane with Br_2.



A **B**


15.29 Draw all resonance structures of the radical that results from abstraction of a hydrogen atom from the antioxidant BHA (**b**utylated **h**ydroxy **a**nisole).



BHA

Radicals and Bond Strength


15.30 With reference to the indicated C – H bonds in 2-methylbutane:




2-methylbutane

a. Rank the C – H bonds in order of increasing bond strength.
b. Draw the radical resulting from cleavage of each C – H bond, and classify it as 1°, 2°, or 3°.
c. Rank the radicals in order of increasing stability.
d. Rank the C – H bonds in order of increasing ease of H abstraction in a radical halogenation reaction.

15.31 Rank the following radicals in order of increasing stability.




15.32 Why is a benzylic C – H bond (labeled in red) unusually weak?






Halogenation of Alkanes




15.33 Rank the indicated hydrogen atoms in order of increasing ease of abstraction in a radical halogenation reaction.



15.34 Draw all constitutional isomers formed by monochlorination of each alkane with Cl_2 and $h\nu$.


a. 
b. 
c. 


15.35 What is the major monobromination product formed by heating each alkane with Br_2?


a. 
b. 
c. 

15.36 Five isomeric alkanes (**A**–**E**) having the molecular formula C_6H_{14} are each treated with $Cl_2 + h\nu$ to give alkyl halides having molecular formula $C_6H_{13}Cl$. **A** yields five constitutional isomers. **B** yields four constitutional isomers. **C** yields two constitutional isomers. **D** yields three constitutional isomers, two of which possess stereogenic centers. **E** yields three constitutional isomers, only one of which possesses a stereogenic center. Identify the structures of **A**–**E.**


15.37 What alkane is needed to make each alkyl halide by radical halogenation?


a. 


b. 

c. 


15.38 Which alkyl halides can be prepared in good yield by radical halogenation of an alkane?

a. 


b. 

c. 

15.39 Draw the products of radical chlorination and bromination of each compound. For which compounds is a single constitutional isomer formed for both reactions? What must be true about the structure of a reactant for both reactions to form a single product?


a. 
b. 
c. 

15.40 Explain why radical bromination of *p*-xylene forms **C** rather than **D.**



p-xylene **C** **D**
NOT formed


15.41 a. What product(s) (excluding stereoisomers) are formed when **Y** is heated with Cl_2?
b. What product(s) (excluding stereoisomers) are formed when **Y** is heated with Br_2?
c. What steps are needed to convert **Y** to the alkene **Z?**



Y **Z**


Resonance

15.42 Draw resonance structures for each radical.




Allylic Halogenation

15.43 Draw the products formed when each alkene is treated with NBS + *h*ν.




15.44 Draw all constitutional isomers formed when **X** is treated with NBS + *h*ν.



X

Reactions

15.45 Draw the organic products formed in each reaction.



15.46 What reagents are needed to convert cyclopentene into (a) bromocyclopentane; (b) *trans*-1,2-dibromocyclopentane; (c) 3-bromocyclopentene?


15.47 Treatment of a hydrocarbon **A** (molecular formula C_9H_{18}) with Br_2 in the presence of light forms alkyl halides **B** and **C,** both having molecular formula $C_9H_{17}Br$. Reaction of either **B** or **C** with $KOC(CH_3)_3$ forms compound **D** (C_9H_{16}) as the major product. Ozonolysis of **D** forms cyclohexanone and acetone. Identify the structures of **A–D.**



cyclohexanone acetone

Stereochemistry and Reactions

15.48 Draw the products formed in each reaction and include the stereochemistry around any stereogenic centers.




15.49 (a) Draw all stereoisomers of molecular formula $C_5H_{10}Cl_2$ formed when (*R*)-2-chloropentane is heated with Cl_2. (b) Assuming that products having different physical properties can be separated into fractions by some physical method (such as fractional distillation), how many different fractions would be obtained? (c) Which of these fractions would be optically active?

15.50 Draw all the monochlorination products that are formed from (*S,S*)-1,2-dimethylcyclopropane.



(*S,S*)-1,2-dimethylcyclopropane

15.51 Draw the six products (including stereoisomers) formed when **A** is treated with NBS + *h*ν.



A


15.52 (a) Draw the products (including stereoisomers) formed when 2-methylhex-2-ene is treated with HBr in the presence of peroxides. (b) Draw the products (including stereoisomers) formed when (*S*)-2,4-dimethylhex-2-ene is treated with HBr and peroxides under similar conditions.

Mechanisms

15.53 Consider the following bromination: $(CH_3)_3CH + Br_2 \xrightarrow{\Delta} (CH_3)_3CBr + HBr$.

a. Calculate $\Delta H°$ for this reaction by using the bond dissociation energies in Table 6.2.
b. Draw out a stepwise mechanism for the reaction, including the initiation, propagation, and termination steps.
c. Calculate $\Delta H°$ for each propagation step.
d. Draw an energy diagram for the propagation steps.
e. Draw the structure of the transition state of each propagation step.

15.54 Draw a stepwise mechanism for the following reaction.




15.55 Although CH_4 reacts with Cl_2 to form CH_3Cl and HCl, the corresponding reaction of CH_4 with I_2 does not occur at an appreciable rate, even though the I – I bond is much weaker than the Cl – Cl bond. Explain why this is so.

15.56 When 3,3-dimethylbut-1-ene is treated with HBr alone, the major product is 2-bromo-2,3-dimethylbutane. When the same alkene is treated with HBr and peroxide, the sole product is 1-bromo-3,3-dimethylbutane. Explain these results by referring to the mechanisms.

Synthesis




15.57 Devise a synthesis of each compound from cyclopentane and any other required organic or inorganic reagents.



15.58 Devise a synthesis of 1-methylcyclohexene oxide from methylcyclohexane. You may use any other required organic or inorganic reagents.

15.59 Devise a synthesis of $CH_3CH_2CH_2CH_2Br$ from HC≡CH. You may use any other required organic or inorganic reagents.

15.60 Devise a synthesis of each compound using CH_3CH_3 as the only source of carbon atoms. You may use any other required organic or inorganic reagents.


a. HC≡CH b.  c.  d. 

15.61 Devise a synthesis of $OHC(CH_2)_4CHO$ from cyclohexane using any required organic or inorganic reagents.

15.62 Devise a synthesis of hexane-2,3-diol from propane as the only source of carbon atoms. You may use any other required organic or inorganic reagents.

Radical Oxidation Reactions

15.63 As described in Section 9.16, the leukotrienes, important components in the asthmatic response, are synthesized from arachidonic acid via the hydroperoxide 5-HPETE. Write a stepwise mechanism for the conversion of arachidonic acid to 5-HPETE with O_2.



arachidonic acid 5-HPETE leukotriene C_4

15.64 Ethers are oxidized with O_2 to form hydroperoxides that decompose violently when heated. Draw a stepwise mechanism for this reaction.




unstable hydroperoxide

15.65 (a) Ignoring stereoisomers, what two allylic hydroperoxides are formed by the oxidation of hex-1-ene with O_2? (b) Draw a stepwise mechanism that shows how these hydroperoxides are formed.


Antioxidants

15.66 Resveratrol is an antioxidant found in the skin of red grapes. Its anticancer, anti-inflammatory, and various cardiovascular effects are under active investigation. (a) Draw all resonance structures for the radical that results from homolysis of the OH bond shown in red. (b) Explain why homolysis of this OH bond is preferred to homolysis of either OH bond in the other benzene ring.



resveratrol

15.67 In cells, vitamin C exists largely as its conjugate base **X. X** is an antioxidant because radicals formed in oxidation processes abstract the labeled H atom, forming a new radical that halts oxidation. Draw the structure of the radical formed by H abstraction, and explain why this H atom is most easily removed.




vitamin C **X**

Polymers and Polymerization


15.68 What monomer is needed to form each polymer?

a.



polyisobutylene
(used to make basketballs)


b.




poly(ethyl acrylate)
(used in latex paints)

15.69 (a) Hard contact lenses, which first became popular in the 1960s, were made by polymerizing methyl methacrylate [$CH_2=C(CH_3)CO_2CH_3$] to form poly(methyl methacrylate) (PMMA). Draw the structure of PMMA. (b) More comfortable softer contact lenses introduced in the 1970s were made by polymerizing hydroxyethyl methacrylate [$CH_2=C(CH_3)CO_2CH_2CH_2OH$] to form poly(hydroxyethyl methacrylate) (poly-HEMA). Draw the structure of poly-HEMA. Since neither polymer allows oxygen from the air to pass through to the retina, newer contact lenses that are both comfortable and oxygen-permeable have now been developed.

15.70 Explain why polystyrene is much more readily oxidized by O_2 in the air than polyethylene is. Which H's in polystyrene are most easily abstracted and why?




polystyrene




polyethylene

15.71 Draw a stepwise mechanism for the following polymerization reaction.



15.72 As we will learn in Chapter 30, styrene derivatives such as **A** can be polymerized by way of cationic rather than radical intermediates. Cationic polymerization is an example of electrophilic addition to an alkene involving carbocations.



A

a. Draw a short segment of the polymer formed by the polymerization of **A.**
b. Why does **A** react faster than styrene ($C_6H_5CH=CH_2$) in a cationic polymerization?


15.73 When two monomers (**X** and **Y**) are polymerized together, a copolymer results. An alternating copolymer is formed when the two monomers **X** and **Y** alternate regularly in the polymer chain. Draw the structure of the alternating copolymer formed when the two monomers, $CH_2=CCl_2$ and $CH_2=CHC_6H_5$, are polymerized together.

Spectroscopy

15.74 **A** and **B,** isomers of molecular formula $C_3H_5Cl_3$, are formed by the radical chlorination of a dihalide **C** of molecular formula $C_3H_6Cl_2$.


a. Identify the structures of **A** and **B** from the following ^1H NMR data:
   Compound **A:** singlet at 2.23 and singlet at 4.04 ppm
   Compound **B:** doublet at 1.69, multiplet at 4.34, and doublet at 5.85 ppm
b. What is the structure of **C?**

15.75 Identify the structure of a minor product formed from the radical chlorination of propane, which has molecular formula $C_3H_6Cl_2$ and exhibits the given ^1H NMR spectrum.




Challenge Problems

15.76 Draw a stepwise mechanism for the following addition reaction to an alkene.



15.77 The triphenylmethyl radical is an unusual persistent radical present in solution in equilibrium with its dimer. For 70 years the dimer was thought to be hexaphenylethane, but in 1970, NMR data showed it to be **A.**



triphenylmethyl radical hexaphenylethane **A**

a. Why is the triphenylmethyl radical more stable than most other radicals?
b. Use curved arrow notation to show how two triphenylmethyl radicals dimerize to form **A.**
c. Propose a reason for the formation of **A** rather than hexaphenylethane.
d. How could ^1H and ^{13}C NMR spectroscopy be used to distinguish between hexaphenylethane and **A?**

15.78 In the presence of a radical initiator (Z·), tributyltin hydride (R_3SnH, R = $CH_3CH_2CH_2CH_2$) reduces alkyl halides to alkanes: R'X + $R_3SnH \rightarrow$ R'H + R_3SnX. The mechanism consists of a radical chain process with an intermediate tin radical:


Initiation: $R_3SnH + Z\cdot \longrightarrow R_3Sn\cdot + HZ$

Propagation:


$R'{-}Br + R_3Sn\cdot \longrightarrow R'\cdot + R_3SnBr$

$R'\cdot + R_3SnH \longrightarrow R'{-}H + R_3Sn\cdot$

This reaction has been employed in many radical cyclization reactions. Draw a stepwise mechanism for the following reaction.




15.79 $PGF_{2\alpha}$ (Section 4.15) is synthesized in cells from arachidonic acid ($C_{20}H_{32}O_2$) using a cyclooxygenase enzyme that catalyzes a multistep radical pathway. Part of this process involves the conversion of radical **A** to PGG_2, an unstable intermediate, which is then transformed to $PGF_{2\alpha}$ and other prostaglandins. Draw a stepwise mechanism for the conversion of **A** to PGG_2. (Hint: The mechanism begins with radical addition to a carbon–carbon double bond to form a resonance-stabilized radical.)



A PGG_2 unstable intermediate $PGF_{2\alpha}$

16 Conjugation, Resonance, and Dienes

16.1 Conjugation
16.2 Resonance and allylic carbocations
16.3 Common examples of resonance
16.4 The resonance hybrid
16.5 Electron delocalization, hybridization, and geometry
16.6 Conjugated dienes
16.7 Interesting dienes and polyenes
16.8 The carbon–carbon σ bond length in buta-1,3-diene
16.9 Stability of conjugated dienes
16.10 Electrophilic addition: 1,2- versus 1,4-addition
16.11 Kinetic versus thermodynamic products
16.12 The Diels–Alder reaction
16.13 Specific rules governing the Diels–Alder reaction
16.14 Other facts about the Diels–Alder reaction
16.15 Conjugated dienes and ultraviolet light



Morphine is an analgesic and narcotic isolated from the opium poppy *Papaver somniferum*. Opium has been widely used as a recreational drug and pain-killing remedy for centuries, and poppy seed tea, which contains morphine, was used as a folk remedy in parts of England until World War II. A key step in a laboratory synthesis of morphine involves the Diels–Alder reaction, a powerful reaction of conjugated dienes discussed in Chapter 16.

Chapter 16 is the first of three chapters that discuss the chemistry of conjugated molecules—molecules with overlapping *p* orbitals on three or more adjacent atoms. Chapter 16 focuses mainly on acyclic conjugated compounds, whereas Chapters 17 and 18 discuss the chemistry of benzene and related compounds that have a *p* orbital on every atom in a ring.

Much of Chapter 16 is devoted to the properties and reactions of 1,3-dienes. To understand these compounds, however, we must first learn about the consequences of having *p* orbitals on three or more adjacent atoms. Because the ability to draw resonance structures is also central to mastering this material, the key aspects of resonance theory are presented in detail.

16.1 Conjugation


***Conjugation* occurs whenever *p* orbitals can overlap on three or more adjacent atoms.** Two common conjugated systems are 1,3-dienes and allylic carbocations.




1,3-diene allylic carbocation

16.1A 1,3-Dienes

1,3-Dienes such as buta-1,3-diene contain two carbon–carbon double bonds joined by a single σ bond. Each carbon atom of a 1,3-diene is bonded to three other atoms and has no nonbonded electron pairs, so each carbon atom is sp^2 hybridized and has one *p* orbital containing an electron. **The four *p* orbitals on adjacent atoms make a 1,3-diene a conjugated system.**




buta-1,3-diene
one σ bond between the double bonds



four adjacent *p* orbitals
Each C is sp^2 hybridized and has a *p* orbital containing one electron.

What is special about conjugation? Having three or more *p* orbitals on adjacent atoms allows *p* orbitals to overlap and electrons to delocalize.

The electron density in the two π bonds is delocalized.



- **When *p* orbitals overlap, the electron density in each of the π bonds is spread out over a larger volume, thus lowering the energy of the molecule and making it more stable.**

Conjugation makes buta-1,3-diene inherently different from penta-1,4-diene, a compound having two double bonds separated by more than one σ bond. The π bonds in penta-1,4-diene are too far apart to be conjugated.

A conjugated diene



buta-1,3-diene

The electrons in the π bonds are **delocalized.**

An isolated diene




penta-1,4-diene

The electrons in the π bonds are **localized.**

Penta-1,4-diene is an **isolated diene.** The electron density in each π bond of an isolated diene is *localized* between two carbon atoms. In buta-1,3-diene, however, the electron density of both π bonds is *delocalized* over the four atoms of the diene. Electrostatic potential maps in Figure 16.1 clearly indicate the difference between these localized and delocalized π bonds.



Figure 16.1
Electrostatic potential plots for a conjugated and an isolated diene



buta-1,3-diene
The red electron-rich region is spread over four adjacent atoms.


penta-1,4-diene
The red electron-rich regions are localized in the π bonds on the two ends of the molecule.

Problem 16.1 Classify each diene as isolated or conjugated.


a. 
b. 
c. 
d. 

16.1B Allylic Carbocations

The **allyl carbocation** is another example of a conjugated system. The three carbon atoms of the allyl carbocation—the positively charged carbon atom and the two that form the double bond—are sp^2 hybridized and have an unhybridized *p* orbital. The *p* orbitals for the double bond carbons each contain an electron, whereas the *p* orbital for the carbocation is empty.




allyl carbocation
Each C is sp^2 hybridized and has a *p* orbital.



three adjacent *p* orbitals




- **Three *p* orbitals on three adjacent atoms, even if one of the *p* orbitals is empty, make the allyl carbocation conjugated.**

Conjugation stabilizes the allyl carbocation because overlap of three adjacent *p* orbitals delocalizes the electron density of the π bond over three atoms.



overlap of adjacent *p* orbitals

Problem 16.2 Which of the following species are conjugated?

a. 
b. 
c. 
d. 
e. 

16.2 Resonance and Allylic Carbocations


The word *resonance* is used in two different contexts. In NMR spectroscopy, a nucleus is ***in resonance*** when it absorbs energy, promoting it to a higher energy state. In drawing molecules, there is ***resonance*** when two different Lewis structures can be drawn for the same arrangement of atoms.

Recall from Section 1.6 that resonance structures are two or more different Lewis structures for the same arrangement of atoms. Being able to draw correct resonance structures is crucial to understanding conjugation and the reactions of conjugated dienes.

- **Two resonance structures differ in the placement of π bonds and nonbonded electrons. The placement of atoms and σ bonds stays the same.**


16.2A The Stability of Allylic Carbocations

We have already drawn resonance structures for the acetate anion (Section 2.5C) and the allyl radical (Section 15.10). The **conjugated allyl carbocation** is another example of a species for which two resonance structures can be drawn. Drawing resonance structures for the allyl carbocation is a way to use Lewis structures to illustrate how conjugation delocalizes electrons.



resonance structures for the allyl carbocation — hybrid

The true structure of the allyl carbocation is a hybrid of the two resonance structures. In the hybrid, the π bond is delocalized over all three atoms. As a result, the positive charge is also delocalized over the two terminal carbons. Delocalizing electron density lowers the energy of the hybrid, thus stabilizing the allyl carbocation and making it more stable than a normal 1° carbocation. Experimental data show that its stability is comparable to a more substituted 2° carbocation.



The electrostatic potential maps in Figure 16.2 compare the resonance-stabilized allyl carbocation with $CH_3CH_2CH_2^+$, a localized 1° carbocation. The electron-deficient region—the site of the positive charge—is concentrated on a single carbon atom in the 1° carbocation $CH_3CH_2CH_2^+$. In the allyl carbocation, however, the electron-poor region is spread out on both terminal carbons.

Problem 16.3 Draw a second resonance structure for each carbocation. Then draw the hybrid.





Figure 16.2
Electrostatic potential maps for a localized and a delocalized carbocation




a localized carbocation — a delocalized carbocation

The electron-deficient region (in blue) of a **1° carbocation** is concentrated on a single carbon atom.

The electron-deficient region (in blue-green) of the **allyl carbocation** is distributed over both terminal carbons.


Problem 16.4 Use resonance theory and the Hammond postulate to explain why 3-chloroprop-1-ene ($CH_2=CHCH_2Cl$) is more reactive than 1-chloropropane ($CH_3CH_2CH_2Cl$) in S_N1 reactions.

16.2B Allylic Carbocations in Biological Reactions



organic diphosphate





diphosphate leaving group

PP_i


Allylic carbocations formed from diphosphates (Section 7.16) are key intermediates in a variety of biological reactions, including the synthesis of geranyl diphosphate from two five-carbon substrates—dimethylallyl diphosphate and isopentenyl diphosphate. Geranyl diphosphate is the precursor of many lipids that occur in plants and animals.



dimethylallyl diphosphate + isopentenyl diphosphate → geranyl diphosphate

This biological process results in the formation of a new carbon–carbon bond and involves two key steps—loss of a good leaving group (diphosphate, $P_2O_7^{4-}$, abbreviated as PP_i) to form an allylic carbocation, followed by nucleophilic attack with an electron-rich double bond. The steps of the mechanism are shown in Mechanism 16.1.

Mechanism 16.1 Biological Formation of Geranyl Diphosphate



1. Loss of the diphosphate leaving group forms an allylic carbocation.
2. Nucleophilic attack of isopentenyl diphosphate on the allylic carbocation forms the new C–C σ bond.
3. Loss of a proton (shown with the general base, B:) forms geranyl diphosphate.

We will learn more about biological reactions involving allylic carbocations derived from diphosphates in Chapter 31.

Problem 16.5 Farnesyl diphosphate is synthesized from isopentenyl diphosphate and **X** by a pathway similar to Mechanism 16.1. Draw the structure of **X.**




isopentenyl diphosphate + **X** → farnesyl diphosphate

16.3 Common Examples of Resonance

When are resonance structures drawn for a molecule or reactive intermediate? Because resonance involves delocalizing π bonds and nonbonded electrons, one or both of these structural features must be present to draw additional resonance forms. There are four common bonding patterns for which more than one Lewis structure can be drawn.

Type [1] The Three Atom "Allyl" System, X=Y–Z*

- **For any group of three atoms having a double bond X=Y and an atom Z that contains a *p* orbital with zero, one, or two electrons, two resonance structures are possible:**




The asterisk [*] corresponds to a charge, a radical, or a lone pair.

* = +, –, •, or ••

This is called **allyl** type resonance because it can be drawn for allylic carbocations, allylic carbanions, and allylic radicals.

X, Y, and Z may all be carbon atoms, as in the case of an allylic carbocation (resonance structures **A** and **B**), or they may be heteroatoms, as in the case of the acetate anion (resonance structures **C** and **D**). The atom Z bonded to the multiple bond can be charged (a net positive or negative charge) or neutral (having zero, one, or two nonbonded electrons). **The two resonance structures differ in the location of the double bond, and either the charge, the radical, or the lone pair, generalized by [*].**




Type [2] Conjugated Double Bonds

Cyclic, completely conjugated rings like benzene have two resonance structures, drawn by moving the electrons in a cyclic manner around the ring. Three resonance structures can be drawn for conjugated dienes, two of which involve charge separation.




two resonance structures for benzene



three resonance structures for buta-1,3-diene

Type [3] Cations Having a Positive Charge Adjacent to a Lone Pair

- **When a lone pair and a positive charge are located on adjacent atoms, two resonance structures can be drawn.**





The overall charge is the same in both resonance structures. Based on formal charge, a neutral X in one structure must bear a (+) charge in the other.

Type [4] Double Bonds Having One Atom More Electronegative Than the Other


- **For a double bond X=Y in which the electronegativity of Y > X, a second resonance structure can be drawn by moving the π electrons onto Y.**



Electronegativity of Y > X.

Sample Problem 16.1 illustrates how to apply these different types of resonance to actual molecules.

Sample Problem 16.1 Draw two more resonance structures for each species.



Solution


Mentally breaking a molecule into two- or three-atom units can make it easier to draw additional resonance structures.

a. Think of the top three atoms of the six-membered ring in **A** as an "allyl" unit. Moving the π bond forms a new "allyl" unit in **B,** and moving the π bond in **B** generates a third resonance structure **C.** No new valid resonance structures are generated by moving electrons in **C.**




A a three-atom "allyl" unit — **B** another three-atom "allyl" unit — **C**

b. Compound **D** contains a carbonyl group, so moving the electron pair in the double bond to the more electronegative oxygen atom separates the charge and generates structure **E. E** now has a three-atom "allyl" unit, so the remaining π bond can be moved to form structure **F.**



D — **E** a three-atom "allyl" unit — **F**

Problem 16.6 Draw additional resonance structures for each ion.



16.4 The Resonance Hybrid

The lower its energy, the more a resonance structure contributes to the overall structure of the hybrid.

Although the resonance hybrid is some combination of all of its valid resonance structures, the **hybrid more closely resembles the best resonance structure.** Recall from Section 1.6C that the best resonance structure is called the **major contributor** to the hybrid, and other resonance structures are called the **minor contributors.** Two identical resonance structures are equal contributors to the hybrid.

Use the following three rules to evaluate the relative energies of two or more valid resonance structures.

Rule [1] **Resonance structures with more bonds and fewer charges are better.**



all neutral atoms
one more bond
better resonance structure

charge separation


Rule [2] **Resonance structures in which every atom has an octet are better.**



All second-row elements have an octet.
better resonance structure

In this example, the resonance structure in which all atoms have octets is better, even though it places a (+) charge on a more electronegative O atom.


Rule [3] **Resonance structures that place a negative charge on a more electronegative atom are better.**



The (–) charge is on the
more electronegative O atom.
better resonance structure

Sample Problem 16.2 illustrates how to determine the relative energy of contributing resonance structures and the hybrid.


Sample Problem 16.2 Draw a second resonance structure for carbocation **A,** as well as the hybrid of both resonance structures. Then use Rules [1]–[3] to rank the relative stability of both resonance structures and the hybrid.



A

Solution

Because **A** contains a positive charge and a lone pair on adjacent atoms, a second resonance structure **B** can be drawn. Because **B** has more bonds and all second-row atoms have octets, **B** is a better resonance structure than **A,** making it the major contributor to the hybrid **C.** Because the hybrid is more stable than either resonance contributor, the order of stability is:




A
minor contributor

B
major contributor

C
hybrid

Increasing stability

Problem 16.7 Draw a second resonance structure and the hybrid for each species, and then rank the two resonance structures and the hybrid in order of increasing stability.




a.

b.

c.

d.


Problem 16.8 Draw all possible resonance structures for the following cation and indicate which structure makes the largest contribution to the resonance hybrid.



16.5 Electron Delocalization, Hybridization, and Geometry

To delocalize nonbonded electrons or electrons in π bonds, there must be *p* orbitals that can overlap. This may mean that the hybridization of an atom is different than would have been predicted using the rules first outlined in Chapter 1.

For example, there are two Lewis structures (**A** and **B**) for the resonance-stabilized anion $(CH_3COCH_2)^-$.




The labeled C is surrounded by four groups—three atoms and one nonbonded electron pair. **Is it sp^3 hybridized?**

The labeled C is surrounded by three groups—three atoms and no nonbonded electron pairs. **Is it sp^2 hybridized?**


Based on structure **A,** the labeled carbon is sp^3 hybridized, with the lone pair of electrons in an sp^3 hybrid orbital. Based on structure **B,** though, it is sp^2 hybridized with the unhybridized *p* orbital forming the π portion of the double bond.

Delocalizing electrons stabilizes a molecule. The electron pair on the carbon atom adjacent to the C=O can only be delocalized, though, if it has a *p* orbital that can overlap with two other *p* orbitals on two adjacent atoms. Thus, the terminal carbon atom is sp^2 hybridized with trigonal planar geometry. **Three adjacent *p* orbitals make the anion conjugated.**



- **In a system X=Y−Z:, Z is generally sp^2 hybridized, and the nonbonded electron pair occupies a *p* orbital to make the system conjugated.**

Sample Problem 16.3 Determine the hybridization around the labeled carbon atom in the following anion.




Solution

Because this is an example of an allyl-type system (X=Y−Z*), a second resonance structure can be drawn that "moves" the lone pair and the π bond. To delocalize the lone pair and make the system conjugated, the labeled carbon atom must be sp^2 hybridized with the lone pair occupying a *p* orbital.



The labeled C atom must be sp^2 hybridized, with the lone pair in a *p* orbital.

Problem 16.9 Determine the hybridization of the labeled atom in each species.




16.6 Conjugated Dienes

Compounds with many π bonds are called **polyenes.**

In the remainder of Chapter 16 we examine **conjugated dienes,** compounds having two double bonds joined by one σ bond. Conjugated dienes are also called **1,3-dienes.** Buta-1,3-diene ($CH_2=CH-CH=CH_2$) is the simplest conjugated diene.

Three stereoisomers are possible for 1,3-dienes with alkyl groups bonded to each end carbon of the diene (RCH=CH−CH=CHR).




both double bonds **trans** — *trans,trans*-1,3-diene or (*E,E*)-1,3-diene

both double bonds **cis** — *cis,cis*-1,3-diene or (*Z,Z*)-1,3-diene

cis,trans-1,3-diene or (*Z,E*)-1,3-diene


Two possible conformations result from rotation around the C−C bond that joins the two double bonds.



s-cis conformation **s-trans** conformation

- The *s*-cis conformation has two double bonds on the *same* side of the single bond.
- The *s*-trans conformation has two double bonds on *opposite* sides of the single bond.


Keep in mind that stereoisomers are discrete molecules, whereas conformations interconvert. Three structures drawn for hexa-2,4-diene illustrate the differences between stereoisomers and conformations in a 1,3-diene:



Problem 16.10 Draw the structure consistent with each description.

a. (2*E*,4*E*)-octa-2,4-diene in the *s*-trans conformation
b. (3*E*,5*Z*)-nona-3,5-diene in the *s*-cis conformation
c. (3*Z*,5*Z*)-4,5-dimethyldeca-3,5-diene. Draw both the *s*-cis and *s*-trans conformations.


Problem 16.11 Neuroprotectin D1 (NPD1) is synthesized in the body from highly unsaturated essential fatty acids. NPD1 is a potent natural anti-inflammatory agent.



NPD1


a. Label each double bond as conjugated or isolated.
b. Label each double bond as *E* or *Z*.
c. For each conjugated system, label the given conformation as *s*-cis or *s*-trans.

16.7 Interesting Dienes and Polyenes




Isoprene is a component of the blue haze seen above forested hillsides, such as Virginia's Blue Ridge Mountains.

Isoprene and **lycopene** are two naturally occurring compounds with conjugated double bonds.



isoprene
(2-methylbuta-1,3-diene)



11 conjugated double bonds shown in red


lycopene

Isoprene, the common name for 2-methylbuta-1,3-diene, is given off by plants as the temperature rises, a process thought to increase a plant's tolerance for heat stress.


Lycopene, a naturally occurring molecule responsible for the red color of tomatoes and other fruits, is an antioxidant like vitamin E. The 11 conjugated double bonds of lycopene cause its red color, a phenomenon discussed in Section 16.15.

Simvastatin and calcitriol are two drugs that contain conjugated double bonds in addition to other functional groups (Figure 16.3). Simvastatin is the generic name of the widely used cholesterol-lowering medicine Zocor. Calcitriol, a biologically active hormone formed from vitamin D_3 obtained in the diet, is responsible for regulating calcium and phosphorus metabolism. Sold under the trade name of Rocaltrol, calcitriol is used to treat patients who are unable to convert vitamin D_3 to the active hormone. Since calcitriol promotes the absorption of calcium ions, it is also used to treat hypocalcemia, the presence of low calcium levels in the blood.

Figure 16.3
Biologically active organic compounds that contain conjugated double bonds



simvastatin
(Zocor)



calcitriol
(Rocaltrol)

16.8 The Carbon–Carbon σ Bond Length in Buta-1,3-diene

Four features distinguish conjugated dienes from isolated dienes.


[1] The C − C single bond joining the two double bonds is unusually short.

[2] Conjugated dienes are more stable than similar isolated dienes.


[3] Some reactions of conjugated dienes are different than reactions of isolated double bonds.

[4] Conjugated dienes absorb longer wavelengths of ultraviolet light.

The bond length of the carbon–carbon double bonds in buta-1,3-diene is similar to an isolated double bond (134 pm), but the central carbon–carbon single bond is shorter than the C−C bond in ethane (148 pm vs. 153 pm).




The observed bond distances can be explained by looking at hybridization. Each carbon atom in buta-1,3-diene is sp^2 hybridized, so the central C−C single bond is formed by the overlap of two sp^2 hybridized orbitals, rather than the sp^3 hybridized orbitals used to form the C−C bond in CH_3CH_3.



Recall from Section 1.11B that increasing percent *s*-character decreases bond length.

- **Based on hybridization, a $C_{sp^2}-C_{sp^2}$ bond should be shorter than a $C_{sp^3}-C_{sp^3}$ bond because it is formed from orbitals having a higher percent *s*-character.**

A resonance argument can also be used to explain the shorter C−C σ bond length of buta-1,3-diene. Buta-1,3-diene can be represented by three resonance structures:




Structures **B** and **C** have charge separation and fewer bonds than **A,** making them less stable resonance structures and only minor contributors to the resonance hybrid. **B** and **C** both contain a double bond between the central carbon atoms, however, so the hybrid must have a partial double bond there. This makes the central C−C bond shorter than a C−C single bond in an alkane.

- **Based on resonance, the central C−C bond in buta-1,3-diene is shorter because it has partial double bond character.**

Problem 16.12 Using hybridization, predict how the bond length of the C−C σ bond in HC≡C−C≡CH should compare with the C−C σ bonds in CH_3CH_3 and $CH_2=CH-CH=CH_2$.


Problem 16.13 Use resonance theory to explain why the labeled C−O bond lengths are equal in the acetate anion.



acetate

16.9 Stability of Conjugated Dienes


In Section 12.3 we learned that hydrogen adds to alkenes to form alkanes, and that the heat released in this reaction, the **heat of hydrogenation,** can be used as a measure of alkene stability.



The relative stability of conjugated and isolated dienes can also be determined by comparing their heats of hydrogenation.


- **When hydrogenation gives the same alkane from two dienes, the more stable diene has the smaller heat of hydrogenation.**

For example, both penta-1,4-diene (an isolated diene) and (*E*)-penta-1,3-diene (a conjugated diene) are hydrogenated to pentane with two equivalents of H_2. Because less energy is released in converting the conjugated diene to pentane, it must be lower in energy (more stable) to begin with. The relative energies of these isomeric pentadienes are illustrated in Figure 16.4.




- **A conjugated diene has a smaller heat of hydrogenation and is more stable than a similar isolated diene.**

Figure 16.4
Relative energies of an isolated and conjugated diene




In Section 16.1, we learned why a conjugated diene is more stable than an isolated diene. A conjugated diene has overlapping *p* orbitals on four adjacent atoms, so its π electrons are delocalized over four atoms, thus stabilizing the diene. This delocalization cannot occur in an isolated diene, so an isolated diene is less stable than a conjugated diene.

Problem 16.14 Which diene in each pair has the larger heat of hydrogenation?



Problem 16.15 Rank the following compounds in order of increasing stability.




16.10 Electrophilic Addition: 1,2- Versus 1,4-Addition

Recall from Chapters 10 and 11 that the characteristic reaction of compounds with π bonds is **addition.** The π bonds in conjugated dienes undergo addition reactions, too, but they differ in two ways from the addition reactions to isolated double bonds.


- **Electrophilic addition in conjugated dienes gives a mixture of products.**
- **Conjugated dienes undergo a unique addition reaction not seen in alkenes or isolated dienes.**

We learned in Chapter 10 that HX adds to the π bond of alkenes to form alkyl halides.



A π bond is broken. alkyl halide


With an **isolated diene,** electrophilic addition of one equivalent of HBr yields *one* product and Markovnikov's rule is followed. The H atom bonds to the less substituted carbon—that is, the carbon atom of the double bond that had more H atoms to begin with.



isolated diene

H bonds to the less substituted carbon.

With a **conjugated diene,** electrophilic addition of one equivalent of HBr affords *two* products.




conjugated diene **1,2-product** **1,4-product**

The ends of the 1,3-diene are called C1 and C4 arbitrarily, without regard to IUPAC numbering.

- The **1,2-addition product** results from Markovnikov addition of HBr across two adjacent carbon atoms (C1 and C2) of the diene.
- The **1,4-addition product** results from addition of HBr to the two end carbons (C1 and C4) of the diene. 1,4-Addition is also called **conjugate addition.**

The mechanism of electrophilic addition of HX involves **two steps:** addition of H^+ (from HX) to form a resonance-stabilized carbocation, followed by nucleophilic attack of X^- at either electrophilic end of the carbocation to form two products. Mechanism 16.2 illustrates the reaction of buta-1,3-diene with HBr.

Mechanism 16.2 Electrophilic Addition of HBr to a 1,3-Diene—1,2- and 1,4-Addition



new bonds shown in red
allylic carbocation

1,4-addition product
new bonds shown in red


1 H^+ of HBr adds to a terminal carbon of the 1,3-diene to form a resonance-stabilized allylic carbocation.

2 Nucleophilic attack of Br^- occurs at either site of the resonance-stabilized carbocation that bears a (+) charge, forming the 1,2- and 1,4-addition products.

Like the electrophilic addition of HX to an alkene, the addition of HBr to a conjugated diene forms the more stable carbocation in Step [1], the rate-determining step. In this case, however, the carbocation is both 2° and **allylic,** and thus two Lewis structures can be drawn for it. In the second step, nucleophilic attack of Br^- can then occur at two different electrophilic sites, forming two different products.


- **Addition of HX to a conjugated diene forms 1,2- and 1,4-products because of the resonance-stabilized allylic carbocation intermediate.**

Sample Problem 16.4 Draw the products of the following reaction.





Solution


Write the steps of the mechanism to determine the structure of the products. Addition of H^+ forms the more stable 2° allylic carbocation, for which two resonance structures can be drawn. Nucleophilic attack of Br^- at either end of the allylic carbocation gives two constitutional isomers, formed by 1,2-addition and 1,4-addition to the diene.



Problem 16.16 Draw the products formed when each diene is treated with one equivalent of HCl.


a.  b.  c.  d. 

Problem 16.17 Draw a stepwise mechanism for the following reaction.




16.11 Kinetic Versus Thermodynamic Products

The amount of 1,2- and 1,4-addition products formed in the electrophilic addition reactions of buta-1,3-diene, a conjugated diene, depends greatly on the reaction conditions.



- **At low temperature the major product is formed by 1,2-addition.**
- **At higher temperature the major product is formed by 1,4-addition.**

Moreover, when a mixture containing predominately the 1,2-product is heated, the 1,4-addition product becomes the major product at equilibrium.



- **The 1,2-product is formed faster because it predominates at low temperature. The product that is formed faster is called the *kinetic product.***
- **The 1,4-product must be more stable because it predominates at equilibrium. The product that predominates at equilibrium is called the *thermodynamic product.***

In many of the reactions we have learned thus far, the more stable product is formed faster—that is, the kinetic and thermodynamic products are the same. The electrophilic addition of HBr to buta-1,3-diene is different, in that **the more stable product is formed more slowly**—that is, the kinetic and thermodynamic products are *different.* Why is the more stable product formed more slowly?

To answer this question, recall that the rate of a reaction is determined by its energy of activation (E_a), whereas the amount of product present at equilibrium is determined by its stability (Figure 16.5). When a single starting material **A** forms two different products (**B** and **C**) by two exothermic pathways, the relative height of the energy barriers determines how fast **B** and **C** are formed, whereas the relative energies of **B** and **C** determine the amount of each at equilibrium. In an exothermic reaction, the relative energies of **B** and **C** do not determine the relative energies of activation to form **B** and **C.**

Why, in the addition of HBr to buta-1,3-diene, is the 1,4-product the more stable thermodynamic product? The 1,4-product (1-bromobut-2-ene) is more stable because it has two alkyl groups bonded to the carbon–carbon double bond, whereas the 1,2-product (3-bromobut-1-ene) has only one.





Figure 16.5
How kinetic and thermodynamic products form in a reaction: **A** → **B** + **C**




- The conversion of **A** → **B** is a faster reaction because the energy of activation leading to **B** is lower. **B** is the kinetic product.
- Because **C** is lower in energy, **C** is the thermodynamic product.

- **The more substituted alkene—1-bromobut-2-ene in this case—is the thermodynamic product.**

The 1,2-product is the kinetic product because of a **proximity effect.** When H^+ (from HBr) adds to the double bond, Br^- is closer to the adjacent carbon (C2) than it is to C4. Even though the resonance-stabilized carbocation bears a partial positive charge on both C2 and C4, attack at C2 is faster simply because Br^- is closer to this carbon.

A **proximity effect** occurs because one species is close to another.




Br^- is closer to C2 than C4.

- **The 1,2-product forms faster because of the proximity of Br^- to C2.**

The overall two-step mechanism for addition of HBr to buta-1,3-diene, forming a 1,2-addition product and 1,4-addition product, is illustrated with the energy diagram in Figure 16.6.


Figure 16.6
Energy diagram for the two-step addition of HBr to $CH_2=CH-CH=CH_2$



Why is the ratio of products temperature dependent?

- At low temperature, the energy of activation is the more important factor. Because most molecules do not have enough kinetic energy to overcome the higher energy barrier at lower temperature, they react by the faster pathway, forming the kinetic product.
- At higher temperature, most molecules have enough kinetic energy to reach either transition state. The two products are in equilibrium with each other, and the more stable compound—which is lower in energy—becomes the major product.


Problem 16.18 Label each product in the following reaction as a 1,2-product or 1,4-product, and decide which is the kinetic product and which is the thermodynamic product.



16.12 The Diels–Alder Reaction

Diels and Alder shared the 1950 Nobel Prize in Chemistry for unraveling the intricate details of this remarkable reaction.


The **Diels–Alder reaction,** named for German chemists Otto Diels and Kurt Alder, is an addition reaction between a **1,3-diene** and an alkene called a **dienophile,** to form a new six-membered ring.



1,3-diene dienophile

The arrows may be drawn in a clockwise or counterclockwise direction to show the flow of electrons in a Diels–Alder reaction.

Three curved arrows are needed to show the cyclic movement of electron pairs because three π bonds break and two σ bonds and one π bond form. Because each new σ bond is ~100 kJ/mol stronger than a π bond that is broken, a typical Diels–Alder reaction releases ~200 kJ/mol of energy. The following equations illustrate two examples of the Diels–Alder reaction.




All Diels–Alder reactions have the following features in common:

[1] They are initiated by heat; that is, the Diels–Alder reaction is a *thermal* reaction.
[2] They form new six-membered rings.
[3] Three π bonds break, and two new C−C σ bonds and one new C−C π bond form.
[4] They are concerted; that is, all bonds are broken and formed in a single step.

The Diels–Alder reaction forms new carbon–carbon bonds, so it can be used to synthesize larger, more complex molecules from smaller ones. For example, Figure 16.7 illustrates a Diels–Alder reaction used in the synthesis of tetrodotoxin, a toxin isolated from many different types of puffer fish.

Figure 16.7
Synthesis of a natural product using the Diels–Alder reaction




- Tetrodotoxin, a complex natural product containing several six-membered rings joined together, is a poison isolated from the ovaries and liver of the puffer fish, so named because the fish inflates itself into a ball when alarmed. One step in the synthesis of tetrodotoxin involves forming a six-membered ring by a Diels–Alder reaction.

Diels–Alder reactions may seem complicated at first, but they are really less complicated than many of the reactions you have already learned, especially those with multistep mechanisms and carbocation intermediates. The key is to learn how to arrange the starting materials to more easily visualize the structure of the product.


How To Draw the Product of a Diels–Alder Reaction

Example **Draw the product of the following Diels–Alder reaction:**




Step [1] **Arrange the 1,3-diene and the dienophile next to each other, with the diene drawn in the *s*-cis conformation.**


- This step is key: **Rotate the diene** so that it is drawn in the ***s*-cis** conformation, and place the end C's of the diene close to the double bond of the dienophile.



Step [2] **Cleave the three π bonds and use arrows to show where the new bonds will be formed.**



Problem 16.19 Draw the product formed when each diene and dienophile react in a Diels–Alder reaction.




16.13 Specific Rules Governing the Diels–Alder Reaction

Several rules govern the course of the Diels–Alder reaction.

16.13A Diene Reactivity

Rule [1] **The diene can react only when it adopts the *s*-cis conformation.**

Both ends of the conjugated diene must be close to the π bond of the dienophile for reaction to occur. Thus, an acyclic diene in the *s*-trans conformation must rotate about the central C−C σ bond to form the *s*-cis conformation before reaction can take place.



This rotation is prevented in cyclic dienes. As a result:


- **When the two double bonds are constrained in the *s*-cis conformation, the diene is unusually *reactive.***
- **When the two double bonds are constrained in the *s*-trans conformation, the diene is *unreactive.***




Zingiberene and β-sesquiphellandrene are trienes obtained from ginger root. Ginger is used as a spice in Indian and Chinese cooking. Ginger candy is sometimes used to treat nausea resulting from seasickness.



Problem 16.20 Rank the following dienes in order of increasing reactivity in a Diels–Alder reaction.



Problem 16.21 Zingiberene and β-sesquiphellandrene, natural products obtained from ginger root, contain conjugated diene units. Which diene reacts faster in the Diels–Alder reaction and why?




16.13B Dienophile Reactivity


Rule [2] **Electron-withdrawing substituents in the dienophile increase the reaction rate.**

In a Diels–Alder reaction, the conjugated diene acts as a nucleophile and the dienophile acts as an electrophile. As a result, electron-withdrawing groups make the dienophile more electrophilic

(and, thus, more reactive) by withdrawing electron density from the carbon–carbon double bond. If Z is an electron-withdrawing group, then the reactivity of the dienophile increases as follows:



A carbonyl group is an effective electron-withdrawing group because the carbonyl carbon bears a partial positive charge (δ+), which withdraws electron density from the carbon–carbon double bond of the dienophile. Common dienophiles that contain a carbonyl group are shown in Figure 16.8.



electron deficient carbonyl carbon

Problem 16.22 Rank the following dienophiles in order of increasing reactivity.





Figure 16.8
Common dienophiles in the Diels–Alder reaction




acrolein | methyl vinyl ketone | methyl acrylate | maleic anhydride | benzoquinone

16.13C Stereospecificity

Rule [3] The stereochemistry of the dienophile is retained in the product.


- **A cis dienophile forms a cis-substituted cyclohexene.**
- **A trans dienophile forms a trans-substituted cyclohexene.**

The two **cis** CO_2H groups of maleic acid become two **cis** substituents in a Diels–Alder adduct. The CO_2H groups can be drawn both above or both below the plane to afford a single achiral **meso** compound. The **trans dienophile** fumaric acid yields two enantiomers with **trans** CO_2H groups.



maleic acid
cis dienophile


an achiral meso compound



fumaric acid
trans dienophile

enantiomers


A **cyclic dienophile** forms a **bicyclic product.** A bicyclic system in which the two rings share a common C−C bond is called a **fused ring system.** The two H atoms at the ring fusion must be cis, because they were cis in the starting dienophile. A bicyclic system of this sort is said to be **cis-fused.**



cyclic dienophile
cis H's in the dienophile

bicyclic product
cis H's in the product


Problem 16.23 Draw the products of each Diels–Alder reaction, and indicate the stereochemistry.



16.13D The Rule of Endo Addition

Rule [4] When endo and exo products are possible, the endo product is preferred.

To understand the rule of endo addition, we must first examine Diels–Alder products that result from cyclic 1,3-dienes. When cyclopentadiene reacts with a dienophile such as ethylene, a new six-membered ring forms, and above the ring there is a one atom "bridge," labeled in green. This carbon atom originated as the sp^3 hybridized carbon of the diene that was not involved in the reaction.



cyclic 1,3-diene


a bridged bicyclic ring system

The product of the Diels–Alder reaction of a cyclic 1,3-diene is bicyclic, but the carbon atoms shared by both rings are *non-adjacent.* Thus, this bicyclic product differs from the fused ring system obtained when the dienophile is cyclic.

- A bicyclic ring system in which the two rings share non-adjacent carbon atoms is called a *bridged* ring system.


Figure 16.9
Fused and bridged bicyclic ring systems compared

a. A fused bicyclic system



- One bond (in red) is shared by two rings.
- The shared C's are adjacent.


b. A bridged bicyclic system



- Two non-adjacent atoms (labeled in blue) are shared by both rings.


Fused and bridged bicyclic ring systems are compared in Figure 16.9.

When cyclopentadiene reacts with a substituted alkene as the dienophile (CH_2=CHZ), the substituent Z can be oriented in one of two ways in the product. The terms **endo** and **exo** are used to indicate the position of Z.




- **A substituent on one bridge is *endo* if it is closer to the *longer* bridge that joins the two carbons common to both rings.**
- **A substituent is *exo* if it is closer to the *shorter* bridge that joins the carbons together.**

In a Diels–Alder reaction, the **endo** product is preferred, as shown in two examples.



To help you distinguish endo and exo, remember that e***n***do is u***n***der the newly formed six-membered ring.




More details on the Diels–Alder reaction are given in Section 27.4.


The Diels–Alder reaction is concerted, and the reaction occurs with the diene and the dienophile arranged one above the other, as shown in Figure 16.10, not side-by-side. In theory, the substituent Z can be oriented either directly under the diene to form the endo product (Pathway [1] in Figure 16.10) or away from the diene to form the exo product (Pathway [2] in Figure 16.10). In practice, though, the **endo product is the major product.** The transition state leading to the endo product allows more interaction between the electron-rich diene and the electron-withdrawing substituent Z on the dienophile, an energetically favorable arrangement.

Figure 16.10
How endo and exo products are formed in the Diels–Alder reaction


Pathway [1] With Z oriented under the diene, the endo product is formed.



Pathway [2] With Z oriented away from the diene, the exo product is formed.



Problem 16.24 Draw the product of each Diels–Alder reaction.




16.14 Other Facts About the Diels–Alder Reaction

16.14A Retrosynthetic Analysis of a Diels–Alder Product

The Diels–Alder reaction is used widely in organic synthesis, so you must be able to look at a compound and determine what conjugated diene and what dienophile were used to make it. To draw the starting materials from a given Diels–Alder adduct:


- Locate the six-membered ring that contains the C=C.
- Draw three arrows around the cyclohexene ring, beginning with the π bond. Each arrow moves two electrons to the adjacent bond, cleaving one π bond and two σ bonds, and forming three π bonds.
- Retain the stereochemistry of substituents on the C=C of the dienophile. Cis substituents on the six-membered ring give a cis dienophile.

Figure 16.11 Finding the diene and dienophile needed for a Diels–Alder reaction




This stepwise retrosynthetic analysis gives the 1,3-diene and dienophile needed for any Diels–Alder reaction, as shown in the two examples in Figure 16.11.

Problem 16.25 What diene and dienophile are needed to prepare each product?




16.14B Retro Diels–Alder Reaction

A reactive diene like cyclopenta-1,3-diene readily undergoes a Diels–Alder reaction with *itself;* that is, **cyclopenta-1,3-diene dimerizes because one molecule acts as the diene and another acts as the dienophile.**



The formation of dicyclopentadiene is so rapid that it takes only a few hours at room temperature for cyclopentadiene to completely dimerize. How, then, can cyclopentadiene be used in a Diels–Alder reaction if it really exists as a dimer?

When heated, dicyclopentadiene undergoes a **retro Diels–Alder reaction,** and two molecules of cyclopentadiene are re-formed. If cyclopentadiene is immediately treated with a different dienophile, it reacts to form a new Diels–Alder adduct with this dienophile.



dicyclopentadiene


two molecules of cyclopentadiene

This diene can now be used with a different dienophile.


16.14C Application: Diels–Alder Reaction in the Synthesis of Steroids

Recall from Section 4.15 that lipids are water-insoluble biomolecules that have diverse structures.

Steroids **are tetracyclic lipids containing three six-membered rings and one five-membered ring.** The four rings are designated as **A, B, C,** and **D.**



steroid skeleton




three-dimensional view from above




carbon skeleton viewed from the side


Steroids exhibit a wide range of biological properties, depending on the substitution pattern of functional groups on the rings. They include **cholesterol** (a component of cell membranes that is implicated in cardiovascular disease), **estrone** (a female sex hormone responsible for the regulation of the menstrual cycle), and **cortisone** (a hormone responsible for the control of inflammation and the regulation of carbohydrate metabolism).



cholesterol




estrone




cortisone

Diels–Alder reactions have been used widely in the laboratory syntheses of steroids. The key Diels–Alder reactions used to prepare the C ring of estrone and the B ring of cortisone are drawn.




diene dienophile Diels–Alder product estrone




diene Diels–Alder product cortisone

Problem 16.26 Draw the product (**A**) of the following Diels–Alder reaction. **A** was a key intermediate in the synthesis of the addicting pain reliever morphine, the chapter-opening molecule.




16.15 Conjugated Dienes and Ultraviolet Light

Recall from Chapter 13 that the absorption of infrared light can promote a molecule from a lower vibrational state to a higher one. In a similar fashion, the absorption of ultraviolet (UV) light can promote an electron from a lower electronic state to a higher one. Ultraviolet light has a slightly shorter wavelength (and, thus, higher frequency) than visible light. The most useful region of UV light for this purpose is **200–400 nm.**



16.15A General Principles

When electrons in a lower energy state (the **ground state**) absorb light having the appropriate energy, an electron is promoted to a higher electronic state (the **excited state**).



The energy difference between the two states depends on the location of the electron. The promotion of electrons in σ bonds and unconjugated π bonds requires light having a wavelength of < 200 nm; that is, it has a shorter wavelength and higher energy than light in the UV region of the electromagnetic spectrum. With conjugated dienes, however, the energy difference between the ground and excited states decreases, so longer wavelengths of light can be used to promote electrons. The wavelength of UV light absorbed by a compound is often referred to as its λ_{max}. Buta-1,3-diene, for example, absorbs UV light at λ_{max} = 217 nm and cyclohexa-1,3-diene has a λ_{max} of 256 nm.




λ_{max} = 217 nm λ_{max} = 256 nm

- **Conjugated dienes and polyenes absorb light in the UV region of the electromagnetic spectrum (200–400 nm).**

A UV spectrum is a plot of the absorbance of UV light versus wavelength. A spectrum consists of very broad bands, and the maximum absorbance corresponds to the λ_{max}, as shown in the UV spectrum of isoprene in Figure 16.12.


Figure 16.12
UV spectrum of isoprene






Lycopene is the red pigment found in tomatoes, watermelon, papaya, guava, and pink grapefruit. Lycopene is not destroyed when fruits and vegetables are processed, so tomato juice and ketchup are high in lycopene.

As the number of conjugated π bonds increases, the energy difference between the ground and excited state decreases, shifting the absorption to longer wavelengths.







With molecules having eight or more conjugated π bonds, the absorption shifts from the UV to the visible region and the compound takes on the color of those wavelengths of visible light it does *not* absorb. For example, lycopene absorbs visible light at λ_{max} = 470 nm, in the blue-green region of the visible spectrum. Because it does not absorb light in the red region, lycopene appears bright red (Figure 16.13).

Figure 16.13 Why lycopene appears red



Problem 16.27 Which compound in each pair absorbs UV light at longer wavelength?

a.  or 

b.  or 


16.15B Sunscreens

Ultraviolet radiation from the sun is high enough in energy to cleave bonds, forming radicals that can prematurely age skin and cause skin cancers. The ultraviolet region is often subdivided, based on the wavelength of UV light: UV-A (320–400 nm), UV-B (290–320 nm), and UV-C (< 290 nm). Fortunately, much of the highest energy UV light (UV-C) is filtered out by the ozone layer, so that only UV light having wavelengths > 290 nm reaches the skin's surface. Much of this UV light is absorbed by **melanin,** the highly conjugated colored pigment in the skin that serves as the body's natural protection against the harmful effects of UV radiation.




Commercial sunscreens are given an **SPF** rating (sun protection factor), according to the amount of sunscreen present. The higher the number, the greater the protection.

Prolonged exposure to the sun can allow more UV radiation to reach your skin than melanin can absorb. A commercial sunscreen can offer added protection, however, because it contains conjugated compounds that absorb UV light, thus shielding your skin (for a time) from the harmful effects of UV radiation. Two sunscreens that have been used for this purpose are *para*-aminobenzoic acid (PABA) and padimate O.




para-aminobenzoic acid (PABA)





padimate O

Many sunscreens contain more than one component to filter out different regions of the UV spectrum. Conjugated compounds generally shield the skin from UV-B radiation, but often have little effect on longer-wavelength UV-A radiation, which does not burn the skin, but can still cause long-term damage to skin cells.

Problem 16.28 Which of the following compounds might be an ingredient in a commercial sunscreen? Explain why or why not.

a. 

c. 

b. 

KEY CONCEPTS


Conjugation, Resonance, and Dienes

Conjugation and Delocalization of Electron Density


- The overlap of *p* orbitals on three or more adjacent atoms allows electron density to delocalize, thus adding stability (16.1).
- An allyl carbocation ($CH_2=CHCH_2^+$) is more stable than a 1° carbocation because of *p* orbital overlap (16.2).
- In a system X=Y–Z:, Z is generally sp^2 hybridized to allow the lone pair to occupy a *p* orbital, making the system conjugated (16.5).

Four Common Examples of Resonance (16.3)


[1] The three-atom "allyl" system:



[2] Conjugated double bonds:



[3] Cations having a positive charge adjacent to a lone pair:



[4] Double bonds involving one atom more electronegative than the other:



Rules on Evaluating the Relative "Stability" of Resonance Structures (16.4)

[1] Structures with more bonds and fewer charges are better.
[2] Structures in which every atom has an octet are better.
[3] Structures that place a negative charge on a more electronegative atom are better.


The Unusual Properties of Conjugated Dienes

[1] The C – C σ bond joining the two double bonds is unusually short (16.8).
[2] Conjugated dienes are more stable than the corresponding isolated dienes. $\Delta H°$ of hydrogenation is smaller for a conjugated diene than for an isolated diene converted to the same product (16.9).
[3] The reactions are unusual:
- Electrophilic addition affords products of 1,2-addition and 1,4-addition (16.10, 16.11).
- Conjugated dienes undergo the Diels–Alder reaction, a reaction that does not occur with isolated dienes (16.12–16.14).

[4] Conjugated dienes absorb UV light in the 200–400 nm region. As the number of conjugated π bonds increases, the absorption shifts to longer wavelength (16.15).

Reactions of Conjugated Dienes


[1] Electrophilic addition of HX (X = halogen) (16.10–16.11)



1,2-product kinetic product **1,4-product thermodynamic product**

- The mechanism has two steps.
- Markovnikov's rule is followed. Addition of H^+ forms the more stable allylic carbocation.
- The 1,2-product is the kinetic product. When H^+ adds to the double bond, X^- adds to the end of the allylic carbocation to which it is closer (C2 not C4). The kinetic product is formed faster at low temperature.
- The thermodynamic product has the more substituted, more stable double bond. The thermodynamic product predominates at equilibrium. With buta-1,3-diene, the thermodynamic product is the 1,4-product.

[2] Diels–Alder reaction (16.12–16.14)




1,3-diene dienophile


- The reaction forms two σ and one π bond in a six-membered ring.
- The reaction is initiated by heat.
- The mechanism is concerted: All bonds are broken and formed in a single step.
- The diene must react in the *s*-cis conformation (16.13A).
- Electron-withdrawing groups in the dienophile increase the reaction rate (16.13B).
- The stereochemistry of the dienophile is retained in the product (16.13C).
- Endo products are preferred (16.13D).

PROBLEMS


Problems Using Three-Dimensional Models


16.29 Name each diene and state whether the ball-and-stick model shows the diene in the *s*-cis or *s*-trans conformation.

a. 

b. 


16.30 What diene and dienophile are needed to prepare each compound by a Diels–Alder reaction?


a. 


b. 


Conjugation

16.31 Which of the following systems are conjugated?

a. 


b. 


c. 


d. 


Resonance and Hybridization


16.32 Draw all reasonable resonance structures for each species.


a. 

b. 


c. 


d. 


e. 


f. 

16.33 For which compounds can a second resonance structure be drawn? Draw an additional resonance structure for each resonance-stabilized compound.


a. 

b. 

c. 


d. 

16.34 Draw all reasonable resonance structures for acetic anhydride.




acetic anhydride

16.35 Explain why the cyclopentadienide anion **A** gives only one signal in its ^{13}C NMR spectrum.



16.36 Why is the bond dissociation energy for the C–C bond in ethane much higher than the bond dissociation energy for the labeled C–C bond in but-1-ene?





Nomenclature and Stereoisomers in Conjugated Dienes



16.37 Draw the structure of each compound.


a. (*Z*)-penta-1,3-diene in the *s*-trans conformation
b. (2*E*,4*Z*)-1-bromo-3-methylhexa-2,4-diene
c. (2*E*,4*E*,6*E*)-octa-2,4,6-triene
d. (2*E*,4*E*)-3-methylhexa-2,4-diene in the *s*-cis conformation

16.38 Draw all possible stereoisomers of hepta-2,4-diene and label each double bond as *E* or *Z*.

16.39 Label each pair of compounds as stereoisomers or conformations.

a.  and 

c.  and 


b.  and 


16.40 Rank the following dienes in order of increasing heat of hydrogenation.




Electrophilic Addition

16.41 Draw the products formed when each compound is treated with one equivalent of HBr.


a. 

b. 


c. 

16.42 Ignoring stereoisomers, draw all products that form by addition of HBr to (*E*)-hexa-1,3,5-triene.


16.43 Treatment of alkenes **A** and **B** with HBr gives the same alkyl halide **C.** Draw a mechanism for each reaction, including all reasonable resonance structures for any intermediate.



16.44 Draw a stepwise mechanism for the following reaction.




16.45 Addition of HCl to alkene **X** forms two alkyl halides **Y** and **Z.**



a. Label **Y** and **Z** as a 1,2-addition product or a 1,4-addition product.
b. Label **Y** and **Z** as the kinetic or thermodynamic product and explain why.
c. Explain why addition of HCl occurs at the indicated C = C (called an exocyclic double bond), rather than the other C = C (called an endocyclic double bond).

16.46 Explain, with reference to the mechanism, why addition of one equivalent of HCl to diene **A** forms only two products of electrophilic addition, even though four constitutional isomers are possible.




16.47 The major product formed by addition of HBr to $(CH_3)_2C=CH-CH=C(CH_3)_2$ is the same at low and high temperature. Draw the structure of the major product and explain why the kinetic and thermodynamic products are the same in this reaction.


Diels–Alder Reaction

16.48 Explain why methyl vinyl ether ($CH_2=CHOCH_3$) is not a reactive dienophile in the Diels–Alder reaction.


16.49 Draw the products of the following Diels–Alder reactions. Indicate stereochemistry where appropriate.




16.50 What diene and dienophile are needed to prepare each Diels–Alder product?




16.51 Give two different ways to prepare the following compound by the Diels–Alder reaction. Explain which method is preferred.



16.52 Compounds containing triple bonds are also Diels–Alder dienophiles. With this in mind, draw the products of each reaction.




16.53 Diels–Alder reaction of a monosubstituted diene (such as $CH_2=CH-CH=CHOCH_3$) with a monosubstituted dienophile (such as $CH_2=CHCHO$) gives a mixture of products, but the 1,2-disubstituted product often predominates. Draw the resonance hybrid for each reactant and use the charge distribution of the hybrids to explain why the 1,2-disubstituted product is the major product.




1,2-disubstituted product **major** | 1,3-disubstituted product **minor**

16.54 The following reactions have been used to synthesize dieldrin and aldrin (named for Diels and Alder), two pesticides having a similar story to DDT (Section 7.4). Identify the lettered compounds in this reaction scheme.




aldrin


16.55 Devise a stepwise synthesis of each compound from dicyclopentadiene using a Diels–Alder reaction as one step. You may also use organic compounds having ≤ 4 C's, and any required organic or inorganic reagents.




16.56 Intramolecular Diels–Alder reactions are possible when a substrate contains both a 1,3-diene and a dienophile, as shown in the following general reaction.



With this in mind, draw the product of each intramolecular Diels–Alder reaction.




16.57 A transannular Diels–Alder reaction is an intramolecular reaction that occurs when the diene and dienophile are contained in one ring, resulting in the formation of a tricyclic ring system. Draw the product formed when the following triene undergoes a transannular Diels–Alder reaction.




General Problems

16.58 (a) Which triene has the largest heat of hydrogenation? (b) Which triene has the smallest heat of hydrogenation? (c) Which triene absorbs the longest wavelength of UV light? (d) Which triene is most reactive in the Diels-Alder reaction?




A **B** **C** **D**


16.59 Draw a stepwise mechanism for the following reaction.



16.60 Draw the products of each reaction. Indicate the stereochemistry of Diels–Alder products.



16.61 Draw a stepwise mechanism for the biological conversion of linalyl diphosphate to limonene.



linalyl diphosphate limonene

16.62 Which benzylic halide reacts faster in an S_N1 reaction? Explain.




A **B**

16.63 Like alkenes, conjugated dienes can be prepared by elimination reactions. Draw a stepwise mechanism for the acid-catalyzed dehydration of 3-methylbut-2-en-1-ol [$(CH_3)_2C=CHCH_2OH$] to isoprene [$CH_2=C(CH_3)CH=CH_2$].


16.64 (a) Draw the two isomeric dienes formed when $CH_2=CHCH_2CH(Cl)CH(CH_3)_2$ is treated with an alkoxide base. (b) Explain why the major product formed in this reaction does not contain the more highly substituted alkene.


Spectroscopy

16.65 The treatment of isoprene [$CH_2=C(CH_3)CH=CH_2$] with one equivalent of mCPBA forms **A** as the major product. **A** gives a molecular ion at 84 in its mass spectrum, and peaks at 2850–3150 cm^{-1} in its IR spectrum. The ^1H NMR spectrum of **A** is given below. What is the structure of **A?**




16.66 The treatment of $(CH_3)_2C=CHCH_2Br$ with H_2O forms **B** (molecular formula $C_5H_{10}O$) as one of the products. Determine the structure of **B** from its ^1H NMR and IR spectra.






UV Absorption

16.67 Which of the following molecules absorbs a longer wavelength of UV light?




A




B

16.68 Explain why both **C** and **D** absorb light in the UV region of the electromagnetic spectrum, despite the fact that they are not 1,3-dienes.




C



D


16.69 Explain why ferulic acid, a natural product found in rice, oats, and other plants, is both an antioxidant and a sunscreen.



ferulic acid

Challenge Problems

16.70 Addition of HBr to allene ($CH_2=C=CH_2$) forms 2-bromoprop-1-ene rather than 3-bromoprop-1-ene, even though 3-bromoprop-1-ene is formed from an allylic carbocation. Considering the arrangement of orbitals in the allene reactant, explain this result.



2-bromoprop-1-ene

3-bromoprop-1-ene


NOT formed

16.71 Determine the hybridization around the N atom in each amine and explain why cyclohexanamine is 10^6 times more basic than aniline.




cyclohexanamine aniline

16.72 Devise a synthesis of **X** from the given starting materials. You may use any organic or inorganic reagents. Account for the stereochemistry observed in **X.**




X


16.73 One step in the synthesis of occidentalol, a natural product isolated from the eastern white cedar tree, involved the following reaction. Identify the structure of **A** and show how **A** is converted to **B.**



16.74 One step in the synthesis of dodecahedrane (Section 4.11) involved reaction of the tetraene **C** with dimethylacetylene dicarboxylate (**D**) to afford two compounds having molecular formula $C_{16}H_{16}O_4$. This reaction has been called a domino Diels–Alder reaction. Identify the two products formed.



16.75 Devise a stepwise mechanism for the conversion of **M** to **N. N** has been converted in several steps to lysergic acid, a naturally occurring precursor of the hallucinogen LSD (Figure 18.4).



M **N** lysergic acid

Benzene and Aromatic Compounds

17



Capsaicin is responsible for the characteristic spicy flavor of jalapeño and habañero peppers. Although it first produces a burning sensation on contact with the mouth or skin, repeated application desensitizes the area to pain. This property has made it the active ingredient in several topical creams for the treatment of chronic pain. Capsaicin has also been used as an animal deterrent in pepper sprays, and as an additive to make birdseed squirrel-proof. Capsaicin is an aromatic compound because it contains a benzene ring. In this chapter, we learn about the characteristics of aromatic compounds like capsaicin.

17.1 Background
17.2 The structure of benzene
17.3 Nomenclature of benzene derivatives
17.4 Spectroscopic properties
17.5 Interesting aromatic compounds
17.6 Benzene's unusual stability
17.7 The criteria for aromaticity—Hückel's rule
17.8 Examples of aromatic compounds
17.9 What is the basis of Hückel's rule?
17.10 The inscribed polygon method for predicting aromaticity
17.11 Buckminsterfullerene—Is it aromatic?


The hydrocarbons we have examined thus far—including the alkanes, alkenes, and alkynes, as well as the conjugated dienes and polyenes of Chapter 16—have been aliphatic hydrocarbons. In Chapter 17, we continue our study of conjugated systems with **aromatic hydrocarbons.**

We begin with **benzene** and then examine other cyclic, planar, and conjugated ring systems to learn the modern definition of what it means to be aromatic. Then, in Chapter 18, we will learn about the reactions of aromatic compounds, highly unsaturated hydrocarbons that do not undergo addition reactions like other unsaturated compounds. An explanation of this behavior relies on an understanding of the structure of aromatic compounds presented in Chapter 17.


17.1 Background

Benzene (C_6H_6) is the simplest aromatic hydrocarbon (or arene). Since its isolation by Michael Faraday from the oily residue remaining in the illuminating gas lines in London in 1825, it has been recognized as an unusual compound. Based on the calculation introduced in Section 10.2, **benzene has four degrees of unsaturation, making it a highly unsaturated hydrocarbon.** But, whereas unsaturated hydrocarbons such as alkenes, alkynes, and dienes readily undergo addition reactions, *benzene does not.* For example, bromine adds to ethylene to form a dibromide, but benzene is inert under similar conditions.

For 6 C's, the maximum number of H's = $2n + 2$ = 2(6) + 2 = 14. Because benzene contains only 6 H's, it has 14 – 6 = 8 H's fewer than the maximum number. This corresponds to 8 H's/2 H's for each degree of unsaturation = **four degrees of unsaturation in benzene.**




ethylene




benzene

Benzene *does* react with bromine, but only in the presence of $FeBr_3$ (a Lewis acid), and the reaction is a **substitution,** *not* an addition.



Thus, any structure proposed for benzene must account for its high degree of unsaturation and its lack of reactivity towards electrophilic addition.

In the last half of the nineteenth century August Kekulé proposed structures that were close to the modern description of benzene. In the Kekulé model, benzene was thought to be a rapidly equilibrating mixture of two compounds, each containing a six-membered ring with three alternating π bonds. These structures are now called **Kekulé structures.** In the Kekulé description, the bond between any two carbon atoms is sometimes a single bond and sometimes a double bond.



Although benzene is still drawn as a six-membered ring with three alternating π bonds, in reality **there is no equilibrium between two different kinds of benzene molecules.** Instead, current descriptions of benzene are based on resonance and electron delocalization due to orbital overlap, as detailed in Section 17.2.

In the nineteenth century, many other compounds having properties similar to those of benzene were isolated from natural sources. Because these compounds possessed strong and characteristic odors, they were called ***aromatic*** **compounds.** It is their chemical properties, though, not their odor that make these compounds special.

- Aromatic compounds resemble benzene—they are unsaturated compounds that do not undergo the addition reactions characteristic of alkenes.

17.2 The Structure of Benzene

Any structure for benzene must account for the following:

- **Benzene contains a six-membered ring and three additional degrees of unsaturation.**
- **Benzene is planar.**
- **All C−C bond lengths are equal.**

Although the Kekulé structures satisfy the first two criteria, they break down with the third, because having three alternating π bonds would mean that benzene should have three short double bonds alternating with three longer single bonds.



This structure implies that the C−C bonds should have **two different lengths.**

- three longer single bonds in red
- three shorter double bonds in black


Resonance

Some texts draw benzene as a hexagon with an inner circle:



The circle represents the **six π electrons,** distributed over the six atoms of the ring.

Benzene is conjugated, so we must use resonance and orbitals to describe its structure. The resonance description of benzene consists of two equivalent Lewis structures, each with three double bonds that alternate with three single bonds.



hybrid


The electrons in the π bonds are **delocalized** around the ring.

The resonance description of benzene matches the Kekulé description with one important exception. **The two Kekulé representations are *not* in equilibrium with each other.** Instead, the true structure of benzene is a resonance **hybrid** of the two Lewis structures, with the dashed lines of the hybrid indicating the position of the π bonds.

We will use one of the two Lewis structures and not the hybrid in drawing benzene, because it is easier to keep track of the electron pairs in the π bonds (the π electrons).

- **Because each π bond has two electrons, benzene has six π electrons.**


The resonance hybrid of benzene explains why all C−C bond lengths are the same. Each C−C bond is single in one resonance structure and double in the other, so the actual bond length (139 pm) is intermediate between a carbon–carbon single bond (153 pm) and a carbon–carbon double bond (134 pm).



The C−C bonds in benzene are equal and intermediate in length.

Hybridization and Orbitals

Each carbon atom in a benzene ring is surrounded by three atoms and no lone pairs of electrons, making it **sp^2 hybridized and trigonal planar with all bond angles 120°.** Each carbon also has a *p* orbital with one electron that extends above and below the plane of the molecule.



The six adjacent *p* orbitals overlap, delocalizing the six electrons over the six atoms of the ring and making benzene a conjugated molecule. Because each *p* orbital has two lobes, one above and one below the plane of the benzene ring, the overlap of the *p* orbitals creates two "doughnuts" of electron density, as shown in Figure 17.1a. The electrostatic potential plot in Figure 17.1b also shows that the electron-rich region is concentrated above and below the plane of the molecule, where the six π electrons are located.

- **Benzene's six π electrons make it electron rich, so it reacts with electrophiles.**


Figure 17.1 Two views of the electron density in a benzene ring

a. View of the *p* orbital overlap




- Overlap of six adjacent *p* orbitals creates two rings of electron density, one above and one below the plane of the benzene ring.

b. Electrostatic potential plot




- The electron-rich region (in red) is concentrated above and below the ring carbons, where the six π electrons are located. (The electron-rich region below the plane is hidden from view.)

Problem 17.1 Draw all possible resonance structures for the antihistamine diphenhydramine, the active ingredient in Benadryl.



diphenhydramine

Problem 17.2 What orbitals are used to form the labeled bonds in the following molecule? Of the labeled C – C bonds, which is the shortest?




17.3 Nomenclature of Benzene Derivatives


Many organic molecules contain a benzene ring with one or more substituents, so we must learn how to name them. Many common names are recognized by the IUPAC system, however, so this complicates the nomenclature of benzene derivatives somewhat.

17.3A Monosubstituted Benzenes


To name a benzene ring with one substituent, **name the substituent and add the word *benzene*.** Carbon substituents are named as alkyl groups.



ethylbenzene




tert-butylbenzene




chlorobenzene


Many monosubstituted benzenes, such as those with methyl (CH_3-), hydroxy ($-OH$), and amino ($-NH_2$) groups, have common names that you must learn, too.



toluene
(methylbenzene)



phenol
(hydroxybenzene)




aniline
(aminobenzene)

17.3B Disubstituted Benzenes


There are three different ways that two groups can be attached to a benzene ring, so a prefix—**ortho, meta,** or **para**—can be used to designate the relative position of the two substituents. Ortho, meta, and para are also abbreviated as ***o, m,*** and ***p,*** respectively.

1,2-Disubstituted benzene
ortho isomer




o-dibromobenzene
or
1,2-dibromobenzene

1,3-Disubstituted benzene
meta isomer



m-dibromobenzene
or
1,3-dibromobenzene


1,4-Disubstituted benzene
para isomer




p-dibromobenzene
or
1,4-dibromobenzene

If the two groups on the benzene ring are different, **alphabetize the names of the substituents** preceding the word *benzene.* If one of the substituents is part of a **common root,** name the **molecule as a derivative of that monosubstituted benzene.**

Alphabetize two different substituent names:




***o*-bromo*c*hloro-benzene**




***m*-fluoro*n*itro-benzene**

Use a common root name:



p*-bromo*toluene



o*-nitro*phenol


17.3C Polysubstituted Benzenes

For three or more substituents on a benzene ring:

[1] Number to give the lowest possible numbers around the ring.


[2] Alphabetize the substituent names.

[3] When substituents are part of common roots, name the molecule as a derivative of that monosubstituted benzene. The substituent that comprises the common root is located at C1.



- Assign the lowest set of numbers.
- Alphabetize the names of all the substituents.

4-chloro-1-ethyl-2-propylbenzene




- Name the molecule as a derivative of the common root **aniline.**
- Designate the position of the NH_2 group as "1," and then assign the lowest possible set of numbers to the other substituents.

2,5-dichloroaniline

17.3D Naming Aromatic Rings as Substituents

A benzene substituent (C_6H_5-) is called a **phenyl group,** and it can be abbreviated in a structure as **Ph–.**




abbreviated as Ph—


phenyl group
C_6H_5-

- A phenyl group (C_6H_5-) is formed by removing one hydrogen from benzene (C_6H_6).

Benzene, therefore, can be represented as PhH, and phenol would be PhOH.




benzene
PhH




phenol
PhOH

The **benzyl** group contains a benzene ring bonded to a CH_2 group. Thus, a benzyl group and a phenyl group differ by the presence of a CH_2 group.




benzyl group
$C_6H_5CH_2-$





phenyl group
C_6H_5-


Finally, substituents derived from benzene, as well as all other substituted aromatic rings, are collectively called **aryl groups,** abbreviated as Ar–.

Problem 17.3 Give the IUPAC name for each compound.

a. 

b. 

c. 

d. 

Problem 17.4 Draw the structure corresponding to each name:



a. isobutylbenzene
b. *o*-dichlorobenzene
c. *cis*-1,2-diphenylcyclohexane
d. *m*-bromoaniline
e. 4-chloro-1,2-diethylbenzene
f. 3-*tert*-butyl-2-ethyltoluene

Problem 17.5 What is the structure of propofol, which has the IUPAC name 2,6-diisopropylphenol? Propofol is an intravenous medication used to induce and maintain anesthesia.

17.4 Spectroscopic Properties

The important IR and NMR absorptions of aromatic compounds are summarized in Table 17.1.


Table 17.1 Characteristic Spectroscopic Absorptions of Benzene Derivatives

Type of spectroscopy	Type of C, H	Absorption
IR absorptions	C_{sp^2}–H C=C (arene)	3150–3000 cm^{-1} 1600, 1500 cm^{-1}
^{1}H NMR absorptions	 (aryl H)	6.5–8 ppm (highly deshielded protons)
	 (benzylic H)	1.5–2.5 ppm (somewhat deshielded C_{sp^3}–H)
^{13}C NMR absorption	C_{sp^2} of arenes	120–150 ppm

The absorption at 6.5–8.0 ppm in the ^{1}H NMR spectrum is particularly characteristic of compounds containing benzene rings. **All aromatic compounds have highly deshielded protons due to the ring current effect of the circulating π electrons,** as discussed in Section 14.4. Observing whether a new compound absorbs in this region of a ^{1}H NMR spectrum is one piece of data used to determine if it is aromatic.

^{13}C NMR spectroscopy is used to determine the substitution patterns in disubstituted benzenes, because each line in a spectrum corresponds to a different kind of carbon atom. For example, *o*-, *m*-, and *p*-dibromobenzene each exhibit a different number of lines in its ^{13}C NMR spectrum, as shown in Figure 17.2.

Figure 17.2 ^{13}C NMR absorptions of the three isomeric dibromobenzenes





o-dibromobenzene	*m*-dibromobenzene	*p*-dibromobenzene
three types of C's **three** ^{13}C NMR signals	**four** types of C's **four** ^{13}C NMR signals	**two** types of C's **two** ^{13}C NMR signals


- The number of signals (lines) in the ^{13}C NMR spectrum of a disubstituted benzene with two identical groups indicates whether they are ortho, meta, or para to each other.

Problem 17.6 What is the structure of a compound of molecular formula $C_{10}H_{14}O_2$ that shows a strong IR absorption at 3150–2850 cm^{-1} and gives the following ^{1}H NMR absorptions: 1.4 (triplet, 6 H), 4.0 (quartet, 4 H), and 6.8 (singlet, 4 H) ppm?

Problem 17.7 How many ^{13}C NMR signals does each compound exhibit?

a. 

b. 


c. 

17.5 Interesting Aromatic Compounds


BTX contains **b**enzene, **t**oluene, and **x**ylene (the common name for dimethylbenzene).

Benzene and **toluene,** the simplest aromatic hydrocarbons obtained from petroleum refining, are useful starting materials for synthetic polymers. They are two components of the **BTX** mixture added to gasoline to boost octane ratings.


benzene


toluene



p-xylene


naphthalene (used in mothballs)

Compounds containing two or more benzene rings that share carbon–carbon bonds are called **polycyclic aromatic hydrocarbons (PAHs).** Naphthalene, the simplest PAH, is present in mothballs.

Benzo[*a*]pyrene, a more complicated PAH shown in Figure 17.3, is formed by the incomplete combustion of organic materials. It is found in cigarette smoke, automobile exhaust, and the fumes from charcoal grills. When ingested or inhaled, benzo[*a*]pyrene and other similar PAHs are oxidized to carcinogenic products, as discussed in Section 9.18.

Figure 17.3
Benzo[*a*]pyrene, a common PAH




- Benzo[*a*]pyrene, produced by the incomplete oxidation of organic compounds in tobacco, is found in cigarette smoke.

Helicene and **twistoflex** are two synthetic PAHs whose unusual shapes are shown in Figure 17.4. Helicene consists of six benzene rings. Because the rings at both ends are not bonded to each other, all of the rings twist slightly, creating a rigid helical shape that prevents the hydrogen atoms on both ends from crashing into each other. Similarly, to reduce steric hindrance between the hydrogen atoms on nearby benzene rings, twistoflex is also nonplanar.


Both helicene and twistoflex are chiral molecules—that is, they are not superimposable on their mirror images, even though neither of them contains a stereogenic center. It's their shape that makes them chiral, not the presence of carbon atoms bonded to four different groups. Each ring system is twisted into a shape that lacks a mirror plane, and each structure is rigid, thus creating the chirality.

Figure 17.4 Helicene and twistoflex—Two synthetic polycyclic aromatic hydrocarbons




Many widely used drugs contain a benzene ring. Six examples are shown in Figure 17.5.


Figure 17.5
Selected drugs that contain a benzene ring




- Trade name **Zoloft**
- Generic name **sertraline**
- Use: a psychotherapeutic drug for depression and panic disorders




- Trade name **Valium**
- Generic name **diazepam**
- Use: a sedative




- Trade name **Novocain**
- Generic name **procaine**
- Use: a local anesthetic



- Trade name **Viracept**
- Generic name **nelfinavir**
- Use: an antiviral drug used to treat HIV



- Trade name **Viagra**
- Generic name **sildenafil**
- Use: a drug used to treat erectile dysfunction




- Trade name **Claritin**
- Generic name **loratadine**
- Use: an antihistamine for seasonal allergies

17.6 Benzene's Unusual Stability

Considering benzene as the hybrid of two resonance structures adequately explains its equal C−C bond lengths, but does not account for its unusual stability and lack of reactivity toward addition.

Heats of hydrogenation, which were used in Section 16.9 to show that conjugated dienes are more stable than isolated dienes, can also be used to estimate the stability of benzene. Equations [1]–[3] compare the heats of hydrogenation of cyclohexene, cyclohexa-1,3-diene, and benzene, all of which give cyclohexane when treated with excess hydrogen in the presence of a metal catalyst.




The relative stability of conjugated dienes versus isolated dienes was first discussed in Section 16.9.

The addition of one mole of H_2 to cyclohexene releases –120 kJ/mol of energy (Equation [1]). If each double bond is worth –120 kJ/mol of energy, then the addition of two moles of H_2 to cyclohexa-1,3-diene (Equation [2]) should release 2 × –120 kJ/mol = –240 kJ/mol of energy. The observed value, however, is –232 kJ/mol. This is slightly smaller than expected because cyclohexa-1,3-diene is a conjugated diene, and conjugated dienes are more stable than two isolated carbon–carbon double bonds.

The hydrogenations of cyclohexene and cyclohexa-1,3-diene occur readily at room temperature, but benzene can be hydrogenated only under forcing conditions, and even then the reaction is extremely slow. If each double bond is worth –120 kJ/mol of energy, then the addition of three moles of H_2 to benzene should release 3 × –120 kJ/mol = –360 kJ/mol of energy. In fact, the observed heat of hydrogenation is only –208 kJ/mol, which is 152 kJ/mol less than predicted and even lower than the observed value for cyclohexa-1,3-diene. Figure 17.6 compares the hypothetical and observed heats of hydrogenation for benzene.


Figure 17.6
A comparison between the observed and hypothetical heats of hydrogenation for benzene




The huge difference between the hypothetical and observed heats of hydrogenation for benzene cannot be explained solely on the basis of resonance and conjugation.

- **The low heat of hydrogenation of benzene means that benzene is especially stable, even more so than the conjugated compounds introduced in Chapter 16. This unusual stability is characteristic of aromatic compounds.**

Benzene's unusual behavior in chemical reactions is not limited to hydrogenation. As mentioned in Section 17.1, **benzene does not undergo addition reactions typical of other highly unsaturated compounds, including conjugated dienes.** Benzene does not react with Br_2 to yield an addition product. Instead, in the presence of a Lewis acid, bromine *substitutes* for a hydrogen atom, thus yielding a product that retains the benzene ring.



An addition product would no longer contain a benzene ring.



A substitution product still contains a benzene ring.

This behavior is characteristic of aromatic compounds. The structural features that distinguish aromatic compounds from the rest are discussed in Section 17.7.

Problem 17.8 Compounds **A** and **B** are both hydrogenated to methylcyclohexane. Which compound has the larger heat of hydrogenation? Which compound is more stable?



A **B**

17.7 The Criteria for Aromaticity—Hückel's Rule


Four structural criteria must be satisfied for a compound to be aromatic:

- **A molecule must be cyclic, planar, completely conjugated, and contain a particular number of π electrons.**

[1] **A molecule must be cyclic.**

- **To be aromatic, each *p* orbital must overlap with *p* orbitals on two adjacent atoms.**


The *p* orbitals on all six carbons of benzene continuously overlap, so benzene is aromatic. Hexa-1,3,5-triene has six *p* orbitals, too, but the two on the terminal carbons cannot overlap with each other, so **hexa-1,3,5-triene is not aromatic.**



benzene

Every *p* orbital overlaps with two neighboring *p* orbitals.

aromatic




hexa-1,3,5-triene

There can be no overlap between the *p* orbitals on the two terminal C's.

not aromatic

[2] A molecule must be planar.


- All adjacent *p* orbitals must be aligned so that the π electron density can be delocalized.



cyclooctatetraene
not aromatic




a tub-shaped,
eight-membered ring



Adjacent *p* orbitals cannot overlap.
Electrons cannot delocalize.

Cyclooctatetraene resembles benzene in that it is a cyclic molecule with alternating double and single bonds. Cyclooctatetraene is tub shaped, however, **not planar,** so overlap between adjacent π bonds is impossible. **Cyclooctatetraene, therefore, is *not* aromatic,** so it undergoes addition reactions like those of other alkenes.




cyclooctatetraene **addition** product

[3] A molecule must be completely conjugated.


- Aromatic compounds must have a *p* orbital on every atom in the ring.



benzene
a *p* orbital on **every** C
aromatic



cyclohexa-1,3-diene
not aromatic



cyclohepta-1,3,5-triene
not aromatic

Both cyclohexa-1,3-diene and cyclohepta-1,3,5-triene contain at least one carbon atom that does not have a *p* orbital, and so they are not completely conjugated and therefore ***not* aromatic.**

[4] A molecule must satisfy Hückel's rule, and contain a particular number of π electrons.

Some compounds satisfy the first three criteria for aromaticity, but still they show none of the stability typical of aromatic compounds. For example, **cyclobutadiene** is so highly reactive that it can only be prepared at extremely low temperatures.



cyclobutadiene

a planar, cyclic, completely conjugated molecule
that is *not* aromatic

Hückel's rule refers to the number of π electrons, *not* the number of atoms in a particular ring.

It turns out that in addition to being cyclic, planar, and completely conjugated, a compound needs a particular number of π electrons to be aromatic. Erich Hückel first recognized in 1931 that the following criterion, expressed in two parts and now known as **Hückel's rule,** had to be satisfied, as well:

- An aromatic compound must contain $4n + 2$ π electrons (n = 0, 1, 2, and so forth).
- Cyclic, planar, and completely conjugated compounds that contain $4n$ π electrons are especially unstable, and are said to be *antiaromatic.*

Table 17.2
The Number of π Electrons That Satisfy Hückel's Rule

n	4*n* + 2
0	2
1	6
2	10
3	14
4, etc.	18

Thus, compounds that contain 2, 6, 10, 14, 18, and so forth π electrons are aromatic, as shown in Table 17.2. **Benzene is aromatic and especially stable because it contains 6 π electrons. Cyclobutadiene is antiaromatic and especially unstable because it contains 4 π electrons.**

Benzene
An aromatic compound



4*n* + 2 = 4(1) + 2 =
6 π electrons
aromatic

Cyclobutadiene
An antiaromatic compound



4*n* = 4(1) =
4 π electrons
antiaromatic

Considering aromaticity, all compounds can be classified in one of three ways:

[1]	Aromatic	• A cyclic, planar, completely conjugated compound with 4*n* + 2 π electrons
[2]	Antiaromatic	• A cyclic, planar, completely conjugated compound with 4*n* π electrons
[3]	Not aromatic or nonaromatic	• A compound that lacks one (or more) of the four requirements to be aromatic or antiaromatic

Note, too, the relationship between each compound type and a similar open-chained molecule having the same number of π electrons.

- An aromatic compound is *more* stable than a similar acyclic compound having the same number of π electrons. Benzene is more stable than hexa-1,3,5-triene.
- An antiaromatic compound is *less* stable than an acyclic compound having the same number of π electrons. Cyclobutadiene is less stable than buta-1,3-diene.
- A compound that is not aromatic is *similar* in stability to an acyclic compound having the same number of π electrons. Cyclohexa-1,3-diene is similar in stability to *cis,cis*-hexa-2,4-diene, so it is not aromatic.


benzene

and


hexa-1,3,5-triene

more stable
aromatic


cyclobutadiene

and


buta-1,3-diene

less stable
antiaromatic



cyclohexa-1,3-diene

and


cis,cis-hexa-2,4-diene



similar stability
nonaromatic

^{1}H NMR spectroscopy readily indicates whether a compound is aromatic. The protons on sp^2 hybridized carbons in aromatic hydrocarbons are highly deshielded and absorb at 6.5–8 ppm, whereas unsaturated hydrocarbons that are not aromatic absorb at 4.5–6 ppm, typical of protons bonded to the C=C of an alkene. Thus, benzene absorbs at 7.3 ppm, whereas cyclooctatetraene, which is not aromatic, absorbs farther upfield, at 5.8 ppm for the protons on its sp^2 hybridized carbons.



Many compounds in addition to benzene are aromatic. Several examples are presented in Section 17.8.

Problem 17.9 Estimate where the protons bonded to the sp^2 hybridized carbons will absorb in the ^{1}H NMR spectrum of each compound.

a.  b.  c. 

17.8 Examples of Aromatic Compounds


In Section 17.8 we look at many different types of aromatic compounds.

17.8A Aromatic Compounds with a Single Ring


Benzene is the most common aromatic compound having a single ring. **Completely conjugated rings larger than benzene are also aromatic if they are planar and have $4n + 2$ π electrons.**

- **Hydrocarbons containing a single ring with alternating double and single bonds are called *annulenes.***

To name an annulene, indicate the number of atoms in the ring in brackets and add the word *annulene*. Thus, benzene is [6]-annulene. Both **[14]-annulene** and **[18]-annulene** are cyclic, planar, completely conjugated molecules that follow Hückel's rule, and so they are aromatic.




[14]-annulene
$4n + 2 = 4(3) + 2 =$
14 π electrons
aromatic



[18]-annulene
$4n + 2 = 4(4) + 2 =$
18 π electrons
aromatic


[10]-Annulene has 10 π electrons, which satisfies Hückel's rule, but a planar molecule would place the two H atoms inside the ring too close to each other, so the ring puckers to relieve this strain. Because **[10]-annulene is not planar,** the 10 π electrons can't delocalize over the entire ring and it is **not aromatic.**




Problem 17.10 Would [16]-, [20]- or [22]-annulene be aromatic if each ring is planar?

17.8B Aromatic Compounds with More Than One Ring

Hückel's rule for determining aromaticity can be applied only to monocyclic systems, but many aromatic compounds containing several benzene rings joined together are also known. Two or more six-membered rings with alternating double and single bonds can be fused together to form **polycyclic aromatic hydrocarbons (PAHs).** Joining two benzene rings together forms **naphthalene.** There are two different ways to join three rings together, forming **anthracene** and **phenanthrene,** and many more complex hydrocarbons are known.



As the number of fused benzene rings increases, the number of resonance structures increases as well. Although two resonance structures can be drawn for benzene, naphthalene is a hybrid of three resonance structures.



Problem 17.11 Draw the four resonance structures for anthracene.


17.8C Aromatic Heterocycles

Recall from Section 9.3 that a **heterocycle** is a ring that contains at least one heteroatom.

Heterocycles containing oxygen, nitrogen, or sulfur—atoms that also have at least one lone pair of electrons—can also be aromatic. With heteroatoms, we must always determine whether the lone pair is localized on the heteroatom or part of the delocalized π system. Two examples, **pyridine** and **pyrrole,** illustrate these different possibilities.

Pyridine


Pyridine is a heterocycle containing a six-membered ring with three π bonds and one nitrogen atom. Like benzene, two resonance structures (with all neutral atoms) can be drawn.



two resonance structures for pyridine
6 π electrons

Pyridine is cyclic, planar, and completely conjugated, because the three single and double bonds alternate around the ring. **Pyridine has six π electrons, two from each π bond, thus satisfying Hückel's rule and making pyridine aromatic.** The nitrogen atom of pyridine also has a nonbonded electron pair, which is localized on the N atom, so it is *not* part of the delocalized π electron system of the aromatic ring.

How is the nitrogen atom of the pyridine ring hybridized? The N atom is surrounded by three groups (two atoms and a lone electron pair), making it ***sp*^{2} hybridized,** and leaving one unhybridized *p* orbital with one electron that overlaps with adjacent *p* orbitals. The lone pair on N resides in an sp^2 hybrid orbital that is perpendicular to the delocalized π electrons.



Pyrrole

Pyrrole contains a five-membered ring with two π bonds and one nitrogen atom. The N atom also has a lone pair of electrons.




pyrrole

Pyrrole is cyclic and planar, with a total of four π electrons from the two π bonds. Is the nonbonded electron pair localized on N or part of a delocalized π electron system? The lone pair on N is *adjacent* to a double bond. Recall the following general rule from Section 16.5:

- In a system X=Y–Z:, Z is generally sp^2 hybridized and the lone pair occupies a *p* orbital to make the system conjugated.

If the lone pair on the N atom occupies a *p* orbital:

- **Pyrrole has a *p* orbital on every adjacent atom, so it is completely conjugated.**
- **Pyrrole has six π electrons—four from the π bonds and two from the lone pair.**




Because pyrrole is cyclic, planar, completely conjugated, and has $4n + 2$ π electrons, **pyrrole is aromatic. The number of electrons—not the size of the ring—determines whether a compound is aromatic.**

Electrostatic potential maps, shown in Figure 17.7 for pyridine and pyrrole, confirm that the **lone pair in pyridine is localized on N,** whereas the **lone pair in pyrrole is part of the delocalized π system.** Thus, a fundamental difference exists between the N atoms in pyridine and pyrrole.


Figure 17.7
Electrostatic potential maps of pyridine and pyrrole

pyridine



- In pyridine, the nonbonded electron pair is localized on the N atom in an sp^2 hybridized orbital, as shown by the region of high electron density (in red) on N.

pyrrole



- In pyrrole, the nonbonded electron pair is in a *p* orbital and is delocalized over the ring, so the entire ring is electron rich (red).


- **When a heteroatom is already part of a double bond (as in the N of pyridine), its lone pair *cannot* occupy a *p* orbital and so it cannot be delocalized over the ring.**
- **When a heteroatom is *not* part of a double bond (as in the N of pyrrole), its lone pair can be located in a *p* orbital and *delocalized* over a ring to make it aromatic.**



Scombroid fish poisoning, associated with facial flushing, hives, and general itching, is caused by the ingestion of inadequately refrigerated fish, typically mahimahi (pictured) and tuna. Bacteria convert the amino acid histidine (Chapter 29) to histamine, which, when consumed in large amounts, results in this clinical syndrome.


Histamine

Histamine, a biologically active amine formed in many tissues, has an aromatic heterocycle with two N atoms, one of which is similar to the N atom of pyridine and one of which is similar to the N atom of pyrrole.



histamine

Histamine has a five-membered ring with two π bonds and two nitrogen atoms, each of which contains a lone pair of electrons. The heterocycle has four π electrons from the two double bonds. The lone pair on the N in red also occupies a *p* orbital, making the heterocycle completely conjugated, and giving it a total of six π electrons. The lone pair on this N atom is thus delocalized over the five-membered ring and the heterocycle is aromatic. The lone pair on the N in blue occupies an sp^2 hybrid orbital perpendicular to the delocalized π electrons.



N: The lone pair resides in a *p* orbital.

N: The lone pair resides in an sp^2 hybrid orbital.


- N (in red) resembles the N atom of pyrrole.
- N (in blue) resembles the N atom of pyridine.

Histamine produces a wide range of physiological effects in the body. Excess histamine is responsible for the runny nose and watery eyes symptomatic of hay fever. It also stimulates the overproduction of stomach acid, and contributes to the formation of hives. These effects result from the interaction of histamine with two different cellular receptors. We will learn more about antihistamines and antiulcer drugs, compounds that block the effects of histamine, in Section 25.6.

Problem 17.12 Which heterocycles are aromatic?


a. 

b. 

c. 

d. 

Problem 17.13 (a) How is each N atom in quinine, an effective antimalarial drug that reduces fever, hybridized? (b) In what type of orbital does the lone pair on each N reside?




quinine



Quinine is isolated from the bark of the cinchona tree native to the Andes Mountains.

Problem 17.14 Januvia, the trade name for sitagliptin, was introduced in 2006 for the treatment of type 2 diabetes. (a) Explain why the five-membered ring in sitagliptin is aromatic. (b) Determine the hybridization of each N atom. (c) In what type of orbital does the lone pair on each N atom reside?



sitagliptin



Januvia increases the body's ability to lower blood sugar levels, so it is used alone or in combination with other drugs to treat type 2 diabetes.

17.8D Charged Aromatic Compounds


Both negatively and positively charged ions can also be aromatic if they satisfy all the necessary criteria.

Cyclopentadienyl Anion

The **cyclopentadienyl anion** is a cyclic and planar anion with two double bonds and a nonbonded electron pair. In this way it resembles pyrrole. The two π bonds contribute four electrons and the lone pair contributes two more, for a total of six. By Hückel's rule, having **six π electrons confers aromaticity.** Like the N atom in pyrrole, the **negatively charged carbon atom must be sp^2 hybridized,** and the **nonbonded electron pair must occupy a *p* orbital** for the ring to be completely conjugated.



cyclopentadienyl anion
all sp^2 hybridized C's
6 π electrons



The lone pair resides in a *p* orbital.

- **The cyclopentadienyl anion is aromatic because it is cyclic, planar, completely conjugated, and has six π electrons.**

We can draw **five equivalent resonance structures for the cyclopentadienyl anion,** delocalizing the negative charge over every carbon atom of the ring.



Although five resonance structures can also be drawn for both the **cyclopentadienyl cation** and **radical,** only the cyclopentadienyl anion has six π electrons, a number that satisfies Hückel's rule. The cyclopentadienyl cation has four π electrons, making it antiaromatic and especially unstable. The cyclopentadienyl radical has five π electrons, so it is neither aromatic nor antiaromatic. Having the "right" number of electrons is necessary for a species to be unusually stable by virtue of aromaticity.



cyclopentadienyl anion
- **6** π electrons
- contains $4n + 2$ π electrons

aromatic



cyclopentadienyl cation
- **4** π electrons
- contains $4n$ π electrons


antiaromatic



cyclopentadienyl radical
- **5** π electrons
- does not contain either $4n$ or $4n + 2$ π electrons

nonaromatic

The cyclopentadienyl anion is readily formed from cyclopentadiene by a Brønsted–Lowry acid–base reaction.




cyclopentadiene
not aromatic
$pK_a = 15$

cyclopentadienyl anion
aromatic
a stabilized conjugate base

Cyclopentadiene itself is not aromatic because it is not fully conjugated. The cyclopentadienyl anion, however, is aromatic, so it is a very stable base. As such, it makes cyclopentadiene more acidic than other hydrocarbons. In fact, the pK_a of cyclopentadiene is 15, much lower (more acidic) than the pK_a of any C−H bond discussed thus far.


- **Cyclopentadiene is more acidic than many hydrocarbons because its conjugate base is aromatic.**

Problem 17.15 Draw the product formed when cyclohepta-1,3,5-triene ($pK_a = 39$) is treated with a strong base. Why is its pK_a so much higher than the pK_a of cyclopentadiene?



cyclohepta-1,3,5-triene
$pK_a = 39$


Problem 17.16 Rank the following compounds in order of increasing acidity.



The cyclopentadienyl anion and the tropylium cation both illustrate an important principle: The **number of π electrons determines aromaticity,** not the number of atoms in a ring or the number of *p* orbitals that overlap. The cyclopentadienyl anion and tropylium cation are aromatic because they each have six π electrons.

Tropylium Cation

The **tropylium cation** is a planar carbocation with three double bonds and a positive charge contained in a seven-membered ring. This carbocation is completely conjugated, because the positively charged carbon is sp^2 hybridized and has a vacant *p* orbital that overlaps with the six *p* orbitals from the carbons of the three double bonds. **Because the tropylium cation has three π bonds and no other nonbonded electron pairs, it contains six π electrons,** thereby satisfying Hückel's rule.



tropylium cation
all sp^2 hybridized C's
6 π electrons

One *p* orbital is vacant.

- **The tropylium cation is aromatic because it is cyclic, planar, completely conjugated, and has six π electrons delocalized over the seven atoms of the ring.**


Problem 17.17 Draw the seven resonance structures for the tropylium cation.

Problem 17.18 Assuming the rings are planar, which ions are aromatic?


a. 

b. 

c. 


d. 

Problem 17.19 Compound **A** exhibits a peak in its ^{1}H NMR spectrum at 7.6 ppm, indicating that it is aromatic. (a) How are the carbon atoms of the triple bonds hybridized? (b) In what type of orbitals are the π electrons of the triple bonds contained? (c) How many π electrons are delocalized around the ring in **A?**

A = 

17.9 What Is the Basis of Hückel's Rule?

Why does the number of π electrons determine whether a compound is aromatic? Cyclobutadiene is cyclic, planar, and completely conjugated, just like benzene, but why is benzene aromatic and cyclobutadiene antiaromatic?



A complete explanation is beyond the scope of an introductory organic chemistry text, but nevertheless, you can better understand the basis of aromaticity by learning more about orbitals and bonding.

17.9A Bonding and Antibonding Orbitals

So far we have used the following basic concepts to describe how bonds are formed:

- Hydrogen uses its 1*s* orbital to form σ bonds with other elements.
- Second-row elements use hybrid orbitals (*sp*, sp^2, or sp^3) to form σ bonds.
- Second-row elements use *p* orbitals to form π bonds.

This description of bonding is called **valence bond theory.** In valence bond theory, a covalent bond is formed by the overlap of two atomic orbitals, and the electron pair in the resulting bond is shared by both atoms. Thus, a carbon–carbon double bond consists of a σ bond, formed by overlap of two sp^2 hybrid orbitals, each containing one electron, and a π bond, formed by overlap of two *p* orbitals, each containing one electron.


This description of bonding works well for most of the organic molecules we have encountered thus far. Unfortunately, it is inadequate for describing systems with many adjacent *p* orbitals that overlap, as there are in aromatic compounds. To more fully explain the bonding in these systems, we must utilize **molecular orbital (MO) theory.**

MO theory describes bonds as the mathematical combination of atomic orbitals that form a new set of orbitals called **molecular orbitals (MOs).** A molecular orbital occupies a region of space *in a molecule* where electrons are likely to be found. When forming molecular orbitals from atomic orbitals, keep in mind:

- A set of *n* atomic orbitals forms *n* molecular orbitals.

If *two* atomic orbitals combine, *two* molecular orbitals are formed. This is fundamentally different than valence bond theory. Because aromaticity is based on *p* orbital overlap, what does MO theory predict will happen when two *p* (atomic) orbitals combine?

The two lobes of each *p* orbital are opposite in phase, with a node of electron density at the nucleus. When two *p* orbitals combine, two molecular orbitals should form. The two *p* orbitals can add together constructively—that is, with like phases interacting—or destructively—that is, with opposite phases interacting.




- When two *p* orbitals of similar phase overlap side-by-side, a π bonding molecular orbital results.
- When two *p* orbitals of opposite phase overlap side-by-side, a π* antibonding molecular orbital results.

A π bonding MO is lower in energy than the two atomic *p* orbitals from which it is formed because a stable bonding interaction results when orbitals of similar phase combine. A bonding interaction holds nuclei together. Similarly, a π* antibonding MO is higher in energy because a destabilizing node results when orbitals of opposite phase combine. A destabilizing interaction pushes nuclei apart.

If two atomic *p* orbitals each have one electron and then combine to form MOs, the two electrons will occupy the lower energy π bonding MO, as shown in Figure 17.8.

Figure 17.8
Combination of two *p* orbitals to form π and π* molecular orbitals



- Two atomic *p* orbitals combine to form two molecular orbitals. The bonding π MO is lower in energy than the two *p* orbitals from which it was formed, and the antibonding π* MO is higher in energy than the two *p* orbitals from which it was formed.
- Two electrons fill the lower energy bonding MO first.


17.9B Molecular Orbitals Formed When More Than Two *p* Orbitals Combine

The molecular orbital description of benzene is much more complex than the two MOs formed in Figure 17.8. Because each of the six carbon atoms of benzene has a *p* orbital, six atomic *p* orbitals combine to form six π molecular orbitals, as shown in Figure 17.9. A description of the exact appearance and energies of these six MOs requires more sophisticated mathematics and understanding of MO theory than is presented in this text. Nevertheless, note that the six MOs are labeled ψ_1–ψ_6, with ψ_1 being the lowest in energy and ψ_6 the highest.

The most important features of the six benzene MOs are as follows:

- **The larger the number of bonding interactions, the lower in energy the MO.** The lowest energy molecular orbital (ψ_1) has all bonding interactions between the *p* orbitals.
- **The larger the number of nodes, the higher in energy the MO.** The highest energy MO (ψ_6*) has all nodes between the *p* orbitals.
- Three MOs are lower in energy than the starting *p* orbitals, making them bonding MOs (ψ_1, ψ_2, and ψ_3), whereas three MOs are higher in energy than the starting *p* orbitals, making them antibonding MOs (ψ_4*, ψ_5*, and ψ_6*).
- The two pairs of MOs (ψ_2 and ψ_3; ψ_4* and ψ_5*) with the same energy are called **degenerate orbitals.**

Figure 17.9
How the six *p* orbitals of benzene overlap to form six molecular orbitals



- Depicted in this diagram are the interactions of the six atomic *p* orbitals of benzene, which form six molecular orbitals. When orbitals of like phase combine, a bonding interaction results. When orbitals of opposite phase combine, a destabilizing node results.

- **The highest energy orbital that contains electrons is called the *highest occupied molecular orbital* (HOMO).** For benzene, the degenerate orbitals ψ_2 and ψ_3 are the HOMOs.
- **The lowest energy orbital that does *not* contain electrons is called the *lowest unoccupied molecular orbital* (LUMO).** For benzene, the degenerate orbitals ψ_4^* and ψ_5^* are the LUMOs.

To fill the MOs, the six electrons are added, two to an orbital, beginning with the lowest energy orbital. As a result, the six electrons completely fill the bonding MOs, leaving the antibonding MOs empty. This is what gives benzene and other aromatic compounds their special stability and this is why six π electrons satisfies Hückel's $4n + 2$ rule.

- **All bonding MOs (and HOMOs) are completely filled in aromatic compounds. No π electrons occupy antibonding MOs.**

17.10 The Inscribed Polygon Method for Predicting Aromaticity

An inscribed polygon is also called a **Frost circle.**


To predict whether a compound has π electrons completely filling bonding MOs, we must know how many bonding molecular orbitals and how many π electrons it has. It is possible to predict the relative energies of cyclic, completely conjugated compounds, without sophisticated math (or knowing what the resulting MOs look like) by using the **inscribed polygon method.**

How To Use the Inscribed Polygon Method to Determine the Relative Energies of MOs for Cyclic, Completely Conjugated Compounds

Example **Plot the relative energies of the MOs of benzene.**

Step [1] **Draw the polygon in question inside a circle with its vertices touching the circle and one of the vertices pointing down. Mark the points at which the polygon intersects the circle.**

- Inscribe a hexagon inside a circle for benzene. The six vertices of the hexagon form six points of intersection, corresponding to the six MOs of benzene. The pattern—a single MO having the lowest energy, two degenerate pairs of MOs, and a single highest energy MO—matches that found in Figure 17.9.




Step [2] **Draw a line horizontally through the center of the circle and label MOs as bonding, nonbonding, or antibonding.**

- **MOs below this line are bonding,** and lower in energy than the *p* orbitals from which they were formed. Benzene has three bonding MOs.
- **MOs at this line are nonbonding,** and equal in energy to the *p* orbitals from which they were formed. Benzene has no nonbonding MOs.
- **MOs above this line are antibonding,** and higher in energy than the *p* orbitals from which they were formed. Benzene has three antibonding MOs.


Step [3] **Add the electrons, beginning with the lowest energy MO.**

- **All the bonding MOs (and the HOMOs) are completely filled in aromatic compounds. No π electrons occupy antibonding MOs.**
- Benzene is aromatic because it has six π electrons that completely fill the bonding MOs.




This method works for all monocyclic, completely conjugated hydrocarbons regardless of ring size. Figure 17.10 illustrates MOs for completely conjugated five- and seven-membered rings using this method. The total number of MOs always equals the number of vertices of the polygon. Because both systems have three bonding MOs, each needs six π electrons to fully occupy them, making the cyclopentadienyl anion and the tropylium cation aromatic, as we learned in Section 17.8D.

Figure 17.10
Using the inscribed polygon method for five- and seven-membered rings



The inscribed polygon method is consistent with Hückel's $4n + 2$ rule; that is, there is always one lowest energy bonding MO that can hold two π electrons and the other bonding MOs come in degenerate pairs that can hold a total of four π electrons. For the compound to be aromatic, these MOs must be completely filled with electrons, so the "magic numbers" for aromaticity fit Hückel's $4n + 2$ rule (Figure 17.11).

Figure 17.11
MO patterns for cyclic, completely conjugated systems



Sample Problem 17.1 Use the inscribed polygon method to show why cyclobutadiene is not aromatic.




cyclobutadiene
4 π electrons

Solution


Cyclobutadiene has four MOs (formed from its four *p* orbitals), to which its four π electrons must be added.

Step [1] Inscribe a square with a vertex down and mark its four points of intersection with the circle.



- The four points of intersection correspond to the four MOs of cyclobutadiene.

Steps [2] and [3] Draw a line through the center of the circle, label the MOs, and add the electrons.



- Cyclobutadiene has four MOs—one bonding, two nonbonding, and one antibonding.
- Adding cyclobutadiene's four π electrons to these orbitals places two in the lowest energy bonding MO and one each in the two nonbonding MOs.
- Separating electrons in two degenerate MOs keeps like charges farther away from each other.

Conclusion: Cyclobutadiene is not aromatic because its HOMOs, two degenerate nonbonding MOs, are not completely filled.


The procedure followed in Sample Problem 17.1 also illustrates why cyclobutadiene is antiaromatic. Having the two unpaired electrons in nonbonding MOs suggests that cyclobutadiene should be a highly unstable diradical. In fact, antiaromatic compounds resemble cyclobutadiene because their HOMOs contain two unpaired electrons, making them especially unstable.

Problem 17.20 Use the inscribed polygon method to show why the following cation is aromatic.



Problem 17.21 Use the inscribed polygon method to show why the cyclopentadienyl cation and radical are not aromatic.

17.11 Buckminsterfullerene—Is It Aromatic?







Diamond and graphite are two elemental forms of carbon.

The two most common elemental forms of carbon are diamond and graphite. Diamond, one of the hardest substances known, is used for industrial cutting tools, whereas graphite, a slippery black substance, is used as a lubricant. Their physical characteristics are so different because their molecular structures are very different.

The structure of diamond consists of a continuous tetrahedral network of sp^3 hybridized carbon atoms, thus creating an infinite array of chair cyclohexane rings. The structure of graphite, on the other hand, consists of parallel sheets of sp^2 hybridized carbon atoms, thus creating an infinite array of benzene rings. The parallel sheets are then held together by weak intermolecular interactions.



diamond
an "infinite" array of six-membered rings, covalently bonded in three dimensions



graphite
an "infinite" array of benzene rings, covalently bonded in two dimensions

Three sheets of graphite, viewed edge-on




Graphite exists in planar sheets of benzene rings, held together by weak intermolecular forces.

Buckminsterfullerene (C_{60}) is a third elemental form of carbon. Its structure consists of 20 hexagons and 12 pentagons of sp^2 hybridized carbon atoms joined in a spherical arrangement. It is completely conjugated because each carbon atom has a *p* orbital with an electron in it.



Buckminsterfullerene (or buckyball) was discovered by Smalley, Curl, and Kroto, who shared the 1996 Nobel Prize in Chemistry for their work. Its unusual name stems from its shape, which resembles the geodesic dome invented by R. Buckminster Fuller. The pattern of five- and six-membered rings also resembles the pattern of rings on a soccer ball.



buckminsterfullerene, C_{60}

20 hexagons + 12 pentagons
of carbon atoms joined together

The 60 C's of buckminsterfullerene are drawn. Each C also contains a *p* orbital with one electron, which is not drawn.

Is C_{60} aromatic? Although it is completely conjugated, it is not planar. Because of its curvature, it is not as stable as benzene. In fact, it undergoes addition reactions with electrophiles in much the same way as ordinary alkenes. Benzene, on the other hand, undergoes substitution reactions with electrophiles, which preserves the unusually stable benzene ring intact. These reactions are the subject of Chapter 18.

Problem 17.22 How many ^{13}C NMR signals does C_{60} exhibit?

KEY CONCEPTS

Benzene and Aromatic Compounds


Comparing Aromatic, Antiaromatic, and Nonaromatic Compounds (17.7)

- **Aromatic compound**
  - A cyclic, planar, completely conjugated compound that contains $4n + 2$ π electrons (n = 0, 1, 2, 3, and so forth)
  - An aromatic compound is more stable than a similar acyclic compound having the same number of π electrons.
- **Antiaromatic compound**
  - A cyclic, planar, completely conjugated compound that contains $4n$ π electrons (n = 0, 1, 2, 3, and so forth)
  - An antiaromatic compound is less stable than a similar acyclic compound having the same number of π electrons.
- **Nonaromatic compound**
  - A compound that lacks one (or more) of the four requirements to be aromatic or antiaromatic

Properties of Aromatic Compounds


- Every atom in an aromatic ring has a *p* orbital to delocalize electron density (17.2).
- Aromatic compounds are unusually stable. $\Delta H°$ for hydrogenation is much less than expected, given the number of degrees of unsaturation (17.6).
- Aromatic compounds do not undergo the usual addition reactions of alkenes (17.6).
- ^1H NMR spectra show highly deshielded protons because of ring currents that reinforce the applied magnetic field (17.4).
- All bonding MOs and HOMOs are completely filled and no electrons occupy antibonding orbitals (17.9).

Examples of Aromatic Compounds with Six π Electrons (17.8)



benzene | pyridine | pyrrole | cyclopentadienyl anion | tropylium cation

Examples of Compounds That Are Not Aromatic (17.8)



not cyclic



not planar





not completely conjugated

PROBLEMS


Problems Using Three-Dimensional Models


17.23 Name each compound and state how many lines are observed in its ^{13}C NMR spectrum.


a. 

b. 

17.24 Classify each compound as aromatic, antiaromatic, or not aromatic.

a. 

b. 

c. 

Benzene Structure and Nomenclature


17.25 Early structural studies on benzene had to explain the following experimental evidence. When benzene was treated with Br_2 (plus a Lewis acid), a single substitution product of molecular formula C_6H_5Br was formed. When this product was treated with another equivalent of Br_2, three different compounds of molecular formula $C_6H_4Br_2$ were formed.


a. Explain why a single Kekulé structure is consistent with the first result, but does not explain the second result.


b. Then explain why a resonance description of benzene is consistent with the results of both reactions.


17.26 Draw all aromatic hydrocarbons that have molecular formula C_8H_{10}. For each compound, determine how many isomers of molecular formula C_8H_9Br would be formed if one H atom on the benzene ring were replaced by a Br atom.


17.27 Give the IUPAC name for each compound.


a. 


c. 


e. 

g. 

b. 

d. 

f. 

h. 





17.28 Draw a structure corresponding to each name.

a. *p*-dichlorobenzene
b. *p*-iodoaniline
c. *o*-bromonitrobenzene
d. 2,6-dimethoxytoluene
e. 2-phenylprop-2-en-1-ol
f. *trans*-1-benzyl-3-phenylcyclopentane




17.29 a. Draw the 14 constitutional isomers of molecular formula C_8H_9Cl that contain a benzene ring.
b. Name all compounds that contain a trisubstituted benzene ring.
c. For which compound(s) are stereoisomers possible? Draw all possible stereoisomers.

Aromaticity





17.30 How many π electrons are contained in each molecule?

a. 
c. 
e. 
b. 
d. 



17.31 Which compounds are aromatic? For any compound that is not aromatic, state why this is so.

a. 
b. 
c. 
d. 


17.32 Which of the following heterocycles are aromatic?

a. 
c. 
e. 
g. 
b. 
d. 
f. 
h. 

17.33 Label each compound as aromatic, antiaromatic, or not aromatic. Assume all completely conjugated rings are planar.


a. 
b. 
c. 
d. 


17.34 Hydrocarbon **A** possesses a significant dipole, even though it is composed of only C – C and C – H bonds. Explain why the dipole arises and use resonance structures to illustrate the direction of the dipole. Which ring is more electron rich?




A


17.35 Pentalene, azulene, and heptalene are conjugated hydrocarbons that do not contain a benzene ring. Which hydrocarbons are especially stable or unstable based on the number of π electrons they contain? Explain your choices.


pentalene


azulene


heptalene


17.36 The purine heterocycle occurs commonly in the structure of DNA.



purine


a. How is each N atom hybridized?
b. In what type of orbital does each lone pair on a N atom reside?
c. How many π electrons does purine contain?
d. Why is purine aromatic?

17.37 Methotrexate, a drug that inhibits the metabolism of folic acid, is used in the treatment of a variety of cancers and autoimmune disorders such as rheumatoid arthritis. (a) GIve the hybridization of each N atom in methotrexate. (b) In what type of orbital does the lone pair of each N reside? (c) Explain why the bicyclic ring system that contains four N atoms is aromatic.



methotrexate


17.38



C


a. How many π electrons does **C** contain?
b. How many π electrons are delocalized in the ring?
c. Explain why **C** is aromatic.

17.39 AZT was the first drug approved to treat HIV, the virus that causes AIDS. Explain why the six-membered ring of AZT is aromatic.




AZT


17.40 Explain the observed rate of reactivity of the following 2° alkyl halides in an S_N1 reaction.



17.41 Draw a stepwise mechanism for the following reaction.



17.42 Explain why α-pyrone reacts with Br_2 to yield a substitution product (like benzene does), rather than an addition product to one of its C=C bonds.



α-pyrone


Resonance

17.43 Draw additional resonance structures for each species.


a. 
cyclopropenyl radical

b. 

pyrrole

c. 
phenanthrene

17.44 The carbon–carbon bond lengths in naphthalene are not equal. Use a resonance argument to explain why bond (a) is shorter than bond (b).



17.45



pyrrole




furan

a. Draw all reasonable resonance structures for pyrrole and explain why pyrrole is less resonance stabilized than benzene.
b. Draw all reasonable resonance structures for furan and explain why furan is less resonance stabilized than pyrrole.

Acidity

17.46 Which compound in each pair is the stronger acid?

a.  or 


b.  or 

17.47 Treatment of indene with $NaNH_2$ forms its conjugate base in a Brønsted–Lowry acid–base reaction. Draw all reasonable resonance structures for indene's conjugate base, and explain why the pK_a of indene is lower than the pK_a of most hydrocarbons.




indene
pK_a = 20

17.48 Considering both 5-methylcyclopenta-1,3-diene (**A**) and 7-methylcyclohepta-1,3,5-triene (**B**), which labeled H atom is most acidic? Which labeled H atom is least acidic? Explain your choices.




A



B

17.49 Draw the conjugate bases of pyrrole and cyclopentadiene. Explain why the sp^3 hybridized C – H bond of cyclopentadiene is more acidic than the N – H bond of pyrrole.


17.50 a. Explain why protonation of pyrrole occurs at C2 to form **A,** rather than on the N atom to form **B.**
b. Explain why **A** is more acidic than **C,** the conjugate acid of pyridine.




pyrrole

pK_a = 0.4

A



B




pK_a = 5.3

C


Inscribed Polygon Method

17.51 Use the inscribed polygon method to show the pattern of molecular orbitals in cyclooctatetraene.




a. Label the MOs as bonding, antibonding, or nonbonding.
b. Indicate the arrangement of electrons in these orbitals for cyclooctatetraene, and explain why cyclooctatetraene is not aromatic.
c. Treatment of cyclooctatetraene with potassium forms a dianion. How many π electrons does this dianion contain?
d. How are the π electrons in this dianion arranged in the molecular orbitals?
e. Classify the dianion of cyclooctatetraene as aromatic, antiaromatic, or not aromatic, and explain why this is so.

17.52 Use the inscribed polygon method to show the pattern of molecular orbitals in cyclonona-1,3,5,7-tetraene and use it to label its cation, radical, and anion as aromatic, antiaromatic, or not aromatic.



Spectroscopy

17.53 How many ^{13}C NMR signals does each compound exhibit?




17.54 Which of the diethylbenzene isomers (ortho, meta, or para) corresponds to each set of ^{13}C NMR spectral data?


[A] ^{13}C NMR signals: 16, 29, 125, 127.5, 128.4, and 144 ppm
[B] ^{13}C NMR signals: 15, 26, 126, 128, and 142 ppm
[C] ^{13}C NMR signals: 16, 29, 128, and 141 ppm

17.55 Propose a structure consistent with each set of data.


a. $C_{10}H_{14}$: IR absorptions at 3150–2850, 1600, and 1500 cm^{-1}



b. C_9H_{12}: ^{13}C NMR signals at 21, 127, and 138 ppm



c. C_8H_{10}: IR absorptions at 3108–2875, 1606, and 1496 cm^{-1}



17.56 Propose a structure consistent with each set of data.

a. Compound **A:**

Molecular formula: $C_8H_{10}O$

IR absorption at 3150–2850 cm^{-1}

^{1}H NMR data: 1.4 (triplet, 3 H), 3.95 (quartet, 2 H), and 6.8–7.3 (multiplet, 5 H) ppm


b. Compound **B:**

Molecular formula: $C_9H_{10}O_2$


IR absorption at 1669 cm^{-1}

^{1}H NMR data: 2.5 (singlet, 3 H), 3.8 (singlet, 3 H), 6.9 (doublet, 2 H), and 7.9 (doublet, 2 H) ppm

17.57 Thymol (molecular formula $C_{10}H_{14}O$) is the major component of the oil of thyme. Thymol shows IR absorptions at 3500–3200, 3150–2850, 1621, and 1585 cm^{-1}. The ^{1}H NMR spectrum of thymol is given below. Propose a possible structure for thymol.



17.58 You have a sample of a compound of molecular formula $C_{11}H_{15}NO_2$, which has a benzene ring substituted by two groups, $(CH_3)_2N-$ and $-CO_2CH_2CH_3$, and exhibits the given ^{13}C NMR. What disubstituted benzene isomer corresponds to these ^{13}C data?



General Problems

17.59 Explain why tetrahydrofuran has a higher boiling point and is much more water soluble than furan, even though both compounds are cyclic ethers containing four carbons.




tetrahydrofuran




furan

17.60 Rizatriptan (trade name Maxalt) is a prescription drug used for the treatment of migraines. (a) How many aromatic rings does rizatriptan contain? (b) Determine the hybridization of each N atom. (c) In what type of orbital does the lone pair on each N reside? (d) Draw all the resonance structures for rizatriptan that contain only neutral atoms. (e) Draw all reasonable resonance structures for the five-membered ring that contains three N atoms.



rizatriptan


17.61 Zolpidem (trade name Ambien) promotes the rapid onset of sleep, making it a widely prescribed drug for treating insomnia.



zolpidem

a. In what type of orbital does the lone pair on each N atom in the heterocycle reside?
b. Explain why the bicyclic ring system that contains both N atoms is aromatic.
c. Draw all reasonable resonance structures for the bicyclic ring system.


17.62 Answer the following questions about curcumin, a yellow pigment isolated from turmeric, a tropical perennial in the ginger family and a principal ingredient in curry powder.



curcumin

a. In Chapter 11 we learned that most enols, compounds that contain a hydroxy group bonded to a C = C, are unstable and tautomerize to carbonyl groups. Draw the keto form of the enol of curcumin, and explain why the enol is more stable than many other enols.
b. Explain why the enol O – H proton is more acidic than an alcohol O – H proton.
c. Why is curcumin colored?
d. Explain why curcumin is an antioxidant.

17.63 Stanozolol is an anabolic steroid that promotes muscle growth. Although stanozolol has been used by athletes and body builders, many physical and psychological problems result from prolonged use and it is banned in competitive sports.




stanozolol


a. Explain why the nitrogen heterocycle—a pyrazole ring—is aromatic.
b. In what type of orbital is the lone pair on each N atom contained?
c. Draw all reasonable resonance structures for stanozolol.
d. Explain why the pK_a of the N – H bond in the pyrazole ring is comparable to the pK_a of the O – H bond, making it considerably more acidic than amines such as CH_3NH_2 (pK_a = 40).

Challenge Problems

17.64 Explain why **A** is aromatic but **B** is not aromatic.




A




B

17.65 Use the observed ^{1}H NMR data to decide whether **C** and its dianion are aromatic, antiaromatic, or not aromatic. **C** shows NMR signals at –4.25 (6 H) and 8.14–8.67 (10 H) ppm. The dianion of **C** shows NMR signals at –3 (10 H) and 21 (6 H) ppm. Why are the signals shifted upfield (or downfield) to such a large extent?




C

17.66 Explain why compound **A** is much more stable than compound **B.**




A




B

17.67 (*R*)-Carvone, the major component of the oil of spearmint, undergoes acid-catalyzed isomerization to carvacrol, a major component of the oil of thyme. Draw a stepwise mechanism and explain why this isomerization occurs.



(*R*)-carvone carvacrol

17.68 Explain why triphenylene resembles benzene in that it does not undergo addition reactions with Br_2, but phenanthrene reacts with Br_2 to yield the addition product drawn. (Hint: Draw resonance structures for both triphenylene and phenanthrene, and use them to determine how delocalized each π bond is.)



triphenylene phenanthrene


17.69 Although benzene itself absorbs at 128 ppm in its ^{13}C NMR spectrum, the carbons of substituted benzenes absorb either upfield or downfield from this value depending on the substituent. Explain the observed values for the carbon ortho to the given substituent in the monosubstituted benzene derivatives **X** and **Y.**



X **Y**

Reactions of Aromatic Compounds

18



Vitamin K_1, phylloquinone, is a fat-soluble vitamin that regulates the synthesis of proteins needed for blood to clot. Dietary sources of vitamin K_1 include cauliflower, broccoli, soy beans, leafy greens, and green tea. A severe deficiency of vitamin K_1 leads to excessive and sometimes fatal bleeding because of inadequate blood clotting. Vitamin K_1 is synthesized by a biological Friedel–Crafts reaction, one of the many examples of electrophilic aromatic substitution, a key reaction of aromatic compounds presented in Chapter 18.

18.1 Electrophilic aromatic substitution
18.2 The general mechanism
18.3 Halogenation
18.4 Nitration and sulfonation
18.5 Friedel–Crafts alkylation and Friedel–Crafts acylation
18.6 Substituted benzenes
18.7 Electrophilic aromatic substitution of substituted benzenes
18.8 Why substituents activate or deactivate a benzene ring
18.9 Orientation effects in substituted benzenes
18.10 Limitations on electrophilic substitution reactions with substituted benzenes
18.11 Disubstituted benzenes
18.12 Synthesis of benzene derivatives
18.13 Nucleophilic aromatic substitution
18.14 Halogenation of alkyl benzenes
18.15 Oxidation and reduction of substituted benzenes
18.16 Multistep synthesis

Chapter 18 discusses the chemical reactions of benzene and other aromatic compounds. Although aromatic rings are unusually stable, making benzene unreactive in most of the reactions discussed so far, benzene acts as a nucleophile with certain electrophiles, yielding substitution products with an intact aromatic ring.

We begin with the basic features and mechanism of electrophilic aromatic substitution (Sections 18.1–18.5), the most prevalent reaction of benzene. Next, we discuss the electrophilic aromatic substitution of substituted benzenes (Sections 18.6–18.12), and conclude with nucleophilic aromatic substitution and other useful reactions of benzene derivatives (Sections 18.13–18.15). The ability to interconvert resonance structures and evaluate their relative energies is crucial to understanding this material.

18.1 Electrophilic Aromatic Substitution

Based on its structure and properties, what kinds of reactions should benzene undergo? Are any of its bonds particularly weak? Does it have electron-rich or electron-deficient atoms?

- **Benzene has six π electrons delocalized in six *p* orbitals that overlap above and below the plane of the ring. These loosely held π electrons make the benzene ring electron rich, so it reacts with electrophiles.**
- **Because benzene's six π electrons satisfy Hückel's rule, benzene is especially stable. Reactions that keep the aromatic ring intact are therefore favored.**

As a result, **the characteristic reaction of benzene is *electrophilic aromatic substitution*—a hydrogen atom is replaced by an electrophile.**

Electrophilic aromatic substitution




aromatic product

As we learned in Section 17.6, benzene does *not* undergo addition reactions like other unsaturated hydrocarbons, because addition would yield a product that is not aromatic. Substitution of a hydrogen, on the other hand, keeps the aromatic ring intact.

Five specific examples of electrophilic aromatic substitution are shown in Figure 18.1. The basic mechanism, discussed in Section 18.2, is the same in all five cases. The reactions differ only in the identity of the electrophile, E^+.

Problem 18.1 Why is benzene less reactive toward electrophiles than an alkene, even though it has more π electrons than an alkene (six versus two)?

Figure 18.1
Five examples of electrophilic aromatic substitution




Friedel–Crafts alkylation and acylation, named for Charles Friedel and James Crafts who discovered the reactions in the nineteenth century, form new carbon–carbon bonds.

18.2 The General Mechanism

No matter what electrophile is used, all electrophilic aromatic substitution reactions occur via a **two-step mechanism:** addition of the electrophile E^+ to form a resonance-stabilized carbocation, followed by deprotonation with base, as shown in Mechanism 18.1.

Mechanism 18.1 General Mechanism—Electrophilic Aromatic Substitution




1. Addition of the electrophile E^+ forms a new C–E bond and a resonance-stabilized carbocation. This step is rate-determining because the aromaticity of the benzene ring is lost.
2. A base removes the proton *on the carbon bonded to the electrophile,* re-forming the aromatic ring. Any resonance structure can be used to draw the product.

The first step in electrophilic aromatic substitution forms a carbocation, for which three resonance structures can be drawn. To help keep track of the location of the positive charge:

- **Always draw in the H atom on the carbon bonded to E. This serves as a reminder that it is the only sp^3 hybridized carbon in the carbocation intermediate.**

- **Notice that the positive charge in a given resonance structure is always located ortho or para to the new C – E bond. In the hybrid, therefore, the charge is delocalized over three atoms of the ring.**




(+) ortho to E **(+) para to E** **(+) ortho to E** **hybrid**

This two-step mechanism for electrophilic aromatic substitution applies to all of the electrophiles in Figure 18.1. **The net result of addition of an electrophile (E^+) followed by elimination of a proton (H^+) is substitution of E for H.**

The energy changes in electrophilic aromatic substitution are shown in Figure 18.2. The mechanism consists of two steps, so the energy diagram has two energy barriers. Because the transition state of the first step is higher in energy, it is rate-determining.

Problem 18.2 In Step [2] of Mechanism 18.1, loss of a proton to form the substitution product was drawn using one resonance structure only. Use curved arrows to show how the other two resonance structures can be converted to the substitution product (PhE) by removal of a proton with :B.

Figure 18.2
Energy diagram for electrophilic aromatic substitution: PhH + E^+ → PhE + H^+




- The mechanism has two steps, so there are two energy barriers.
- Step [1] is rate-determining; its transition state is at higher energy.


18.3 Halogenation

The general mechanism outlined in Mechanism 18.1 can now be applied to each of the five specific examples of electrophilic aromatic substitution shown in Figure 18.1. For each mechanism we must learn how to generate a specific electrophile. This step is *different* with each electrophile. Then, the electrophile reacts with benzene by the two-step process of Mechanism 18.1. These two steps are the *same* for all five reactions.

In **halogenation,** benzene reacts with Cl_2 or Br_2 in the presence of a Lewis acid catalyst, such as $FeCl_3$ or $FeBr_3$, to give the **aryl halides** chlorobenzene or bromobenzene, respectively. Analogous reactions with I_2 and F_2 are not synthetically useful because I_2 is too unreactive and F_2 reacts too violently.




chlorobenzene



bromobenzene


In bromination (Mechanism 18.2), the Lewis acid $FeBr_3$ reacts with Br_2 to form a Lewis acid–base complex that weakens and polarizes the Br−Br bond, making it more electrophilic. This reaction is Step [1] of the mechanism for the bromination of benzene. The remaining two steps follow directly from the general mechanism for electrophilic aromatic substitution: addition of the electrophile (Br^+ in this case) forms a resonance-stabilized carbocation, and loss of a proton regenerates the aromatic ring.

Mechanism 18.2 Bromination of Benzene




1. Lewis acid–base reaction of Br_2 with $FeBr_3$ forms a species with a weakened Br−Br bond that serves as source of Br^+.
2. Addition of the electrophile forms a new C−Br bond and a resonance-stabilized carbocation.
3. $FeBr_4^-$ removes the proton *on the carbon bonded to the electrophile,* re-forming the aromatic ring. The Lewis acid catalyst $FeBr_3$ is regenerated for another reaction cycle.

Figure 18.3
Examples of biologically active aryl chlorides





Generic name **bupropion**
Trade names **Wellbutrin, Zyban**
antidepressant,
also used to reduce nicotine cravings




chlorpheniramine
antihistamine

Herbicides were used extensively during the Vietnam War to defoliate dense jungle areas. The concentration of certain herbicide by-products in the soil remains high today.





2,4-D
2,4-dichlorophenoxy-acetic acid
herbicide



2,4,5-T
2,4,5-trichlorophenoxy-acetic acid
herbicide


the active components in **Agent Orange,**
a defoliant used in the Vietnam War


Chlorination proceeds by a similar mechanism. Reactions that introduce a halogen substituent on a benzene ring are widely used, and many halogenated aromatic compounds with a range of biological activity have been synthesized, as shown in Figure 18.3.

Problem 18.3 Draw a detailed mechanism for the chlorination of benzene using Cl_2 and $FeCl_3$.

18.4 Nitration and Sulfonation

Nitration and **sulfonation** of benzene introduce two different functional groups on an aromatic ring. Nitration is an especially useful reaction because a nitro group can then be reduced to an NH_2 group, a common benzene substituent, in a reaction discussed in Section 18.15.






benzenesulfonic acid


Generation of the electrophile in both nitration and sulfonation requires strong acid. In **nitration,** the electrophile is $^{+}NO_2$ (the **nitronium ion**), formed by protonation of HNO_3 followed by loss of water (Mechanism 18.3).

Mechanism 18.3 Formation of the Nitronium Ion ($^{+}NO_2$) for Nitration




In **sulfonation,** protonation of sulfur trioxide, SO_3, forms a positively charged sulfur species ($^+SO_3H$) that acts as an electrophile (Mechanism 18.4).

Mechanism 18.4 Formation of the Electrophile $^+SO_3H$ for Sulfonation



These steps illustrate how to generate the electrophile E^+ for nitration and sulfonation, the process that begins any mechanism for electrophilic aromatic substitution. To complete either of these mechanisms, you must replace the electrophile E^+ by either $^+NO_2$ or $^+SO_3H$ in the general mechanism (Mechanism 18.1). Thus, **the two-step sequence that replaces H by E is the same regardless of E^+.** This is shown in Sample Problem 18.1 using the reaction of benzene with the nitronium ion.

Sample Problem 18.1 Draw a stepwise mechanism for the nitration of a benzene ring.




nitrobenzene


Solution

We must first generate the electrophile and then write the two-step mechanism for electrophilic aromatic substitution using it.

Part [1] Generation of the electrophile $^+NO_2$




Part [2] Two-step mechanism for electrophilic aromatic substitution



Any species with a lone pair of electrons can be used to remove the proton in the last step. In this case, the mechanism is drawn with HSO_4^-, formed when $^+NO_2$ is generated as the electrophile.

Problem 18.4 Draw a stepwise mechanism for the sulfonation of an alkyl benzene such as **A** to form a substituted benzenesulfonic acid **B.** Treatment of **B** with base forms a sodium salt **C** that can be used as a synthetic detergent to clean away dirt (see Problem 3.22).



A **B** synthetic detergent **C**


18.5 Friedel–Crafts Alkylation and Friedel–Crafts Acylation

Friedel–Crafts alkylation and **Friedel–Crafts acylation** form new carbon–carbon bonds.

18.5A General Features


In **Friedel–Crafts alkylation,** treatment of benzene with an alkyl halide and a Lewis acid ($AlCl_3$) forms an alkyl benzene. This reaction is an **alkylation** because it results in transfer of an alkyl group from one atom to another (from Cl to benzene).

Friedel–Crafts alkylation




In **Friedel–Crafts acylation,** a benzene ring is treated with an **acid chloride** (RCOCl) and $AlCl_3$ to form a ketone. Because the new group bonded to the benzene ring is called an **acyl group,** the transfer of an acyl group from one atom to another is an **acylation.**

Friedel–Crafts acylation




Acid chlorides are also called **acyl chlorides.**

Problem 18.5 What product is formed when benzene is treated with each organic halide in the presence of $AlCl_3$?



Problem 18.6 What acid chloride would be needed to prepare each of the following ketones from benzene using a Friedel–Crafts acylation?




18.5B Mechanism

The mechanisms of alkylation and acylation proceed in a manner analogous to those for halogenation, nitration, and sulfonation. The unique feature in each reaction is how the electrophile is generated.


In **Friedel–Crafts alkylation,** the Lewis acid $AlCl_3$ reacts with the alkyl chloride to form a Lewis acid–base complex, illustrated with CH_3CH_2Cl and $(CH_3)_3CCl$ as alkyl chlorides. The identity of the alkyl chloride determines the exact course of the reaction as shown in Mechanism 18.5.

Mechanism 18.5 Formation of the Electrophile in Friedel–Crafts Alkylation—Two Possibilities

Possibility [1] For CH_3Cl and 1° RCl




Possibility [2] For 2° and 3° RCl



- **For CH_3Cl and 1° RCl, the Lewis acid–base complex itself serves as the electrophile for electrophilic aromatic substitution.**
- **With 2° and 3° RCl, the Lewis acid–base complex reacts further to give a 2° or 3° carbocation, which serves as the electrophile. Carbocation formation occurs only with 2° and 3° alkyl chlorides, because they afford more stable carbocations.**

In either case, the electrophile goes on to react with benzene in the two-step mechanism characteristic of electrophilic aromatic substitution, illustrated in Mechanism 18.6 using the 3° carbocation, $(CH_3)_3C^+$.

Mechanism 18.6 Friedel–Crafts Alkylation Using a 3° Carbocation




1 Addition of the carbocation electrophile forms a new carbon−carbon bond.

2 $AlCl_4^-$ removes a proton on the carbon bearing the new substituent to re-form the aromatic ring.

In **Friedel–Crafts acylation,** the Lewis acid $AlCl_3$ ionizes the carbon–halogen bond of the acid chloride, thus forming a positively charged carbon electrophile called an **acylium ion,** which is resonance stabilized (Mechanism 18.7). The positively charged carbon atom of the acylium ion then goes on to react with benzene in the two-step mechanism of electrophilic aromatic substitution.


Mechanism 18.7 Formation of the Electrophile in Friedel–Crafts Acylation



To complete the mechanism for acylation, insert the electrophile into the general mechanism and draw the last two steps, as illustrated in Sample Problem 18.2.

Sample Problem 18.2


Draw a stepwise mechanism for the following Friedel–Crafts acylation.




Solution

First generate the acylium ion, and then write the two-step mechanism for electrophilic aromatic substitution using it for the electrophile.

Part [1] Generation of the electrophile $(CH_3CO)^+$



Part [2] Two-step mechanism for electrophilic aromatic substitution




18.5C Other Facts About Friedel–Crafts Alkylation


Three additional facts about Friedel–Crafts alkylations must be kept in mind.


[1] Vinyl halides and aryl halides do *not* react in Friedel–Crafts alkylation.


Most Friedel–Crafts reactions involve carbocation electrophiles. Because the carbocations derived from vinyl halides and aryl halides are highly unstable and do not readily form, these organic halides do *not* undergo Friedel–Crafts alkylation.




Problem 18.7 Which halides are unreactive in a Friedel–Crafts alkylation reaction?

a. 

b. 


c. 

d. 


[2] Rearrangements can occur.

The Friedel–Crafts reaction can yield products having rearranged carbon skeletons when 1° and 2° alkyl halides are used as starting materials, as shown in Equations [1] and [2]. In both reactions, the carbon atom bonded to the halogen in the starting material (labeled in blue) is not bonded to the benzene ring in the product, thus indicating that a rearrangement has occurred.

Recall from Section 9.9 that a 1,2-shift converts a less stable carbocation to a more stable carbocation by shift of a hydrogen atom or an alkyl group.

[1] 

2° halide


[2] 

1° halide

The result in Equation [1] is explained by a carbocation rearrangement involving a 1,2-hydride shift: **the less stable 2° carbocation (formed from the 2° halide) rearranges to a more stable 3° carbocation,** as illustrated in Mechanism 18.8.

Mechanism 18.8 Friedel–Crafts Alkylation Involving Carbocation Rearrangement

Part [1] Formation of a 2° carbocation and rearrangement




2° carbocation + $AlCl_4^-$

3° carbocation

1 – 2 Lewis acid–base reaction of the alkyl chloride with $AlCl_3$ and cleavage of the C–Cl bond form a 2° carbocation.

3 1,2-Hydride shift converts a 2° carbocation to a more stable 3° carbocation.

Part [2] Two-step mechanism for electrophilic aromatic substitution



3° carbocation

[+ two more resonance structures]


+ HCl + $AlCl_3$

4 Addition of the 3° carbocation forms a new carbon–carbon bond and a resonance-stabilized carbocation.


5 $AlCl_4^-$ removes a proton on the carbon bearing the new substituent to re-form the aromatic ring.

Rearrangements can occur even when no free carbocation is formed initially. For example, the 1° alkyl chloride in Equation [2] forms a complex with $AlCl_3$, which does *not* decompose to an unstable 1° carbocation, as shown in Mechanism 18.9. Instead, a **1,2-hydride shift** forms a 2° carbocation, which then serves as the electrophile in the two-step mechanism for electrophilic aromatic substitution.

Mechanism 18.9 A Rearrangement Reaction Beginning with a 1° Alkyl Chloride



Problem 18.8 Draw a stepwise mechanism for the following reaction.





[3] Other functional groups that form carbocations can also be used as starting materials.

Although Friedel–Crafts alkylation works well with alkyl halides, any compound that readily forms a carbocation can be used instead. The two most common alternatives are alkenes and alcohols, both of which afford carbocations in the presence of strong acid.


- **Protonation of an alkene forms a carbocation, which can then serve as an electrophile in a Friedel–Crafts alkylation.**

- **Protonation of an alcohol, followed by loss of water, likewise forms a carbocation.**






Each carbocation can then go on to react with benzene to form a product of electrophilic aromatic substitution. For example:




Problem 18.9 Draw the product of each reaction.



18.5D Intramolecular Friedel–Crafts Reactions

All of the Friedel–Crafts reactions discussed thus far have resulted from intermolecular reaction of a benzene ring with an electrophile. Starting materials that contain both units are capable of **intramolecular reaction,** and this forms a new ring. Treatment of compound **A,** which contains both a benzene ring and an acid chloride, with $AlCl_3$, forms α-tetralone by an intramolecular Friedel–Crafts acylation reaction.




A

α-tetralone

new C–C bond in red

Such an intramolecular Friedel–Crafts acylation was a key step in the synthesis of the hallucinogen LSD, as shown in Figure 18.4.

Figure 18.4
Intramolecular Friedel–Crafts acylation in the synthesis of LSD




LSD
lysergic acid diethyl amide



Ergot-infected grain, the source of lysergic acid

- Intramolecular Friedel–Crafts acylation at the labeled carbons formed a product containing a new six-membered ring, which was converted to LSD in several steps.
- LSD was first prepared by Swiss chemist Albert Hoffman in 1938 from a related organic compound isolated from the ergot fungus that attacks rye and other grains. Ergot has a long history as a dreaded poison, affecting individuals who become ill from eating ergot-contaminated bread. The hallucinogenic effects of LSD were first discovered when Hoffman accidentally absorbed a small amount of the drug through his fingertips.

Problem 18.10 Draw a stepwise mechanism for the intramolecular Friedel–Crafts acylation of compound **A** to form **B.** **B** can be converted in one step to the antidepressant sertraline.




A **B** sertraline (Zoloft)



Sertraline (trade name Zoloft) is an effective antidepressant because it increases the concentration of the neurotransmitter serotonin in the brain.


Problem 18.11 Intramolecular reactions are also observed in Friedel–Crafts alkylation. Draw the intramolecular alkylation product formed from each of the following reactants. (Watch out for rearrangements!)



18.5E Biological Friedel–Crafts Reactions

Biological Friedel–Crafts reactions occur as well. As we learned in Section 16.2, allylic diphosphates contain a good leaving group, so they can serve as a source of allylic carbocations. A key step in the biological synthesis of vitamin K_1, the chapter-opening molecule, involves Friedel–Crafts reaction of 1,4-dihydroxynaphthoic acid with phytyl diphosphate to form **X,** which is converted to vitamin K_1 in several steps, as shown in Figure 18.5.

Figure 18.5 Friedel–Crafts reaction in the synthesis of vitamin K_1



Problem 18.12 (a) Draw resonance structures for the carbocation formed after loss of a leaving group from phytyl diphosphate. (b) Draw the two-step mechanism for Friedel–Crafts alkylation of 1,2-dihydroxynaphthoic acid with this carbocation to form **X.**

18.6 Substituted Benzenes

Many substituted benzene rings undergo electrophilic aromatic substitution. Common substituents include halogens, OH, NH_2, alkyl, and many functional groups that contain a carbonyl. Each substituent either increases or decreases the electron density in the benzene ring, and this affects the course of electrophilic aromatic substitution, as we will learn in Section 18.7.

What makes a substituent on a benzene ring electron donating or electron withdrawing? The answer is **inductive effects** and **resonance effects,** both of which can add or remove electron density.



Inductive Effects

Inductive effects stem from the **electronegativity** of the atoms in the substituent and the **polarizability** of the substituent group.

Inductive and resonance effects were first discussed in Sections 2.5B and 2.5C, respectively.

- **Atoms more electronegative than carbon—including N, O, and X—pull electron density away from carbon and thus exhibit an electron-withdrawing inductive effect.**
- **Polarizable alkyl groups donate electron density, and thus exhibit an electron-donating inductive effect.**

Considering inductive effects *only,* an NH_2 group withdraws electron density and CH_3 donates electron density.

Electron-withdrawing inductive effect	Electron-donating inductive effect
	
• N is **more electronegative** than C. • N inductively withdraws electron density.	• Alkyl groups are **polarizable,** making them electron-donating groups.


Problem 18.13 Which substituents have an electron-withdrawing and which have an electron-donating inductive effect: (a) $CH_3CH_2CH_2CH_2-$; (b) $Br-$; (c) CH_3CH_2O-?

Resonance Effects

Resonance effects can either donate or withdraw electron density, depending on whether they place a positive or negative charge on the benzene ring.

- **A resonance effect is electron donating when resonance structures place a negative charge on carbons of the benzene ring.**
- **A resonance effect is electron withdrawing when resonance structures place a positive charge on carbons of the benzene ring.**

An electron-donating resonance effect is observed whenever an atom Z having a lone pair of electrons is directly bonded to a benzene ring (general structure—$\mathbf{C_6H_5-Z:}$). Common examples of Z include N, O, and halogen. For example, five resonance structures can be drawn for aniline ($C_6H_5NH_2$). Because three of them place a negative charge on a carbon atom of the benzene ring, an **NH_2 group donates electron density to a benzene ring by a resonance effect.**




aniline

Three resonance structures place a (–) charge on atoms in the ring.

In contrast, **an electron-withdrawing resonance effect is observed in substituted benzenes having the general structure $\mathbf{C_6H_5-Y{=}Z}$,** where Z is more electronegative than Y. For


example, seven resonance structures can be drawn for benzaldehyde (C_6H_5CHO). Because three of them place a positive charge on a carbon atom of the benzene ring, a CHO group withdraws electron density from a benzene ring by a resonance effect.



benzaldehyde

Three resonance structures place a (+) charge on atoms in the ring.

Problem 18.14 Draw all resonance structures for each compound and use the resonance structures to determine if the substituent has an electron-donating or electron-withdrawing resonance effect.



Considering Both Inductive and Resonance Effects

To predict whether a substituted benzene is more or less electron rich than benzene itself, we must consider the **net balance of *both* the inductive and the resonance effects.** Alkyl groups, for instance, donate electrons by an inductive effect, but they have no resonance effect because they lack nonbonded electron pairs or π bonds. As a result,

- **An alkyl group is an electron-donating group and an alkyl benzene is more electron rich than benzene.**

When electronegative atoms, such as N, O, or halogen, are bonded to the benzene ring, they inductively *withdraw* electron density from the ring. All of these groups also have a nonbonded pair of electrons, so they *donate* electron density to the ring by resonance. **The *identity of the element* determines the net balance of these opposing effects.**



Z = N, O, X

Induction and resonance have opposite effects.
- Z inductively *withdraws* electron density.
- Z *donates* electron density by resonance.


- **When a neutral O or N atom is bonded directly to a benzene ring, the resonance effect dominates and the net effect is electron donation.**
- **When a halogen X is bonded to a benzene ring, the inductive effect dominates and the net effect is electron withdrawal.**

Thus, **NH_2 and OH are electron-donating groups** because the resonance effect predominates, whereas **Cl and Br are electron-withdrawing groups** because the inductive effect predominates.

Finally, the inductive and resonance effects in compounds having the general structure **$C_6H_5-Y=Z$** (with Z more electronegative than Y) are **both electron withdrawing;** in other words, the two effects *reinforce* each other. This is true for benzaldehyde (C_6H_5CHO) and all other compounds that contain a carbonyl group bonded directly to the benzene ring.

Thus, on balance, an **NH_2 group is electron donating,** so the benzene ring of aniline ($C_6H_5NH_2$) has more electron density than benzene. An **aldehyde group (CHO), on the other hand, is**


Figure 18.6
The effect of substituents on the electron density in substituted benzenes



- The NH_2 group donates electron density, making the benzene ring more electron rich (redder), whereas the CHO group withdraws electron density, making the benzene ring less electron rich (greener).

electron withdrawing, so the benzene ring of benzaldehyde (C_6H_5CHO) has less electron density than benzene. These effects are illustrated in the electrostatic potential maps in Figure 18.6. These compounds represent examples of the general structural features in electron-donating and electron-withdrawing substituents:

Electron-donating groups **Electron-withdrawing groups**




R = alkyl Z = N or O X = halogen


- **Common electron-donating groups are alkyl groups or groups with an N or O atom (with a lone pair) bonded to the benzene ring.**

- **Common electron-withdrawing groups are halogens or groups with an atom Y bearing a full or partial positive charge (+ or δ+) bonded to the benzene ring.**

The net effect of electron donation and withdrawal on the reactions of substituted aromatics is discussed in Sections 18.7–18.9.


Sample Problem 18.3 Classify each substituent as electron donating or electron withdrawing.

a. 

b. 


Solution

If necessary, draw out the atoms and bonds of the substituent to clearly see lone pairs and multiple bonds. **Always look at the atom bonded directly to the benzene ring** to determine electron-donating or electron-withdrawing effects. An O or N atom with a lone pair of electrons makes a substituent electron donating. A halogen or an atom with a partial positive charge makes a substituent electron withdrawing.

a. 

- An O atom with a lone pair is bonded directly to the benzene ring.


an electron-donating group


b. 


- An atom with a partial (+) charge is bonded directly to the benzene ring.

an electron-withdrawing group

Problem 18.15 Classify each substituent as electron donating or electron withdrawing.

a. 

b. 

c. 

18.7 Electrophilic Aromatic Substitution of Substituted Benzenes


Electrophilic aromatic substitution is a general reaction of *all* aromatic compounds, including polycyclic aromatic hydrocarbons, heterocycles, and substituted benzene derivatives. A substituent affects two aspects of electrophilic aromatic substitution:

- **The rate of reaction:** A substituted benzene reacts faster or slower than benzene itself.
- **The orientation:** The new group is located either ortho, meta, or para to the existing substituent. The identity of the first substituent determines the position of the second substituent.

Toluene ($C_6H_5CH_3$) and nitrobenzene ($C_6H_5NO_2$) illustrate two possible outcomes.


[1] Toluene

Toluene reacts **faster** than benzene in all substitution reactions. Thus, its **electron-donating CH_3 group *activates* the benzene ring** to electrophilic attack. Although three products are possible, compounds with the new group ortho or para to the CH_3 group predominate. The CH_3 group is therefore called an **ortho, para director.**



[2] Nitrobenzene


Nitrobenzene reacts **more slowly** than benzene in all substitution reactions. Thus, its **electron-withdrawing NO_2 group *deactivates* the benzene ring** to electrophilic attack. Although three products are possible, the compound with the new group meta to the NO_2 group predominates. The NO_2 group is called a **meta director.**



Substituents either activate or deactivate a benzene ring towards electrophiles, and direct selective substitution at specific sites on the ring. **All substituents can be divided into three general types.**

[1] Ortho, para directors and activators

- **Substituents that *activate* a benzene ring and direct substitution ortho and para.**




[2] Ortho, para deactivators

- **Substituents that *deactivate* a benzene ring and direct substitution ortho and para.**




[3] Meta directors

- **Substituents that direct substitution meta.**
- **All meta directors *deactivate* the ring.**




To learn these lists: **Keep in mind that the halogens are in a class by themselves.** Then learn the general structures for each type of substituent.

- **All ortho, para directors are R groups or have a nonbonded electron pair on the atom bonded to the benzene ring.**




Z = N or O ⟶ The ring is **activated.**
Z = halogen → The ring is **deactivated.**

- **All meta directors have a full or partial positive charge on the atom bonded to the benzene ring.**



Sample Problem 18.4 shows how this information can be used to predict the products of electrophilic aromatic substitution reactions.


Sample Problem 18.4 Draw the products of each reaction and state whether the reaction is faster or slower than a similar reaction with benzene.




Solution

To draw the products:

- Draw the Lewis structure for the substituent to see if it has a lone pair or partial positive charge on the atom bonded to the benzene ring.
- Classify the substituent—ortho, para activating; ortho, para deactivating; or meta deactivating—and draw the products.




The lone pair on N makes this group an **ortho, para activator. This compound reacts faster than benzene.**



The δ+ on the C bonded to the benzene ring makes the group a **meta deactivator. This compound reacts more slowly than benzene.**

Problem 18.16 Draw the products formed when each compound is treated with HNO_3 and H_2SO_4. State whether the reaction occurs faster or slower than a similar reaction with benzene.




18.8 Why Substituents Activate or Deactivate a Benzene Ring

- **Why do substituents activate or deactivate a benzene ring?**
- **Why are particular orientation effects observed?** Why are some groups ortho, para directors and some groups meta directors?

To understand why some substituents make a benzene ring react *faster* than benzene itself (activators), whereas others make it react *slower* (deactivators), we must evaluate the rate-determining step (the first step) of the mechanism. Recall from Section 18.2 that the first step in electrophilic aromatic substitution is the addition of an electrophile (E^+) to form a resonance-stabilized carbocation. The Hammond postulate (Section 7.15) makes it possible to predict the relative rate of the reaction by looking at the stability of the carbocation intermediate.

- **The more stable the carbocation, the lower in energy the transition state that forms it, and the faster the reaction.**



Stabilizing the carbocation makes the reaction faster.


The principles of inductive effects and resonance effects, first introduced in Section 18.6, can now be used to predict carbocation stability.

- **Electron-donating groups stabilize the carbocation and *activate* a benzene ring toward electrophilic attack. All activators are R groups or they have an N or O atom with a lone pair bonded directly to the benzene ring.**

- **Electron-withdrawing groups destabilize the carbocation and *deactivate* a benzene ring toward electrophilic attack. All deactivators are either halogens or they have an atom with a full or partial positive charge bonded directly to the benzene ring.**


The energy diagrams in Figure 18.7 illustrate the effect of electron-donating and electron-withdrawing groups on the energy of the transition state of the rate-determining step in electrophilic aromatic substitution.

Figure 18.7 Energy diagrams comparing the rate of electrophilic aromatic substitution of substituted benzenes




- Electron-donor groups **D** stabilize the carbocation intermediate, lower the energy of the transition state, and increase the rate of reaction.
- Electron-withdrawing groups **W** destabilize the carbocation intermediate, raise the energy of the transition state, and decrease the rate of reaction.

Problem 18.17 Label each compound as more or less reactive than benzene in electrophilic aromatic substitution.



Problem 18.18 Rank the following compounds in order of increasing reactivity in electrophilic aromatic substitution.



18.9 Orientation Effects in Substituted Benzenes

To understand why particular orientation effects arise, you must keep in mind the general structures for ortho, para directors and for meta directors already given in Section 18.7. There are two general types of ortho, para directors and one general type of meta director:


- **All ortho, para directors are R groups or have a nonbonded electron pair on the atom bonded to the benzene ring.**

- **All meta directors have a full or partial positive charge on the atom bonded to the benzene ring.**

To evaluate the directing effects of a given substituent, we can follow a stepwise procedure.

How To Determine the Directing Effects of a Particular Substituent

Step [1] **Draw all resonance structures for the carbocation formed from attack of an electrophile E^+ at the ortho, meta, and para positions of a substituted benzene (C_6H_5-A).**



- There are at least three resonance structures for each site of reaction.
- Each resonance structure places a positive charge ortho or para to the new C – E bond.


Step [2] **Evaluate the stability of the intermediate resonance structures. The electrophile attacks at those positions that give the most stable carbocation.**

Sections 18.9A–C show how this two-step procedure can be used to determine the directing effects of the CH_3 group in toluene, the NH_2 group in aniline, and the NO_2 group in nitrobenzene, respectively.

18.9A The CH_3 Group—An ortho, para Director

To determine why a **CH_3 group directs electrophilic aromatic substitution to the ortho and para positions,** first draw all resonance structures that result from electrophilic attack at the ortho, meta, and para positions to the CH_3 group.

Always draw in the H atom at the site of electrophilic attack. This will help you keep track of where the charges go.




The positive charge in all resonance structures is always ortho or para to the new C−E bond. It is *not* necessarily ortho or para to the CH_3 group.

To evaluate the stability of the resonance structures, determine whether any are especially stable or unstable. In this example, **attack ortho or para to CH_3 generates a resonance structure that places a positive charge on a carbon atom with the CH_3 group.** The electron-donating CH_3 group *stabilizes* the adjacent positive charge. In contrast, attack meta to the CH_3 group does *not* generate any resonance structure stabilized by electron donation. Other alkyl groups are ortho, para directors for the same reason.

- The CH_3 group directs electrophilic attack ortho and para to itself because an electron-donating inductive effect stabilizes the carbocation intermediate.

18.9B The NH_2 Group—An ortho, para Director

To determine why an **amino group (NH_2) directs electrophilic aromatic substitution to the ortho and para positions,** follow the same procedure.




Attack at the meta position generates the usual three resonance structures. Because of the lone pair on the N atom, attack at the ortho and para positions generates a fourth resonance structure, which is stabilized because **every atom has an octet of electrons. This additional resonance structure can be drawn for all substituents that have an N, O, or halogen atom bonded directly to the benzene ring.**

- The NH_2 group directs electrophilic attack ortho and para to itself because the carbocation intermediate has additional resonance stabilization.

18.9C The NO_2 Group—A meta Director

To determine why a **nitro group (NO_2) directs electrophilic aromatic substitution to the meta position,** follow the same procedure.



Attack at each position generates three resonance structures. One resonance structure resulting from attack at the ortho and para positions is especially *destabilized,* because it contains a positive charge on two adjacent atoms. Attack at the meta position does not generate any particularly unstable resonance structures.

- **With the NO_2 group (and all meta directors), meta attack occurs because attack at the ortho or para position gives a destabilized carbocation intermediate.**

Problem 18.19 Draw all resonance structures for the carbocation formed by ortho attack of the electrophile $^+NO_2$ on each starting material. Label any resonance structures that are especially stable or unstable.





Figure 18.8 summarizes the reactivity and directing effects of the common substituents on benzene rings.

Figure 18.8
The reactivity and directing effects of common substituted benzenes




In summary:
[1] All ortho, para directors except the halogens activate the benzene ring.
[2] All meta directors deactivate the benzene ring.
[3] The halogens deactivate the benzene ring.

18.10 Limitations on Electrophilic Substitution Reactions with Substituted Benzenes


Although electrophilic aromatic substitution works well with most substituted benzenes, halogenation and the Friedel–Crafts reactions have some additional limitations that must be kept in mind.

18.10A Halogenation of Activated Benzenes


Considering all electrophilic aromatic substitution reactions, halogenation occurs the most readily. As a result, benzene rings activated by strong electron-donating groups—OH, NH_2, and their alkyl derivatives (OR, NHR, and NR_2)—undergo **polyhalogenation** when treated with X_2 and FeX_3. Aniline ($C_6H_5NH_2$) and phenol (C_6H_5OH) both give a tribromo derivative when treated with Br_2 and $FeBr_3$. **Substitution occurs at all hydrogen atoms ortho and para to the NH_2 and OH groups.**



Monosubstitution of H by Br occurs with Br_2 *alone* without added catalyst to form a mixture of ortho and para products.




Problem 18.20 Draw the products of each reaction.




18.10B Limitations in Friedel–Crafts Reactions

Friedel–Crafts reactions are the most difficult electrophilic aromatic substitution reactions to carry out in the laboratory. They do not occur when the benzene ring is substituted with NO_2 (a strong deactivator) or with NH_2, NHR, or NR_2 (strong activators).


A benzene ring deactivated by a strong electron-withdrawing group—that is, any of the meta directors—is not electron rich enough to undergo Friedel–Crafts reactions.



Friedel–Crafts reactions also do not occur with NH_2 groups, which are strong activating groups. NH_2 groups are strong Lewis bases (due to the nonbonded electron pair on N), so they react with $AlCl_3$, the Lewis acid needed for alkylation or acylation. The resulting product contains a positive charge adjacent to the benzene ring, so the ring is now strongly deactivated and therefore unreactive in Friedel–Crafts reactions.




Problem 18.21 Which of the following compounds undergo Friedel–Crafts alkylation with CH_3Cl and $AlCl_3$? Draw the products formed when a reaction occurs.




Another limitation of the Friedel–Crafts alkylation arises because of **polyalkylation.** Treatment of benzene with an alkyl halide and $AlCl_3$ places an electron-donor R group on the ring. Because R groups activate a ring, the alkylated product (C_6H_5R) is now *more reactive* than benzene itself toward further substitution, and it reacts again with RCl to give products of polyalkylation.

To minimize polyalkylation a large excess of benzene is used relative to the amount of alkyl halide.



Polysubstitution does not occur with Friedel–Crafts acylation, because the product now has an electron-withdrawing group that deactivates the ring toward another electrophilic substitution.




18.11 Disubstituted Benzenes

What happens in electrophilic aromatic substitution when a disubstituted benzene ring is used as starting material? **To predict the products, look at the directing effects of both substituents and then determine the net result,** using the following three guidelines.


Rule [1] **When the directing effects of two groups reinforce, the new substituent is located on the position directed by both groups.**

The CH_3 group in *p*-nitrotoluene is an ortho, para director and the NO_2 group is a meta director. These two effects reinforce each other so that one product is formed on treatment with Br_2 and $FeBr_3$. The position para to the CH_3 group is "blocked" by a nitro group, so no substitution can occur on that carbon.



Rule [2] **If the directing effects of two groups oppose each other, the more powerful activator "wins out."**


In compound **A,** the $NHCOCH_3$ group activates its two ortho positions, and the CH_3 group activates its two ortho positions to reaction with electrophiles. Because the $NHCOCH_3$ is a stronger activator, substitution occurs ortho to it.



A


Rule [3] **No substitution occurs between two meta substituents because of crowding.**

For example, no substitution occurs at the carbon atom between the two CH_3 groups in *m*-xylene, even though two CH_3 groups activate that position.




m-xylene
(1,3-dimethylbenzene)

Sample Problem 18.5 Draw the products formed from nitration of each compound.




Solution


a. Both the OH and CH_3 groups are ortho, para directors. Because the OH group is a stronger activator, substitution occurs ortho to it.



b. Both the OH and CH_3 groups are ortho, para directors whose directing effects reinforce each other in this case. No substitution occurs between the two meta substituents, however, so two products result.




Problem 18.22 Draw the products formed when each compound is treated with HNO_3 and H_2SO_4.



18.12 Synthesis of Benzene Derivatives

To synthesize benzene derivatives with more than one substituent, we must always take into account the directing effects of each substituent. In a disubstituted benzene, for example, **the directing effects indicate which substituent must be added to the ring first.**

For example, the Br group in *p*-bromonitrobenzene is an ortho, para director and the NO_2 group is a meta director. Because the two substituents are para to each other, the ortho, para director must be introduced *first* when synthesizing this compound from benzene.




p-bromonitrobenzene

With two para substituents, add the ortho, para director first.


Thus, Pathway [1], in which bromination precedes nitration, yields the desired para product, whereas Pathway [2], in which nitration precedes bromination, yields the undesired meta isomer.

Pathway [1] Bromination before nitration: The desired para product is formed.



desired product


Pathway [2] Nitration before bromination: The undesired meta isomer is formed.



meta director meta isomer

Pathway [1] yields both the desired para product as well as the undesired ortho isomer. Because these compounds are constitutional isomers, they are separable. Obtaining such a mixture of ortho and para isomers is often unavoidable.


Sample Problem 18.6 Devise a synthesis of *o*-nitrotoluene from benzene.



o-nitrotoluene


Solution

The CH_3 group in *o*-nitrotoluene is an ortho, para director and the NO_2 group is a meta director. Because the two substituents are ortho to each other, the **ortho, para director must be introduced first.** The synthesis thus involves two steps: Friedel–Crafts alkylation followed by nitration.




o-nitrotoluene

Problem 18.23 Devise a synthesis of each compound from the indicated starting material.



18.13 Nucleophilic Aromatic Substitution

Although most reactions of aromatic compounds occur by way of electrophilic aromatic substitution, aryl halides undergo a limited number of substitution reactions with strong nucleophiles.




X = F, Cl, Br, I
A = H or electron-withdrawing group

- **Nucleophilic aromatic substitution results in the substitution of a halogen X on a benzene ring by a nucleophile (:Nu⁻).**

As we learned in Section 7.17, these reactions *cannot* occur by an S_N1 or S_N2 mechanism, which take place only at sp^3 hybridized carbons. Instead, two different mechanisms are proposed to explain the results: **addition–elimination** (Section 18.13A) and **elimination–addition** (Section 18.13B).

18.13A Nucleophilic Aromatic Substitution by Addition–Elimination


Aryl halides with strong electron-withdrawing groups (such as NO_2) on the ortho or para positions react with nucleophiles to afford substitution products. Treatment of *p*-chloronitrobenzene with hydroxide (^-OH) affords *p*-nitrophenol by replacement of Cl by OH.



p-chloronitrobenzene *p*-nitrophenol

Nucleophilic aromatic substitution occurs with a variety of strong nucleophiles, including ^-OH, ^-OR, $^-NH_2$, ^-SR, and in some cases, neutral nucleophiles such as NH_3 and RNH_2. The mechanism of these reactions has two steps: **addition of the nucleophile** to form a resonance-stabilized carbanion, followed by **elimination of the halogen leaving group.** Mechanism 18.10 is drawn with an aryl chloride containing a general electron-withdrawing group W.

Mechanism 18.10 Nucleophilic Aromatic Substitution by Addition–Elimination



1 Addition of the nucleophile forms a resonance-stabilized carbanion and a new C–Nu bond in the rate-determining step.


2 Loss of the leaving group re-forms the aromatic ring.

In nucleophilic aromatic substitution, the following trends in reactivity are observed.


- **Increasing the number of electron-withdrawing groups increases the reactivity of the aryl halide. Electron-withdrawing groups stabilize the intermediate carbanion, and by the Hammond postulate, lower the energy of the transition state that forms it.**


- **Increasing the electronegativity of the halogen increases the reactivity of the aryl halide. A more electronegative halogen stabilizes the intermediate carbanion by an inductive effect, making aryl fluorides (ArF) much *more* reactive than other aryl halides, which contain less electronegative halogens.**

Thus, aryl chloride **B** is more reactive than *o*-chloronitrobenzene (**A**) because it contains two electron-withdrawing NO_2 groups. Aryl fluoride **C** is more reactive than **B** because **C** contains the more electronegative halogen, fluorine.




The location of the electron-withdrawing group greatly affects the rate of nucleophilic aromatic substitution. When a nitro group is located ortho or para to the halogen, the negative charge of the intermediate carbanion can be delocalized onto the NO_2 group, thus stabilizing it. With a meta NO_2 group, no such additional delocalization onto the NO_2 group occurs.






Thus, nucleophilic aromatic substitution by an addition–elimination mechanism occurs only with aryl halides that contain electron-withdrawing substituents at the ortho or para position.

Problem 18.24 Draw the products of each reaction.




Problem 18.25 Draw a stepwise mechanism for the following reaction that forms ether **D. D** can be converted to the antidepressant fluoxetine (trade name Prozac) in a single step.




18.13B Nucleophilic Aromatic Substitution by Elimination–Addition: Benzyne

Aryl halides that do not contain an electron-withdrawing group generally do not react with nucleophiles. Under extreme reaction conditions, however, nucleophilic aromatic substitution can occur with aryl halides. For example, heating chlorobenzene with NaOH above 300 °C and 170 atmospheres of pressure affords phenol.



The mechanism proposed to explain this result involves formation of a **benzyne** intermediate (C_6H_4) by elimination–addition. As shown in Mechanism 18.11, benzyne is a highly reactive, unstable intermediate formed by elimination of HX from an aryl halide.


Mechanism 18.11 Nucleophilic Aromatic Substitution by Elimination–Addition: Benzyne




1 – 2 Elimination of H and X from two adjacent atoms forms a reactive benzyne intermediate.

3 – 4 Nucleophilic attack and protonation form the substitution product.

Formation of a benzyne intermediate explains why substituted aryl halides form mixtures of products. **Nucleophilic aromatic substitution by an elimination–addition mechanism affords substitution on the carbon directly bonded to the leaving group and the carbon adjacent to it.** As an example, treatment of *p*-chlorotoluene with $NaNH_2$ forms para- and meta-substitution products.




This result is explained by the fact that nucleophilic attack on the benzyne intermediate may occur at either C3 to form *m*-methylaniline, or C4 to form *p*-methylaniline.




As you might expect, the triple bond in benzyne is unusual. Each carbon of the six-membered ring is sp^2 hybridized, and as a result, the σ bond and two π bonds of the triple bond are formed with the following orbitals.

- **The σ bond is formed by overlap of two sp^2 hybrid orbitals.**
- **One π bond is formed by overlap of two *p* orbitals perpendicular to the plane of the molecule.**
- **The second π bond is formed by overlap of two sp^2 hybrid orbitals.**



Thus, the second π bond of benzyne differs from all other π bonds seen thus far, because **it is formed by the side-by-side overlap of *sp^2 hybrid orbitals,* not *p* orbitals.** This π bond, located in the plane of the molecule, is extremely weak.

Problem 18.26 Draw the products of each reaction.




Problem 18.27 Draw all products formed when *m*-chlorotoluene is treated with KNH_2 in NH_3.

18.14 Halogenation of Alkyl Benzenes

Radical halogenation of alkanes was discussed in Chapter 15. The mechanism of radical halogenation at an allylic carbon was given in Section 15.10.


We finish Chapter 18 by learning some additional reactions of substituted benzenes that greatly expand the ability to synthesize benzene derivatives. In Section 18.14 we return to radical halogenation, and in Section 18.15 we examine useful oxidation and reduction reactions.

Benzylic C−H bonds are weaker than most other sp^3 hybridized C−H bonds, because homolysis forms a resonance-stabilized benzylic radical.



The bond dissociation energy for a benzylic C – H bond (356 kJ/mol) is even less than the bond dissociation energy for a 3° C – H bond (381 kJ/mol).


As a result, an alkyl benzene undergoes selective bromination at the weak benzylic C – H bond under radical conditions to form a **benzylic halide.** For example, radical bromination of ethylbenzene using either Br_2 (in the presence of light or heat) or *N*-bromosuccinimide (NBS, in the presence of light or peroxides) forms a benzylic bromide as the sole product.



The mechanism for halogenation at the benzylic position resembles other radical halogenation reactions, and so it involves initiation, propagation, and termination. Mechanism 18.12 illustrates the radical bromination of ethylbenzene using Br_2 ($h\nu$ or Δ).


Mechanism 18.12 Benzylic Bromination

Part [1] Initiation



1 Homolysis of the Br–Br bond requires light or heat and forms two bromine radicals.

Part [2] Propagation



2 Abstraction of a benzylic hydrogen by Br· forms the resonance-stabilized benzylic radical.


3 The benzylic radical abstracts a bromine atom to form the benzylic bromide. Because Br· is also formed, Steps [2] and [3] can occur repeatedly without additional initiation.

Part [3] Termination



4 Termination of the chain occurs when any two radicals combine to form a bond.

Thus, an alkyl benzene undergoes two different reactions with Br_2, depending on the reaction conditions.




- With Br_2 and $FeBr_3$ (**ionic conditions**), electrophilic aromatic substitution occurs, resulting in replacement of H by Br on the aromatic ring to form ortho and para isomers.
- With Br_2 and light or heat (**radical conditions**), substitution of H by Br occurs at the *benzylic* carbon of the alkyl group.

Problem 18.28 Explain why $C_6H_5CH_2CH_2Br$ is not formed during the radical bromination of $C_6H_5CH_2CH_3$.


The radical bromination of alkyl benzenes is a useful reaction because the resulting benzylic halide can serve as starting material for a variety of substitution and elimination reactions, thus making it possible to form many new substituted benzenes. Sample Problem 18.7 illustrates one possibility.

Sample Problem 18.7 Design a synthesis of styrene from ethylbenzene.




Solution

The double bond can be introduced by a two-step reaction sequence: bromination at the benzylic position under radical conditions, followed by elimination of HBr with strong base to form the π bond.



Problem 18.29 How could you use ethylbenzene to prepare each compound? More than one step is required.




18.15 Oxidation and Reduction of Substituted Benzenes


Oxidation and reduction reactions are valuable tools for preparing many other benzene derivatives. Because the mechanisms are complex and do not have general applicability, reagents and reactions are presented only, without reference to the detailed mechanism.

18.15A Oxidation of Alkyl Benzenes

Arenes containing at least one benzylic C−H bond are oxidized with $KMnO_4$ to benzoic acid, a carboxylic acid with the carboxy group (COOH) bonded directly to the benzene ring. With some alkyl benzenes, this also results in the cleavage of carbon–carbon bonds, so the product has fewer carbon atoms than the starting material.




Substrates with more than one alkyl group are oxidized to dicarboxylic acids. Compounds without a benzylic C−H bond are inert to oxidation.



18.15B Reduction of Aryl Ketones to Alkyl Benzenes


Ketones formed as products in Friedel–Crafts acylation can be reduced to alkyl benzenes by two different methods.



- The **Clemmensen reduction** uses zinc and mercury in the presence of strong acid.
- The **Wolff–Kishner reduction** uses hydrazine (NH_2NH_2) and strong base (KOH).

Because both C−O bonds in the starting material are converted to C−H bonds in the product, the reduction is difficult and the reaction conditions must be harsh.

Clemmensen reduction



Wolff–Kishner reduction





Figure 18.9
Two methods to prepare an alkyl benzene




We now know two different ways to introduce an alkyl group on a benzene ring (Figure 18.9):


- **A one-step method using Friedel–Crafts alkylation**
- **A two-step method using Friedel–Crafts acylation to form a ketone, followed by reduction**


Although the two-step method seems more roundabout, it must be used to synthesize certain alkyl benzenes that cannot be prepared by the one-step Friedel–Crafts alkylation because of rearrangements.

Recall from Section 18.5C that propylbenzene cannot be prepared by a Friedel–Crafts alkylation. Instead, when benzene is treated with 1-chloropropane and $AlCl_3$, isopropylbenzene is formed by a rearrangement reaction. Propylbenzene can be made, however, by a two-step procedure using Friedel–Crafts acylation followed by reduction.



Problem 18.30 Write out the two-step sequence that converts benzene to each compound.

a. 

b. 


Problem 18.31 What steps are needed to convert benzene into *p*-isobutylacetophenone, a synthetic intermediate used in the synthesis of the anti-inflammatory agent ibuprofen.



p-isobutylacetophenone ibuprofen

18.15C Reduction of Nitro Groups


A nitro group (NO_2) is easily introduced on a benzene ring by nitration with strong acid (Section 18.4). This process is useful because the nitro group is readily reduced to an amino group (NH_2) under a variety of conditions. The most common methods use H_2 and a catalyst, or a metal (such as Fe or Sn) and a strong acid like HCl.





Benzocaine is the active ingredient in the over-the-counter topical anesthetic Orajel.


For example, reduction of ethyl *p*-nitrobenzoate with H_2 and a palladium catalyst forms ethyl *p*-aminobenzoate, a local anesthetic commonly called benzocaine.



ethyl *p*-nitrobenzoate — ethyl *p*-aminobenzoate (benzocaine)

Sample Problem 18.8 illustrates the utility of this process in a short synthesis.

Sample Problem 18.8 Design a synthesis of *m*-bromoaniline from benzene.



m-bromoaniline


Solution

To devise a retrosynthetic plan, keep in mind:

- The NH_2 group cannot be introduced directly on the ring by electrophilic aromatic substitution. It must be added by a two-step process: nitration followed by reduction.
- Both the Br and NH_2 groups are ortho, para directors, but they are located meta to each other on the ring. However, an NO_2 group (from which an NH_2 group is made) *is* a meta director, and we can use this fact to our advantage.

Retrosynthetic Analysis

Working backward gives the following **three-step retrosynthetic analysis:**




m-bromoaniline


- [1] Form the NH_2 group by reduction of NO_2.
- [2] Introduce the Br group meta to the NO_2 group by halogenation.
- [3] Add the NO_2 group by nitration.


Synthesis


The synthesis involves three steps, and the order is crucial for success. Halogenation (Step [2] of the synthesis) must occur *before* reduction (Step [3]) in order to form the meta substitution product.



Problem 18.32 Synthesize each compound from benzene.

a. 


b. 

c. 

18.16 Multistep Synthesis

The reactions learned in Chapter 18 make it possible to synthesize a wide variety of substituted benzenes, as shown in Sample Problems 18.9–18.11.

Sample Problem 18.9 Synthesize *p*-nitrobenzoic acid from benzene.




p-nitrobenzoic acid

Solution

Both groups on the ring (NO_2 and COOH) are meta directors. To place these two groups para to each other, remember that the COOH group is prepared by oxidizing an alkyl group, which is an ortho, para director.

Retrosynthetic Analysis




p-nitrobenzoic acid

Working backwards:

- [1] Form the COOH group by oxidation of an alkyl group.
- [2] Introduce the NO_2 group para to the CH_3 group (an ortho, para director) by nitration.
- [3] Add the CH_3 group by Friedel–Crafts alkylation.


Synthesis



- Friedel–Crafts alkylation with CH_3Cl and $AlCl_3$ forms toluene in Step [1]. Because CH_3 is an ortho, para director, nitration yields the desired para product, which can be separated from its ortho isomer (Step [2]).
- Oxidation with $KMnO_4$ converts the CH_3 group into a COOH group, giving the desired product in Step [3].

Sample Problem 18.10

Synthesize *p*-chlorostyrene from benzene.




p-chlorostyrene

Solution

Both groups on the ring are ortho, para directors located para to each other. To introduce the double bond in the side chain, we must follow the two-step sequence in Sample Problem 18.7.

Retrosynthetic Analysis




p-chlorostyrene

Working backwards:


- [1] Form the double bond by two steps: benzylic halogenation followed by elimination.
- [2] Introduce the CH_3CH_2 group by Friedel–Crafts alkylation.
- [3] Add the Cl atom by chlorination.

Synthesis



- Chlorination in Step [1] followed by Friedel–Crafts alkylation in Step [2] forms the desired para product, which can be separated from its ortho isomer.
- Benzylic bromination followed by elimination with strong base [$KOC(CH_3)_3$] (Steps [3] and [4]) forms the double bond of the target compound, *p*-chlorostyrene.

Sample Problem 18.11 Synthesize the trisubstituted benzene **A** from benzene.




Solution


Two groups (CH_3CO and NO_2) in **A** are meta directors located meta to each other, and the third substituent, an alkyl group, is an ortho, para director.

Retrosynthetic Analysis

With three groups on the benzene ring, begin by determining the possible disubstituted benzenes that are immediate precursors of the target compound, and then eliminate any that cannot be converted to the desired product. For example, three different disubstituted benzenes (**B**–**D**) can theoretically be precursors to **A.** However, conversion of compounds **B** or **D** to **A** would require a Friedel–Crafts reaction on a deactivated benzene ring, a reaction that does not occur. Thus, only **C** is a feasible precursor of **A.**




To complete the retrosynthetic analysis, prepare **C** from benzene:




- [1] Add the ketone by Friedel–Crafts acylation.
- [2] Add the alkyl group by the two-step process—Friedel–Crafts acylation followed by reduction. It is not possible to prepare butylbenzene by a one-step Friedel–Crafts alkylation because of a rearrangement reaction (Section 18.15B).

Synthesis



- Friedel–Crafts acylation followed by reduction with Zn(Hg), HCl yields butylbenzene (Steps [1]–[2]).
- Friedel–Crafts acylation gives the para product **C,** which can be separated from its ortho isomer (Step [3]).
- Nitration in Step [4] introduces the NO_2 group ortho to the alkyl group (an ortho, para director) and meta to the CH_3CO group (a meta director).

Problem 18.33 Synthesize each compound from benzene.




KEY CONCEPTS

Reactions of Aromatic Compounds

Mechanism of Electrophilic Aromatic Substitution (18.2)

- Electrophilic aromatic substitution follows a two-step mechanism. Reaction of the aromatic ring with an electrophile forms a carbocation, and loss of a proton regenerates the aromatic ring.
- The first step is rate-determining.
- The intermediate carbocation is stabilized by resonance; a minimum of three resonance structures can be drawn. The positive charge is always located ortho or para to the new C – E bond.







Three Rules Describing the Reactivity and Directing Effects of Common Substituents (18.7–18.9)

[1] All ortho, para directors except the halogens activate the benzene ring.

[2] All meta directors deactivate the benzene ring.


[3] The halogens deactivate the benzene ring and direct ortho, para.

Summary of Substituent Effects in Electrophilic Aromatic Substitution (18.6–18.9)

	Substituent	Inductive effect	Resonance effect	Reactivity	Directing effect
[1]	 R = alkyl	donating	none	activating	ortho, para
[2]	 Z = N or O	withdrawing	donating	activating	ortho, para
[3]	 X = halogen	withdrawing	donating	deactivating	ortho, para
[4]		withdrawing	withdrawing	deactivating	meta


Five Examples of Electrophilic Aromatic Substitution

[1] Halogenation—Replacement of H by Cl or Br (18.3)




- Polyhalogenation occurs on benzene rings substituted by OH and NH_2 (and related substituents) (18.10A).


[2] Nitration—Replacement of H by NO_2 (18.4)



[3] Sulfonation—Replacement of H by SO_3H (18.4)




[4] Friedel–Crafts alkylation—Replacement of H by R (18.5)




Variations:

[1] with alcohols




[2] with alkenes



- Rearrangements can occur.
- Vinyl halides and aryl halides are unreactive.
- The reaction does not occur on benzene rings substituted by meta deactivating groups or NH_2 groups (18.10B).
- Polyalkylation can occur.


[5] Friedel–Crafts acylation—Replacement of H by RCO (18.5)



- The reaction does not occur on benzene rings substituted by meta deactivating groups or NH_2 groups (18.10B).

Nucleophilic Aromatic Substitution (18.13)


[1] Nucleophilic substitution by an addition–elimination mechanism



X = F, Cl, Br, I
A = electron-withdrawing group

- The mechanism has two steps.
- Strong electron-withdrawing groups at the ortho or para positions are required.
- Increasing the number of electron-withdrawing groups increases the rate.
- Increasing the electronegativity of the halogen increases the rate.

[2] Nucleophilic substitution by an elimination–addition mechanism




X = halogen


- Reaction conditions are harsh.
- Benzyne is formed as an intermediate.
- Product mixtures may result.

Other Reactions of Benzene Derivatives

[1] Benzylic halogenation (18.14)




[2] Oxidation of alkyl benzenes (18.15A)




- A benzylic C – H bond is needed for reaction.

[3] Reduction of ketones to alkyl benzenes (18.15B)




[4] Reduction of nitro groups to amino groups (18.15C)




PROBLEMS

Problems Using Three-Dimensional Models

18.34 Draw the products formed when **A** and **B** are treated with each of the following reagents: (a) Br_2, $FeBr_3$; (b) HNO_3, H_2SO_4; (c) CH_3CH_2COCl, $AlCl_3$.




A



B

18.35 What is the major product formed by an intramolecular Friedel–Crafts acylation of the following compound?





Reactions


18.36 Draw the products formed when phenol (C_6H_5OH) is treated with each set of reagents.

a. [1] HNO_3, H_2SO_4; [2] Sn, HCl
b. [1] $(CH_3CH_2)_2CHCOCl$, $AlCl_3$; [2] Zn(Hg), HCl
c. [1] CH_3CH_2Cl, $AlCl_3$; [2] Br_2, $h\nu$
d. [1] $(CH_3)_2CHCl$, $AlCl_3$; [2] $KMnO_4$


18.37 Draw the products formed when each compound is treated with CH_3CH_2COCl, $AlCl_3$.


a. 


b. 


c. 


18.38 Draw the products of each reaction.


a. 

b. 


c. 


d. 


e. 

f. 


18.39 What products are formed when benzene is treated with each alkyl chloride and $AlCl_3$?


a. 


b. 


c. 


18.40 Draw the products of each reaction.


a. 


c. 




b. 





d. 




18.41 You have learned two ways to make an alkyl benzene: Friedel–Crafts alkylation, and Friedel–Crafts acylation followed by reduction. Although some alkyl benzenes can be prepared by both methods, it is often true that only one method can be used to prepare a given alkyl benzene. Which method(s) can be used to prepare each of the following compounds from benzene? Show the steps that would be used.


a. 

b. 

c. 

18.42 Draw the structure of **A,** an intermediate in the synthesis of the antipsychotic drug risperidone. Explain why three rings in risperidone are considered aromatic.






A


$C_{12}H_{13}FN_2O$

several steps




risperidone

18.43 One step in the synthesis of pioglitazone (trade name Actos), a drug used to treat type 2 diabetes, involves the reaction of **A** with **B** in the presence of NaH to afford **C.** What is the structure of **C?**




A **B**

18.44 **D** is an intermediate in the synthesis of rosiglitazone (trade name Avandia), a drug used for type 2 diabetes. Suggest two different methods to prepare the ether in **D** by substitution reactions.




D



rosiglitazone

18.45 Explain why each of the following reactions will not form the given product. Then, design a synthesis of **A** from benzene and **B** from phenol (C_6H_5OH).




Substituent Effects

18.46 Rank the compounds in each group in order of increasing reactivity in electrophilic aromatic substitution: (a) C_6H_6, C_6H_5Cl, C_6H_5CHO; (b) $C_6H_5CH_3$, $C_6H_5NH_2$, $C_6H_5CH_2NH_2$.


18.47 For each of the following substituted benzenes: [1] C_6H_5Br; [2] C_6H_5CN; [3] $C_6H_5OCOCH_3$:

a. Does the substituent donate or withdraw electron density by an inductive effect?
b. Does the substituent donate or withdraw electron density by a resonance effect?
c. On balance, does the substituent make a benzene ring more or less electron rich than benzene itself?
d. Does the substituent activate or deactivate the benzene ring in electrophilic aromatic substitution?


18.48 Consider the tetracyclic aromatic compound drawn below, with rings labeled as **A, B, C,** and **D.** (a) Which of the four rings is ***most*** reactive in electrophilic aromatic substitution? (b) Which of the four rings is ***least*** reactive in electrophilic aromatic substitution? (c) What are the major product(s) formed when this compound is treated with one equivalent of Br_2?



18.49 For each N-substituted benzene, predict whether the compound reacts faster than, slower than, or at a similar rate to benzene in electrophilic aromatic substitution. Then draw the major product(s) formed when each compound reacts with a general electrophile E^+.



18.50 What is the major product of electrophilic addition of HBr to the following alkene? Explain your choice.



18.51 Using resonance structures, explain why a nitroso group (−NO) is an ortho, para director that deactivates a benzene ring toward electrophilic attack.


18.52 Explain the following observation. Ethyl 3-phenylpropanoate ($C_6H_5CH_2CH_2CO_2CH_2CH_3$) reacts with electrophiles to afford ortho- and para-disubstituted arenes, but ethyl 3-phenylprop-2-enoate ($C_6H_5CH{=}CHCO_2CH_2CH_3$) reacts with electrophiles to afford meta-disubstituted arenes.

18.53 Rank the aryl halides in each group in order of increasing reactivity in nucleophilic aromatic substitution by an addition–elimination mechanism.


a. chlorobenzene, *p*-fluoronitrobenzene, *m*-fluoronitrobenzene
b. 1-fluoro-2,4-dinitrobenzene, 1-fluoro-3,5-dinitrobenzene, 1-fluoro-3,4-dinitrobenzene
c. 1-fluoro-2,4-dinitrobenzene, 4-chloro-3-nitrotoluene, 4-fluoro-3-nitrotoluene

Mechanisms


18.54 Draw a stepwise mechanism for the following reaction.



18.55 Draw a stepwise, detailed mechanism for the following intramolecular reaction.




18.56 Draw a stepwise, detailed mechanism for the following reaction.



18.57 Friedel–Crafts alkylation of benzene with (*R*)-2-chlorobutane and $AlCl_3$ affords *sec*-butylbenzene.


a. How many stereogenic centers are present in the product?
b. Would you expect the product to exhibit optical activity? Explain, with reference to the mechanism.

18.58 Draw a stepwise mechanism for the following substitution. Explain why 2-chloropyridine reacts faster than chlorobenzene in this type of reaction.




2-chloropyridine

18.59 Draw a stepwise mechanism for the following reaction.




18.60 Although two products (**A** and **B**) are possible when naphthalene undergoes electrophilic aromatic substitution, only **A** is formed. Draw resonance structures for the intermediate carbocation to explain why this is observed.



naphthalene **A** **B**

This product is formed. This product is *not* formed.

18.61 Draw a stepwise mechanism for the following reaction, which results in the synthesis of bisphenol F (R = H), an additive used in a variety of packaging materials. Bisphenol F is related to BPA (bisphenol A, R = CH_3), a reagent used to harden some plastics, now removed from certain baby products because of its estrogen-like activity that can disrupt endocrine pathways.





bisphenol F
R = H


18.62 Benzyl bromide ($C_6H_5CH_2Br$) reacts rapidly with CH_3OH to afford benzyl methyl ether ($C_6H_5CH_2OCH_3$). Draw a stepwise mechanism for the reaction, and explain why this 1° alkyl halide reacts rapidly with a weak nucleophile under conditions that favor an S_N1 mechanism. Would you expect the para-substituted benzylic halides $CH_3OC_6H_4CH_2Br$ and $O_2NC_6H_4CH_2Br$ to each be more or less reactive than benzyl bromide in this reaction? Explain your reasoning.


Synthesis

18.63 Synthesize each compound from benzene and any other organic or inorganic reagents.


a.



b.



c.



d.


(PABA)
sunscreen component


e.



f.



g.


h.



18.64 Synthesize each compound from toluene ($C_6H_5CH_3$) and any other organic or inorganic reagents.


a.


b.



c.



18.65 Devise a synthesis of each compound from phenol (C_6H_5OH) and any other organic or inorganic reagents.


a.



b.


18.66 Use the reactions in this chapter along with those learned in Chapters 11 and 12 to synthesize each compound. You may use benzene, acetylene (HC≡CH), two-carbon alcohols, ethylene oxide, and any inorganic reagents.

a. 


c. 

b. 

d. 


18.67 Ibufenac, a para-disubstituted arene with the structure $HO_2CCH_2C_6H_4CH_2CH(CH_3)_2$, is a much more potent analgesic than aspirin, but it was never sold commercially because it caused liver toxicity in some clinical trials. Devise a synthesis of ibufenac from benzene and organic halides having fewer than five carbons.


18.68 Carboxylic acid **X** is an intermediate in the multistep synthesis of proparacaine, a local anesthetic. Devise a synthesis of **X** from phenol and any needed organic or inorganic reagents.



Spectroscopy


18.69 Identify the structures of isomers **A** and **B** (molecular formula C_8H_9Br).






18.70 Propose a structure of compound **C** (molecular formula $C_{10}H_{12}O$) consistent with the following data. **C** is partly responsible for the odor and flavor of raspberries.


Compound **C:** IR absorption at 1717 cm^{-1}



18.71 Compound **X** (molecular formula $C_{10}H_{12}O$) was treated with NH_2NH_2, ^-OH to yield compound **Y** (molecular formula $C_{10}H_{14}$). Based on the ^1H NMR spectra of **X** and **Y** given below, what are the structures of **X** and **Y?**




18.72 Reaction of *p*-cresol with two equivalents of 2-methylprop-1-ene affords BHT, a preservative with molecular formula $C_{15}H_{24}O$. BHT gives the following ^1H NMR spectral data: 1.4 (singlet, 18 H), 2.27 (singlet, 3 H), 5.0 (singlet, 1 H), and 7.0 (singlet, 2 H) ppm. What is the structure of BHT? Draw a stepwise mechanism illustrating how it is formed.



p-cresol 2-methylprop-1-ene (2 equiv)

Challenge Problems

18.73 Devise a synthesis of optically active (*S*)-fluoxetine (trade name Prozac) from the given starting materials and any other needed reagents.



(*S*)-fluoxetine

18.74 The ^1H NMR spectrum of phenol (C_6H_5OH) shows three absorptions in the aromatic region: 6.70 (2 ortho H's), 7.14 (2 meta H's), and 6.80 (1 para H) ppm. Explain why the ortho and para absorptions occur at lower chemical shift than the meta absorption.


18.75 Explain the reactivity and orientation effects observed in each heterocycle.




a. Pyridine is less reactive than benzene in electrophilic aromatic substitution and yields 3-substituted products.

b. Pyrrole is more reactive than benzene in electrophilic aromatic substitution and yields 2-substituted products.


18.76 Draw a stepwise, detailed mechanism for the dienone–phenol rearrangement, a reaction that forms alkyl-substituted phenols from cyclohexadienones.




18.77 Draw a stepwise mechanism for the following intramolecular reaction, which is used in the synthesis of the female sex hormone estrone.



18.78 The bicyclic heterocycles quinoline and indole undergo electrophilic aromatic substitution to give the products shown. (a) Explain why electrophilic substitution occurs on the ring without the N atom for quinoline, but occurs on the ring with the N atom in indole. (b) Explain why electrophilic substitution occurs more readily at C8 than C7 in quinoline. (c) Explain why electrophilic substitution occurs more readily at C3 rather than C2 of indole.




18.79 Devise a stepwise mechanism for the following reaction. The reaction does not take place by direct electrophilic aromatic substitution at C2. (Hint: The mechanism begins with addition of an electrophile at C3.)



Carboxylic Acids and the Acidity of the O–H Bond

19



Lysine is an essential amino acid that is needed for protein synthesis. Because lysine cannot be synthesized by humans and it is not stored in the body, lysine must be ingested on a regular basis. Common food sources of lysine are meat, beans, peas, soy, and peanuts. Although most grains are low in lysine, quinoa is relatively high in lysine content and a good source of essential amino acids for a vegetarian diet. Like other amino acids, lysine contains both a carboxylic acid and an amine base. In Chapter 19 we learn about carboxylic acids and the acid–base properties of amino acids.

19.1 Structure and bonding
19.2 Nomenclature
19.3 Physical properties
19.4 Spectroscopic properties
19.5 Interesting carboxylic acids
19.6 Aspirin, arachidonic acid, and prostaglandins
19.7 Preparation of carboxylic acids
19.8 Reactions of carboxylic acids—General features
19.9 Carboxylic acids—Strong organic Brønsted–Lowry acids
19.10 Inductive effects in aliphatic carboxylic acids
19.11 Substituted benzoic acids
19.12 Extraction
19.13 Sulfonic acids
19.14 Amino acids


Chapter 19 serves as a transition between the preceding discussion of resonance and aromaticity, and the subsequent treatment of carbonyl chemistry. We pause to study the chemistry of the OH group by examining **carboxylic acids (RCO_2H)**, and to a lesser extent, **phenols (PhOH)** and **alcohols (ROH).**

In Chapter 19 we concentrate on the acidity of carboxylic acids, and revisit some of the factors that determine acidity, a topic first discussed in Chapter 2. Then, in Chapters 20 and 22 we will learn other reactions of carboxylic acids that occur at the carbonyl group.


19.1 Structure and Bonding

The word **carboxy** (for a COOH group) is derived from ***carb***onyl (C=O) + hydr***oxy*** (OH).


Carboxylic acids **are organic compounds containing a carboxy group (COOH).** Although the structure of a carboxylic acid is often abbreviated as **RCOOH** or **RCO_2H,** keep in mind that the central carbon atom of the functional group is doubly bonded to one oxygen atom and singly bonded to another.



The carbon atom of a carboxy group is surrounded by three groups, making it ***sp*^{2} hybridized** and **trigonal planar,** with bond angles of approximately 120°. The C=O of a carboxylic acid is *shorter* than its C−O.




The C−O single bond of a carboxylic acid is shorter than the C−O single bond of an alcohol. This can be explained by looking at the hybridization of the respective carbon atoms. In the alcohol, the carbon is sp^3 hybridized, whereas in the carboxylic acid the carbon is sp^2 hybridized. As a result, the higher percent *s*-character in the sp^2 hybrid orbital shortens the C−O bond in the carboxylic acid.



Because oxygen is more electronegative than either carbon or hydrogen, **the C−O and O−H bonds are polar.** The electrostatic potential plot of acetic acid in Figure 19.1 shows that the carbon and hydrogen atoms are electron poor and the oxygen atoms are electron rich.

Figure 19.1
Electrostatic potential plot of acetic acid (CH_3COOH)



Acetic acid contains two electron-rich oxygen atoms (in red). Its carbonyl carbon and hydroxy hydrogen are both electron deficient.

19.2 Nomenclature

Both IUPAC and common names are used for carboxylic acids.

19.2A IUPAC System


In IUPAC nomenclature, carboxylic acids are identified by a suffix added to the parent name of the longest chain, and two different endings are used depending on whether the carboxy group is bonded to a chain or ring.


To name a carboxylic acid using the IUPAC system:

[1] If the COOH is bonded to a chain of carbons, find the longest chain containing the COOH group, and change the ***-e*** ending of the parent alkane to the suffix ***-oic acid.*** If the COOH group is bonded to a ring, name the ring and add the words ***carboxylic acid.***

[2] Number the carbon chain or ring to put the **COOH group at C1,** but omit this number from the name. Apply all of the other usual rules of nomenclature.


Sample Problem 19.1 Give the IUPAC name of each compound.

a. 

b. 

Solution


a. [1] Find and name the longest chain containing COOH:



hexane ⟶ **hexano*ic acid***
(6 C's)

The COOH contributes one C to the longest chain.


[2] Number and name the substituents:



two methyl substituents on C4 and C5


Answer: 4,5-dimethylhexanoic acid

b. [1] Find and name the ring bonded to COOH.



cyclohexane + carboxylic acid
(6 C's)


[2] Number and name the substituents:





Number to put COOH at C1 and give the second substituent (CH_3) the lower number (C2).


Answer: 2,5,5-trimethylcyclohexanecarboxylic acid

Problem 19.1 Give the IUPAC name for each compound.

a. 

c. 

b. 

d. 

Problem 19.2 Give the structure corresponding to each IUPAC name.

a. 2-bromobutanoic acid
b. 2,3-dimethylpentanoic acid
c. 3,3,4-trimethylheptanoic acid
d. 2-*sec*-butyl-4,4-diethylnonanoic acid
e. 3,4-diethylcyclohexanecarboxylic acid
f. 1-isopropylcyclobutanecarboxylic acid

19.2B Common Names

Most simple carboxylic acids have common names that are more widely used than their IUPAC names.

- **A common name is formed by using a common parent name followed by the suffix *-ic acid.***

Table 19.1 lists common parent names for some simple carboxylic acids. These parent names are used in the nomenclature of many other compounds with carbonyl groups (Chapters 21 and 22).








Greek letters are used to designate the location of substituents in common names.

- **The carbon adjacent to the COOH is called the α carbon.**
- **The carbon bonded to the α carbon is the β carbon, followed by the γ (gamma) carbon, the δ (delta) carbon, and so forth down the chain. The last carbon in the chain is sometimes called the Ω (omega) carbon.**




Caproic acid, the common name for hexanoic acid, is a foul-smelling carboxylic acid found in the fleshy coat of ginkgo seeds, giving the seeds an unpleasant and even repulsive odor.

Table 19.1 Common Names for Some Simple Carboxylic Acids

Number of C atoms	Structure	Parent name	Common name
1		**form-**	formic acid
2		**acet-**	acetic acid
3		**propion-**	propionic acid
4		**butyr-**	butyric acid
5		**valer-**	valeric acid
6		**capro-**	caproic acid
		**benzo-**	benzoic acid

The α carbon in the common system is numbered C2 in the IUPAC system.




IUPAC numbering begins at the C=O.
Greek lettering begins at the C bonded to the C=O.

Problem 19.3 Draw the structure corresponding to each common name:


a. α-methoxyvaleric acid
b. β-phenylpropionic acid
c. α,β-dimethylcaproic acid
d. α-chloro-β-methylbutyric acid

19.2C Other Nomenclature Facts


Many compounds containing two carboxy groups are also known. In the IUPAC system, **diacids** are named by adding the suffix ***-dioic acid*** to the name of the parent alkane. The three simplest diacids are most often identified by their common names, as shown.



oxalic acid
(ethanedioic acid)



malonic acid
(propanedioic acid)




succinic acid
(butanedioic acid)


Metal salts of carboxylate anions are formed from carboxylic acids in many reactions in Chapter 19. To name the **metal salt of a carboxylate anion,** change the *-ic acid* ending of the carboxylic acid to the suffix ***-ate*** and put three parts together:


name of the metal cation	+	parent	+	suffix
		common or IUPAC		***-ate***

Two examples are shown in Figure 19.2.

Problem 19.4 Give the IUPAC name for each metal salt of a carboxylate anion.

a. 

c. 

b. 



d. 


Figure 19.2
Naming the metal salts of carboxylate anions



sodium cation

parent + suffix
acet- -ate

sodium acetate



potassium cation

parent + suffix
propano- -ate


potassium propanoate

Problem 19.5 Depakote, a drug used to treat seizures and bipolar disorder, consists of a mixture of valproic acid [$(CH_3CH_2CH_2)_2CHCO_2H$] and its sodium salt. Give IUPAC names for each of these compounds.

19.3 Physical Properties


Carboxylic acids exhibit **dipole–dipole** interactions because they have polar C−O and O−H bonds. They also exhibit intermolecular **hydrogen bonding** because they possess a hydrogen atom bonded to an electronegative oxygen atom. Carboxylic acids often exist as **dimers,** held together by *two* intermolecular hydrogen bonds between the carbonyl oxygen atom of one molecule and the OH hydrogen atom of another molecule (Figure 19.3). Carboxylic acids are the **most polar** organic compounds we have studied so far.

Figure 19.3
Two molecules of acetic acid (CH_3COOH) held together by two hydrogen bonds




How these intermolecular forces affect the physical properties of carboxylic acids is summarized in Table 19.2.

Table 19.2 Physical Properties of Carboxylic Acids

Property	Observation
Boiling point and melting point	• Carboxylic acids have higher boiling points and melting points than other compounds of comparable molecular weight.  VDW — MW = 58, bp 0 °C VDW, DD — MW = 58, bp 48 °C VDW, DD, HB — MW = 60, bp 97 °C VDW, DD, two HB — MW = 60, bp 118 °C **Increasing strength of intermolecular forces** **Increasing boiling point**
Solubility	• Carboxylic acids are soluble in organic solvents regardless of size. • Carboxylic acids having ≤ 5 C's are water soluble because they can hydrogen bond with H_2O (Section 3.4C). • Carboxylic acids having > 5 C's are water insoluble because the nonpolar alkyl portion is too large to dissolve in the polar H_2O solvent. These "fatty" acids dissolve in a nonpolar fat-like environment but do not dissolve in water.

Key: VDW = van der Waals, DD = dipole–dipole, HB = hydrogen bonding, MW = molecular weight

Problem 19.6 Rank the following compounds in order of increasing boiling point. Which compound is the most water soluble? Which compound is the least water soluble?




19.4 Spectroscopic Properties

Carboxylic acids have very characteristic IR and NMR absorptions. In the IR, carboxylic acids show two strong absorptions.

- The **C=O group absorbs at about 1710 cm^{-1},** in the usual region for a carbonyl.
- The **O–H absorption** occurs from **2500–3500 cm^{-1}.** This very broad absorption sometimes obscures the C–H peak at 3000 cm^{-1}.

The IR spectrum of butanoic acid in Figure 19.4 illustrates these characteristic peaks.

Figure 19.4
The IR spectrum of butanoic acid, $CH_3CH_2CH_2COOH$




- A strong C=O absorption occurs at 1712 cm^{-1}.
- The broad O–H absorption (2500–3500 cm^{-1}) nearly obscures the C–H peak at ~3000 cm^{-1}.

Carboxylic acids have two noteworthy ^1H NMR absorptions and one noteworthy ^{13}C NMR absorption.

- **The highly deshielded OH proton absorbs in the ^1H NMR spectrum somewhere between 10 and 12 ppm,** farther *downfield* than all other absorptions of common organic compounds. Like the OH signal of an alcohol, the exact location depends on the degree of hydrogen bonding and the concentration of the sample.
- The protons on the α carbon to the carboxy group are somewhat deshielded, absorbing at 2–2.5 ppm.
- In the ^{13}C NMR spectrum, the carbonyl absorption is highly deshielded, appearing at 170–210 ppm.

Sometimes the OH absorption of a carboxylic acid is very broad, so that it is almost buried in the baseline of the ^1H NMR spectrum, making it difficult to see (Figure 19.5).


Figure 19.5 The ^{1}H and ^{13}C NMR spectra of propanoic acid



- **^{1}H NMR spectrum:** There are three signals due to three different kinds of H atoms. The H_a and H_b signals are split into a triplet and quartet, respectively. The H_c signal, a singlet, is due to the highly deshielded OH proton.
- **^{13}C NMR spectrum:** There are three signals due to three different kinds of carbon atoms. The carbonyl carbon is highly deshielded.


Figure 19.5 illustrates the ^{1}H and ^{13}C NMR spectra of propanoic acid.

Problem 19.7 Explain how you could use IR spectroscopy to distinguish among the following three compounds.




19.5 Interesting Carboxylic Acids

Several simple carboxylic acids have characteristic odors and flavors.




formic acid

- **Formic acid (HCOOH),** a carboxylic acid with an acrid odor and a biting taste, is responsible for the sting of some types of ants. The name is derived from the Latin word formica, meaning "ant."



acetic acid


- **Acetic acid (CH_3COOH)** is the sour-tasting component of vinegar. The name comes from the Latin *acetum,* meaning "vinegar." The air oxidation of ethanol to acetic acid is the process that makes "bad" wine taste sour. Acetic acid is an industrial starting material for polymers used in paints and adhesives. Pure acetic acid is often called *glacial* acetic acid, because it freezes just below room temperature (mp = 17 °C), forming white crystals reminiscent of the ice in a glacier.



butanoic acid

- **Butanoic acid ($CH_3CH_2CH_2COOH$)** is an oxidation product that contributes to the disagreeable smell of body odor. Its common name, butyric acid, is derived from the Latin word *butyrum,* meaning "butter," because butyric acid gives rancid butter its peculiar odor and taste.

Oxalic acid and **lactic acid** are simple carboxylic acids quite prevalent in nature. Oxalic acid occurs naturally in spinach and rhubarb. Lactic acid gives sour milk its distinctive taste.



oxalic acid lactic acid



Although oxalic acid is toxic, you would have to eat about nine pounds of spinach at one time to ingest a fatal dose.


4-Hydroxybutanoic acid, known by its common name γ-hydroxybutyric acid (GHB), is an illegal recreational drug that depresses the central nervous system and results in intoxication. GHB is highly addictive and widely abused, and because its taste can be easily masked in an alcoholic beverage, it has been used as a "date rape" drug.



4-hydroxybutanoic acid
GHB


Salts of carboxylic acids are commonly used as preservatives. Sodium benzoate, a fungal growth inhibitor, is a preservative used in soft drinks, and potassium sorbate is an additive that prolongs the shelf-life of baked goods and other foods.

Soaps, the sodium salts of fatty acids, were discussed in Section 3.6.



sodium benzoate potassium sorbate

Problem 19.8 Isotretinoin, a fatty acid, is a medication used to treat severe acne that has not responded to other drugs. Because it causes fetal abnormalities, it cannot be taken by pregnant women, and its prescription and use are carefully monitored. (a) Label each C=C as *E* or *Z*. (b) Label each σ bond joining two C=C's as *s*-cis or *s*-trans.




isotretinoin

19.6 Aspirin, Arachidonic Acid, and Prostaglandins



The word *aspirin* is derived from the prefix ***a-*** for *acetyl* + ***spir*** from the Latin name *spirea* for the meadowsweet plant.

Recall from Chapter 2 that **aspirin (acetylsalicylic acid)** is a synthetic carboxylic acid, similar in structure to **salicin,** a naturally occurring compound isolated from willow bark, and **salicylic acid,** found in meadowsweet.



aspirin (acetylsalicylic acid)

salicin (isolated from willow bark)

salicylic acid (isolated from meadowsweet)


sodium salicylate (sweet carboxylate salt)

Both salicylic acid and sodium salicylate (its sodium salt) were widely used analgesics in the nineteenth century, but both had undesirable side effects. Salicylic acid irritated the mucous membranes of the mouth and stomach, and sodium salicylate was too sweet for most patients. Aspirin, a synthetic compound, was first sold in 1899 after Felix Hoffmann, a German chemist at Bayer Company,

developed a feasible commercial synthesis. Hoffmann's work was motivated by personal reasons; his father suffered from rheumatoid arthritis and was unable to tolerate the sweet taste of sodium salicylate.


How does aspirin relieve pain and reduce inflammation? Aspirin blocks the synthesis of **prostaglandins,** 20-carbon fatty acids with a five-membered ring that are responsible for pain, inflammation, and a wide variety of other biological functions. **$PGF_{2\alpha}$** contains the typical carbon skeleton of a prostaglandin.

Aspirin is the most widely used pain reliever and anti-inflammatory agent in the world, yet its mechanism of action remained unknown until the 1970s. John Vane, Bengt Samuelsson, and Sune Bergstrom shared the 1982 Nobel Prize in Physiology or Medicine for unraveling the details of its mechanism.




$PGF_{2\alpha}$
a prostaglandin


Prostaglandins are not stored in cells. Rather they are synthesized from arachidonic acid, a polyunsaturated fatty acid having four cis double bonds. Unlike hormones, which are transported in the bloodstream to their sites of action, prostaglandins act where they are synthesized. Aspirin acts by blocking the synthesis of prostaglandins from arachidonic acid. Aspirin inactivates cyclooxygenase, an enzyme that converts arachidonic acid to PGG_2, an unstable precursor of $PGF_{2\alpha}$ and other prostaglandins. **Aspirin lessens pain and decreases inflammation because it prevents the synthesis of prostaglandins, the compounds responsible for both of these physiological responses.**



Although prostaglandins have a wide range of biological activity, their inherent instability often limits their usefulness as drugs. Consequently, more stable analogues with useful medicinal properties have been synthesized. For example, latanoprost (trade name Xalatan) and bimatoprost (trade name Lumigan) are prostaglandin analogues used to reduce eye pressure in individuals with glaucoma.



latanoprost



bimatoprost

Problem 19.9 How many tetrahedral stereogenic centers does $PGF_{2\alpha}$ contain? Draw its enantiomer. How many of its double bonds can exhibit cis-trans isomerism? Considering both its double bonds and its tetrahedral stereogenic centers, how many stereoisomers are possible for $PGF_{2\alpha}$?

19.7 Preparation of Carboxylic Acids


Our discussion of the reactions involving carboxylic acids begins with a brief list of reactions that synthesize them. In these reactions, the carboxy group is formed in the *product,* and many different functional groups serve as starting materials. Reactions that produce a particular functional group are called **preparations.**

In the remainder of Chapter 19 (and Chapters 20 and 22) we discuss reactions in which a carboxylic acid is a *starting material* that may be converted to a variety of different products. Keep in mind that **reactions of a particular functional group follow a common theme.** For example, alkenes undergo addition reactions. As a result, these reactions are easier to learn than the list of preparations, in which vastly different functional groups undergo a wide variety of reactions to form the same kind of product.

Where have we encountered carboxylic acids as reaction products before? The carbonyl carbon is highly oxidized, because it has three C−O bonds, so **carboxylic acids are typically prepared by oxidation reactions.** Three oxidation methods are summarized below. Two other useful methods to prepare carboxylic acids are presented in Chapter 20.

[1] By oxidation of 1° alcohols (Section 12.12B)


1° Alcohols are converted to carboxylic acids with $Na_2Cr_2O_7$, $K_2Cr_2O_7$, or CrO_3 in the presence of H_2O and H_2SO_4.



1° alcohol

[2] By oxidation of alkyl benzenes (Section 18.15A)


Alkyl benzenes having at least one benzylic C−H bond are oxidized with $KMnO_4$ to benzoic acid.



Benzoic acid is *always* the product regardless of the alkyl benzene used as starting material.


[3] By oxidative cleavage of alkynes (Section 12.11)

Both internal and terminal alkynes are oxidatively cleaved with ozone to give carboxylic acids.




With internal alkynes two carboxylic acids are formed as products. With terminal alkynes, the *sp* hybridized C−H bond is converted to CO_2.

Problem 19.10 What alcohol can be oxidized to each carboxylic acid?




Problem 19.11 Identify **A–D** in the following reactions.



19.8 Reactions of Carboxylic Acids—General Features


The polar C−O and O−H bonds, nonbonded electron pairs on oxygen, and the π bond give a carboxylic acid many reactive sites, complicating its chemistry somewhat. By far, **the most important reactive feature of a carboxylic acid is its polar O−H bond, which is readily cleaved with base.**

- Carboxylic acids react as Brønsted–Lowry acids—that is, as proton donors.




Much of the rest of Chapter 19 is devoted to the acidity of carboxylic acids, as well as some related acid–base reactions. Two other structural features are less important in the reactions of carboxylic acids, but they play a role in the reactions of Chapters 20 and 22.

The nonbonded electron pairs on oxygen create electron-rich sites that can be protonated by strong acids (H−A). Protonation occurs at the carbonyl oxygen because the resulting conjugate acid is resonance stabilized. As a result, **carboxylic acids are weakly basic—they react with strong acids by protonation of the carbonyl oxygen.** This reaction plays an important role in several mechanisms in Chapter 22.



three resonance structures for the conjugate acid

Finally, the polar C−O bonds make the carboxy carbon electrophilic, so carboxylic acids react with nucleophiles. Nucleophilic attack occurs at an sp^2 hybridized carbon atom, so it results in the cleavage of the π bond, as well. This reaction is also discussed in Chapter 22.



Problem 19.12 Draw the cation that results when a carboxylic acid is protonated on the oxygen of the OH group. Explain why protonation of the carbonyl oxygen is preferred to protonation of the hydroxy oxygen.

19.9 Carboxylic Acids—Strong Organic Brønsted–Lowry Acids

Carboxylic acids are strong organic acids, and as such, readily react with Brønsted–Lowry bases to form carboxylate anions.

Recall from Section 2.3 that **the lower the pK_a, the stronger the acid.**




carboxylate anion


What bases are used to deprotonate a carboxylic acid? As we learned in Section 2.3, equilibrium favors the products of an acid–base reaction when the weaker base and acid are formed. Because a weaker acid has a higher pK_a, the following general rule results:

- **An acid can be deprotonated by a base that has a conjugate acid with a higher pK_a.**

Because the pK_a values of many carboxylic acids are ~5, bases that have conjugate acids with pK_a values higher than 5 are strong enough to deprotonate them. Thus, acetic acid (pK_a = 4.8) and benzoic acid (pK_a = 4.2) can be deprotonated with NaOH and $NaHCO_3$, as shown in the following equations.



acetic acid
stronger acid
pK_a = 4.8




benzoic acid
stronger acid
pK_a = 4.2

Table 19.3 lists common bases that can be used to deprotonate carboxylic acids. It is noteworthy that even a weak base like $NaHCO_3$ is strong enough to remove a proton from RCOOH.

Table 19.3 Common Bases Used to Deprotonate Carboxylic Acids

	Base	Conjugate acid (pK_a)
Increasing basicity ↓	$Na^+ HCO_3^-$	H_2CO_3 (6.4)
	NH_3	NH_4^+ (9.4)
	Na_2CO_3	HCO_3^- (10.2)
	$Na^+ {}^-OCH_3$	CH_3OH (15.5)
	$Na^+ {}^-OH$	H_2O (15.7)
	$Na^+ {}^-OCH_2CH_3$	CH_3CH_2OH (16)
	$Na^+ H^-$	H_2 (35)


Why are carboxylic acids such strong organic acids? Remember that a strong acid has a weak, stabilized conjugate base. **Deprotonation of a carboxylic acid forms a resonance-stabilized conjugate base—a carboxylate anion.** Two equivalent resonance structures can be drawn for acetate (the conjugate base of acetic acid), both of which place a negative charge on an electronegative O atom. In the resonance hybrid, therefore, the negative charge is delocalized over two oxygen atoms.



acetic acid — two resonance structures for acetate, the conjugate base — hybrid


How resonance affects acidity was first discussed in Section 2.5C.

Experimental data support this resonance description of acetate. **The acetate anion has two C–O bonds of equal length** (127 pm) and intermediate between the length of a C–O single bond (136 pm) and C=O (121 pm).



acetate hybrid

Resonance stabilization accounts for why carboxylic acids are more acidic than other compounds with O–H bonds—namely, alcohols and phenols. For example, the pK_a values of ethanol (CH_3CH_2OH) and phenol (C_6H_5OH) are 16 and 10, respectively, both higher than the pK_a of acetic acid (4.8).



To understand the relative acidity of ethanol, phenol, and acetic acid, we must compare the stability of their conjugate bases and use the following rule:

- **Anything that stabilizes a conjugate base A:⁻ makes the starting acid H–A more acidic.**


Ethoxide, the conjugate base of ethanol, bears a negative charge on an oxygen atom, but there are no additional factors to further stabilize the anion. Because ethoxide is less stable than acetate, **ethanol is a weaker acid than acetic acid.**



ethanol — ethoxide

The resonance hybrid of phenoxide illustrates that its negative charge is dispersed over four atoms—three C atoms and one O atom.

Like acetate, **phenoxide** ($C_6H_5O^-$, the conjugate base of phenol) is also resonance stabilized. In the case of phenoxide, however, there are *five* resonance structures that disperse the negative charge over a total of *four* different atoms (three different carbons and the oxygen).




hybrid — phenol — 1 — 2 — 3 — 4 — 5

Phenoxide is more stable than ethoxide, but less stable than acetate, because acetate has two electronegative oxygen atoms upon which to delocalize the negative charge, whereas phenoxide has only one. Additionally, phenoxide resonance structures **2–4** have the negative charge on a carbon, a less electronegative element than oxygen. As a result, structures **2–4** are less stable than structures **1** and **5,** which have the negative charge on oxygen.

Moreover, resonance structures **1** and **5** have intact aromatic rings, whereas structures **2–4** do not. This, too, makes structures **2–4** less stable than **1** and **5.** Figure 19.6 summarizes this information about phenoxide by displaying the approximate relative energies of its five resonance structures and its hybrid.

Figure 19.6
The relative energies of the five resonance structures for phenoxide and its hybrid



As a result, resonance stabilization of the conjugate base is important in determining acidity, but **the absolute number of resonance structures alone is not what's important.** We must evaluate their relative contributions to predict the relative stability of the conjugate bases.

Keep in mind that although carboxylic acids are strong organic acids, they are still much weaker than strong inorganic acids like HCl and H_2SO_4, which have pK_a values < 0.

- **Because of their O–H bond, RCOOH, ROH, and C_6H_5OH are more acidic than most organic hydrocarbons.**

- **A carboxylic acid is a stronger acid than an alcohol or phenol because its conjugate base is most effectively resonance stabilized.**

The relationship between acidity and stability of the conjugate base is summarized for acetic acid, phenol, and ethanol in Figure 19.7.

Because alcohols and phenols are weaker acids than carboxylic acids, stronger bases are needed to deprotonate them. To deprotonate C_6H_5OH (pK_a = 10), a base whose conjugate acid has a pK_a > 10 is needed. Thus, of the bases listed in Table 19.3, $NaOCH_3$, NaOH, $NaOCH_2CH_3$, and NaH are strong enough. To deprotonate CH_3CH_2OH (pK_a = 16), only NaH is strong enough.

Problem 19.13 Draw the products of each acid–base reaction.





Figure 19.7


Summary: The relationship between acidity and conjugate base stability for acetic acid, phenol, and ethanol



- **Acetate is the most stable conjugate base** because it has two equivalent resonance structures, both of which place a negative charge on an O atom.
- **Phenoxide** has only one O atom to accept the negative charge. The two resonance structures that contain an intact aromatic ring and place a negative charge on an O atom are major contributors to the hybrid. Resonance stabilizes phenoxide but not as much as resonance stabilizes acetate.
- **Ethoxide is the least stable conjugate base** because it has no additional resonance stabilization.

Problem 19.14 Given the pK_a values in Appendix A, which of the following bases are strong enough to deprotonate CH_3COOH: (a) F^-; (b) $(CH_3)_3CO^-$; (c) CH_3^-; (d) $^-NH_2$; (e) Cl^-?

Problem 19.15 Rank the labeled protons (H_a–H_c) in mandelic acid, a naturally occurring carboxylic acid in plums and peaches, in order of increasing acidity. Explain in detail why you chose this order.



mandelic acid


19.10 Inductive Effects in Aliphatic Carboxylic Acids

The pK_a of a carboxylic acid is affected by nearby groups that inductively donate or withdraw electron density.

- **Electron-withdrawing groups stabilize a conjugate base, making a carboxylic acid more acidic.**
- **Electron-donating groups destabilize the conjugate base, making a carboxylic acid less acidic.**

The relative acidity of CH_3COOH, $ClCH_2COOH$, and $(CH_3)_3CCOOH$ illustrates these principles in the following equations.


We first learned about inductive effects and acidity in Section 2.5B.




- $ClCH_2COOH$ is more acidic ($pK_a = 2.8$) than CH_3COOH ($pK_a = 4.8$) because its conjugate base is stabilized by the electron-withdrawing inductive effect of the electronegative Cl.
- $(CH_3)_3CCOOH$ is less acidic ($pK_a = 5.1$) than CH_3COOH because the three polarizable CH_3 groups donate electron density and destabilize the conjugate base.

The number, electronegativity, and location of substituents also affect acidity.


- **The larger the number of electronegative substituents, the stronger the acid.**



- **The more electronegative the substituent, the stronger the acid.**




- **The closer the electron-withdrawing group to the COOH, the stronger the acid.**



Problem 19.16 Match each of the following pK_a values (3.2, 4.9, and 0.2) to the appropriate carboxylic acid: (a) CH_3CH_2COOH; (b) CF_3COOH; (c) ICH_2COOH.

Problem 19.17 Rank the compounds in each group in order of increasing acidity.




19.11 Substituted Benzoic Acids

Recall from Chapter 18 that substituents on a benzene ring either donate or withdraw electron density, depending on the balance of their inductive and resonance effects. These same effects also determine the acidity of substituted benzoic acids. There are two rules to keep in mind.

Rule [1] **Electron-donor groups destabilize a conjugate base, making an acid less acidic.**


An electron-donor group destabilizes a conjugate base by donating electron density onto a negatively charged carboxylate anion. A benzoic acid substituted by an electron-donor group has a higher pK_a than benzoic acid (pK_a = 4.2).



Rule [2] **Electron-withdrawing groups stabilize a conjugate base, making an acid more acidic.**

An electron-withdrawing group stabilizes a conjugate base by removing electron density from the negatively charged carboxylate anion. A benzoic acid substituted by an electron-withdrawing group has a lower pK_a than benzoic acid (pK_a = 4.2).

W = Electron-withdrawing group



This acid is more acidic than benzoic acid.

$pK_a < 4.2$

W stabilizes the carboxylate anion.

How do we know which groups are electron donating or electron withdrawing on a benzene ring? We already learned the characteristics of electron-donating and electron-withdrawing groups in Chapter 18, and how they affect the rate of electrophilic aromatic substitution. These principles can now be extended to substituted benzoic acids.

Electron-donor groups D → **• activate benzene to electrophilic attack • make a benzoic acid less acidic**

Electron-withdrawing groups W → **• deactivate benzene to electrophilic attack • make a benzoic acid more acidic**

Figure 19.8 illustrates how common electron-donating and electron-withdrawing groups affect both the rate of reaction of a benzene ring towards electrophiles and the acidity of substituted benzoic acids.

Figure 19.8


How common substituents affect the reactivity of a benzene ring towards electrophiles and the acidity of substituted benzoic acids

	Substituent	Effect in electrophilic substitution	Effect on acidity of substituted benzoic acids
electron-donating groups	$-\ddot{N}H_2$ [$\ddot{N}HR$, $\ddot{N}R_2$]	**activating groups**	These groups make a benzoic acid **less acidic.**
	$-\ddot{O}H$		
	$-\ddot{O}R$		
	$-\ddot{N}HCOR$		
	$-R$		
electron-withdrawing groups	$-\ddot{X}:$ [X = F, Cl, Br, I]	**deactivating groups**	These groups make a benzoic acid **more acidic.**
	$-CHO$		
	$-COR$		
	$-COOR$		
	$-COOH$		
	$-CN$		
	$-SO_3H$		
	$-NO_2$		
	$-\overset{+}{N}R_3$		

Increasing acidity ↓


- **Groups that donate electron density activate a benzene ring towards electrophilic attack and make a benzoic acid *less* acidic.** Common electron-donating groups are R groups, or groups that have an N or O atom (with a lone pair) bonded to the benzene ring.
- **Groups that withdraw electron density deactivate a benzene ring towards electrophilic attack, and make a benzoic acid *more* acidic.** Common electron-withdrawing groups are the halogens, or groups with an atom Y (with a full or partial positive charge) bonded to the benzene ring.

Sample Problem 19.2 Rank the following three carboxylic acids in order of increasing acidity.




Solution

***p*-Methoxybenzoic acid (B):** The CH_3O group is an electron-donor group because its electron-donating resonance effect is stronger than its electron-withdrawing inductive effect (Section 18.6). This destabilizes the conjugate base by donating electron density to the negatively charged carboxylate anion, making **B** less acidic than benzoic acid **A.** Two of the possible resonance structures for **B**'s conjugate base are drawn.




***p*-Nitrobenzoic acid (C):** The NO_2 group is an electron-withdrawing group because of both inductive effects and resonance (Section 18.6). This stabilizes the conjugate base by removing electron density from the negatively charged carboxylate anion, making **C** more acidic than benzoic acid **A.** Two of the possible resonance structures for **C**'s conjugate base are drawn.




By this analysis, the order of acidity is **B < A < C.**

Problem 19.18 Rank the compounds in each group in order of increasing acidity.



Problem 19.19 Substituted phenols show substituent effects similiar to substituted benzoic acids. Should the pK_a of phenol **A,** one of the naturally occurring phenols called urushiols isolated from poison ivy, be higher or lower than the pK_a of phenol (C_6H_5OH, pK_a = 10)? Explain.



A



Poison ivy contains the irritant urushiol.

Extraction has long been and remains the first step in isolating a natural product from its source.

19.12 Extraction


An organic chemist in the laboratory must separate and purify mixtures of compounds. One particularly useful technique is **extraction,** which uses solubility differences and acid–base principles to separate and purify compounds.

Two solvents are used in extraction: water or an aqueous solution such as 10% $NaHCO_3$ or 10% NaOH; and an organic solvent such as dichloromethane (CH_2Cl_2), diethyl ether, or hexane. **Compounds are separated by their solubility differences in an aqueous and organic solvent.**

An item of glassware called a **separatory funnel,** depicted in Figure 19.9, is used for the extraction. When two insoluble liquids are added to the separatory funnel, two layers form, with the less dense liquid on top and the more dense liquid on the bottom.

Suppose a mixture of benzoic acid (C_6H_5COOH) and NaCl is added to a separatory funnel containing H_2O and CH_2Cl_2. The benzoic acid would dissolve in the organic layer and the NaCl would dissolve in the water layer. Separating the organic and aqueous layers and placing them in different flasks separates the benzoic acid and NaCl from each other.

Figure 19.9
Using a separatory funnel for extraction




- When two insoluble liquids are added to a separatory funnel, two layers are visible, and the less dense liquid forms the upper layer.
- To separate the layers, the lower layer can be drained from the bottom of the separatory funnel by opening the stopcock. The top layer can then be poured out the top neck of the funnel.

How could we separate a mixture of benzoic acid and cyclohexanol? Both compounds are organic, and as a result, both are soluble in an organic solvent such as CH_2Cl_2 and insoluble in water. If a mixture of benzoic acid and cyclohexanol were added to a separatory funnel with CH_2Cl_2 and water, both would dissolve in the CH_2Cl_2 layer, and the two compounds would *not*


be separated from each other. Is it possible to use extraction to separate two compounds of this sort that have similar solubility properties?

Recall from Tables 9.1 and 19.2 that alcohols and carboxylic acids having more than five carbons are water insoluble.




If a carboxylic acid is one of the compounds, the answer is *yes,* because we can use acid–base chemistry to change its solubility properties.

When benzoic acid (a strong organic acid) is treated with aqueous NaOH, benzoic acid is deprotonated, forming sodium benzoate. Because sodium benzoate is ionic, it is soluble in water, but insoluble in organic solvents.



A similar acid–base reaction does *not* occur when cyclohexanol is treated with NaOH because organic alcohols are much weaker organic acids, so they can be deprotonated only by a *very strong base* such as NaH. NaOH is not strong enough to form significant amounts of the sodium alkoxide.




This difference in acid–base chemistry can be used to separate benzoic acid and cyclohexanol by the stepwise extraction procedure illustrated in Figure 19.10. This extraction scheme relies on two principles:

- **Extraction can separate only compounds having different solubility properties. One compound must dissolve in the aqueous layer and one must dissolve in the organic layer.**

- **A carboxylic acid can be separated from other organic compounds by converting it to a water-soluble carboxylate anion by an acid–base reaction.**



Thus, the water-soluble salt, $C_6H_5CO_2^-Na^+$ (derived from $C_6H_5CO_2H$ by an acid–base reaction) can be separated from water-insoluble cyclohexanol by an extraction procedure.



Figure 19.10 Separation of benzoic acid and cyclohexanol by an extraction procedure




- Both compounds dissolve in the organic solvent CH_2Cl_2.
- Adding 10% aqueous NaOH solution forms two layers. When the two layers are mixed, the NaOH deprotonates $C_6H_5CO_2H$ to form $C_6H_5CO_2^-Na^+$, which dissolves in the aqueous layer.
- The cyclohexanol remains in the CH_2Cl_2 layer.
- Draining the lower layer out the bottom stopcock separates the two layers, and the separation process is complete.
- Cyclohexanol (dissolved in CH_2Cl_2) is in one flask. The sodium salt of benzoic acid, $C_6H_5CO_2^-Na^+$ (dissolved in water) is in another flask.

Problem 19.20 Which of the following pairs of compounds can be separated from each other by an extraction procedure?

a.  and 

b.  and 


c.  and NaCl

d. NaCl and KCl


19.13 Sulfonic Acids

Although much less common than carboxylic acids, **sulfonic acids** constitute a useful group of organic acids. Sulfonic acids have the general structure **RSO_3H.** The most widely used sulfonic acid, ***p*-toluenesulfonic acid,** was first discussed in Section 2.6.

Recall from Section 9.13 that **$CH_3C_6H_4SO_2$–** is called a **tosyl group,** abbreviated by the letters **Ts.** For this reason, *p*-toluenesulfonic acid (also called tosic acid) is abbreviated as **TsOH.**




sulfonic acid



p-toluenesulfonic acid

TsOH

Sulfonic acids are very strong acids (pK_a values ≈ –7) because their conjugate bases are resonance stabilized, and all the resonance structures delocalize a negative charge on oxygen. The conjugate base of a sulfonic acid is called a **sulfonate anion.**



strong acid
$pK_a \approx -7$

three resonance structures
All have a negative charge on oxygen.

Because sulfonate anions are such weak bases, they make **good leaving groups** in nucleophilic substitution reactions, as we learned in Section 9.13.

Problem 19.21 Two other commonly used sulfonic acids are methanesulfonic acid (CH_3SO_3H) and trifluoromethanesulfonic acid (CF_3SO_3H). Which has the weaker conjugate base? Which conjugate base is the better leaving group? Which of these acids has the higher pK_a?

19.14 Amino Acids

Chapter 29 discusses the synthesis of amino acids and their conversion to proteins.


Amino acids, one of four kinds of small biomolecules that have important biological functions in the cell (Section 3.9), also undergo proton transfer reactions.

19.14A Introduction




Humans can synthesize only 10 of the 20 amino acids needed for protein synthesis. The remaining 10, called **essential amino acids,** must be obtained from the diet and consumed on a regular, almost daily basis. Vegetarian diets must be carefully balanced to obtain all the essential amino acids. Grains—wheat, rice, and corn—are low in lysine, and legumes—beans, peas, and peanuts—are low in methionine, but a combination of these foods provides all the needed amino acids. Thus, a diet of corn tortillas and beans, or rice and tofu, provides all essential amino acids. A peanut butter sandwich on wheat bread does, too.

Amino acids contain two functional groups—an amino group (NH_2) and a carboxy group (COOH). In most naturally occurring amino acids, the amino group is bonded to the α carbon, and so they are called **α-amino acids.** Amino acids are the building blocks of proteins, biomolecules that comprise muscle, hair, fingernails, and many other biological tissues.



α-amino acid

The 20 amino acids that occur naturally in proteins differ in the identity of the R group bonded to the α carbon. **The simplest amino acid, called glycine, has R = H.** When the R group is any other substituent, **the α carbon is a stereogenic center,** and there are two possible enantiomers.



glycine
no stereogenic centers


L amino acid
Only this isomer occurs in proteins.

D amino acid

Amino acids exist in nature as only one of these enantiomers. Except when the R group is CH_2SH, the stereogenic center on the α carbon has the *S* configuration. An older system of nomenclature names the **naturally occurring enantiomer of an amino acid as the L isomer, and its unnatural enantiomer the D isomer.**

The R group of an amino acid can be H, alkyl, aryl, or an alkyl chain containing an N, O, or S atom. Representative examples are listed in Table 19.4. All amino acids have common names, which are abbreviated by a three-letter or one-letter designation. For example, glycine is often written as the three-letter abbreviation **Gly,** or the one-letter abbreviation **G.** These abbreviations are also given in Table 19.4. A complete list of the 20 naturally occurring amino acids is found in Figure 29.2.

Table 19.4 Representative Amino Acids

General structure: 

R group	Name	Three-letter abbreviation	One-letter abbreviation
H	glycine	Gly	G
CH_3	alanine	Ala	A
$CH_2C_6H_5$	phenylalanine	Phe	F
CH_2OH	serine	Ser	S
CH_2SH	cysteine	Cys	C
$CH_2CH_2SCH_3$	methionine	Met	M
CH_2CH_2COOH	glutamic acid	Glu	E
$(CH_2)_4NH_2$	lysine	Lys	K

Problem 19.22 Draw both enantiomers of each amino acid and label them as *R* or *S:* (a) phenylalanine; (b) methionine.


19.14B Acid–Base Properties

An amino acid is both an acid and a base.

- **The NH_2 group has a nonbonded electron pair, making it a base.**
- **The COOH group has an acidic proton, making it an acid.**

Amino acids are never uncharged neutral compounds. They exist as salts, so they have very high melting points and are very soluble in water.

- **Proton transfer from the acidic carboxy group to the basic amino group forms a salt called a *zwitterion,* which contains both a positive and a negative charge.**




This neutral form of an amino acid does *not* exist.

zwitterion
This salt is the neutral form of an amino acid.


In actuality, an amino acid can exist in three different forms, depending on the pH of the aqueous solution in which it is dissolved.

When the pH of a solution is ~6, alanine (R = CH_3) exists in its zwitterionic form (**A**), having no net charge. In this form the carboxy group bears a negative charge—it is a **carboxylate anion**—and the amino group bears a net positive charge (an **ammonium cation**).




alanine
A
a neutral zwitterion
This form exists at pH ≈ 6.

When strong acid is added to lower the pH (≤ 2), the carboxylate anion is protonated and the **amino acid has a net positive charge** (form **B**).



A **B**
overall (+1) charge
This form exists at pH ≤ 2.

When strong base is added to **A** to raise the pH (≥ 10), the ammonium cation is deprotonated and the **amino acid has a net negative charge** (form **C**).




A **C**
overall (–1) charge
This form exists at pH ≥ 10.

Thus, **alanine exists in one of three different forms depending on the pH of the solution in which it is dissolved.** If the pH of a solution is gradually increased from 2 to 10, the following process occurs.

- At low pH alanine has a net (+) charge (form **B**).
- As the pH is increased to ~6, the carboxy group is deprotonated, and the amino acid exists as a zwitterion with no overall charge (form **A**).
- At high pH, the ammonium cation is deprotonated, and the amino acid has a net (–) charge (form **C**).

These reactions are summarized in Figure 19.11.

Figure 19.11
Summary of the acid–base reactions of alanine



Problem 19.23 Explain why amino acids, unlike most other organic compounds, are insoluble in organic solvents like diethyl ether.

Problem 19.24 Draw the positively charged, neutral, and negatively charged forms for the amino acid glycine. Which species predominates at pH 11? Which species predominates at pH 1?

19.14C Isoelectric Point

Because a protonated amino acid has at least two different protons that can be removed, a pK_a value is reported for each of these protons. For example, the pK_a of the carboxy proton of alanine is 2.35 and the pK_a of the ammonium proton is 9.87. Table 29.1 lists these values for all 20 amino acids.

- **The pH at which the amino acid exists primarily in its neutral form is called its *isoelectric point,* abbreviated as p*I*.**

More information on the isoelectric point can be found in Section 29.1.

Generally, the isoelectric point is the average of both pK_a values of an amino acid:

$$\textbf{Isoelectric point} = \textbf{p}I = \frac{\text{p}K_a\,(\text{COOH}) + \text{p}K_a\,(\text{NH}_3^+)}{2}$$

$$\text{For alanine:}\quad \textbf{p}I = \frac{2.35 + 9.87}{2} = \frac{6.12}{\text{p}I\ \text{(alanine)}}$$

Problem 19.25 The pK_a values for the carboxy and ammonium protons of phenylalanine are 2.58 and 9.24, respectively. What is the isoelectric point of phenylalanine? Draw the structure of phenylalanine at its isoelectric point.

Problem 19.26 Explain why the pK_a of the COOH group of glycine is much lower than the pK_a of the COOH of acetic acid (2.35 compared to 4.8).

KEY CONCEPTS

Carboxylic Acids and the Acidity of the O–H Bond


General Facts

- Carboxylic acids contain a carboxy group (COOH). The central carbon is sp^2 hybridized and trigonal planar (19.1).
- Carboxylic acids are identified by the suffixes *-oic acid, carboxylic acid,* or *-ic acid* (19.2).
- Carboxylic acids are polar compounds that exhibit hydrogen bonding interactions (19.3).

Summary of Spectroscopic Absorptions (19.4)

IR absorptions	C=O	~1710 cm^{-1}
	O–H	3500–2500 cm^{-1} (very broad and strong)
^{1}H NMR absorptions	O–H	10–12 ppm (highly deshielded proton)
	C–H α to COOH	2–2.5 ppm (somewhat deshielded C_{sp^3}–H)
^{13}C NMR absorption	C=O	170–210 ppm (highly deshielded carbon)

General Acid–Base Reaction of Carboxylic Acids (19.9)




p$K_a \approx 5$ carboxylate anion

- Carboxylic acids are especially acidic because carboxylate anions are resonance stabilized.
- For equilibrium to favor the products, the base must have a conjugate acid with a p$K_a > 5$. Common bases are listed in Table 19.3.

Factors That Affect Acidity

Resonance effects

- A carboxylic acid is more acidic than an alcohol or phenol because its conjugate base is more effectively stabilized by resonance (19.9).




Inductive effects

- Acidity increases with the presence of electron-withdrawing groups (like the electronegative halogens) and decreases with the presence of electron-donating groups (like polarizable alkyl groups) (19.10).

Substituted benzoic acids

- Electron-donor groups **(D)** make a substituted benzoic acid less acidic than benzoic acid.
- Electron-withdrawing groups **(W)** make a substituted benzoic acid more acidic than benzoic acid.




Other Facts

- Extraction is a useful technique for separating compounds having different solubility properties. Carboxylic acids can be separated from other organic compounds by extraction, because aqueous base converts a carboxylic acid into a water-soluble carboxylate anion (19.12).
- A sulfonic acid (RSO_3H) is a strong acid because it forms a weak, resonance-stabilized conjugate base on deprotonation (19.13).
- Amino acids have an amino group on the α carbon to the carboxy group [$RCH(NH_2)COOH$]. Amino acids exist as zwitterions at pH ≈ 6. Adding acid forms a species with a net (+1) charge [$RCH(NH_3)COOH]^+$. Adding base forms a species with a net (–1) charge [$RCH(NH_2)COO]^-$ (19.14).

PROBLEMS

Problems Using Three-Dimensional Models


19.27 Answer each question for **A** and **B** depicted in the ball-and-stick models.



A **B**


a. What is the IUPAC name for each compound?
b. What product is formed when each compound is treated with NaOH?
c. Name the products formed in part (b).
d. Draw the structure of an isomer that is at least 10^5 times less acidic than each compound.


19.28 Rank the carboxylic acids in order of increasing acidity.





Nomenclature


19.29 Give the IUPAC name for each compound.


a. 

b. 

c. 

d. 

e. 

f. 


19.30 Draw the structure corresponding to each name.

a. 3,3-dimethylpentanoic acid
b. 4-chloro-3-phenylheptanoic acid
c. (*R*)-2-chloropropanoic acid
d. *m*-hydroxybenzoic acid
e. potassium acetate
f. sodium α-bromobutyrate
g. 2,2-dichloropentanedioic acid
h. 4-isopropyl-2-methyloctanedioic acid

19.31 Give an IUPAC and common name for each of the following naturally occurring carboxylic acids: (a) $CH_3CH(OH)CO_2H$ (lactic acid); (b) $HOCH_2CH_2C(OH)(CH_3)CH_2CO_2H$ (mevalonic acid).


Physical Properties


19.32 Rank the following compounds in order of increasing boiling point.





Preparation of Carboxylic Acids

19.33 Draw the organic products formed in each reaction.


a. 


b. 


c. 

d. 

19.34 Identify the lettered compounds in each reaction sequence.


a. 


b. 


c. 

Acid–Base Reactions; General Questions on Acidity


19.35 Using the pK_a table in Appendix A, determine whether each of the following bases is strong enough to deprotonate the three compounds listed below. Bases: [1] ^-OH; [2] $CH_3CH_2^-$; [3] $^-NH_2$; [4] NH_3; [5] $HC{\equiv}C^-$.


a. 
pK_a = 4.3


b. 
pK_a = 9.4


c. 
pK_a = 18

19.36 Draw the products of each acid–base reaction, and using the pK_a table in Appendix A, determine if equilibrium favors the reactants or products.


a. 


b. 

c. 


d. 


19.37 Which compound in each pair has the lower pK_a? Which compound in each pair has the stronger conjugate base?

a. 


b. 


19.38 Rank the compounds in each group in order of increasing acidity.

a. 

b. 


19.39 Rank the compounds in each group in order of increasing basicity.

a. 

b. 

19.40 Match the pK_a values to the appropriate compound. pK_a values: 0.28, 1.24, 2.66, 2.86, and 3.12. Compounds: (a) FCH_2COOH; (b) CF_3COOH; (c) $F_2CHCOOH$; (d) ICH_2COOH; (e) $BrCH_2COOH$.

19.41 Although codeine occurs in low concentration in the opium poppy, most of the codeine used in medicine is prepared from morphine (the principal component of opium) by the following reaction. Explain why selective methylation occurs at only one OH in morphine to give codeine. Codeine is a less potent and less addictive analgesic than morphine.



morphine codeine

19.42 Which carboxylic acid has the lower pK_a, pyruvic acid (CH_3COCO_2H) or acetoacetic acid ($CH_3COCH_2CO_2H$)? Explain your choice.


19.43 Explain each statement.

a. The pK_a of *p*-nitrophenol is lower than the pK_a of phenol (7.2 vs. 10).
b. The pK_a of *p*-nitrophenol is lower than the pK_a of *m*-nitrophenol (7.2 vs. 8.3).

19.44 Explain the following statement. Although 2-methoxyacetic acid (CH_3OCH_2COOH) is a stronger acid than acetic acid (CH_3COOH), *p*-methoxybenzoic acid ($CH_3OC_6H_4COOH$) is a weaker acid than benzoic acid (C_6H_5COOH).


19.45 The pK_a of *p*-methylthiophenol ($CH_3SC_6H_4OH$) is 9.53. Is *p*-methylthiophenol more or less reactive in electrophilic aromatic substitution than phenol?

19.46 Explain why the pK_a of compound **A** is lower than the pK_a's of both compounds **B** and **C.**




A pK_a = 3.2 **B** pK_a = 3.9 **C** pK_a = 4.4

19.47 Rank the following compounds in order of increasing acidity and explain in detail your choice of order.



C **D** **E**


19.48 Phthalic acid and isophthalic acid have protons on two carboxy groups that can be removed with base. (a) Explain why the pK_a for loss of the first proton (pK_{a1}) is lower for phthalic acid than isophthalic acid. (b) Explain why the pK_a for loss of the second proton (pK_{a2}) is higher for phthalic acid than isophthalic acid.



phthalic acid
pK_{a1} = 2.9
pK_{a2} = 5.4


isophthalic acid
pK_{a1} = 3.7
pK_{a2} = 4.6

19.49 Explain the following result. Acetic acid (CH_3COOH), labeled at its OH oxygen with the uncommon ^{18}O isotope (shown in red), was treated with aqueous base, and then the solution was acidified. Two products having the ^{18}O label at different locations were formed.




19.50 Draw all resonance structures of the conjugate bases formed by removal of the labeled protons (H_a, H_b, and H_c) in cyclohexane-1,3-dione and acetanilide. For each compound, rank these protons in order of increasing acidity and explain the order you chose.

a.



cyclohexane-1,3-dione


b.



acetanilide

19.51 As we will see in Chapter 23, C–H bonds are sometimes more acidic than O–H bonds. Explain why the pK_a of $CH_2(CHO)_2$ is lower than the pK_a of $HO(CH_2)_3OH$ (9 vs. 16).

19.52 Identify **X** in the following equation, and explain how hexanoic acid (Section 19.2B) is formed by this stepwise reaction sequence.




hexanoic acid

19.53 The pK_a of acetamide (CH_3CONH_2) is 16. Draw the structure for its conjugate base and explain why acetamide is less acidic than CH_3COOH.

Extraction

19.54 Write out the steps needed to separate hydrocarbon **A** and carboxylic acid **B** by using an extraction procedure.



A **B**

19.55 Because phenol (C_6H_5OH) is less acidic than a carboxylic acid, it can be deprotonated by NaOH but not by the weaker base $NaHCO_3$. Using this information, write out an extraction sequence that can be used to separate C_6H_5OH, benzoic acid, and cyclohexanol. Show what compound is present in each layer at each stage of the process, and if it is present in its neutral or ionic form.

19.56 Can octane and octan-1-ol be separated using an aqueous extraction procedure? Explain why or why not.

Spectroscopy

19.57 Identify each compound from its spectral data.


a. Molecular formula: $C_3H_5ClO_2$
IR: 3500–2500 cm^{-1}, 1714 cm^{-1}
^1H NMR data: 2.87 (triplet, 2 H), 3.76 (triplet, 2 H), and 11.8 (singlet, 1 H) ppm


b. Molecular formula: $C_8H_8O_3$
IR: 3500–2500 cm^{-1}, 1688 cm^{-1}
^1H NMR data: 3.8 (singlet, 3 H), 7.0 (doublet, 2 H), 7.9 (doublet, 2 H), and 12.7 (singlet, 1 H) ppm

c. Molecular formula: $C_8H_8O_3$
IR: 3500–2500 cm^{-1}, 1710 cm^{-1}
^1H NMR data: 4.7 (singlet, 2 H), 6.9–7.3 (multiplet, 5 H), and 11.3 (singlet, 1 H) ppm


19.58 Use the ^{1}H NMR and IR spectra given below to identify the structures of two isomers (**A** and **B**) having molecular formula $C_4H_8O_2$.


Compound **A:**






Compound **B:**






19.59 An unknown compound **C** (molecular formula $C_4H_8O_3$) exhibits IR absorptions at 3600–2500 and 1734 cm^{-1}, as well as the following ^{1}H NMR spectrum. What is the structure of **C?**



19.60 Propose a structure for **D** (molecular formula $C_9H_9ClO_2$) consistent with the given spectroscopic data.

^{13}C NMR signals at 30, 36, 128, 130, 133, 139, and 179 ppm




19.61 Match the ^{13}C NMR data to the appropriate structure.

Spectrum [1]: signals at 14, 22, 27, 34, 181 ppm

Spectrum [2]: signals at 27, 39, 186 ppm


Spectrum [3]: signals at 22, 26, 43, 180 ppm



19.62 γ-Butyrolactone ($C_4H_6O_2$, GBL) is a biologically inactive compound that is converted to the biologically active recreational drug GHB (Section 19.5) by a lactonase enzyme in the body. Since γ-butyrolactone is more fat soluble than GHB, it is more readily absorbed by tissues and thus produces a faster onset of physiological symptoms. γ-Butyrolactone shows an absorption in its IR spectrum at 1770 cm^{-1} and the following ^1H NMR spectral data: 2.28 (multiplet, 2 H), 2.48 (triplet, 2 H), and 4.35 (triplet, 2 H) ppm. What is the structure of γ-butyrolactone?

Amino Acids


19.63 Threonine is a naturally occurring amino acid that has two stereogenic centers.



threonine

a. Draw the four possible stereoisomers using wedges and dashes.
b. The naturally occurring amino acid has the 2*S*,3*R* configuration at its two stereogenic centers. Which structure does this correspond to?

19.64 Proline is an unusual amino acid because its N atom on the α carbon is part of a five-membered ring.



proline

a. Draw both enantiomers of proline.
b. Draw proline in its zwitterionic form.


19.65 For each amino acid [$RCH(NH_2)COOH$], draw its neutral, positively charged, and negatively charged forms. Which form predominates at pH = 1, 6, and 11? What is the structure of each amino acid at its isoelectric point?

a. methionine (R = $CH_2CH_2SCH_3$)
b. serine (R = CH_2OH)


19.66 Calculate the isoelectric point for each amino acid.

a. cysteine: pK_a (COOH) = 2.05; pK_a (α-NH_3^+) = 10.25
b. methionine: pK_a (COOH) = 2.28; pK_a (α-NH_3^+) = 9.21

19.67 Lysine and tryptophan are two amino acids that contain an additional N atom in the R group bonded to the α carbon. While lysine is classified as a basic amino acid because it contains an additional basic N atom, tryptophan is classified as a neutral amino acid. Explain why this difference in classification occurs.




lysine



tryptophan

19.68 Glutamic acid is a naturally occurring α-amino acid that contains a carboxy group in its R group side chain (Table 19.4). (Glutamic acid is drawn in its neutral form with no charged atoms, a form that does not actually exist at any pH.)




glutamic acid


a. What form of glutamic acid exists at pH = 1?
b. If the pH is gradually increased, what form of glutamic acid exists after one equivalent of base is added? After two equivalents? After three equivalents?
c. Propose a structure of monosodium glutamate, the common flavor enhancer known as MSG.

Challenge Problems


19.69 Explain why using one or two equivalents of NaH results in different products in the following reactions.



19.70 Although *p*-hydroxybenzoic acid is less acidic than benzoic acid, *o*-hydroxybenzoic acid is slightly more acidic than benzoic acid. Explain this result.




p-hydroxybenzoic acid




o-hydroxybenzoic acid

19.71 2-Hydroxybutanedioic acid occurs naturally in apples and other fruits. Rank the labeled protons (H_a–H_e) in order of increasing acidity and explain in detail the order you chose.



2-hydroxybutanedioic acid


19.72 Although it was initially sold as a rat poison, warfarin is an effective anticoagulant used to prevent blood clots. Label the most acidic proton in warfarin, and explain why its pK_a is comparable to the pK_a of a carboxylic acid.



warfarin

20 Introduction to Carbonyl Chemistry; Organometallic Reagents; Oxidation and Reduction

20.1 Introduction
20.2 General reactions of carbonyl compounds
20.3 A preview of oxidation and reduction
20.4 Reduction of aldehydes and ketones
20.5 The stereochemistry of carbonyl reduction
20.6 Enantioselective carbonyl reductions
20.7 Reduction of carboxylic acids and their derivatives
20.8 Oxidation of aldehydes
20.9 Organometallic reagents
20.10 Reaction of organometallic reagents with aldehydes and ketones
20.11 Retrosynthetic analysis of Grignard products
20.12 Protecting groups
20.13 Reaction of organometallic reagents with carboxylic acid derivatives
20.14 Reaction of organometallic reagents with other compounds
20.15 α,β-Unsaturated carbonyl compounds
20.16 Summary—The reactions of organometallic reagents
20.17 Synthesis



Ciguatoxin CTX3C is a potent neurotoxin found in more than 400 species of warm-water fish. Thousands of people contract ciguatera seafood poisoning each year from ingesting tropical reef fish containing this neurotoxin. Interest in providing a practical supply of ciguatoxin CTX3C for biological testing led to its laboratory synthesis in 2001. One step in the multistep synthesis involved a selective reduction, one of the many reactions presented in Chapter 20.

Chapters 20 through 24 of this text discuss carbonyl compounds—aldehydes, ketones, acid halides, esters, amides, and carboxylic acids. **The carbonyl group is perhaps the most important functional group in organic chemistry,** because its electron-deficient carbon and easily broken π bond make it susceptible to a wide variety of useful reactions.

We begin by examining the similarities and differences between two broad classes of carbonyl compounds. We will then spend the remainder of Chapter 20 on reactions that are especially important in organic synthesis. Chapters 21 and 22 present specific reactions that occur at the carbonyl carbon, and Chapters 23 and 24 concentrate on reactions occurring at the α carbon to the carbonyl group.

Although Chapter 20 is "jam-packed" with reactions, most of them follow one of two general pathways, so they can be classified in a well-organized fashion, provided you remember a few basic principles. Keep in mind the following fundamental themes about reactions:

- Nucleophiles attack electrophiles.
- π Bonds are easily broken.
- Bonds to good leaving groups are easily cleaved.

20.1 Introduction

Two broad classes of compounds contain a ***carbonyl group:***




carbonyl group

[1] Compounds that have only carbon and hydrogen atoms bonded to the carbonyl group




aldehyde



ketone

- An **aldehyde** has at least one H atom bonded to the carbonyl group.
- A **ketone** has two alkyl or aryl groups bonded to the carbonyl group.


[2] Compounds that contain an electronegative atom bonded to the carbonyl group




carboxylic acid



acid chloride



ester



amide

These include **carboxylic acids, acid chlorides, esters,** and **amides,** as well as other similar compounds discussed in Chapter 22. Each of these compounds contains an atom (Cl, O, or N) more electronegative than carbon, capable of acting as a **leaving group.** Acid chlorides, esters, and amides are often called **carboxylic acid derivatives,** because they can be synthesized from carboxylic acids (Chapter 22). Since each compound contains an acyl group (RCO−), they are also called **acyl derivatives.**


- The presence or absence of a leaving group on the carbonyl carbon determines the type of reactions these compounds undergo (Section 20.2).

The carbonyl carbon atom is sp^2 hybridized and trigonal planar, and all bond angles are ~120°. The double bond of a carbonyl group consists of one σ bond and one π bond. The π bond is




The aldehyde α-sinensal (Problem 20.1) is the major compound responsible for the orange-like odor of mandarin oil, obtained from the mandarin tree in southern China.

formed by the overlap of two *p* orbitals, and extends above and below the plane. In these features the carbonyl group resembles the trigonal planar, sp^2 hybridized carbons of a C–C double bond.




In one important way, though, a C=O and C=C are very different. **The electronegative oxygen atom in the carbonyl group means that the bond is polarized, making the carbonyl carbon electron deficient.** Using a resonance description, the carbonyl group is represented by two resonance structures, with a charge-separated resonance structure a minor contributor to the hybrid. An electrostatic potential plot for formaldehyde, the simplest aldehyde, is shown in Figure 20.1. It clearly indicates the polarized carbonyl group.



the major contributor to the hybrid | a minor contributor to the hybrid | hybrid polarized carbonyl


Problem 20.1



α-sinensal

a. What orbitals are used to form the indicated bonds in α-sinensal?
b. In what type of orbitals do the lone pairs on O reside?


Figure 20.1
Electrostatic potential map of formaldehyde, $CH_2=O$



- An electrostatic potential map shows the electron-deficient carbon atom and the electron-rich oxygen atom of the carbonyl group.

20.2 General Reactions of Carbonyl Compounds


With what types of reagents should a carbonyl group react? The electronegative oxygen makes the carbonyl carbon electrophilic, and because it is trigonal planar, a carbonyl carbon is uncrowded. Moreover, a carbonyl group has an easily broken π bond.




uncrowded
sp^2 hybridized carbon

As a result, **carbonyl compounds react with nucleophiles.** The outcome of nucleophilic attack, however, depends on the identity of the carbonyl starting material.

- **Aldehydes and ketones undergo nucleophilic addition.**



- **Carbonyl compounds that contain leaving groups undergo nucleophilic substitution.**




Let's examine each of these general reactions individually.

20.2A Nucleophilic Addition to Aldehydes and Ketones

Aldehydes and ketones react with nucleophiles to form addition products by the two-step process shown in Mechanism 20.1: **nucleophilic attack** followed by **protonation.**

Mechanism 20.1 Nucleophilic Addition—A Two-Step Process



1. The nucleophile attacks the electrophilic carbonyl. The π bond is broken, moving an electron pair out on oxygen and forming an sp^3 hybridized carbon.
2. Protonation of the negatively charged oxygen by H_2O forms the addition product.

More examples of nucleophilic addition to aldehydes and ketones are discussed in Chapter 21.

The net result is that the π bond is broken, two new σ bonds are formed, and the elements of H and Nu are added across the π bond. Nucleophilic addition with two different nucleophiles—**hydride (H:⁻)** and **carbanions (R:⁻)**—is discussed in Chapter 20.

Aldehydes are more reactive than ketones towards nucleophilic attack for both steric and electronic reasons.



aldehyde
- less crowded
- less stable
- more reactive


ketone
- more crowded
- more stable
- less reactive

- **The two R groups bonded to the ketone carbonyl group make it more crowded, so nucleophilic attack is more difficult.**
- **The two electron-donor R groups stabilize the partial charge on the carbonyl carbon of a ketone, making it more stable and less reactive.**

20.2B Nucleophilic Substitution of RCOZ (Z = Leaving Group)

Carbonyl compounds with leaving groups react with nucleophiles to form substitution products by the two-step process shown in Mechanism 20.2: **nucleophilic attack,** followed by **loss of the leaving group.**

Mechanism 20.2 Nucleophilic Substitution—A Two-Step Process



Z = OH, Cl, OR', NH_2

1 The nucleophile attacks the electrophilic carbonyl. The π bond is broken, moving an electron pair out on oxygen and forming an sp^3 hybridized carbon.

2 An electron pair on oxygen re-forms the π bond and Z comes off as a leaving group with the electron pair in the C–Z bond.


The net result is that Nu replaces Z—a nucleophilic substitution reaction. This reaction is often called **nucleophilic *acyl* substitution** to distinguish it from the nucleophilic substitution reactions at sp^3 hybridized carbons discussed in Chapter 7. Nucleophilic substitution with two different nucleophiles—**hydride (H:⁻)** and **carbanions (R:⁻)**—is discussed in Chapter 20. Other nucleophiles are examined in Chapter 22.

Carboxylic acid derivatives differ greatly in their reactivity towards nucleophiles. The order in which they react parallels the leaving group ability of the group Z bonded to the carbonyl carbon.

- **The better the leaving group Z, the more reactive RCOZ is in nucleophilic acyl substitution.**

Recall from Section 7.7 that the weaker the base, the better the leaving group.


Thus, the following trends result:



- **Acid chlorides (RCOCl), which have the best leaving group (Cl^-), are the most reactive carboxylic acid derivatives, and amides ($RCONH_2$), which have the worst leaving group ($^-NH_2$), are the least reactive.**
- **Carboxylic acids (RCOOH) and esters (RCOOR'), which have leaving groups of similar basicity (^-OH and $^-OR'$), fall in the middle.**


Nucleophilic addition and nucleophilic acyl substitution involve the *same* first step—**nucleophilic attack on the electrophilic carbonyl group** to form a tetrahedral intermediate. The difference between them is what then happens to this intermediate. Aldehydes and ketones cannot undergo substitution because they have no leaving group bonded to the newly formed sp^3 hybridized carbon. Nucleophilic substitution with an aldehyde, for example, would form $H:^-$, an extremely strong base and therefore a very poor (and highly unlikely) leaving group.

An aldehyde does not undergo nucleophilic substitution...




...because $H:^-$ is a very poor leaving group.

Problem 20.2 Which carbonyl groups in the anticancer drug taxol (Section 5.5) will undergo nucleophilic addition and which will undergo nucleophilic substitution?



taxol

Problem 20.3 Which compound in each pair is more reactive towards nucleophilic attack?




To show how these general principles of nucleophilic substitution and addition apply to carbonyl compounds, we are going to discuss oxidation and reduction reactions, and reactions with organometallic reagents—compounds that contain carbon–metal bonds. We begin with reduction to build on what you learned previously in Chapter 12.

20.3 A Preview of Oxidation and Reduction

Recall the definitions of oxidation and reduction presented in Section 12.1:


- **Oxidation results in an increase in the number of C – Z bonds (usually C – O bonds) or a decrease in the number of C – H bonds.**
- **Reduction results in a decrease in the number of C – Z bonds (usually C – O bonds) or an increase in the number of C – H bonds.**

Carbonyl compounds are either reactants or products in many of these reactions, as illustrated in the accompanying diagram. For example, because aldehydes fall in the middle of this scheme, they can be both oxidized and reduced. Carboxylic acids and their derivatives (RCOZ), on the other hand, are already highly oxidized, so their only useful reaction is reduction.




The three most useful oxidation and reduction reactions of carbonyl starting materials can be summarized as follows:

[1] Reduction of aldehydes and ketones to alcohols (Sections 20.4–20.6)




Aldehydes and ketones are reduced to 1° and 2° alcohols, respectively.

[2] Reduction of carboxylic acid derivatives (Section 20.7)



The reduction of carboxylic acids and their derivatives gives a variety of products, depending on the identity of Z and the nature of the reducing agent. The usual products are aldehydes or 1° alcohols.

[3] Oxidation of aldehydes to carboxylic acids (Section 20.8)




The most useful oxidation reaction of carbonyl compounds is the oxidation of aldehydes to carboxylic acids.

We begin with reduction, because the mechanisms of reduction reactions follow directly from the general mechanisms for nucleophilic addition and substitution.

20.4 Reduction of Aldehydes and Ketones

$LiAlH_4$ and $NaBH_4$ serve as a source of $H:^-$, but there are no free $H:^-$ ions present in reactions with these reagents.


The most useful reagents for reducing aldehydes and ketones are the metal hydride reagents (Section 12.2). The two most common metal hydride reagents are **sodium borohydride ($NaBH_4$)** and **lithium aluminum hydride ($LiAlH_4$).** These reagents contain a polar metal–hydrogen bond that serves as a source of the nucleophile hydride, **$H:^-$.** $LiAlH_4$ is a stronger reducing agent than $NaBH_4$, because the Al−H bond is more polar than the B−H bond.



sodium borohydride | lithium aluminum hydride | a polar metal–hydrogen bond

20.4A Reduction with Metal Hydride Reagents

Treating an aldehyde or a ketone with $NaBH_4$ or $LiAlH_4$, followed by water or some other proton source, affords an **alcohol.** This is an addition reaction because **the elements of H_2 are added across the π bond**, but it is also a reduction because the product alcohol has fewer C−O bonds than the starting carbonyl compound.




R' = H or alkyl

aldehyde or ketone

1° or 2° alcohol

$LiAlH_4$ reductions must be carried out under anhydrous conditions, because water reacts violently with the reagent. Water is added to the reaction mixture (to serve as a proton source) *after* the reduction with $LiAlH_4$ is complete.


The product of this reduction reaction is a **1° alcohol** when the starting carbonyl compound is an aldehyde, and a **2° alcohol** when it is a ketone.





aldehyde | 1° alcohol | ketone | 2° alcohol

$NaBH_4$ selectively reduces aldehydes and ketones in the presence of most other functional groups. Reductions with $NaBH_4$ are typically carried out in CH_3OH as solvent. $LiAlH_4$ reduces aldehydes and ketones and many other functional groups as well (Sections 12.6 and 20.7).


Problem 20.4 What alcohol is formed when each compound is treated with $NaBH_4$ in CH_3OH?


a. 


b. 

c. 

Problem 20.5 What aldehyde or ketone is needed to prepare each alcohol by metal hydride reduction?

a. 

b. 


c. 

Problem 20.6 Why can't 1-methylcyclohexanol be prepared from a carbonyl compound by reduction?

20.4B The Mechanism of Hydride Reduction

Hydride reduction of aldehydes and ketones occurs via the general mechanism of nucleophilic addition—that is, **nucleophilic attack** followed by **protonation.** Mechanism 20.3 is shown using $LiAlH_4$, but an analogous mechanism can be written for $NaBH_4$.

Mechanism 20.3 $LiAlH_4$ Reduction of RCHO and $R_2C{=}O$




1 The nucleophile (AlH_4^-) donates $H:^-$ to the carbonyl group, breaking the π bond and moving an electron pair out on oxygen. This forms a new C – H bond.

2 Protonation of the negatively charged oxygen by H_2O (or CH_3OH) forms the reduction product with a new O – H bond.

- **The net result of adding $H:^-$ (from $NaBH_4$ or $LiAlH_4$) and H^+ (from H_2O) is the addition of the elements of H_2 to the carbonyl π bond.**

20.4C Catalytic Hydrogenation of Aldehydes and Ketones


Catalytic hydrogenation also reduces aldehydes and ketones to 1° and 2° alcohols, respectively, using H_2 and Pd-C (or another metal catalyst). H_2 adds to the C=O in much the same way that it adds to the C=C of an alkene (Section 12.3). The metal catalyst (Pd-C) provides a surface that binds the carbonyl starting material and H_2, and two H atoms are sequentially transferred with cleavage of the π bond.



When a compound contains both a carbonyl group and a carbon–carbon double bond, selective reduction of one functional group can be achieved by proper choice of reagent.

- **A C=C is reduced faster than a C=O with H_2 (Pd-C).**
- **A C=O is readily reduced with $NaBH_4$ and $LiAlH_4$, but a C=C is inert.**

Thus, cyclohex-2-enone, a compound that contains both a carbon–carbon double bond and a carbonyl group, can be reduced to three different compounds—an allylic alcohol, a carbonyl compound, or an alcohol—depending on the reagent.




Problem 20.7 Draw the products formed when $CH_3COCH_2CH_2CH{=}CH_2$ is treated with each reagent: (a) $LiAlH_4$, then H_2O; (b) $NaBH_4$ in CH_3OH; (c) H_2 (1 equiv), Pd-C; (d) H_2 (excess), Pd-C; (e) $NaBH_4$ (excess) in CH_3OH; (f) $NaBD_4$ in CH_3OH.

The reduction of aldehydes and ketones is a common reaction used in the synthesis of many useful natural products. Two examples are shown in Figure 20.2.

Figure 20.2
$NaBH_4$ reductions used in organic synthesis



The male musk deer, a small antlerless deer found in the mountain regions of China and Tibet, has long been hunted for its musk, a strongly scented liquid used in early medicine and later in perfumery.




- **Muscone** is the major compound in musk, one of the oldest known ingredients in perfumes. Musk was originally isolated from the male musk deer, but it can now be prepared synthetically in the laboratory in a variety of ways.

20.5 The Stereochemistry of Carbonyl Reduction


Recall from Section 9.15 that an achiral starting material gives a racemic mixture when a new stereogenic center is formed.

The stereochemistry of carbonyl reduction follows the same principles we have previously learned. Reduction converts a **planar sp^2 hybridized carbonyl carbon to a tetrahedral sp^3 hybridized carbon.** What happens when a new stereogenic center is formed in this process? With an achiral reagent like $NaBH_4$ or $LiAlH_4$, a racemic product is obtained. For example, $NaBH_4$ in CH_3OH solution reduces butan-2-one, an achiral ketone, to butan-2-ol, an alcohol that contains a new stereogenic center. Both enantiomers of butan-2-ol are formed in equal amounts.




Why is a racemic mixture formed? Because the carbonyl carbon is sp^2 hybridized and planar, hydride can approach the double bond with equal probability from both sides of the plane,

forming two alkoxides, which are **enantiomers** of each other. Protonation of the alkoxides gives an equal amount of two alcohols, which are also **enantiomers.**



- **Conclusion: Hydride reduction of an achiral ketone with $LiAlH_4$ or $NaBH_4$ gives a racemic mixture of two alcohols when a new stereogenic center is formed.**

Problem 20.8 Draw the products formed (including stereoisomers) when each compound is reduced with $NaBH_4$ in CH_3OH.



20.6 Enantioselective Carbonyl Reductions

20.6A CBS Reagents

One enantiomer can be formed selectively from the reduction of a carbonyl group, provided a **chiral reducing agent** is used. This strategy is identical to that employed in the Sharpless asymmetric epoxidation reaction (Section 12.15). A reduction that forms one enantiomer predominantly or exclusively is an **enantioselective** or **asymmetric reduction.**


Many different chiral reducing agents have now been prepared for this purpose. One such reagent, formed by reacting borane ($\mathbf{BH_3}$) with a heterocycle called an **oxazaborolidine,** has one stereogenic center (and thus two enantiomers).



(*S*)-CBS reagent **(*R*)-CBS reagent**

These reagents are called the **(*S*)-CBS reagent** and the **(*R*)-CBS reagent,** named for *C*orey, *B*akshi, and *S*hibata, the chemists who developed these versatile reagents. One B−H bond of BH_3 serves as the source of hydride in this reduction. The stereochemistry of the new stereogenic center in the product is often predictable. For ketones having the general structure C_6H_5COR, draw the starting material with the aryl group on the left side of the carbonyl, as shown with acetophenone. Then, to draw the product, keep in mind:

- The (*S*)-CBS reagent delivers hydride (H:⁻) from the *front* side of the C=O. This generally affords the *R* alcohol as the major product.
- The (*R*)-CBS reagent delivers hydride (H:⁻) from the *back* side of the C=O. This generally affords the *S* alcohol as the major product.



These reagents are highly enantioselective. Treatment of propiophenone with the (***S***)-CBS reagent forms the ***R*** alcohol in 97% enantiomeric excess (*ee*). Enantioselective reductions are key steps in the synthesis of several widely used drugs, including salmeterol, a long-acting bronchodilator shown in Figure 20.3. This new technology provides access to single enantiomers of biologically active compounds, often previously available only as a racemic mixture.





Figure 20.3
Enantioselective reduction—A key step in the synthesis of salmeterol




- (*R*)-Salmeterol is a long-acting bronchodilator used for the treatment of asthma.
- In this example, the (*R*)-CBS reagent adds the new H atom from behind, the same result observed with acetophenone and propiophenone. In this case, however, alcohol **A** has the *R* configuration using the rules for assigning priority in Chapter 5.



Ezetimibe is sold as a single medication under the trade name of Zetia, or in combination with simvastatin, another cholesterol-lowering medication, and marketed as Vytorin. These drugs are prescribed for individuals who cannot tolerate or derive no benefit from other cholesterol-lowering medications.


Problem 20.9 What carbonyl compound and CBS reagent are needed to prepare **X,** an intermediate in the synthesis of ezetimibe (trade name Zetia), a drug that lowers cholesterol levels by inhibiting its absorption in the intestines.



X ezetimibe


20.6B Enantioselective Biological Reduction

Although laboratory reduction reactions often do not proceed with 100% enantioselectivity, biological reductions that occur in cells *always* proceed with complete selectivity, forming a single enantiomer. In cells, the reducing agent is **NADH,** nicotinamide adenine dinucleotide (reduced form), the coenzyme introduced in Section 12.14.




nicotinamide adenine dinucleotide
(reduced form)
NADH

In biological reduction, NADH donates H:⁻, in much the same way as a metal hydride reagent. Nucleophilic attack of hydride and protonation thus form an alcohol from a carbonyl group, and NADH is converted to NAD^+.



Pyruvic acid is formed during the metabolism of glucose. During periods of strenuous exercise, when there is insufficient oxygen to metabolize pyruvic acid to CO_2, pyruvic acid is reduced to lactic acid. The tired feeling of sore muscles is a result of lactic acid accumulation.


This reaction is completely enantioselective. Reduction of pyruvic acid with NADH catalyzed by lactate dehydrogenase affords a single enantiomer of lactic acid with the *S* configuration. NADH reduces a variety of different carbonyl compounds in biological systems. The configuration of the product (*R* or *S*) depends on the enzyme used to catalyze the process.





Niacin can be obtained from foods such as soybeans, which contain it naturally, and from breakfast cereals, which are fortified with it to help people consume their recommended daily allowance of this B vitamin.

As we learned in Section 12.4, **NAD^+, the oxidized form of NADH, is a biological oxidizing agent** capable of oxidizing alcohols to carbonyl compounds, forming NADH in the process. NAD^+ is synthesized from the vitamin niacin, which can be obtained from soybeans among other dietary sources.



niacin
vitamin B_3


20.7 Reduction of Carboxylic Acids and Their Derivatives

The reduction of carboxylic acids and their derivatives (RCOZ) is complicated because the products obtained depend on the identity of both the leaving group (Z) and the reducing agent. Metal hydride reagents are the most useful reducing reagents. **Lithium aluminum hydride is a strong reducing agent that reacts with *all* carboxylic acid derivatives.** Two other related but more selective reducing agents are also used.

[1] **Diisobutylaluminum hydride, $[(CH_3)_2CHCH_2]_2AlH$,** abbreviated as **DIBAL-H,** has two bulky isobutyl groups, which make this reagent less reactive than $LiAlH_4$.

[2] **Lithium tri-*tert*-butoxyaluminum hydride, $LiAlH[OC(CH_3)_3]_3$,** has three electronegative oxygen atoms bonded to aluminum, which make this reagent less nucleophilic than $LiAlH_4$.


$LiAlH_4$ is a strong, nonselective reducing agent. DIBAL-H and $LiAlH[OC(CH_3)_3]_3$ are milder, more selective reducing agents.




In both reagents, the single H atom bonded to Al is donated as $H:^-$ in hydride reductions.

20.7A Reduction of Acid Chlorides and Esters


Acid chlorides and esters can be reduced to either aldehydes or alcohols, depending on the reagent.



- $LiAlH_4$ converts RCOCl and RCOOR' to alcohols.
- A milder reducing agent (DIBAL-H or $LiAlH[OC(CH_3)_3]_3$) converts RCOCl or RCOOR' to RCHO at low temperatures.




In the reduction of an acid chloride, Cl^- comes off as the leaving group.



In the reduction of the ester, CH_3O^- comes off as the leaving group, which is then protonated by H_2O to form CH_3OH.

Mechanism 20.4 illustrates why two different products are possible. It can be conceptually divided into two parts: **nucleophilic substitution** to form an aldehyde (Steps [1] and [2]), followed by **nucleophilic addition** to the aldehyde to form an alcohol (Steps [3] and [4]). A general mechanism is drawn using $LiAlH_4$ as reducing agent.


Mechanism 20.4 Reduction of RCOCl and RCOOR' with a Metal Hydride Reagent




1. Nucleophilic attack of $H:^-$ forms a tetrahedral intermediate with a leaving group Z.
2. The π bond is re-formed and the leaving group Z departs. The overall result of addition of $H:^-$ and elimination of $Z:^-$ is substitution of H for Z.
3. Nucleophilic attack of $H:^-$ forms an alkoxide with no leaving group.
4. Protonation of the alkoxide by H_2O forms the alcohol reduction product. The overall result of Steps [3] and [4] is addition of H_2.

With less nucleophilic reducing agents such as DIBAL-H and $LiAlH[OC(CH_3)_3]_3$, the process stops after reaction with one equivalent of $H:^-$ and the aldehyde is formed as product. With a stronger reducing agent like $LiAlH_4$, two equivalents of $H:^-$ are added and an alcohol is formed.

Problem 20.10 Draw a stepwise mechanism for the following reaction.




Problem 20.11 Draw the structure of both an acid chloride and an ester that can be used to prepare each compound by reduction.



Selective reductions are routinely used in the synthesis of highly complex natural products such as **ciguatoxin CTX3C,** a potent neurotoxin introduced in the chapter opener. One reaction in a synthesis of ciguatoxin CTX3C involved the reduction of an ester to an aldehyde using DIBAL-H, as shown in Figure 20.4.


Figure 20.4
The DIBAL-H reduction of an ester to an aldehyde in the synthesis of the marine neurotoxin ciguatoxin CTX3C




- One step in a lengthy synthesis of ciguatoxin CTX3C involved selective reduction of an ester to an aldehyde using DIBAL-H.

20.7B Reduction of Carboxylic Acids and Amides


Carboxylic acids are reduced to alcohols with $LiAlH_4$. $LiAlH_4$ is too strong a reducing agent to stop the reaction at the aldehyde stage, but milder reagents are not strong enough to initiate the reaction in the first place, so this is the only useful reduction reaction of carboxylic acids.



Unlike the $LiAlH_4$ reduction of all other carboxylic acid derivatives, which affords alcohols, the **$LiAlH_4$ reduction of amides forms amines.**



Both C–O bonds are reduced to C–H bonds by $LiAlH_4$, and any H atom or R group bonded to the amide nitrogen atom remains bonded to it in the product. Because $^-NH_2$ (or ^-NHR or $^-NR_2$) is a poorer leaving group than Cl^- or ^-OR, $^-NH_2$ is never lost during reduction, and therefore it forms an amine in the final product.




Imines and related compounds are discussed in Chapter 21.

The mechanism, illustrated in Mechanism 20.5 with $RCONH_2$ as starting material, is somewhat different than the previous reductions of carboxylic acid derivatives. Amide reduction proceeds

Mechanism 20.5 Reduction of an Amide to an Amine with $LiAlH_4$


Part [1] Reduction of an amide to an imine



1 – 2 AlH_4^- removes a proton from the amide to form a Lewis base that complexes with AlH_3 in Step [2].

3 – 4 Nucleophilic attack of $H:^-$ and loss of a leaving group, $(OAlH_3)^{2-}$, form an imine.


Part [2] Reduction of an imine to an amine





5 – 6 Nucleophilic addition of $H:^-$ and protonation form the amine.


with formation of an intermediate ***imine,*** **a compound containing a C–N double bond,** which is then further reduced to an amine.

Problem 20.12 Draw the products formed from $LiAlH_4$ reduction of each compound.


a. 


b. 


c. 

d. 

Problem 20.13 What amide(s) will form each of the following amines on treatment with $LiAlH_4$?

a. 


b. 


c. 

20.7C A Summary of the Reagents for Reduction

The many available metal hydride reagents reduce a wide variety of functional groups. Keep in mind that $LiAlH_4$ is such a strong reducing agent that it *nonselectively* reduces most polar functional groups. All other metal hydride reagents are more selective, and each has its particular reactions that best utilize its reduced reactivity. The reagents and their uses are summarized in Table 20.1.

Problem 20.14 What product is formed when each compound is treated with either $LiAlH_4$ (followed by H_2O), or $NaBH_4$ in CH_3OH?

a. 

b. 



c. 

Table 20.1 A Summary of Metal Hydride Reducing Agents

	Reagent	Starting material	→	Product
Strong reagent	$LiAlH_4$	RCHO	→	RCH_2OH
		R_2CO	→	R_2CHOH
		RCOOH	→	RCH_2OH
		RCOOR'	→	RCH_2OH
		RCOCl	→	RCH_2OH
		$RCONH_2$	→	RCH_2NH_2
Milder reagents	$NaBH_4$	RCHO	→	RCH_2OH
		R_2CO	→	R_2CHOH
	$LiAlH[OC(CH_3)_3]_3$	RCOCl	→	RCHO
	DIBAL-H	RCOOR'	→	RCHO




Aldehydes give a positive Tollens test; that is, they react with Ag^+ to form RCOOH and Ag. When the reaction is carried out in a glass flask, a silver mirror is formed on its walls. Other functional groups give a negative Tollens test, because no silver mirror forms.

20.8 Oxidation of Aldehydes


The most common oxidation reaction of carbonyl compounds is the oxidation of **aldehydes to carboxylic acids.** A variety of oxidizing agents can be used, including CrO_3, $Na_2Cr_2O_7$, $K_2Cr_2O_7$, and $KMnO_4$. Cr^{6+} reagents are also used to oxidize 1° and 2° alcohols, as discussed in Section 12.12. Because ketones have no H on the carbonyl carbon, they do not undergo this oxidation reaction.

Aldehydes are oxidized selectively in the presence of other functional groups using **silver(I) oxide in aqueous ammonium hydroxide (Ag_2O in NH_4OH).** This is called **Tollens reagent.** Oxidation with Tollens reagent provides a distinct color change, because the Ag^+ reagent is reduced to silver metal (Ag), which precipitates out of solution.




Only the aldehyde is oxidized.

Problem 20.15 What product is formed when each compound is treated with either Ag_2O, NH_4OH or $Na_2Cr_2O_7$, H_2SO_4, H_2O?



Problem 20.16 Review the oxidation reactions using Cr^{6+} reagents in Section 12.12. Then, draw the product formed when compound **B** is treated with each reagent.




B

a. $NaBH_4$, CH_3OH
b. [1] $LiAlH_4$; [2] H_2O
c. PCC
d. Ag_2O, NH_4OH
e. CrO_3, H_2SO_4, H_2O

20.9 Organometallic Reagents

We will now discuss the reactions of carbonyl compounds with organometallic reagents, another class of nucleophiles.

- *Organometallic reagents* **contain a carbon atom bonded to a metal.**



organometallic reagent

= R—M

M = metal

Most common metals:
M = Li, Mg, Cu

Lithium, magnesium, and copper are the most commonly used metals in organometallic reagents, but others (such as Sn, Si, Tl, Al, Ti, and Hg) are known. General structures of the three common organometallic reagents are shown. R can be alkyl, aryl, allyl, benzyl, sp^2 hybridized, and with

M = Li or Mg, *sp* hybridized. Because metals are *more electropositive* (less electronegative) than carbon, they donate electron density towards carbon, so that **carbon bears a partial negative charge.**

R—Li	R—Mg—X	R—Cu⁻ Li⁺ (R above Cu)
organolithium reagents	organomagnesium reagents or Grignard reagents	organocopper reagents or organocuprates


- **The more polar the carbon–metal bond, the more reactive the organometallic reagent.**

Because both Li and Mg are very electropositive metals, **organolithium (RLi)** and **organomagnesium reagents (RMgX)** contain very polar carbon–metal bonds and are therefore *very reactive* reagents. Organomagnesium reagents are called **Grignard reagents,** after Victor Grignard, who received the Nobel Prize in Chemistry in 1912 for his work with them.

Electronegativity values for carbon and the common metals in R – M reagents are C (2.5), Li (1.0), Mg (1.3), and Cu (1.8).

Organocopper reagents (R_2CuLi), also called **organocuprates,** have a less polar carbon–metal bond and are therefore *less reactive*. Although organocuprates contain two alkyl groups bonded to copper, only one R group is utilized in a reaction.

Regardless of the metal, organometallic reagents are useful synthetically because they react as if they were free carbanions; that is, carbon bears a partial *negative* charge, so the **reagents react as bases and nucleophiles.**




carbanion


a base and a nucleophile

20.9A Preparation of Organometallic Reagents

Organolithium and Grignard reagents are typically prepared by reaction of an organic halide with the corresponding metal, as shown in the accompanying equations.





With lithium, the halogen and metal exchange to form the organolithium reagent. With magnesium, the metal inserts in the carbon–halogen bond, forming the Grignard reagent. Grignard reagents are usually prepared in diethyl ether ($CH_3CH_2OCH_2CH_3$) as solvent. It is thought that two ether oxygen atoms complex with the magnesium atom, stabilizing the reagent.



Two molecules of diethyl ether complex with the Mg atom of the Grignard reagent.

Organocuprates are prepared from organolithium reagents by reaction with a Cu^+ salt, often CuI.







Problem 20.17 Write the step(s) needed to convert CH_3CH_2Br to each reagent: (a) CH_3CH_2Li; (b) CH_3CH_2MgBr; (c) $(CH_3CH_2)_2CuLi$.

20.9B Acetylide Anions

The **acetylide anions** discussed in Chapter 11 are another example of organometallic compounds. These reagents are prepared by an acid–base reaction of an alkyne with a base such as $NaNH_2$ or NaH. We can think of these compounds as **organosodium** reagents. Because sodium is even more electropositive (less electronegative) than lithium, the C−Na bond of these organosodium compounds is best described as **ionic,** rather than polar covalent.



An acid–base reaction can also be used to prepare *sp* hybridized organolithium compounds. Treatment of a terminal alkyne with CH_3Li affords a lithium acetylide. Equilibrium favors the products because the *sp* hybridized C−H bond of the terminal alkyne is more acidic than the sp^3 hybridized conjugate acid, CH_4, that is formed.




Problem 20.18 Oct-1-yne ($HC{\equiv}CCH_2CH_2CH_2CH_2CH_2CH_3$) reacts rapidly with NaH, forming a gas that bubbles out of the reaction mixture, as one product. Oct-1-yne also reacts rapidly with CH_3MgBr, and a different gas is produced. Write balanced equations for both reactions and identify the gases formed.

Problem 20.19 Which of the following species represent organometallic compounds:
(a) $BrMgC{\equiv}CCH_2CH_3$; (b) $NaOCH_2CH_3$; (c) $KOC(CH_3)_3$; (d) PhLi?


20.9C Reaction as a Base

- **Organometallic reagents are strong bases that readily abstract a proton from water to form hydrocarbons.**


The electron pair in the carbon–metal bond is used to form a new bond to the proton. Equilibrium favors the products of this acid–base reaction because H_2O is a much stronger acid than the alkane product.




Similar reactions occur for the same reason with the O–H proton in alcohols and carboxylic acids, and the N–H protons of amines.



Because organolithium and Grignard reagents are themselves prepared from alkyl halides, a two-step method converts an alkyl halide into an alkane (or another hydrocarbon).




Problem 20.20 Draw the product formed when each organometallic reagent is treated with H_2O.



20.9D Reaction as a Nucleophile


Organometallic reagents are also strong nucleophiles that react with electrophilic carbon atoms to form new carbon–carbon bonds. These reactions are very valuable in forming the carbon skeletons of complex organic molecules. The following reactions of organometallic reagents are examined in Sections 20.10, 20.13, and 20.14:

[1] Reaction of R–M with aldehydes and ketones to afford alcohols (Section 20.10)




Aldehydes and ketones are converted to 1°, 2°, or 3° alcohols with R"Li or R"MgX.

[2] Reaction of R–M with carboxylic acid derivatives (Section 20.13)



Acid chlorides and esters can be converted to ketones or 3° alcohols with organometallic reagents. The identity of the product depends on the identity of R"–M and the leaving group Z.


[3] Reaction of R – M with other electrophilic functional groups (Section 20.14)



Organometallic reagents also react with CO_2 to form carboxylic acids and with epoxides to form alcohols.

20.10 Reaction of Organometallic Reagents with Aldehydes and Ketones


Treatment of an aldehyde or ketone with either an organolithium or Grignard reagent followed by water forms an alcohol with a new carbon–carbon bond. This reaction is an **addition reaction** because the elements of R" and H are added across the π bond.



20.10A General Features

This reaction follows the general mechanism for nucleophilic addition (Section 20.2A)—that is, **nucleophilic attack** by a carbanion followed by **protonation.** Mechanism 20.6 is shown using R"MgX, but the same steps occur with organolithium reagents and acetylide anions.


Mechanism 20.6 Nucleophilic Addition of R"MgX to RCHO and $R_2C{=}O$



1 The nucleophile $(R")^-$ attacks the carbonyl group, breaking the π bond and yielding an alkoxide. This forms a new carbon–carbon bond.


2 Protonation of the alkoxide by H_2O forms the addition product with a new O – H bond. The overall result is addition of R" and H to the carbonyl group.

This reaction is used to prepare 1°, 2°, and 3° alcohols, depending on the number of alkyl groups bonded to the carbonyl carbon of the aldehyde or ketone.




[1] Addition of R"MgX to formaldehyde ($CH_2=O$) forms a 1° alcohol.
[2] Addition of R"MgX to all other aldehydes forms a 2° alcohol.
[3] Addition of R"MgX to ketones forms a 3° alcohol.

Each reaction results in addition of one new alkyl group to the carbonyl carbon, and forms one new carbon–carbon bond. The reaction is general for all organolithium and Grignard reagents, and works for acetylide anions as well, as illustrated in Equations [1]–[3].



Because organometallic reagents are strong bases that rapidly react with H_2O (Section 20.9C), the addition of the new alkyl group must be carried out under anhydrous conditions to prevent traces of water from reacting with the reagent, thus reducing the yield of the desired alcohol. Water is added after the addition to protonate the alkoxide.


Problem 20.21 Draw the product of each reaction.



20.10B Stereochemistry


Like reduction, addition of organometallic reagents converts an sp^2 hybridized carbonyl carbon to a tetrahedral sp^3 hybridized carbon. Addition of R−M always occurs from both sides of the trigonal planar carbonyl group. **When a new stereogenic center is formed from an achiral starting material, an equal mixture of enantiomers results,** as shown in Sample Problem 20.1.

Sample Problem 20.1 Draw all stereoisomers formed in the following reaction.




Solution

The Grignard reagent adds from both sides of the trigonal planar carbonyl group, forming two alkoxides, each containing a new stereogenic center labeled in blue. Protonation with water yields **an equal amount of two enantiomers—a racemic mixture.**




Problem 20.22 Draw the products (including stereochemistry) of the following reactions.



20.10C Applications in Synthesis


Many syntheses of useful compounds utilize the nucleophilic addition of a Grignard or organolithium reagent to form carbon–carbon bonds. For example, a key step in the synthesis of ethynylestradiol (Section 11.4), an oral contraceptive component, is the addition of lithium acetylide to a ketone, as shown in Figure 20.5.

Figure. 20.5
The synthesis of ethynylestradiol



The synthesis of C_{18} juvenile hormone, a member of a group of structurally related molecules that regulate the complex life cycle of an insect, is another example. The last steps of the synthesis are outlined in Figure 20.6.

Figure 20.6
C_{18} juvenile hormone






Juvenile hormones regulate the life cycle of the cecropia moth.


- Addition of CH_3MgX to ketone **A** gives an alkoxide, **B,** which is protonated with H_2O to form 3° alcohol **C.** Although the ester group ($-COOCH_3$) can also react with the Grignard reagent (Section 20.13), it is less reactive than the ketone carbonyl. Thus, with control of reaction conditions, nucleophilic addition occurs selectively at the ketone.
- Treatment of halohydrin **C** with K_2CO_3 forms the C_{18} juvenile hormone in one step. Conversion of a halohydrin to an epoxide was discussed in Section 9.6.

Juvenile hormones maintain the juvenile stage of an insect until it is ready for adulthood. This property has been exploited to control mosquitoes and other insects infecting livestock and crops. Although juvenile hormone itself is too unstable in light and too expensive to synthesize for use in controlling insect populations, related compounds, called **juvenile hormone *mimics,*** have been used effectively. Application of these synthetic hormones to an egg or larva of an insect prevents maturation. With no sexually mature adults to propagate the next generation, the insect population is reduced. The best known example of a synthetic juvenile hormone is called **methoprene,** sold under such trade names as Altocid, Precor, and Diacon. Methoprene is used in cattle salt blocks to control hornflies, in stored tobacco to control pests, and on dogs and cats to control fleas.



methoprene
juvenile hormone mimic

=





20.11 Retrosynthetic Analysis of Grignard Products

To use the Grignard addition in synthesis, you must be able to determine what carbonyl and Grignard components are needed to prepare a given compound—that is, **you must work backwards, in the retrosynthetic direction.** This involves a two-step process:

Step [1] Find the carbon bonded to the OH group in the product.

Step [2] Break the molecule into two components: One alkyl group bonded to the carbon with the OH group comes from the organometallic reagent. The rest of the molecule comes from the carbonyl component.






pentan-3-ol


To synthesize pentan-3-ol [$(CH_3CH_2)_2CHOH$] by a Grignard reaction, locate the carbon bonded to the OH group, and then break the molecule into two components at this carbon. Thus, retrosynthetic analysis shows that one of the ethyl groups on this carbon comes from a Grignard reagent (CH_3CH_2MgX), and the rest of the molecule comes from the carbonyl component, a three-carbon aldehyde.

Retrosynthetic analysis




Then, writing the reaction in the synthetic direction—that is, from starting material to product—shows whether the analysis is correct. In this example, a three-carbon aldehyde reacts with CH_3CH_2MgBr to form an alkoxide, which can then be protonated by H_2O to form pentan-3-ol, the desired alcohol.

In the synthetic direction:



There is often more than one way to synthesize a 2° alcohol by Grignard addition, as shown in Sample Problem 20.2.

Sample Problem 20.2 Show two different methods to synthesize butan-2-ol using a Grignard reaction.



butan-2-ol

Solution


Because butan-2-ol has two different alkyl groups bonded to the carbon bearing the OH group, there are two different ways to form a new carbon–carbon bond by Grignard addition.

Possibility [1] Use CH_3MgX and a three-carbon aldehyde.


Possibility [2] Use CH_3CH_2MgX and a two-carbon aldehyde.




Both methods give the desired product, butan-2-ol, as can be seen by writing the reactions from starting material to product.




Problem 20.23 What Grignard reagent and carbonyl compound are needed to prepare each alcohol? As shown in part (d), 3° alcohols with three different R groups on the carbon bonded to the OH group can be prepared by three different Grignard reactions.



Problem 20.24 Linalool (the Chapter 9 opening molecule) and lavandulol are two of the major components of lavender oil. (a) What organolithium reagent and carbonyl compound can be used to make each alcohol? (b) How might lavandulol be formed by reduction of a carbonyl compound? (c) Why can't linalool be prepared by a similar pathway?



Problem 20.25 What Grignard reagent and carbonyl compound can be used to prepare the antidepressant venlafaxine (trade name Effexor)?



venlafaxine


20.12 Protecting Groups


Although the addition of organometallic reagents to carbonyls is a very versatile reaction, it cannot be used with molecules that contain both a carbonyl group and N−H or O−H bonds.

Rapid acid–base reactions occur between organometallic reagents and all of the following functional groups: ROH, RCOOH, RNH_2, R_2NH, $RCONH_2$, RCONHR, and RSH.

- **Carbonyl compounds that also contain N−H or O−H bonds undergo an acid–base reaction with organometallic reagents, not nucleophilic addition.**

Suppose, for example, that you wanted to add methylmagnesium chloride (CH_3MgCl) to the carbonyl group of 5-hydroxypentan-2-one to form a diol. Nucleophilic addition will *not* occur with this substrate. Instead, **because Grignard reagents are strong bases and proton transfer reactions are fast, CH_3MgCl removes the O−H proton before nucleophilic addition takes place.** The stronger acid and base react to form the weaker conjugate acid and conjugate base, as we learned in Section 20.9C.





products of proton transfer

Solving this problem requires a three-step strategy:


Step [1] Convert the OH group into another functional group that does not interfere with the desired reaction. This new blocking group is called a **protecting group,** and the reaction that creates it is called ***protection.***

Step [2] Carry out the desired reaction.

Step [3] Remove the protecting group. This reaction is called ***deprotection.***


Application of the general strategy to the Grignard addition of CH_3MgCl to 5-hydroxypentan-2-one is illustrated in Figure 20.7.

Figure 20.7
General strategy for using a protecting group




- In Step [1], the OH proton in 5-hydroxypentan-2-one is replaced with a protecting group, written as **PG.** Because the product of Step [1] no longer has an OH proton, it can now undergo nucleophilic addition.
- In Step [2], CH_3MgCl adds to the carbonyl group to yield a 3° alcohol after protonation with water.
- Removal of the protecting group in Step [3] forms the desired product, 4-methylpentane-1,4-diol.

A common OH protecting group is a **silyl ether.** A silyl ether has a new O−Si bond in place of the O−H bond of the alcohol. The most widely used silyl ether protecting group is the ***tert*-butyldimethylsilyl ether,** abbreviated as **TBDMS.**




tert-Butyldimethylsilyl ethers are prepared from alcohols by reaction with *tert*-butyldimethylsilyl chloride and an amine base, usually imidazole.



imidazole



The silyl ether is typically removed with a fluoride salt, usually **tetrabutylammonium fluoride** $(CH_3CH_2CH_2CH_2)_4N^+\ F^-$, drawn as $Bu_4N^+F^-$ (Bu = butyl).




The use of a *tert*-butyldimethylsilyl ether as a protecting group makes possible the synthesis of 4-methylpentane-1,4-diol by a three-step sequence.



- **Step [1] Protect the OH group** as a *tert*-butyldimethylsilyl ether by reaction with *tert*-butyldimethylsilyl chloride and imidazole.
- **Step [2] Carry out nucleophilic addition** by using CH_3MgCl, followed by protonation.
- **Step [3] Remove the protecting group** with tetrabutylammonium fluoride to form the desired addition product.

Protecting groups block interfering functional groups, and in this way, a wider variety of reactions can take place with a particular substrate. For more on protecting groups, see the discussion of acetals in Section 21.15.


Problem 20.26 Using protecting groups, show how estrone can be converted to ethynylestradiol, a widely used oral contraceptive.



estrone ethynylestradiol

20.13 Reaction of Organometallic Reagents with Carboxylic Acid Derivatives

Organometallic reagents react with carboxylic acid derivatives (RCOZ) to form two different products, depending on the identity of both the leaving group Z and the reagent R−M. The most useful reactions are carried out with esters and acid chlorides, forming either **ketones** or **3° alcohols.**




Z = Cl or OR' ketone 3° alcohol


- Keep in mind that RLi and RMgX are very reactive reagents, whereas R_2CuLi is much less reactive. This reactivity difference makes selective reactions possible.

20.13A Reaction of RLi and RMgX with Esters and Acid Chlorides


Both esters and acid chlorides form 3° alcohols when treated with two equivalents of either Grignard or organolithium reagents. Two new carbon–carbon bonds are formed in the product.



Two examples using Grignard reagents are shown.




Problem 20.27 Draw the product formed when each compound is treated with two equivalents of $CH_3CH_2CH_2CH_2MgBr$ followed by H_2O.



The mechanism for this addition reaction resembles the mechanism for the metal hydride reduction of acid chlorides and esters discussed in Section 20.7A. The mechanism is conceptually divided into two parts: **nucleophilic substitution** to form a ketone (Steps [1] and [2]), followed by **nucleophilic addition** to form a 3° alcohol (Steps [3] and [4]), as shown in Mechanism 20.7.

Mechanism 20.7 Reaction of R"MgX or R"Li with RCOCl and RCOOR'




1. Nucleophilic attack of $(R")^-$ forms a tetrahedral intermediate with a leaving group Z.
2. The π bond is re-formed and the leaving group Z departs to form a ketone. The overall result of addition of $(R")^-$ and elimination of $Z:^-$ is substitution of R" for Z.
3. Nucleophilic attack of $(R")^-$ forms an alkoxide with no leaving group.
4. Protonation of the alkoxide by H_2O forms a 3° alcohol.

Organolithium and Grignard reagents afford 3° alcohols when they react with esters and acid chlorides. As soon as the ketone forms by addition of one equivalent of reagent to RCOZ (Steps [1] and [2] of the mechanism), it reacts with a second equivalent of reagent to form the 3° alcohol.


This reaction is more limited than the Grignard addition to aldehydes and ketones, because only 3° alcohols having two identical alkyl groups can be prepared. Nonetheless, it is still a valuable reaction because it forms two new carbon–carbon bonds.

Sample Problem 20.3 What ester and Grignard reagent are needed to prepare the following alcohol?




Solution


A 3° alcohol formed from an ester and Grignard reagent must have **two identical R groups,** and these R groups come from RMgX. The remainder of the molecule comes from the ester. The carbon (labeled in blue) bonded to the OH group comes from the carbonyl carbon.



Checking in the synthetic direction:




Problem 20.28 What ester and Grignard reagent are needed to prepare each alcohol?






20.13B Reaction of R_2CuLi with Acid Chlorides

To form a ketone from a carboxylic acid derivative, a less reactive organometallic reagent—namely an **organocuprate**—is needed. **Acid chlorides, which have the best leaving group (Cl^-) of the carboxylic acid derivatives, react with R'_2CuLi, to give a ketone as product.** Esters, which contain a poorer leaving group (^-OR), do *not* react with R'_2CuLi.







This reaction results in nucleophilic substitution of an alkyl group R' for the leaving group Cl, forming one new carbon–carbon bond.

Problem 20.29 What organocuprate reagent is needed to convert CH_3CH_2COCl to each ketone?

a. 
b. 
c. 


Problem 20.30 What reagent is needed to convert $(CH_3)_2CHCH_2COCl$ into each compound?

a. 
b. 

c. 
d. 

A ketone with two different R groups bonded to the carbonyl carbon can be made by two different methods, as illustrated in Sample Problem 20.4.

Sample Problem 20.4 Show two different ways to prepare pentan-2-one from an acid chloride and an organocuprate reagent.




pentan-2-one


Solution

In each case, one alkyl group comes from the organocuprate and one comes from the acid chloride.



Possibility [1] Use $(CH_3)_2CuLi$ and a four-carbon acid chloride.



Possibility [2] Use $(CH_3CH_2CH_2)_2CuLi$ and a two-carbon acid chloride.



Problem 20.31 Draw two different ways to prepare each ketone from an acid chloride and an organocuprate reagent.


a. 
b. 

20.14 Reaction of Organometallic Reagents with Other Compounds

Because organometallic reagents are strong nucleophiles, they react with many other electrophiles in addition to carbonyl groups. Because these reactions always lead to the formation of new carbon–carbon bonds, they are also valuable in organic synthesis. In Section 20.14, we examine the reactions of organometallic reagents with **carbon dioxide** and **epoxides.**


20.14A Reaction of Grignard Reagents with Carbon Dioxide

Grignard reagents react with CO_2 to give carboxylic acids after protonation with aqueous acid. This reaction, called **carboxylation,** forms a carboxylic acid with one more carbon atom than the Grignard reagent from which it is prepared.




carboxylic acid

Because Grignard reagents are made from alkyl or aryl halides, an organic halide can be converted to a carboxylic acid having one more carbon atom by a two-step reaction sequence: **formation of a Grignard reagent,** followed by **reaction with CO_2.**




The mechanism resembles earlier reactions of nucleophilic Grignard reagents with carbonyl groups, as shown in Mechanism 20.8.

Mechanism 20.8 Carboxylation—Reaction of RMgX with CO_2




1. The nucleophilic Grignard reagent attacks the electrophilic carbon of CO_2, cleaving the π bond and forming a new carbon–carbon bond.
2. Protonation of the carboxylate anion with aqueous acid forms the carboxylic acid.

Problem 20.32 What carboxylic acid is formed from each alkyl halide on treatment with [1] Mg; [2] CO_2; [3] H_3O^+?




20.14B Reaction of Organometallic Reagents with Epoxides

Like other strong nucleophiles, **organometallic reagents**—RLi, RMgX, and R_2CuLi—**open epoxide rings to form alcohols.**




alcohol




The opening of epoxide rings with negatively charged nucleophiles was discussed in Section 9.15A.

The reaction follows the same two-step process as the opening of epoxide rings with other negatively charged nucleophiles—that is, **nucleophilic attack from the back side of the epoxide ring, followed by protonation of the resulting alkoxide.** In unsymmetrical epoxides, nucleophilic attack occurs at the less substituted carbon atom.




Problem 20.33 What epoxide is needed to convert CH_3CH_2MgBr to each of the following alcohols, after quenching with water?



20.15 α,β-Unsaturated Carbonyl Compounds


α,β-Unsaturated carbonyl compounds are conjugated molecules containing a carbonyl group and a carbon–carbon double bond, separated by a single σ bond.



α,β-unsaturated carbonyl compound

Both functional groups of α,β-unsaturated carbonyl compounds have π bonds, but individually, they react with very different kinds of reagents. Carbon–carbon double bonds react with electrophiles (Chapter 10) and carbonyl groups react with nucleophiles (Section 20.2). What happens, then, when these two functional groups having opposite reactivity are in close proximity?


Because the two π bonds are conjugated, the electron density in an α,β-unsaturated carbonyl compound is delocalized over four atoms. Three resonance structures show that the carbonyl carbon and the β carbon bear a partial positive charge. This means that **α,β-unsaturated carbonyl compounds can react with nucleophiles at two different sites.**



three resonance structures for an α,β-unsaturated carbonyl compound


hybrid two electrophilic sites

- **Addition of a nucleophile to the carbonyl carbon, called 1,2-addition, adds the elements of H and Nu across the C=O, forming an allylic alcohol.**



allylic alcohol

- **Addition of a nucleophile to the β carbon, called 1,4-addition or conjugate addition, forms a carbonyl compound.**




Both 1,2- and 1,4-addition result in nucleophilic addition of the elements of H and Nu.

20.15A The Mechanisms for 1,2-Addition and 1,4-Addition

The steps for the mechanism of 1,2-addition are exactly the same as those for the nucleophilic addition to an aldehyde or ketone—that is, **nucleophilic attack,** followed by **protonation** (Section 20.2A), as shown in Mechanism 20.9.

Mechanism 20.9 1,2-Addition to an α,β-Unsaturated Carbonyl Compound




1 The nucleophile attacks the electrophilic carbonyl. The π bond is broken, moving an electron pair out on oxygen.

2 Protonation of the negatively charged oxygen by H_2O forms the addition product. H and Nu are added to the carbonyl group.

The mechanism for 1,4-addition also begins with nucleophilic attack, and then protonation and tautomerization add the elements of H and Nu to the α and β carbons of the carbonyl compound, as shown in Mechanism 20.10.

Mechanism 20.10 1,4-Addition to an α,β-Unsaturated Carbonyl Compound



1 Nucleophilic attack at the electrophilic β carbon forms a resonance-stabilized enolate anion, which can react on either carbon or oxygen in the second step.


2a Protonation of the carbon end of the enolate forms the 1,4-addition product directly.

2b – 3 Protonation of the oxygen end of the enolate forms an enol, which undergoes tautomerization by the two-step process described in Section 11.9. This forms the same 1,4-addition product that results from protonation on carbon.


20.15B Reaction of α,β-Unsaturated Carbonyl Compounds with Organometallic Reagents


The identity of the metal in an organometallic reagent determines whether it reacts with an α,β-unsaturated aldehyde or ketone by 1,2-addition or 1,4-addition.

Why is conjugate addition also called 1,4-addition? If the atoms of the enol are numbered beginning with the O atom, then the elements of H and Nu are bonded to atoms "1" and "4," respectively.





- **Organolithium and Grignard reagents form 1,2-addition products.**






- **Organocuprate reagents form 1,4-addition products.**





Sample Problem 20.5 Draw the products of each reaction.




Solution


The characteristic reaction of α,β-unsaturated carbonyl compounds is nucleophilic addition. The reagent determines the mode of addition (1,2- or 1,4-).

a. **Grignard reagents undergo 1,2-addition.** CH_3CH_2MgBr adds a new CH_3CH_2 group at the carbonyl carbon.

b. **Organocuprate reagents undergo 1,4-addition.** The cuprate reagent adds a new vinyl group ($CH_2{=}CH$) at the β carbon.




Problem 20.34 Draw the product when each compound is treated with either $(CH_3)_2CuLi$, followed by H_2O, or $HC{\equiv}CLi$, followed by H_2O.



20.16 Summary—The Reactions of Organometallic Reagents


We have now seen many different reactions of organometallic reagents with a variety of functional groups, and you may have some difficulty keeping them all straight. Rather than memorizing them all, keep in mind the following three concepts:

[1] Organometallic reagents (R – M) attack electrophilic carbon atoms, especially the carbonyl carbon.



[2] After an organometallic reagent adds to a carbonyl group, the fate of the intermediate depends on the presence or absence of a leaving group.

- Without a leaving group, the characteristic reaction is *nucleophilic addition.*
- With a leaving group, it is *nucleophilic substitution.*



[3] The polarity of the R – M bond determines the reactivity of the reagents.

- RLi and RMgX are very reactive reagents.
- R_2CuLi is much less reactive.

20.17 Synthesis

The reactions learned in Chapter 20 have proven extremely useful in organic synthesis. Oxidation and reduction reactions interconvert two functional groups that differ in oxidation state. Organometallic reagents form new carbon–carbon bonds.

Synthesis is perhaps the most difficult aspect of organic chemistry. It requires you to remember both the new reactions you've just learned, and the ones you've encountered in previous chapters. In a successful synthesis, you must also put these reactions in a logical order. Don't be discouraged. Learn the basic reactions and then practice them over and over again with synthesis problems.

In Sample Problems 20.6–20.8 that follow, keep in mind that the products formed by the reactions of Chapter 20 can themselves be transformed into many other functional groups. For example, hexan-2-ol, the product of Grignard addition of butylmagnesium chloride to acetaldehyde, can be transformed into a variety of other compounds, as shown in Figure 20.8.






Figure 20.8
Conversion of hexan-2-ol into other compounds




Before proceeding with Sample Problems 20.6–20.8, you should review the stepwise strategy for designing a synthesis found in Section 11.12.

Sample Problem 20.6 Synthesize 1-methylcyclohexene from cyclohexanone and any organic alcohol.



Retrosynthetic Analysis




Thinking backwards:

- [1] Form the double bond by dehydration of an alcohol.
- [2] Make the 3° alcohol by Grignard addition of CH_3MgX.
- [3] Prepare the Grignard reagent stepwise from an alcohol.

Synthesis

Four steps are needed:




major product
trisubstituted alkene

- Conversion of CH_3OH to the Grignard reagent CH_3MgBr requires two steps: formation of an alkyl halide (Step [1]), followed by reaction with Mg (Step [2]).
- Addition of CH_3MgBr to cyclohexanone followed by protonation forms an alcohol in Step [3].
- Acid-catalyzed elimination of water in Step [4] forms a mixture of alkenes, with the desired trisubstituted alkene as the major product.


Sample Problem 20.7

Synthesize 2,4-dimethylhexan-3-one from four-carbon alcohols.



2,4-dimethylhexan-3-one

Retrosynthetic Analysis




Synthesize each of these components.

Thinking backwards:


- [1] Form the ketone by oxidation of a 2° alcohol.
- [2] Make the 2° alcohol by Grignard addition to an aldehyde. Both of these compounds have 4 C's, and each must be synthesized from an alcohol.

Synthesis

First, make both components needed for the Grignard reaction.




Then, complete the synthesis with Grignard addition, followed by oxidation of the alcohol to the ketone.



new C–C bond in red


Sample Problem 20.8

Synthesize isopropylcyclopentane from alcohols having ≤ 5 C's.



isopropylcyclopentane

Retrosynthetic Analysis




Thinking backwards:


- [1] Form the alkane by hydrogenation of an alkene.
- [2] Introduce the double bond by dehydration of an alcohol.
- [3] Form the 3° alcohol by Grignard addition to a ketone. Both components of the Grignard reaction must then be synthesized.

Synthesis

First make both components needed for the Grignard reaction.





Complete the synthesis with Grignard addition, dehydration, and hydrogenation.





major product
tetrasubstituted
double bond


Problem 20.35 Synthesize each compound from cyclohexanol, ethanol, and any other needed reagents.

a. 

b. 

c. 

d. 


e. 

KEY CONCEPTS


Introduction to Carbonyl Chemistry; Organometallic Reagents; Oxidation and Reduction

Reduction Reactions


[1] Reduction of aldehydes and ketones to 1° and 2° alcohols (20.4)



[2] Reduction of α,β-unsaturated aldehydes and ketones (20.4C)




[3] Enantioselective ketone reduction (20.6)




- A single enantiomer is formed.

[4] Reduction of acid chlorides (20.7A)




- $LiAlH_4$, a strong reducing agent, reduces an acid chloride to an alcohol.
- With $LiAlH[OC(CH_3)_3]_3$, a milder reducing agent, reduction stops at the aldehyde stage.

[5] Reduction of esters (20.7A)




- $LiAlH_4$, a strong reducing agent, reduces an ester to an alcohol.
- With DIBAL-H, a milder reducing agent, reduction stops at the aldehyde stage.

[6] Reduction of carboxylic acids to 1° alcohols (20.7B)




[7] Reduction of amides to amines (20.7B)



Oxidation Reactions

Oxidation of aldehydes to carboxylic acids (20.8)




- All Cr^{6+} reagents except PCC oxidize RCHO to RCOOH.
- Tollens reagent (Ag_2O + NH_4OH) oxidizes RCHO only. Primary (1°) and 2° alcohols do not react with Tollens reagent.

Preparation of Organometallic Reagents (20.9)


[1] Organolithium reagents:




[2] Grignard reagents:





[3] Organocuprate reagents:





[4] Lithium and sodium acetylides:






Reactions with Organometallic Reagents

[1] Reaction as a base (20.9C)



- RM = RLi, RMgX, R_2CuLi
- This acid–base reaction occurs with H_2O, ROH, RNH_2, R_2NH, RSH, RCOOH, $RCONH_2$, and RCONHR.


[2] Reaction with aldehydes and ketones to form 1°, 2°, and 3° alcohols (20.10)



R' = H or alkyl


1°, 2°, or 3° alcohol

[3] Reaction with esters to form 3° alcohols (20.13A)




3° alcohol

[4] Reaction with acid chlorides (20.13B)




- More reactive organometallic reagents—R"Li and R"MgX—add two equivalents of R" to an acid chloride to form a 3° alcohol with two identical R" groups.
- Less reactive organometallic reagents—R'_2CuLi—add only one equivalent of R' to an acid chloride to form a ketone.

[5] Reaction with carbon dioxide—Carboxylation (20.14A)




carboxylic acid

[6] Reaction with epoxides (20.14B)



alcohol


[7] Reaction with α,β-unsaturated aldehydes and ketones (20.15B)



- More reactive organometallic reagents—R'Li and R'MgX—react with α,β-unsaturated carbonyls by 1,2-addition.
- Less reactive organometallic reagents—R'_2CuLi—react with α,β-unsaturated carbonyls by 1,4-addition.


Protecting Groups (20.12)

[1] Protecting an alcohol as a *tert*-butyldimethylsilyl ether



tert-butyldimethylsilyl ether


[2] Deprotecting a *tert*-butyldimethylsilyl ether to re-form an alcohol



PROBLEMS

Problems Using Three-Dimensional Models


20.36 Draw the products formed when **A** or **B** is treated with each reagent. In some cases, no reaction occurs.





A **B**


a. $NaBH_4$, CH_3OH
b. [1] $LiAlH_4$; [2] H_2O
c. [1] CH_3MgBr (excess); [2] H_2O
d. [1] C_6H_5Li (excess); [2] H_2O
e. $Na_2Cr_2O_7$, H_2SO_4, H_2O

20.37 Devise a synthesis of each alcohol from organic alcohols having one or two carbons and any required reagents.

a. 
(+ enantiomer)

b. 

c. 


d. 


Reactions and Reagents


20.38 Draw the product formed when pentanal ($CH_3CH_2CH_2CH_2CHO$) is treated with each reagent. With some reagents, no reaction occurs.


a. $NaBH_4$, CH_3OH
b. [1] $LiAlH_4$; [2] H_2O
c. H_2, Pd-C
d. PCC
e. $Na_2Cr_2O_7$, H_2SO_4, H_2O
f. Ag_2O, NH_4OH
g. [1] CH_3MgBr; [2] H_2O
h. [1] C_6H_5Li; [2] H_2O
i. [1] $(CH_3)_2CuLi$; [2] H_2O
j. [1] $HC{\equiv}CNa$; [2] H_2O
k. [1] $CH_3C{\equiv}CLi$; [2] H_2O
l. The product in (a), then TBDMS−Cl, imidazole

20.39 Draw the product formed when $(CH_3CH_2CH_2CH_2)_2CuLi$ is treated with each compound. In some cases, no reaction occurs.

a. 


b. 

c.  , then H_2O

d.  , then H_2O


20.40 The stereochemistry of the products of reduction depends on the reagent used, as you learned in Sections 20.5 and 20.6. With this in mind, how would you convert 3,3-dimethylbutan-2-one [$CH_3COC(CH_3)_3$] to: (a) racemic 3,3-dimethylbutan-2-ol [$CH_3CH(OH)C(CH_3)_3$]; (b) only (*R*)-3,3-dimethylbutan-2-ol; (c) only (*S*)-3,3-dimethylbutan-2-ol?


20.41 Draw the product formed when the α,β-unsaturated ketone **A** is treated with each reagent.



A


a. $NaBH_4$, CH_3OH
b. H_2 (1 equiv), Pd-C
c. H_2 (excess), Pd-C
d. [1] CH_3Li; [2] H_2O
e. [1] CH_3CH_2MgBr; [2] H_2O
f. [1] $(CH_2{=}CH)_2CuLi$; [2] H_2O

20.42 Draw the products of each reduction reaction.


a. 

b. 


c. 

d. 


20.43 Draw the products of the following reactions with organometallic reagents.



20.44 Draw all stereoisomers formed in each reaction.




20.45 Which carbon is most electrophilic in **Y?** Explain your choice.



Y

20.46 Treatment of ketone **A** with ethynyllithium (HC≡CLi) followed by D_3O^+ afforded a compound **B** of molecular formula $C_{12}H_{13}DO_3$, which gave an IR absorption at approximately 1715 cm^{-1}. What is the structure of **B** and how is it formed?



A


20.47 Explain why metal hydride reduction gives an endo alcohol as the major product in one reaction and an exo alcohol as the major product in the other reaction.



endo OH group


exo OH group

20.48 A student tried to carry out the following reaction sequence, but none of diol **A** was formed. Explain what was wrong with this plan, and design a successful stepwise synthesis of **A.**




A

20.49 Identify the lettered compounds in the following reaction scheme. Compounds **F, G,** and **K** are isomers of molecular formula $C_{13}H_{18}O$. How could ^1H NMR spectroscopy distinguish these three compounds from each other?




20.50 Fill in the lettered products (**A**–**D**) in the synthesis of (*R*)-isoproterenol, a drug that increases heart rate and dilates lung passages.




(*R*)-isoproterenol
trade name Isuprel

Mechanism


20.51 Draw a stepwise mechanism for the following reaction. Your mechanism must show how both organic products are formed.




20.52 Draw a stepwise mechanism for the following reaction.



20.53 Tertiary alcohols can be formed by the reaction of dimethyl carbonate [$(CH_3O)_2C=O$] with excess Grignard reagent. Draw a stepwise mechanism for the following transformation.




20.54 Draw a stepwise mechanism for the following reduction.




Synthesis

20.55 What Grignard reagent and aldehyde (or ketone) are needed to prepare each alcohol? Show all possible routes.




20.56 Procyclidine is a drug that has been used to treat the uncontrolled body movements associated with Parkinson's disease. Draw three different methods to prepare procyclidine using a Grignard reagent.




procyclidine

20.57 What ester and Grignard reagent are needed to synthesize each alcohol?




20.58 What organolithium reagent and carbonyl compound can be used to prepare each of the following compounds? You may use aldehydes, ketones, or esters as carbonyl starting materials.



(two ways) (three ways)


20.59 What epoxide and organometallic reagent are needed to synthesize each alcohol?





20.60 Propose at least three methods to convert $C_6H_5CH_2CH_2Br$ to $C_6H_5CH_2CH_3$.


20.61 Propose two different methods to synthesize oct-1-en-3-ol [$CH_3(CH_2)_4CH(OH)CH=CH_2$] using a Grignard reagent and a carbonyl compound. Oct-1-en-3-ol is commonly called matsutake alcohol because it was first isolated from the Japanese matsutake mushroom.

20.62 Synthesize each compound from cyclohexanol using any other organic or inorganic compounds.

a. 


b. 


c. 

d. 


(Each cyclohexane ring must come from cyclohexanol.)


20.63 Convert propan-2-ol [$(CH_3)_2CHOH$] into each compound. You may use any other organic or inorganic compounds.

a. 


b. 


20.64 Convert benzene into each compound. You may also use any inorganic reagents and organic alcohols having three carbons or fewer. One step of the synthesis must use a Grignard reagent.


a. 

b. 


20.65 Design a synthesis of each compound from alcohols having four carbons or fewer as the only organic starting materials. You may use any other inorganic reagents you choose.

a. 


b. 

c. 


20.66 Synthesize each compound from the given starting material. You may use any other required inorganic reagents.

a. 


+ organic halides

b. 

20.67 Devise a synthesis of the given alcohol from benzene, organic alcohols having four or fewer carbons, and any needed inorganic reagents.




20.68 Devise a synthesis of (*E*)-tetradec-11-enal, a sex pheromone of the spruce budworm, a pest that destroys fir and spruce forests, from acetylene, $Br(CH_2)_{10}OH$, and any needed organic compounds or inorganic reagents.



(*E*)-tetradec-11-enal

Spectroscopy

20.69 An unknown compound **A** (molecular formula $C_7H_{14}O$) was treated with $NaBH_4$ in CH_3OH to form compound **B** (molecular formula $C_7H_{16}O$). Compound **A** has a strong absorption in its IR spectrum at 1716 cm^{-1}. Compound **B** has a strong absorption in its IR spectrum at 3600–3200 cm^{-1}. The ^1H NMR spectra of **A** and **B** are given. What are the structures of **A** and **B?**



20.70 Treatment of compound **C** (molecular formula C_4H_8O) with C_6H_5MgBr, followed by H_2O, affords compound **D** (molecular formula $C_{10}H_{14}O$). Compound **D** has a strong peak in its IR spectrum at 3600–3200 cm^{-1}. The ^1H NMR spectral data of **C** and **D** are given. What are the structures of **C** and **D?**

Compound **C** signals at 1.3 (singlet, 6 H) and 2.4 (singlet, 2 H) ppm


Compound **D** signals at 1.2 (singlet, 6 H), 1.6 (singlet, 1 H), 2.7 (singlet, 2 H), and 7.2 (multiplet, 5 H) ppm

20.71 Treatment of compound **E** (molecular formula $C_4H_8O_2$) with excess CH_3CH_2MgBr yields compound **F** (molecular formula $C_6H_{14}O$) after protonation with H_2O. **E** shows a strong absorption in its IR spectrum at 1743 cm^{-1}. **F** shows a strong IR absorption at 3600–3200 cm^{-1}. The ^1H NMR spectral data of **E** and **F** are given. What are the structures of **E** and **F?**


Compound **E** signals at 1.2 (triplet, 3 H), 2.0 (singlet, 3 H), and 4.1 (quartet, 2 H) ppm

Compound **F** signals at 0.9 (triplet, 6 H), 1.1 (singlet, 3 H), 1.5 (quartet, 4 H), and 1.55 (singlet, 1 H) ppm

20.72 Reaction of butanenitrile ($CH_3CH_2CH_2CN$) with methylmagnesium bromide (CH_3MgBr), followed by treatment with aqueous acid, forms compound **G. G** has a molecular ion in its mass spectrum at *m/z* = 86 and a base peak at *m/z* = 43. **G** exhibits a strong absorption in its IR spectrum at 1721 cm^{-1} and has the ^1H NMR spectrum given below. What is the structure of **G?** We will learn about the details of this reaction in Chapter 22.




20.73 Treatment of isobutene [$(CH_3)_2C=CH_2$] with $(CH_3)_3CLi$ forms a carbanion that reacts with $CH_2=O$ to form **H** after water is added to the reaction mixture. **H** has a molecular ion in its mass spectrum at m/z = 86, and shows fragments at 71 and 68. **H** exhibits absorptions in its IR spectrum at 3600–3200 and 1651 cm^{-1}, and has the ^1H NMR spectrum given below. What is the structure of **H?**




Challenge Problems


20.74 Draw a stepwise mechanism for the following reaction.




20.75 Design a synthesis of (*R*)-salmeterol (Figure 20.3) from the following starting materials.



20.76 Lithium tri-*sec*-butylborohydride, also known as L-selectride, is a metal hydride reagent that contains three *sec*-butyl groups bonded to boron. When this reagent is used to reduce cyclic ketones, one stereoisomer often predominates as product. Explain why the reduction of 4-*tert*-butylcyclohexanone with L-selectride forms the cis alcohol as the major product.




20.77 Explain why the β carbon of an α,β-unsaturated carbonyl compound absorbs farther downfield in the ^{13}C NMR spectrum than the α carbon, even though the α carbon is closer to the electron-withdrawing carbonyl group. For example, the β carbon of mesityl oxide absorbs at 150.5 ppm, while the α carbon absorbs at 122.5 ppm.




mesityl oxide


20.78 Identify **X** and **Y,** two of the intermediates in a synthesis of the antidepressant venlafaxine (trade name Effexor), in the following reaction scheme. Write a mechanism for the formation of **X** from **W.**




20.79 Reaction of benzylmagnesium chloride with formaldehyde yields alcohols **N** and **P** after protonation. Draw a stepwise mechanism that shows how both products are formed.



20.80 Draw a stepwise mechanism for the following reaction. (Hint: Conjugate addition can occur with heteroatoms as well as carbon nucleophiles.)




20.81 Draw a stepwise mechanism for the following reaction of a Grignard reagent with a cyclic amide.



Aldehydes and Ketones—Nucleophilic Addition

21



21.1 Introduction
21.2 Nomenclature
21.3 Physical properties
21.4 Spectroscopic properties
21.5 Interesting aldehydes and ketones
21.6 Preparation of aldehydes and ketones
21.7 Reactions of aldehydes and ketones—General considerations
21.8 Nucleophilic addition of H^- and R^-—A review
21.9 Nucleophilic addition of ^-CN
21.10 The Wittig reaction
21.11 Addition of 1° amines
21.12 Addition of 2° amines
21.13 Addition of H_2O—Hydration
21.14 Addition of alcohols—Acetal formation
21.15 Acetals as protecting groups
21.16 Cyclic hemiacetals
21.17 An introduction to carbohydrates

The natural product **digoxin** has been prescribed since the 1960s for patients with congestive heart failure, a condition that results when fluid builds up in the body because the heart's pumping action is weak. Unlike many commercial medications that are synthesized from simple precursors, digoxin is still obtained by extraction of the leaves of the woolly foxglove plant, which is grown in the Netherlands and shipped to the United States for processing. One thousand kilograms of dried leaves yield one kilogram of digoxin, sold under the trade name of Lanoxin. Digoxin contains three acetal units, which are formed by addition reactions to carbonyl groups. In Chapter 21, we learn about nucleophilic addition, the characteristic reaction of aldehydes and ketones.


In Chapter 21 we continue the study of carbonyl compounds with a detailed look at **aldehydes** and **ketones.** We will first learn about the nomenclature, physical properties, and spectroscopic absorptions that characterize aldehydes and ketones. The remainder of Chapter 21 is devoted to **nucleophilic addition** reactions. Although we have already learned two examples of this reaction in Chapter 20, nucleophilic addition to aldehydes and ketones is a general reaction that occurs with many nucleophiles, forming a wide variety of products.

Every new reaction in Chapter 21 involves nucleophilic addition, so the challenge lies in learning the specific reagents and mechanisms that characterize each reaction.

21.1 Introduction


An aldehyde is often written as **RCHO.** Remember that the **H atom is bonded to the carbon atom,** *not* the oxygen. Likewise, a ketone is written as **RCOR,** or if both alkyl groups are the same, **R_2CO.** Each structure must contain a C=O for every atom to have an octet.

As we learned in Chapter 20, **aldehydes and ketones contain a carbonyl group.** An aldehyde contains at least one H atom bonded to the carbonyl carbon, whereas a ketone has two alkyl or aryl groups bonded to it.



carbonyl group **aldehyde** **ketone**

Two structural features determine the chemistry and properties of aldehydes and ketones.




trigonal planar electrophilic carbon


- The carbonyl group is sp^2 hybridized and trigonal planar, making it relatively uncrowded.
- The electronegative oxygen atom polarizes the carbonyl group, making the carbonyl carbon electrophilic.

As a result, **aldehydes and ketones react with nucleophiles.** The relative reactivity of the carbonyl group is determined by the number of R groups bonded to it. **As the number of R groups around the carbonyl carbon increases, the reactivity of the carbonyl compound decreases,** resulting in the following order of reactivity:

Increasing the number of alkyl groups on the carbonyl carbon decreases reactivity for both steric and electronic reasons, as discussed in Section 20.2B.



Problem 21.1 Rank the following compounds in order of increasing reactivity towards nucleophilic attack.



Problem 21.2 Explain why benzaldehyde is less reactive than cyclohexanecarbaldehyde towards nucleophilic attack.



benzaldehyde cyclohexanecarbaldehyde

21.2 Nomenclature

Both IUPAC and common names are used for aldehydes and ketones.

21.2A Naming Aldehydes in the IUPAC System


In IUPAC nomenclature, aldehydes are identified by a suffix added to the parent name of the longest chain. Two different suffixes are used, depending on whether the CHO group is bonded to a chain or a ring.


To name an aldehyde using the IUPAC system:

[1] If the CHO is bonded to a chain of carbons, find the longest chain containing the CHO group, and change the ***-e*** ending of the parent alkane to the suffix ***-al.*** If the CHO group is bonded to a ring, name the ring and add the suffix ***-carbaldehyde.***

[2] Number the chain or ring to put the CHO group at C1, but omit this number from the name. Apply all of the other usual rules of nomenclature.


Sample Problem 21.1 Give the IUPAC name for each compound.

a. 

b. 


Solution

a. [1] Find and name the longest chain containing the CHO:




butane ⟶ butan*al*
(4 C's)

[2] Number and name substituents:




Answer: 2,3-dimethylbutanal

b. [1] Find and name the ring bonded to the CHO group:




cyclohexane + carbaldehyde
(6 C's)


[2] Number and name substituents:




Answer: 2-ethylcyclohexanecarbaldehyde

Problem 21.3 Give the IUPAC name for each aldehyde.

a. 

b. 

c. 

Problem 21.4 Give the structure corresponding to each IUPAC name.


a. 2-isobutyl-3-isopropylhexanal
b. *trans*-3-methylcyclopentanecarbaldehyde
c. 1-methylcyclopropanecarbaldehyde
d. 3,6-diethylnonanal

21.2B Common Names for Aldehydes

Like carboxylic acids, many simple aldehydes have common names that are widely used.

- **A common name for an aldehyde is formed by taking the common parent name and adding the suffix *-aldehyde.***


The common parent names are similar to those used for carboxylic acids, listed in Table 19.1. The common names **formaldehyde, acetaldehyde,** and **benzaldehyde** are virtually always used instead of their IUPAC names.



formaldehyde (methanal) | acetaldehyde (ethanal) | benzaldehyde (benzenecarbaldehyde)

(IUPAC names are in parentheses.)


Greek letters are used to designate the location of substituents in common names. **The carbon adjacent to the CHO group is the α carbon,** and so forth down the chain.



IUPAC numbering begins at the C=O.
Greek lettering begins at the C bonded to the C=O.

Figure 21.1 gives the common and IUPAC names for three aldehydes.

Figure 21.1
Three examples of aldehyde nomenclature



2-chloropropanal (α-chloropropionaldehyde) | 3-methylpentanal (β-methylvaleraldehyde) | phenylethanal (phenylacetaldehyde)

(Common names are in parentheses.)

21.2C Naming Ketones in the IUPAC System

- **In the IUPAC system all ketones are identified by the suffix *-one.***


To name an acyclic ketone using IUPAC rules:


[1] Find the longest chain containing the carbonyl group, and change the ***-e*** ending of the parent alkane to the suffix ***-one.***

[2] Number the carbon chain to give the carbonyl carbon the lower number. Apply all of the other usual rules of nomenclature.

With cyclic ketones, numbering always begins at the carbonyl carbon, but the "1" is usually omitted from the name. The ring is then numbered clockwise or counterclockwise to give the first substituent the lower number.


Sample Problem 21.2 Give the IUPAC name for each ketone.

a. 

b. 


Solution

a. [1] Find and name the longest chain containing the carbonyl group:




pentane → pentan*one*
(5 C's)

[2] Number and name substitutents:




Answer: 3-methylpentan-2-one

b. [1] Name the ring:




cyclohexane → cyclohexan*one*
(6 C's)


[2] Number and name substitutents:




Answer:
3-isopropyl-4-methylcyclohexanone

Problem 21.5 Give the IUPAC name for each ketone.


a. 

b. 


c. 

21.2D Common Names for Ketones

Most common names for ketones are formed by **naming both alkyl groups** on the carbonyl carbon, **arranging them alphabetically,** and adding the word ***ketone.*** Using this method, the common name for butan-2-one becomes ethyl methyl ketone.



IUPAC name: **butan-2-one**




Common name: **ethyl methyl ketone**


Three widely used common names for some simple ketones do not follow this convention:



acetone




acetophenone




benzophenone

Figure 21.2 gives acceptable names for two ketones.

Figure 21.2
Two examples of ketone nomenclature



IUPAC name: 2-methylpentan-3-one
Common name: ethyl isopropyl ketone




m-bromoacetophenone
or
3-bromoacetophenone


21.2E Additional Nomenclature Facts

Do not confuse a **benzyl** group with a **benzoyl** group.

Sometimes acyl groups (RCO−) must be named as substituents. To name an acyl group, take either the IUPAC or common parent name and add the suffix ***-yl*** or ***-oyl.*** The three most common acyl groups are drawn below.




benz*yl* group



formyl group




acetyl group




benzoyl group

Compounds containing both a C−C double bond and an aldehyde are named as **enals,** and compounds that contain both a C−C double bond and a ketone are named as **enones.** The chain is numbered to give the carbonyl group the lower number.



2,2-dimethylbut-3-enal




4-methylpent-3-en-2-one

Problem 21.6 Give the structure corresponding to each name: (a) *sec*-butyl ethyl ketone; (b) methyl vinyl ketone; (c) *p*-ethylacetophenone; (d) 3-benzoyl-2-benzylcyclopentanone; (e) 6,6-dimethylcyclohex-2-enone; (f) 3-ethylhex-5-enal.


Problem 21.7 Give the IUPAC name (including any *E,Z* designation) for each unsaturated aldehyde. Neral is obtained from lemon grass, and cucumber aldehyde (Problem 1.30) contributes to the aroma of a fresh mango.

a.



neral

b.



cucumber aldehyde

21.3 Physical Properties

Aldehydes and ketones exhibit dipole–dipole interactions because of their polar carbonyl group. Because they have no O−H bond, two molecules of RCHO or RCOR are incapable of intermolecular hydrogen bonding, making them less polar than alcohols and carboxylic acids. How these intermolecular forces affect the physical properties of aldehydes and ketones is summarized in Table 21.1.

Problem 21.8 The boiling point of butan-2-one (80 °C) is significantly higher than the boiling point of diethyl ether (35 °C), even though both compounds exhibit dipole–dipole interactions and have comparable molecular weights. Offer an explanation.

Table 21.1 Physical Properties of Aldehydes and Ketones

Property	Observation
Boiling point and melting point	• For compounds of comparable molecular weight, bp's and mp's follow the usual trend: The stronger the intermolecular forces, the higher the bp or mp. VDW MW = 72 bp 36 °C VDW, DD MW = 72 bp 76 °C VDW, DD MW = 72 bp 80 °C VDW, DD, HB MW = 74 bp 118 °C **Increasing strength of intermolecular forces** **Increasing boiling point**
Solubility	• RCHO and RCOR are soluble in organic solvents regardless of size. • RCHO and RCOR having ≤ 5 C's are H_2O soluble because they can hydrogen bond with H_2O (Section 3.4C). • RCHO and RCOR having > 5 C's are H_2O insoluble because the nonpolar alkyl portion is too large to dissolve in the polar H_2O solvent.

Key: VDW = van der Waals, DD = dipole–dipole, HB = hydrogen bonding, MW = molecular weight

21.4 Spectroscopic Properties

The presence of the carbonyl group in aldehydes and ketones gives them characteristic absorptions in their IR and NMR spectra.


21.4A IR Spectra

Aldehydes and ketones exhibit the following characteristic IR absorptions:

- Like all carbonyl compounds, **aldehydes and ketones give a strong peak at ~1700 cm^{-1} due to the C=O.**
- **The sp^2 hybridized C–H bond of an aldehyde shows one or two peaks at ~2700–2830 cm^{-1}.**

The IR spectrum of propanal in Figure 21.3 illustrates these characteristic peaks.


Figure 21.3
The IR spectrum of propanal, CH_3CH_2CHO



- A strong C=O occurs at 1739 cm^{-1}.
- The sp^2 C–H of the CHO appears as two peaks at 2813 and 2716 cm^{-1}.

The exact position of the carbonyl absorption often provides additional information about a compound. For example, most aldehydes have a C=O peak around **1730 cm^{-1},** whereas for ketones, it is typically around **1715 cm^{-1}.** Two other structural features—**ring size** (for cyclic ketones) and **conjugation**—affect the location of the carbonyl absorption in a predictable manner.

[1] The carbonyl absorption of cyclic ketones shifts to higher wavenumber as the size of the ring decreases and the ring strain increases.



[2] Conjugation of the carbonyl group with a C=C or a benzene ring shifts the absorption to lower wavenumber by ~30 cm^{-1}.

The effect of conjugation on the frequency of the C=O absorption is explained by **resonance.** An α,β-unsaturated carbonyl compound can be written as three resonance structures, two of which place a single bond between the carbon and oxygen atoms of the carbonyl group. Thus, the π bond of the carbonyl group is delocalized, giving the conjugated carbonyl group some single bond character, and making it somewhat **weaker** than an unconjugated C=O. **Weaker bonds absorb at lower frequency (lower wavenumber) in an IR spectrum.**






Figure 21.4 illustrates the effects of conjugation on the location of the carbonyl absorption in some representative compounds.

Figure 21.4
The effect of conjugation on the carbonyl absorption in an IR spectrum



Problem 21.9 Which carbonyl group in each pair absorbs at a higher frequency?




21.4B NMR Spectra

Aldehydes and ketones exhibit the following characteristic ^1H and ^{13}C NMR absorptions:

- **The sp^2 hybridized C−H proton of an aldehyde is highly deshielded and absorbs far downfield at 9–10 ppm.** Splitting occurs with protons on the α carbon, but the coupling constant is often very small (J = 1–3 Hz).
- **Protons on the α carbon to the carbonyl group absorb at 2–2.5 ppm.** Methyl ketones, for example, give a characteristic singlet at ~2.1 ppm.
- In a ^{13}C NMR spectrum, **the carbonyl carbon is highly deshielded, appearing in the 190–215 ppm region.**

The ^1H and ^{13}C NMR spectra of propanal are illustrated in Figure 21.5.

Figure 21.5 The ^1H and ^{13}C NMR spectra of propanal, CH_3CH_2CHO




- **^1H NMR:** There are three signals due to the three different kinds of hydrogens, labeled H_a, H_b, and H_c. The deshielded CHO proton occurs downfield at 9.8 ppm. The H_c signal is split into a triplet by the adjacent CH_2 group, but the coupling constant is small.
- **^{13}C NMR:** There are three signals due to the three different kinds of carbons, labeled C_a, C_b, and C_c. The deshielded carbonyl carbon absorbs downfield at 203 ppm.

Problem 21.10 Draw the structure of all constitutional isomers that contain a ketone and have molecular formula $C_5H_{10}O$. Give the IUPAC name for each isomer and state how ^{13}C NMR spectroscopy could be used to distinguish these isomers.

21.5 Interesting Aldehydes and Ketones



formaldehyde
$CH_2{=}O$



acetone
$(CH_3)_2C{=}O$

Because it is a starting material for the synthesis of many resins and plastics, billions of pounds of **formaldehyde** are produced annually in the United States by the oxidation of methanol (CH_3OH). Formaldehyde is also sold as a 37% aqueous solution called **formalin,** which has been used as a disinfectant, antiseptic, and preservative for biological specimens. Formaldehyde, a product of the incomplete combustion of coal and other fossil fuels, is partly responsible for the irritation caused by smoggy air.


Acetone is an industrial solvent and a starting material in the synthesis of some organic polymers. Acetone is produced in vivo during the breakdown of fatty acids. In diabetes, a common endocrine disease in which normal metabolic processes are altered because of the inadequate secretion of insulin, individuals often have unusually high levels of acetone in their bloodstreams. The characteristic odor of acetone can be detected on the breath of diabetic patients when their disease is poorly controlled.

Many aldehydes with characteristic odors occur in nature, as shown in Figure 21.6.

Figure 21.6 Some naturally occurring aldehydes with strong odors



Many steroid hormones contain a carbonyl along with other functional groups. **Cortisone** and **prednisone** are two anti-inflammatory steroids with closely related structures. Cortisone is secreted by the body's adrenal gland, whereas prednisone is a synthetic analogue used in the treatment of inflammatory diseases such as arthritis and asthma.




21.6 Preparation of Aldehydes and Ketones

Aldehydes and ketones can be prepared by a variety of methods. Because these reactions are needed for many multistep syntheses, Section 21.6 briefly summarizes earlier reactions that synthesize an aldehyde or ketone.

21.6A Common Methods to Synthesize Aldehydes


Aldehydes are prepared from 1° alcohols, esters, acid chlorides, and alkynes.

- **By oxidation of 1° alcohols with PCC**




(Section 12.12B)

- **By reduction of esters and acid chlorides**



(Section 20.7A)

- **By hydroboration–oxidation of an alkyne**




(Section 11.10)

21.6B Common Methods to Synthesize Ketones


Ketones are prepared from 2° alcohols, acid chlorides, and alkynes.

- **By oxidation of 2° alcohols with Cr^{6+} reagents**




(Section 12.12A)

- **By reaction of acid chlorides with organocuprates**




(Section 20.13)

- **By Friedel–Crafts acylation**




(Section 18.5)

- **By hydration of an alkyne**





(Section 11.9)


Aldehydes and ketones are also both obtained as products of the oxidative cleavage of alkenes (Section 12.10).




Problem 21.11 What reagents are needed to convert each compound into butanal ($CH_3CH_2CH_2CHO$)?

a. 


b. 


c. 

d. 


Problem 21.12 What reagents are needed to convert each compound into acetophenone ($C_6H_5COCH_3$)?

a. 

b. 

c. 

Problem 21.13 What alkene would yield 2,2-dimethoxycyclopentane-1,3-dicarbaldehyde on treatment with O_3 followed by $(CH_3)_2S$?




2,2-dimethoxycyclopentane-1,3-dicarbaldehyde

21.7 Reactions of Aldehydes and Ketones—General Considerations

Let's begin our discussion of carbonyl reactions by looking at the two general kinds of reactions that aldehydes and ketones undergo.

[1] Reaction at the carbonyl carbon

Recall from Chapter 20 that the uncrowded, electrophilic carbonyl carbon makes aldehydes and ketones susceptible to **nucleophilic addition** reactions.




The elements of H and Nu are added to the carbonyl group. In Chapter 20 you learned about this reaction with hydride ($H:^-$) and carbanions ($R:^-$) as nucleophiles. In Chapter 21, we will discuss similar reactions with other nucleophiles.

[2] Reaction at the α carbon

A second general reaction of aldehydes and ketones involves reaction at the **α carbon.** A C−H bond on the α carbon to a carbonyl group is more acidic than many other C−H bonds, because reaction with base forms a resonance-stabilized enolate anion.

- **Enolates are nucleophiles, and so they react with electrophiles (E^+) to form new bonds on the α carbon.**




Chapters 23 and 24 are devoted to reactions at the α carbon to a carbonyl group.

- **Aldehydes and ketones react with nucleophiles at the carbonyl carbon.**
- **Aldehydes and ketones form enolates that react with electrophiles at the α carbon.**

21.7A The General Mechanism of Nucleophilic Addition

Two general mechanisms are usually drawn for nucleophilic addition, depending on the nucleophile (negatively charged versus neutral) and the presence or absence of an acid catalyst. With negatively charged nucleophiles, nucleophilic addition follows the two-step process first discussed in Chapter 20—**nucleophilic attack** followed by **protonation,** as shown in Mechanism 21.1.

Mechanism 21.1 General Mechanism—Nucleophilic Addition



R' = H or alkyl


1 The nucleophile attacks the electrophilic carbonyl. The π bond is broken, moving an electron pair out on oxygen and forming an sp^3 hybridized carbon.

2 Protonation of the negatively charged oxygen by H_2O forms the addition product.

In this mechanism **nucleophilic attack *precedes* protonation.** This process occurs with strong neutral or negatively charged nucleophiles.

With some neutral nucleophiles, however, nucleophilic addition does not occur unless an acid catalyst is added. The general mechanism for this reaction consists of three steps (not two), but the same product results because H and Nu add across the carbonyl π bond. In this mechanism, **protonation *precedes* nucleophilic attack.** Mechanism 21.2 is shown with the neutral nucleophile H−Nu: and a general acid H−A.

Mechanism 21.2 General Mechanism—Acid-Catalyzed Nucleophilic Addition




R' = H or alkyl

resonance-stabilized cation

1 Protonation of the carbonyl oxygen forms a resonance-stabilized cation.

2 – 3 Nucleophilic attack and deprotonation form the neutral addition product. The overall result is addition of H and Nu to the carbonyl group.

The effect of protonation is to convert a neutral carbonyl group to one having a net positive charge. **This protonated carbonyl group is much more electrophilic,** and much more susceptible to attack by a nucleophile. This step is unnecessary with strong nucleophiles like hydride ($H:^-$) that were used in Chapter 20. With weaker nucleophiles, however, nucleophilic attack does not occur unless the carbonyl group is first protonated.




This step is a specific example of a general phenomenon.

- **Any reaction involving a carbonyl group and a strong acid begins with the same first step—protonation of the carbonyl oxygen.**

21.7B The Nucleophile

What nucleophiles add to carbonyl groups? This cannot be predicted solely on the trends in nucleophilicity learned in Chapter 7. Only *some* of the nucleophiles that react well in nucleophilic substitution at sp^3 hybridized carbons give reasonable yields of nucleophilic addition products.

Cl^-, Br^-, and I^- are good nucleophiles in substitution reactions at sp^3 hybridized carbons, but they are ineffective nucleophiles in addition. Addition of Cl^- to a carbonyl group, for example, would cleave the C–O π bond, forming an alkoxide. Because Cl^- is a much weaker base than the alkoxide formed, equilibrium favors the starting materials (the weaker base, Cl^-), *not* the addition product.



The situation is further complicated because some of the initial nucleophilic addition adducts are unstable and undergo elimination to form a stable product. For example, amines (RNH_2) add to carbonyl groups in the presence of mild acid to form unstable **carbinolamines,** which readily lose water to form **imines.** This addition–elimination sequence replaces a C=O by a C=N. The details of this process are discussed in Section 21.11.





Figure 21.7 lists nucleophiles that add to a carbonyl group, as well as the products obtained from nucleophilic addition using cyclohexanone as a representative ketone. These reactions are discussed in the remaining sections of Chapter 21. In cases in which the initial addition adduct is unstable, it is enclosed within brackets, followed by the final product.


Figure 21.7
Specific examples of nucleophilic addition




21.8 Nucleophilic Addition of H^- and R^-—A Review

We begin our study of nucleophilic additions to aldehydes and ketones by briefly reviewing nucleophilic addition of hydride and carbanions, two reactions examined in Sections 20.4 and 20.10, respectively.


Treatment of an aldehyde or ketone with either $NaBH_4$ or $LiAlH_4$ followed by protonation forms a 1° or 2° alcohol. $NaBH_4$ and $LiAlH_4$ serve as a source of **hydride, $H:^-$—the nucleophile**—and the reaction results in addition of the elements of H_2 across the C−O π bond. Addition of H_2 reduces the carbonyl group to an alcohol.



Hydride reduction of aldehydes and ketones occurs via the two-step mechanism of nucleophilic addition—that is, **nucleophilic attack of H:⁻ followed by protonation**—shown previously in Section 20.4B.




Treatment of an aldehyde or ketone with either an organolithium (R"Li) or Grignard reagent (R"MgX) followed by water forms a 1°, 2°, or 3° alcohol containing a new carbon–carbon bond. R"Li and R"MgX serve as a source of a **carbanion (R")⁻—the nucleophile**—and the reaction results in addition of the elements of R" and H across the C−O π bond.



The stereochemistry of hydride reduction and Grignard addition was discussed previously in Sections 20.5 and 20.10B, respectively.


The nucleophilic addition of carbanions to aldehydes and ketones occurs via the two-step mechanism of nucleophilic addition—that is, **nucleophilic attack of (R")⁻ followed by protonation**—shown previously in Section 20.10.



In both reactions, the nucleophile—either hydride or a carbanion—attacks the trigonal planar sp^2 hybridized carbonyl from both sides, so that when a new stereogenic center is formed, a mixture of stereoisomers results, as shown in Sample Problem 21.3.

Sample Problem 21.3


Draw the products (including the stereochemistry) formed in the following reaction.




(*R*)-3-methylcyclopentanone

Solution

The Grignard reagent adds CH_3^- from both sides of the trigonal planar carbonyl group, yielding a mixture of 3° alcohols after protonation with water. In this example, the starting ketone and both alcohol products are chiral. The two products, which contain two stereogenic centers, are stereoisomers but not mirror images—that is, they are **diastereomers.**




Problem 21.14 Draw the products of each reaction. Include all stereoisomers formed.




21.9 Nucleophilic Addition of ⁻CN

Treatment of an aldehyde or ketone with NaCN and a strong acid such as HCl adds the elements of HCN across the carbon–oxygen π bond, forming a **cyanohydrin.**



This reaction adds one carbon to the aldehyde or ketone, forming a new carbon–carbon bond.




acetaldehyde cyanohydrin
new C–C bond in red

21.9A The Mechanism

The mechanism of cyanohydrin formation involves the usual two steps of nucleophilic addition: **nucleophilic attack followed by protonation** as shown in Mechanism 21.3.

Mechanism 21.3 Nucleophilic Addition of ⁻CN—Cyanohydrin Formation




R' = H or alkyl

new C–C bond in red

1. Nucleophilic attack of ⁻CN forms a new carbon–carbon bond with cleavage of the C–O π bond.
2. Protonation of the negatively charged oxygen by HCN forms the addition product. The HCN used in this step is formed by the acid–base reaction of ⁻CN with the strong acid, HCl.


This reaction does not occur with HCN alone. The **cyanide anion** makes addition possible because it is a strong nucleophile that attacks the carbonyl group.

Cyanohydrins can be reconverted to carbonyl compounds by treatment with base. This process is just the reverse of the addition of HCN: **deprotonation followed by elimination of ⁻CN.**





Note the difference between two similar terms. **Hydration** results in *adding* water to a compound. **Hydrolysis** results in *cleaving bonds* with water.

The cyano group (CN) of a cyanohydrin is readily hydrolyzed to a carboxy group (COOH) by heating with aqueous acid or base. **Hydrolysis replaces the three C–N bonds by three C–O bonds.**



Problem 21.15 Draw the products of each reaction.

a. 


b. 

21.9B Application: Naturally Occurring Cyanohydrin Derivatives




Peach and apricot pits are a natural source of the cyanohydrin derivative amygdalin.


Although the cyanohydrin is an uncommon functional group, **linamarin** and **amygdalin** are two naturally occurring cyanohydrin derivatives. Both contain a carbon atom bonded to both an oxygen atom and a cyano group, analogous to a cyanohydrin.



linamarin



amygdalin




laetrile



Cassava is a widely grown root crop, first introduced to Africa by Portuguese traders from Brazil in the sixteenth century. The peeled root is eaten after boiling or roasting. If the root is eaten without processing, illness and even death can result from high levels of HCN.

Linamarin is isolated from cassava, a woody shrub grown as a root crop in the humid tropical regions of South America and Africa. **Amygdalin** is present in the seeds and pits of apricots, peaches, and wild cherries. Amygdalin and the related synthetic compound **laetrile** were once touted as anticancer drugs, although their effectiveness is unproven.

Linamarin, amygdalin, and laetrile are toxic compounds because they are metabolized to cyanohydrins, which are hydrolyzed to carbonyl compounds and toxic HCN gas, a cellular poison with a characteristic almond odor. This second step is merely the reconversion of a cyanohydrin to a carbonyl compound, a process that occurs with base in reactions run in the laboratory (Section 21.9A). If cassava root is processed with care, linamarin is enzymatically metabolized by this reaction sequence and the toxic HCN is released before the root is ingested, making it safe to eat.



linamarin

cyanohydrin derivative

acetone cyanohydrin


toxic by-product

Problem 21.16 What cyanohydrin and carbonyl compound are formed when amygdalin is metabolized in a similar manner to linamarin?


21.10 The Wittig Reaction

The additions of H^-, R^-, and ^-CN all involve the same two steps—nucleophilic attack followed by protonation. Other examples of nucleophilic addition in Chapter 21 are somewhat different. Although they still involve attack of a nucleophile, the initial addition adduct is converted to another product by one or more reactions.

The first reaction in this category is the **Wittig reaction,** named for German chemist Georg Wittig, who was awarded the Nobel Prize in Chemistry in 1979 for its discovery. The Wittig reaction uses a carbon nucleophile, the **Wittig reagent,** to form **alkenes.** When a carbonyl compound is treated with a Wittig reagent, the carbonyl oxygen atom is replaced by the negatively charged alkyl group bonded to the phosphorus—that is, **the C=O is converted to a C=C.**




- **A Wittig reaction forms two new carbon–carbon bonds—one new σ bond and one new π bond—as well as a phosphorus by-product, $Ph_3P=O$ (triphenylphosphine oxide).**



21.10A The Wittig Reagent


A **Wittig reagent** is an **organophosphorus reagent**—a reagent that contains a carbon–phosphorus bond. A typical Wittig reagent has a phosphorus atom bonded to three phenyl groups, plus another alkyl group that bears a negative charge.



Phosphorus ylides are also called **phosphoranes.**


A Wittig reagent is an ***ylide,*** **a species that contains two oppositely charged atoms bonded to each other, and both atoms have octets.** In a Wittig reagent, a negatively charged carbon atom is bonded to a positively charged phosphorus atom.

Because phosphorus is a third-row element, it can be surrounded by more than eight electrons. As a result, a second resonance structure can be drawn that places a double bond between carbon and phosphorus. Regardless of which resonance structure is drawn, a **Wittig reagent has no net charge.** In one resonance structure, though, the carbon atom bonded to phosphorus (labeled in blue) bears a net negative charge, so it is nucleophilic.



Wittig reagents are synthesized by a two-step procedure.


Step [1] **S_N2 reaction of triphenylphosphine with an alkyl halide forms a phosphonium salt.**



Because phosphorus is located below nitrogen in the periodic table, a neutral phosphorus atom with three bonds also has a lone pair of electrons.

Triphenylphosphine (Ph_3P:), which contains a lone pair of electrons on P, is the nucleophile. Because the reaction follows an S_N2 mechanism, it works best with unhindered CH_3X and 1° alkyl halides (RCH_2X). Secondary alkyl halides (R_2CHX) can also be used, although yields are often lower.

Step [2] **Deprotonation of the phosphonium salt with a strong base (:B) forms the ylide.**






Section 20.9C discussed the reaction of organometallic reagents as strong bases.


Because removal of a proton from a carbon bonded to phosphorus generates a resonance-stabilized carbanion (the ylide), this proton is somewhat more acidic than other protons on an alkyl group in the phosphonium salt. Very strong bases are still needed, though, to favor the products of this acid–base reaction. Common bases used for this reaction are the organolithium reagents such as butyllithium, $CH_3CH_2CH_2CH_2Li$, abbreviated as BuLi.

To synthesize the Wittig reagent, $Ph_3P{=}CH_2$, use these two steps:




- **Step [1]** Form the phosphonium salt by S_N2 reaction of Ph_3P: and CH_3Br.
- **Step [2]** Form the ylide by removal of a proton using BuLi as a strong base.

Problem 21.17 Draw the products of the following Wittig reactions.



Problem 21.18 Outline a synthesis of each Wittig reagent from Ph_3P and an alkyl halide.



21.10B Mechanism of the Wittig Reaction


The currently accepted mechanism of the Wittig reaction involves two steps. Like other nucleophiles, the Wittig reagent attacks an electrophilic carbonyl carbon, but then the initial addition adduct undergoes elimination to form an alkene. Mechanism 21.4 is drawn using $Ph_3P{=}CH_2$.

Mechanism 21.4 The Wittig Reaction




1. The negatively charged carbon of the ylide attacks the carbonyl carbon as the carbonyl oxygen attacks the positively charged P atom. This step forms two bonds and generates a four-membered ring called an oxaphosphetane.
2. Elimination of triphenylphosphine oxide forms two new π bonds. The formation of the strong P=O provides the driving force for the Wittig reaction.

One limitation of the Wittig reaction is that a mixture of alkene stereoisomers sometimes forms. For example, reaction of propanal (CH_3CH_2CHO) with a Wittig reagent forms the mixture of *E* and *Z* isomers shown.



Because the Wittig reaction forms two carbon–carbon bonds in a single reaction, it has been used to synthesize many natural products, including β-carotene, shown in Figure 21.8.


Figure 21.8
A Wittig reaction used to synthesize β-carotene





β-carotene
orange pigment found in carrots
(vitamin A precursor)

- The more stable *E* alkene is the major product in this Wittig reaction.

Problem 21.19 Draw the products (including stereoisomers) formed when benzaldehyde (C_6H_5CHO) is treated with each Wittig reagent.

a. 

b. 


c. 

21.10C Retrosynthetic Analysis

To use the Wittig reaction in synthesis, you must be able to determine what carbonyl compound and Wittig reagent are needed to prepare a given compound—that is, **you must work backwards, in the retrosynthetic direction.** There can be two different Wittig routes to a given alkene, but one is often preferred on steric grounds.


How To Determine the Starting Materials for a Wittig Reaction Using Retrosynthetic Analysis

Example What starting materials are needed to synthesize alkene X by a Wittig reaction?



X


Step [1] Cleave the carbon–carbon double bond into two components.




- Part of the molecule becomes the carbonyl component and the other part becomes the Wittig reagent.

There are usually two routes to a given alkene using a Wittig reaction:

Possibility [1]




Possibility [2]



Step [2] Compare the Wittig reagents. The preferred pathway uses a Wittig reagent derived from an unhindered alkyl halide—CH_3X or RCH_2X.


Determine what alkyl halide is needed to prepare each Wittig reagent:

Possibility [1]



1° halide preferred path




Possibility [2]



2° halide

Because the synthesis of the Wittig reagent begins with an S_N2 reaction, **the preferred pathway begins with an unhindered methyl halide or 1° alkyl halide.** In this example, retrosynthetic analysis of both Wittig reagents indicates that only one of them ($Ph_3P{=}CHCH_3$) can be synthesized from a 1° alkyl halide, making Possibility [1] the preferred pathway.

Problem 21.20 What starting materials are needed to prepare each alkene by a Wittig reaction? When there are two possible routes, indicate which route, if any, is preferred.

a.  b.  c. 

21.10D Comparing Methods of Alkene Synthesis

An advantage in using the Wittig reaction over other elimination methods to synthesize alkenes is that **you always know the location of the double bond.** Whereas other methods of alkene synthesis often give a mixture of constitutional isomers, the Wittig reaction always gives a single constitutional isomer.


For example, two methods can be used to convert cyclohexanone into alkene **B** (methylenecyclohexane): a two-step method consisting of Grignard addition followed by dehydration, or a one-step Wittig reaction.



cyclohexanone **B**


Recall from Section 9.8 that the major product formed in acid-catalyzed dehydration of an alcohol is the more substituted alkene.

In a two-step method, treatment of cyclohexanone with CH_3MgBr forms a 3° alcohol after protonation. Dehydration of the alcohol with H_2SO_4 forms a mixture of alkenes, in which the desired disubstituted alkene is the minor product.




cyclohexanone | 3° alcohol | trisubstituted C=C **major product** | disubstituted C=C **minor product**


By contrast, reaction of cyclohexanone with $Ph_3P{=}CH_2$ affords the desired alkene as the only product. The newly formed double bond always joins the carbonyl carbon with the negatively charged carbon of the Wittig reagent. In other words, **the position of the double bond is always unambiguous in the Wittig reaction.** This makes the Wittig reaction an especially attractive method for preparing many alkenes.



cyclohexanone **B** only product


Problem 21.21 Show two methods to synthesize each alkene: a one-step method using a Wittig reagent, and a two-step method that forms a carbon–carbon bond with an organometallic reagent in one of the steps.

a. 

b. 

21.11 Addition of 1° Amines


We now move on to the reaction of aldehydes and ketones with nitrogen and oxygen heteroatoms. **Amines are organic nitrogen compounds that contain a nonbonded electron pair on the N atom.** As we learned in Section 3.2, amines are classified as 1°, 2°, or 3° by the number of alkyl groups bonded to the *nitrogen* atom.




Both 1° and 2° amines react with aldehydes and ketones. We begin by examining the reaction of aldehydes and ketones with 1° amines.

21.11A Formation of Imines

Treatment of an aldehyde or ketone with a 1° amine affords an **imine** (also called a **Schiff base**). Nucleophilic attack of the 1° amine on the carbonyl group forms an unstable **carbinolamine,** which loses water to form an imine. The overall reaction results in **replacement of C=O by C=NR.**




Because the N atom of an imine is surrounded by three groups (two atoms and a lone pair), it is sp^2 hybridized, making the C−N−R" bond angle ~120° (*not* 180°). Imine formation is fastest when the reaction medium is weakly acidic.



The mechanism of imine formation (Mechanism 21.5) can be divided into two distinct parts: **nucleophilic addition of the 1° amine (Steps [1] and [2]), followed by elimination of H_2O (Steps [3]–[5]).** Each step involves a reversible equilibrium, so that the reaction is driven to completion by removing H_2O.

Imine formation is most rapid at pH 4–5. Mild acid is needed for protonation of the hydroxy group in Step [3] to form a **good leaving group.** Under strongly acidic conditions, the reaction rate decreases because the amine nucleophile is protonated. With no free electron pair, it is no longer a nucleophile, and so nucleophilic addition cannot occur.

Mechanism 21.5 Imine Formation from an Aldehyde or Ketone




1 – 2 Nucleophilic attack of the amine followed by proton transfer forms the carbinolamine.

3 Protonation of the OH group forms a good leaving group.


4 Loss of H_2O forms a resonance-stabilized iminium ion.

5 Loss of a proton forms the imine.


Problem 21.22 Draw the product formed when $CH_3CH_2CH_2CH_2NH_2$ reacts with each carbonyl compound in the presence of mild acid.


a. 

b. 

c. 


Problem 21.23 What 1° amine and carbonyl compound are needed to prepare each imine?

a. 

b. 


21.11B Application: Retinal, Rhodopsin, and the Chemistry of Vision

Many imines play vital roles in biological systems. A key molecule in the chemistry of vision is the highly conjugated imine **rhodopsin,** which is synthesized in the rod cells of the eye from **11-*cis*-retinal** and a 1° amine in the protein **opsin.**



11-*cis*-retinal

Figure 21.9
The key reaction in the chemistry of vision



- Rhodopsin is a light-sensitive compound located in the membrane of the rod cells in the retina of the eye. Rhodopsin contains the protein opsin bonded to 11-*cis*-retinal via an imine linkage. When light strikes this molecule, the crowded 11-cis double bond isomerizes to the 11-trans isomer, and a nerve impulse is transmitted to the brain by the optic nerve.


The central role of rhodopsin in the visual process was delineated by Nobel Laureate George Wald of Harvard University.

The complex process of vision centers around this imine derived from retinal (Figure 21.9). The 11-cis double bond in rhodopsin creates crowding in the rather rigid side chain. When light strikes the rod cells of the retina, it is absorbed by the conjugated double bonds of rhodopsin, and the 11-cis double bond is isomerized to the 11-trans arrangement. This isomerization is accompanied by a drastic change in shape in the protein, altering the concentration of Ca^{2+} ions moving across the cell membrane, and sending a nerve impulse to the brain, which is then processed into a visual image.

21.12 Addition of 2° Amines


21.12A Formation of Enamines

A 2° amine reacts with an aldehyde or ketone to give an **enamine.** ***Enamines*** **have a nitrogen atom bonded to a double bond** (alk***ene*** **+** ***amine*** **=** ***enamine***).




R' = H or alkyl carbinolamine enamine

Like imines, enamines are also formed by the addition of a nitrogen nucleophile to a carbonyl group followed by elimination of water. In this case, however, elimination occurs across two adjacent *carbon* atoms to form a new carbon–carbon π bond.



The mechanism for enamine formation (Mechanism 21.6) is identical to the mechanism for imine formation except for the last step, involving formation of the π bond. The mechanism can be divided into two distinct parts: **nucleophilic addition of the 2° amine (Steps [1] and [2]), followed by elimination of H_2O (Steps [3]–[5]).** Each step involves a reversible equilibrium once again, so that the reaction is driven to completion by removing H_2O.

Mechanism 21.6 Enamine Formation from an Aldehyde or Ketone



1 – 2 Nucleophilic attack of the amine followed by proton transfer forms the carbinolamine.


3 Protonation of the OH group forms a good leaving group.

4 Loss of H_2O forms a resonance-stabilized iminium ion.

5 Loss of a proton from the adjacent C–H bond forms the enamine.

The mechanisms illustrate why **the reaction of 1° amines with carbonyl compounds forms *imines*, but the reaction with 2° amines forms *enamines*.** In Figure 21.10, the last step of both mechanisms is compared using cyclohexanone as starting material. The position of the double bond depends on which proton is removed in the last step. Removal of an N–H proton forms a C=N, whereas removal of a C–H proton forms a C=C.

Figure 21.10
The formation of imines and enamines compared




- With a **1° amine,** the intermediate iminium ion still has a proton on the N atom that may be removed to form a C=N.
- With a **2° amine,** the intermediate iminium ion has no proton on the N atom. A proton must be removed from an adjacent C–H bond, and this forms a C=C.


21.12B Imine and Enamine Hydrolysis

Because imines and enamines are formed by a set of reversible reactions, both can be converted back to carbonyl compounds by hydrolysis with mild acid. The mechanism of these reactions is exactly the *reverse* of the mechanism written for the formation of imines and enamines. In the hydrolysis of enamines, the carbonyl carbon in the product comes from the sp^2 hybridized carbon bonded to the N atom in the starting material.

- **Hydrolysis of imines and enamines forms aldehydes and ketones.**




imine





enamine

Problem 21.24 What two enamines are formed when 2-methylcyclohexanone is treated with $(CH_3)_2NH$?


Problem 21.25 What carbonyl compound and amine are formed by the hydrolysis of each compound?

a. 

b. 


c. 

Problem 21.26 Draw a stepwise mechanism for the following imine hydrolysis.



21.13 Addition of H_2O—Hydration

Treatment of a carbonyl compound with H_2O in the presence of an acid or base catalyst **adds the elements of H and OH across the carbon–oxygen π bond,** forming a ***gem*-diol** or **hydrate.**



R' = H or alkyl

gem-diol (hydrate)

Hydration of a carbonyl group gives a good yield of *gem*-diol only with an unhindered aldehyde like formaldehyde, and with aldehydes containing nearby electron-withdrawing groups.




formaldehyde — formaldehyde hydrate — chloral — chloral hydrate

21.13A The Thermodynamics of Hydrate Formation

Whether addition of H_2O to a carbonyl group affords a good yield of the *gem*-diol depends on the relative energies of the starting material and the product. With less stable carbonyl starting materials, equilibrium favors the hydrate product, whereas with more stable carbonyl starting materials, equilibrium favors the carbonyl starting material. Because **alkyl groups stabilize a carbonyl group** (Section 20.2B):

- **Increasing the number of alkyl groups on the carbonyl carbon decreases the amount of hydrate at equilibrium.**

This can be illustrated by comparing the amount of hydrate formed from formaldehyde, acetaldehyde, and acetone.




Formaldehyde, the least stable carbonyl compound, forms the largest percentage of hydrate. On the other hand, acetone and other ketones, which have two electron-donor R groups, form < 1% of the hydrate at equilibrium. Other electronic factors come into play as well.

- **Electron-donating groups near the carbonyl carbon stabilize the carbonyl group, decreasing the amount of the hydrate at equilibrium.**
- **Electron-withdrawing groups near the carbonyl carbon destabilize the carbonyl group, increasing the amount of hydrate at equilibrium.**

This explains why chloral (trichloroacetaldehyde) forms a large amount of hydrate at equilibrium. Three electron-withdrawing Cl atoms place a partial positive charge on the α carbon to the carbonyl, destabilizing the carbonyl group, and therefore increasing the amount of hydrate at equilibrium.


Chloral hydrate, a sedative sometimes administered to calm a patient prior to a surgical procedure, has also been used for less reputable purposes. Adding it to an alcoholic beverage makes a so-called knock-out drink, causing an individual who drinks it to pass out. Because it is addictive and care must be taken in its administration, it is a controlled substance.



chloral

Adjacent like charges (δ+) *destabilize* the carbonyl and *increase* the amount of hydrate.


Problem 21.27 Which compound in each pair forms the higher percentage of *gem*-diol at equilibrium?



21.13B The Kinetics of Hydrate Formation

Although H_2O itself adds slowly to a carbonyl group, both acid and base catalyze the addition. In base, the nucleophile is ^-OH, and the mechanism follows the usual two steps for nucleophilic addition: **nucleophilic attack followed by protonation,** as shown in Mechanism 21.7.

Mechanism 21.7 Base-Catalyzed Addition of H_2O to a Carbonyl Group




R' = H or alkyl

1 The nucleophile (^-OH) attacks the carbonyl, breaking the π bond and moving an electron pair out on oxygen.

2 Protonation of the negatively charged oxygen by H_2O forms the hydration product.

The acid-catalyzed addition follows the general mechanism presented in Section 21.7A. For a poorer nucleophile like H_2O to attack a carbonyl group, the **carbonyl must be protonated by acid first; thus, protonation *precedes* nucleophilic attack.** The overall mechanism has three steps, as shown in Mechanism 21.8.

Mechanism 21.8 Acid-Catalyzed Addition of H_2O to a Carbonyl Group



R' = H or alkyl

1 Protonation of the carbonyl oxygen forms a resonance-stabilized cation.


2 – 3 Nucleophilic attack and deprotonation form the *gem-diol*. The overall result is addition of H and OH to the carbonyl group.

Acid and base increase the rate of reaction for different reasons.

- **Base converts H_2O into ^-OH, a *stronger nucleophile.***
- **Acid protonates the carbonyl group, making it *more electrophilic* towards nucleophilic attack.**

These catalysts increase the rate of the reaction, but they do not affect the equilibrium constant. Starting materials that give a low yield of *gem*-diol do so whether or not a catalyst is present. Because these reactions are reversible, the conversion of *gem*-diols to aldehydes and ketones is also catalyzed by acid and base, and the steps of the mechanism are reversed.


Problem 21.28 Draw a stepwise mechanism for the following reaction.



21.14 Addition of Alcohols—Acetal Formation

The term *acetal* refers to any compound derived from an aldehyde or ketone, having two OR groups bonded to a single carbon. The term *ketal* is sometimes used when the starting carbonyl compound is a ketone; that is, the carbon bonded to the alkoxy groups is *not* bonded to a H atom and the general structure is $R_2C(OR')_2$. Because ketals are considered a subclass of acetals in the IUPAC system, we will use the single general term *acetal* for any compound having two OR groups on a carbon atom.


Aldehydes and ketones react with *two* equivalents of alcohol to form acetals. In an acetal, the carbonyl carbon from the aldehyde or ketone is now singly bonded to two OR" (alkoxy) groups.



R' = H or alkyl


acetal

This reaction differs from other additions we have seen thus far, because **two equivalents of alcohol are added to the carbonyl group,** and two new C−O σ bonds are formed. Acetal formation is catalyzed by acids, commonly *p*-toluenesulfonic acid (TsOH).




acetal

When a diol such as ethylene glycol is used in place of two equivalents of ROH, a cyclic acetal is formed. Both oxygen atoms in the cyclic acetal come from the diol.



a cyclic acetal


Acetals are *not* ethers, even though both functional groups contain a C−O σ bond. Having two C−O σ bonds on the *same* carbon atom makes an acetal very different from an ether.



acetal ether

Like *gem*-diol formation, the synthesis of acetals is reversible, and often the equilibrium favors reactants, not products. In acetal synthesis, however, water is formed as a by-product, so the equilibrium can be driven to the right by removing the water as it is formed. This can be done in a variety of ways in the laboratory. A drying agent can be added that reacts with the water, or more commonly, the water can be distilled from the reaction mixture as it

Figure 21.11
A Dean–Stark trap for removing water




- **A Dean–Stark trap is an apparatus used for removing water from a reaction mixture.** To use a Dean–Stark trap to convert a carbonyl compound to an acetal:

  The carbonyl compound, an alcohol, and an acid are dissolved in benzene. As the mixture is heated, the carbonyl compound is converted to the acetal with water as a by-product. Benzene and water co-distill from the reaction mixture. When the hot vapors reach the cold condenser, they condense, forming a liquid that then collects in the glass tube below. Water, the more dense liquid, forms the lower layer, so that as it collects, it can be drained through the stopcock into a flask. In this way, water can be removed from a reaction mixture, driving the equilibrium.


is formed by using a Dean–Stark trap, as pictured in Figure 21.11. Driving an equilibrium to the right by removing one of the products is an application of Le Châtelier's principle (see Section 9.8).

Problem 21.29 Draw the products of each reaction.



21.14A The Mechanism

The mechanism for acetal formation can be divided into two parts: **the addition of one equivalent of alcohol** to form a **hemiacetal,** followed by the **conversion of the hemiacetal** to the **acetal.** A **hemiacetal** has a carbon atom bonded to one OH group and one OR group.




Like *gem*-diols, hemiacetals are often higher in energy than their carbonyl starting materials, making the direction of equilibrium unfavorable for hemiacetal formation. The elimination of H_2O, which can be removed from the reaction mixture to drive the equilibrium to favor product, occurs during the conversion of the hemiacetal to the acetal. This explains why two equivalents of ROH react with a carbonyl compound, forming the acetal as product.

Mechanism 21.9 is written in two parts with a general acid HA.

Mechanism 21.9 Acetal Formation


Part [1] Formation of a hemiacetal



1 Protonation of the carbonyl oxygen forms a resonance-stabilized cation.

2 – 3 Nucleophilic attack by R'OH and deprotonation form the hemiacetal. The overall result is addition of H and OR' to the carbonyl group.

Part [2] Formation of an acetal




4 Protonation of the OH group of the hemiacetal forms a good leaving group.

5 Loss of H_2O forms a resonance-stabilized cation.


6 – 7 Nucleophilic attack by R'OH followed by loss of a proton forms the acetal. The overall result of Part [2] is the addition of a second OR' group to the carbonyl.

Although this mechanism is lengthy—there are seven steps altogether—there are only three different kinds of reactions: **addition of a nucleophile, elimination of a leaving group, and proton transfer.** Steps [2] and [6] involve nucleophilic attack and Step [5] eliminates H_2O. The other four steps in the mechanism shuffle protons from one oxygen atom to another, to make a better leaving group or a more electrophilic carbonyl group.

Problem 21.30 Label each compound as an acetal, a hemiacetal, or an ether.




Problem 21.31 Draw a stepwise mechanism for the following reaction.




21.14B Hydrolysis of Acetals

Conversion of an aldehyde or ketone to an acetal is a **reversible reaction,** so **an acetal can be hydrolyzed to an aldehyde or ketone by treatment with aqueous acid.** Because this reaction is also an equilibrium process, it is driven to the right by using a large excess of water for hydrolysis.



The mechanism for this reaction is the reverse of acetal synthesis, as illustrated in Sample Problem 21.4.

Sample Problem 21.4 Draw a stepwise mechanism for the following reaction.




Solution

The mechanism is the reverse of acetal formation and involves two parts—conversion of the acetal to a hemiacetal, followed by conversion of the hemiacetal to the carbonyl compound.


Part [1] Conversion of the acetal to a hemiacetal

To convert this acetal to a hemiacetal, one molecule of CH_3OH must be eliminated and one molecule of H_2O must be added.



Part [2] Conversion of the hemiacetal to the carbonyl compound


To convert the hemiacetal to a carbonyl compound, one molecule of CH_3OH must be eliminated and the C–O π bond must be formed.




Steps [2] and [6] involve loss of the leaving group (CH_3OH), and Step [3] involves nucleophilic attack of H_2O. The other four steps in the mechanism shuffle protons from one oxygen atom to another. Steps [2] and [6] form resonance-stabilized carbocations, but only one resonance structure is drawn.


Acetal hydrolysis requires a strong acid to make a good leaving group (ROH). In Sample Problem 21.4, H_2SO_4 converts CH_3O^- into CH_3OH, a weak base and neutral leaving group. Acetal hydrolysis does not occur in base.

Problem 21.32 Draw the products of each reaction.

a. 

b. 

Problem 21.33 Safrole is a naturally occurring acetal isolated from sassafras plants. Once used as a common food additive in root beer and other beverages, it is now banned because it is carcinogenic. What compounds are formed when safrole is hydrolyzed with aqueous acid?




safrole



Sassafras, source of safrole

Problem 21.34 Identify the acetal in oleandrin, and draw the products formed by acid-catalyzed hydrolysis of the acetal.



oleandrin




Oleandrin, a toxic cardiac glycoside used in some countries for the treatment of congestive heart failure, is obtained from the sap of the oleander plant (*Nerium oleander*), a common ornamental plant in tropical and subtropical regions.

21.15 Acetals as Protecting Groups

Just as the *tert*-butyldimethylsilyl ethers are used as protecting groups for alcohols (Section 20.12), **acetals are valuable protecting groups for aldehydes and ketones.**

Suppose a starting material **A** contains both a ketone and an ester, and it is necessary to selectively reduce the ester to an alcohol (6-hydroxyhexan-2-one), leaving the ketone untouched. Such a selective reduction is *not* possible in one step. Because ketones are more readily reduced, methyl 5-hydroxyhexanoate is formed instead.




To solve this problem we can use a protecting group to block the more reactive ketone carbonyl group. The overall process requires three steps.

[1] Protect the interfering functional group—the ketone carbonyl.

[2] Carry out the desired reaction—reduction.

[3] Remove the protecting group.


The following three-step sequence using a cyclic acetal leads to the desired product.



- **Step [1]** The ketone carbonyl is protected as a cyclic acetal by reaction of the starting material with $HOCH_2CH_2OH$ and TsOH.
- **Step [2]** Reduction of the ester is then carried out with $LiAlH_4$, followed by treatment with H_2O.
- **Step [3]** The acetal is then converted back to a ketone carbonyl group with aqueous acid.

Acetals are widely used protecting groups for aldehydes and ketones because they are easy to add and easy to remove, and they are stable to a wide variety of reaction conditions. Acetals do not react with base, oxidizing agents, reducing agents, or nucleophiles. Good protecting groups must survive a variety of reaction conditions that take place at other sites in a molecule, but they must also be selectively removed under mild conditions when needed.


Problem 21.35 How would you use a protecting group to carry out the following transformation?



21.16 Cyclic Hemiacetals

Cyclic hemiacetals are also called **lactols.**

Although acyclic hemiacetals are generally unstable and therefore not present in appreciable amounts at equilibrium, **cyclic hemiacetals containing five- and six-membered rings are stable compounds** that are readily isolated.



hemiacetal

One C is bonded to:
- an **OH** group
- an **OR** group




cyclic hemiacetals

Each C is bonded to:
- an **OH** group
- an **OR** group that is part of a ring

21.16A Forming Cyclic Hemiacetals


All hemiacetals are formed by nucleophilic addition of a hydroxy group to a carbonyl group. In the same way, cyclic hemiacetals are formed by **intramolecular cyclization of hydroxy aldehydes.**



[Equilibrium proportions of each compound are given.]


Such intramolecular reactions to form five- and six-membered rings are faster than the corresponding intermolecular reactions. The two reacting functional groups, in this case OH and C=O, are held in close proximity, increasing the probability of reaction.

Problem 21.36 What lactol (cyclic hemiacetal) is formed from intramolecular cyclization of each hydroxy aldehyde?



Hemiacetal formation is catalyzed by both acid and base. The acid-catalyzed mechanism is identical to Part [1] of Mechanism 21.9, except that the reaction occurs in an **intramolecular** fashion, as shown for the acid-catalyzed cyclization of 5-hydroxypentanal to form a six-membered cyclic hemiacetal in Mechanism 21.10.


Mechanism 21.10 Acid-Catalyzed Cyclic Hemiacetal Formation



1 – 2 Protonation of the carbonyl oxygen followed by intramolecular nucleophilic attack forms the six-membered ring.


3 Deprotonation forms the neutral cyclic hemiacetal.

Intramolecular cyclization of a hydroxy aldehyde forms a **hemiacetal with a new stereogenic center, so that an equal amount of two enantiomers** results.




21.16B The Conversion of Hemiacetals to Acetals

Cyclic hemiacetals can be converted to acetals by treatment with an alcohol and acid. This reaction converts the OH group that is part of the hemiacetal to an OR group.



Mechanism 21.11, which is similar to Part [2] of Mechanism 21.9, illustrates the conversion of a cyclic hemiacetal to an acetal.

Mechanism 21.11 A Cyclic Acetal from a Cyclic Hemiacetal



1 Protonation of the OH group of the hemiacetal forms a good leaving group.


2 Loss of H_2O forms a resonance-stabilized cation.

3 – 4 Nucleophilic attack by CH_3OH followed by loss of a proton forms the acetal.


The overall result of this reaction is the **replacement of the hemiacetal OH group by an OCH_3 group.** This substitution reaction readily occurs because the carbocation formed in Step [2] is stabilized by resonance. This fact makes the OH group of a hemiacetal different from the hydroxy group in other alcohols.

Thus, when a compound that contains both an alcohol OH group and a hemiacetal OH group is treated with an alcohol and acid, only the hemiacetal OH group reacts to form an acetal. The alcohol OH group does *not* react.


The conversion of cyclic hemiacetals to acetals is an important reaction in carbohydrate chemistry, as discussed in Chapter 28.



Problem 21.37 Two naturally occurring compounds that contain stable cyclic hemiacetals and acetals are monensin and digoxin, the chapter-opening molecule. Monensin, a polyether antibiotic produced by *Streptomyces cinamonensis,* is used as an additive in cattle feed. Digoxin is a widely prescribed cardiac drug used to increase the force of heart contractions. Label each acetal, hemiacetal, and ether in both compounds.




monensin



digoxin

Problem 21.38 Draw the products of each reaction.




21.17 An Introduction to Carbohydrates


Carbohydrates, commonly referred to as sugars and starches, are polyhydroxy aldehydes and ketones, or compounds that can be hydrolyzed to them. Along with proteins, fatty acids, and nucleotides, they form one of the four main groups of biomolecules responsible for the structure and function of all living cells.

Many carbohydrates contain cyclic acetals or hemiacetals. Examples include **glucose,** the most common simple sugar, and **lactose,** the principal carbohydrate in milk. Hemiacetal carbons are labeled in blue, whereas the acetal carbon is labeled in red.


Glucose is the carbohydrate that is transported in the blood to individual cells. The hormone insulin regulates the level of glucose in the blood. Diabetes is a common disease that results from a deficiency of insulin, resulting in increased glucose levels in the blood and other metabolic abnormalities. Insulin injections control glucose levels.



3-D structure



β-D-glucose
(one form of glucose)




lactose

3-D structure

Hemiacetals in sugars are formed in the same way that other hemiacetals are formed—that is, by **cyclization of hydroxy aldehydes.** Thus, the hemiacetal of glucose is formed by cyclization

of an acyclic *poly*hydroxy aldehyde **A,** as shown in the accompanying equation. This process illustrates two important features.




- When the OH group on C5 is the nucleophile, **cyclization yields a six-membered ring,** and this ring size is preferred.
- **Cyclization forms a new stereogenic center** (labeled in blue), exactly analogous to the cyclization of the simpler hydroxy aldehyde (5-hydroxypentanal) in Section 21.16A. **The new OH group of the hemiacetal can occupy either the equatorial or axial position.**

For glucose, this results in two cyclic forms, called **β-D-glucose** (having an equatorial OH group) and **α-D-glucose** (having an axial OH group). Because β-D-glucose has the new OH group in the more roomy equatorial position, this cyclic form of glucose is the major product. At equilibrium, only a trace of the acyclic hydroxy aldehyde **A** is present.

Many more details on this process and other aspects of carbohydrate chemistry are presented in Chapter 28.

Problem 21.39



α-D-galactose

a. How many stereogenic centers are present in α-D-galactose?
b. Label the hemiacetal carbon in α-D-galactose.
c. Draw the structure of β-D-galactose.
d. Draw the structure of the polyhydroxy aldehyde that cyclizes to α- and β-D-galactose.
e. From what you learned in Section 21.16B, what product(s) is (are) formed when α-D-galactose is treated with CH_3OH and an acid catalyst?

KEY CONCEPTS

Aldehydes and Ketones—Nucleophilic Addition

General Facts


- Aldehydes and ketones contain a carbonyl group bonded to only H atoms or R groups. The carbonyl carbon is sp^2 hybridized and trigonal planar (21.1).
- Aldehydes are identified by the suffix *-al,* whereas ketones are identified by the suffix *-one* (21.2).
- Aldehydes and ketones are polar compounds that exhibit dipole–dipole interactions (21.3).

Summary of Spectroscopic Absorptions of RCHO and R_2CO (21.4)

IR absorptions	C=O	~1715 cm^{-1} for ketones • Frequency increases with decreasing ring size. ~1730 cm^{-1} for aldehydes • For both RCHO and R_2CO, the frequency decreases with conjugation.
	C_{sp^2}–H of CHO	~2700–2830 cm^{-1} (one or two peaks)
^{1}H NMR absorptions	CHO	9–10 ppm (highly deshielded proton)
	C–H α to C=O	2–2.5 ppm (somewhat deshielded C_{sp^3}–H)
^{13}C NMR absorption	C=O	190–215 ppm

Nucleophilic Addition Reactions


[1] Addition of hydride (H^-) (21.8)



R' = H or alkyl — 1° or 2° alcohol

- The mechanism has two steps.
- $H:^-$ adds to the planar C=O from both sides.


[2] Addition of organometallic reagents (R^-) (21.8)



R' = H or alkyl — 1°, 2°, or 3° alcohol

- The mechanism has two steps.
- $(R'')^-$ adds to the planar C=O from both sides.


[3] Addition of cyanide (^-CN) (21.9)



R' = H or alkyl — cyanohydrin

- The mechanism has two steps.
- ^-CN adds to the planar C=O from both sides.


[4] Wittig reaction (21.10)



R' = H or alkyl — Wittig reagent — alkene

- The reaction forms a new C–C σ bond and a new C–C π bond.
- $Ph_3P{=}O$ is formed as by-product.


[5] Addition of 1° amines (21.11)



R' = H or alkyl — imine

- The reaction is fastest at pH 4–5.
- The intermediate carbinolamine is unstable, and loses H_2O to form the C=N.


[6] Addition of 2° amines (21.12)



R' = H or alkyl — enamine

- The reaction is fastest at pH 4–5.
- The intermediate carbinolamine is unstable, and loses H_2O to form the C=C.


[7] Addition of H_2O—Hydration (21.13)



R' = H or alkyl — *gem*-diol

- The reaction is reversible. Equilibrium favors the product only with less stable carbonyl compounds (e.g., H_2CO and Cl_3CCHO).
- The reaction is catalyzed by either H^+ or ^-OH.

[8] Addition of alcohols (21.14)




R' = H or alkyl — acetal

- The reaction is reversible.
- The reaction is catalyzed by acid.
- Removal of H_2O drives the equilibrium to favor the products.


Other Reactions

[1] Synthesis of Wittig reagents (21.10A)




- Step [1] is best with CH_3X and RCH_2X because the reaction follows an S_N2 mechanism.
- A strong base is needed for proton removal in Step [2].

[2] Conversion of cyanohydrins to aldehydes and ketones (21.9)



- This reaction is the reverse of cyanohydrin formation.


[3] Hydrolysis of nitriles (21.9)



R' = H or alkyl

α-hydroxy carboxylic acid

[4] Hydrolysis of imines and enamines (21.12)




R' = H or alkyl
imine

enamine

aldehyde or ketone

[5] Hydrolysis of acetals (21.14)



R' = H or alkyl


aldehyde or ketone

- The reaction is acid catalyzed and is the reverse of acetal synthesis.
- A large excess of H_2O drives the equilibrium to favor the products.


PROBLEMS

Problems Using Three-Dimensional Models

21.40 (a) Give the IUPAC name for **A** and **B.** (b) Draw the product formed when **A** or **B** is treated with each reagent: [1] $NaBH_4$, CH_3OH; [2] CH_3MgBr, then H_2O; [3] $Ph_3P{=}CHOCH_3$; [4] $CH_3CH_2CH_2NH_2$, mild acid; [5] $HOCH_2CH_2CH_2OH$, H^+.





A




B


21.41 Rank the following compounds in order of increasing reactivity in nucleophilic addition.









21.42 What carbonyl compound and diol are needed to prepare each compound?


a. 


b. 


Nomenclature


21.43 Give the IUPAC name for each compound.


a. 

c. 

e. 

b. 

d. 


f. 


21.44 Give the structure corresponding to each name.

a. 2-methyl-3-phenylbutanal
b. 3,3-dimethylcyclohexanecarbaldehyde
c. 3-benzoylcyclopentanone
d. 2-formylcyclopentanone
e. (*R*)-3-methylheptan-2-one
f. *m*-acetylbenzaldehyde
g. 2-*sec*-butylcyclopent-3-enone
h. 5,6-dimethylcyclohex-1-enecarbaldehyde


Reactions


21.45 Draw the products formed in each reaction sequence.


a. 


b. 


21.46 Draw the products of each reaction.


a. 


b. 


c. 

d. 


e. 


f. 


g. 

h. 


21.47 What carbonyl compound and alcohol are formed by hydrolysis of each acetal?


a. 


b. 


c. 

21.48 Draw all stereoisomers formed in each reaction.

a. 

b. 


c. 


d. 

21.49 Hydroxy aldehydes **A** and **B** readily cyclize to form hemiacetals. Draw the stereoisomers formed in this reaction from both **A** and **B.** Explain why this process gives an optically inactive product mixture from **A,** and an optically active product mixture from **B.**




21.50 What products are formed when each acetal is hydrolyzed with aqueous acid?


a. 

b. 

21.51 What hydrolysis products are formed when the following compound is treated with aqueous acid?



21.52 Etoposide, sold as a phosphate derivative with the trade name of Etopophos, is used for the treatment of lung cancer, testicular cancer, and lymphomas. (a) Locate the acetals in etoposide. (b) What products are formed when all of the acetals are hydrolyzed with aqueous acid?



Properties of Aldehydes and Ketones




21.53 Which compound forms the higher concentration of hydrate at equilibrium, PhCOCHO or $PhCH_2CHO$? Explain your reasoning.

21.54 Consider the para-substituted aromatic ketones, $NO_2C_6H_4COCH_3$ (*p*-nitroacetophenone) and $CH_3OC_6H_4COCH_3$ (*p*-methoxyacetophenone).


a. Which carbonyl compound is more stable?
b. Which compound forms the higher percentage of hydrate at equilibrium?
c. Which compound exhibits a carbonyl absorption at higher wavenumber in its IR spectrum? Explain your reasoning in each part.


Synthesis


21.55 What Wittig reagent and carbonyl compound are needed to prepare each alkene? When two routes are possible, indicate which route, if any, is preferred.


a. 
b. 
c. 

21.56 What carbonyl compound and amine or alcohol are needed to prepare each product?

a. 

b. 



c. 

d. 


21.57 Show two different methods to carry out the following transformation: a one-step method using a Wittig reagent, and a two-step method using a Grignard reagent. Which route, if any, is preferred?




21.58 Devise a synthesis of each alkene using a Wittig reaction to form the double bond. You may use benzene and organic alcohols having four or fewer carbons as starting materials and any required reagents.


a. 
(+ *Z* isomer)
b. 

21.59 Devise a synthesis of each compound from cyclohexene and organic alcohols. You may use any other required organic or inorganic reagents.

a. 


b. 

21.60 Devise a synthesis of each compound from the given starting materials. You may also use organic alcohols having four or fewer carbons, and any organic or inorganic reagents.


a. 


(+ *Z* isomer)

b. 

c. 

21.61 Devise a synthesis of each compound from ethanol (CH_3CH_2OH) as the only source of carbon atoms. You may use any other organic or inorganic reagents you choose.

a. 

b. 


Protecting Groups

21.62 Design a stepwise synthesis to convert cyclopentanone and 4-bromobutanal to hydroxy aldehyde **A.**



cyclopentanone 4-bromobutanal **A**

21.63 Besides the *tert*-butyldimethylsilyl ethers introduced in Chapter 20, there are many other widely used alcohol protecting groups. For example, an alcohol such as cyclohexanol can be converted to a **m**eth**o**xy **m**ethyl ether (a MOM protecting group) by treatment with base and chloromethyl methyl ether, $ClCH_2OCH_3$. The protecting group can be removed by treatment with aqueous acid.




cyclohexanol methoxy methyl ether


a. Write a stepwise mechanism for the formation of a MOM ether from cyclohexanol.
b. What functional group comprises a MOM ether?
c. Besides cyclohexanol, what other products are formed by aqueous hydrolysis of the MOM ether? Draw a stepwise mechanism that accounts for formation of each product.

Mechanism

21.64 Draw a stepwise mechanism for the following reaction.




21.65 Draw a stepwise mechanism for the following reaction, a key step in the synthesis of the anti-inflammatory drug celecoxib (trade name Celebrex).




celecoxib


21.66 Treatment of $(HOCH_2CH_2CH_2CH_2)_2CO$ with acid forms a product of molecular formula $C_9H_{16}O_2$ and a molecule of water. Draw the structure of the product and explain how it is formed.

21.67 Draw a stepwise mechanism for each reaction.


a. 

b. 

21.68 Draw a stepwise mechanism for the following reaction that converts a dicarbonyl compound to a furan.




21.69 Draw a stepwise mechanism for the following reaction, a key step in the synthesis of ticlopidine, a drug that inhibits platelet aggregation. Ticlopidine has been used to reduce the risk of stroke in patients who cannot tolerate aspirin.




ticlopidine

21.70 Salsolinol is a naturally occurring compound found in bananas, chocolate, and several foods derived from plant sources. Salsolinol is also formed in the body when acetaldehyde, an oxidation product of the ethanol ingested in an alcoholic beverage, reacts with dopamine, a neurotransmitter. Draw a stepwise mechanism for the formation of salsolinol in the following reaction.




dopamine acetaldehyde salsolinol

21.71 (a) Explain how $NaBH_4$ in CH_3OH can reduce hemiacetal **A** to butane-1,4-diol ($HOCH_2CH_2CH_2CH_2OH$). (b) What product is formed when **A** is treated with $Ph_3P{=}CHCH_2CH(CH_3)_2$? (c) The drug isotretinoin is formed by reaction of **X** and **Y.** What is the structure of isotretinoin? Although isotretinoin (trade name Accutane or Roaccutane) is used for the treatment of severe acne, it is dispensed under strict controls because it also causes birth defects.



A **X** **Y**

21.72 Reaction of 5,5-dimethoxypentan-2-one with methylmagnesium iodide followed by treatment with aqueous acid forms cyclic hemiacetal **Y.** Draw a stepwise mechanism that illustrates how **Y** is formed.



5,5-dimethoxypentan-2-one **Y**

Spectroscopy

21.73 Although the carbonyl absorption of cyclic ketones generally shifts to higher wavenumber with decreasing ring size, the C=O of cyclopropenone absorbs at lower wavenumber in its IR spectrum than the C=O of cyclohex-2-enone. Explain this observation by using the principles of aromaticity learned in Chapter 17.




cyclopropenone (1640 cm^{-1}) cyclohex-2-enone (1685 cm^{-1})


21.74 Use the ^1H NMR and IR data to determine the structure of each compound.


Compound **A**	Molecular formula:	$C_5H_{10}O$
	IR absorptions at	1728, 2791, and 2700 cm^{-1}
	^1H NMR data:	1.08 (singlet, 9 H) and 9.48 (singlet, 1 H) ppm
Compound **B**	Molecular formula:	$C_5H_{10}O$
	IR absorption at	1718 cm^{-1}
	^1H NMR data:	1.10 (doublet, 6 H), 2.14 (singlet, 3 H), and 2.58 (septet, 1 H) ppm
Compound **C**	Molecular formula:	$C_{10}H_{12}O$
	IR absorption at	1686 cm^{-1}
	^1H NMR data:	1.21 (triplet, 3 H), 2.39 (singlet, 3 H), 2.95 (quartet, 2 H), 7.24 (doublet, 2 H), and 7.85 (doublet, 2 H) ppm
Compound **D**	Molecular formula:	$C_{10}H_{12}O$
	IR absorption at	1719 cm^{-1}
	^1H NMR data:	1.02 (triplet, 3 H), 2.45 (quartet, 2 H), 3.67 (singlet, 2 H), and 7.06–7.48 (multiplet, 5 H) ppm


21.75 A solution of acetone [$(CH_3)_2C=O$] in ethanol (CH_3CH_2OH) in the presence of a trace of acid was allowed to stand for several days, and a new compound of molecular formula $C_7H_{16}O_2$ was formed. The IR spectrum showed only one major peak in the functional group region around 3000 cm^{-1}, and the ^1H NMR spectrum is given here. What is the structure of the product?




21.76 Compounds **A** and **B** have molecular formula $C_9H_{10}O$. Identify their structures from the ^{1}H NMR and IR spectra given.











21.77 An unknown compound **C** of molecular formula $C_6H_{12}O_3$ exhibits a strong absorption in its IR spectrum at 1718 cm^{-1} and the given ^{1}H NMR spectrum. What is the structure of **C?**





21.78 An unknown compound **D** exhibits a strong absorption in its IR spectrum at 1692 cm^{-1}. The mass spectrum of **D** shows a molecular ion at *m/z* = 150 and a base peak at 121. The ^1H NMR spectrum of **D** is shown below. What is the structure of **D?**




Carbohydrates

21.79 Draw the structure of the acyclic polyhydroxy aldehyde that cyclizes to each hemiacetal.

a. 

b. 


21.80 β-D-Glucose, a hemiacetal, can be converted to a mixture of acetals on treatment with CH_3OH in the presence of acid. Draw a stepwise mechanism for this reaction. Explain why two acetals are formed from a single starting material.



β-D-glucose

Challenge Problems


21.81 Draw a stepwise mechanism for the following reaction.



21.82 Brevicomin, the aggregation pheromone of the western pine bark beetle, contains a bicyclic bridged ring system and is prepared by the acid-catalyzed cyclization of 6,7-dihydroxynonan-2-one.

a. Suggest a structure for brevicomin.


b. Devise a synthesis of 6,7-dihydroxynonan-2-one from 6-bromohexan-2-one. You may also use three-carbon alcohols and any required organic or inorganic reagents.




6-bromohexan-2-one

6,7-dihydroxynonan-2-one

21.83 Draw a stepwise mechanism for the following reaction.




21.84 Maltose is a carbohydrate present in malt, the liquid obtained from barley and other grains. Although maltose has numerous functional groups, its reactions are explained by the same principles we have already encountered.




a. Label the acetal and hemiacetal carbons.
b. What products are formed when maltose is treated with each of the following reagents: [1] H_3O^+; [2] CH_3OH and HCl; [3] excess NaH, then excess CH_3I?
c. Draw the products formed when the compound formed in Reaction [3] of part (b) is treated with aqueous acid.

The reactions in parts (b) and (c) are used to determine structural features of carbohydrates like maltose. We will learn much more about maltose and similar carbohydrates in Chapter 28.

21.85 Identify **R** and **S** in the following reaction sequence, and draw a mechanism for the conversion of **R** to **S** (molecular formula $C_6H_{10}O_3$). **S** was used in the synthesis of darunavir (trade name Prezista), used to treat HIV.



21.86 Draw a stepwise mechanism for the following reaction, a key step in the synthesis of conivaptan (trade name Vaprisol), a drug used in the treatment of low sodium levels.



22 Carboxylic Acids and Their Derivatives—Nucleophilic Acyl Substitution

22.1 Introduction
22.2 Structure and bonding
22.3 Nomenclature
22.4 Physical properties
22.5 Spectroscopic properties
22.6 Interesting esters and amides
22.7 Introduction to nucleophilic acyl substitution
22.8 Reactions of acid chlorides
22.9 Reactions of anhydrides
22.10 Reactions of carboxylic acids
22.11 Reactions of esters
22.12 Application: Lipid hydrolysis
22.13 Reactions of amides
22.14 Application: The mechanism of action of β-lactam antibiotics
22.15 Summary of nucleophilic acyl substitution reactions
22.16 Natural and synthetic fibers
22.17 Biological acylation reactions
22.18 Nitriles




Ginkgolide B, a major constituent of the extracts of the ginkgo tree *Ginkgo biloba,* is a complex molecule whose structure was determined in 1967. Although its compact, rigid ring system of 20 carbons made it a challenging molecule to prepare in the laboratory, Nobel Laureate E. J. Corey and co-workers reported its synthesis in 1988. Ginkgo extracts are widely used herbal supplements, taken to enhance memory and treat dementia. Recent findings of the National Institutes of Health, however, have cast doubt on their efficacy in providing long-term improvement in cognitive function. In Chapter 22 we learn about the reactions of esters, three of the many functional groups in ginkgolide B.

Chapter 22 continues the study of carbonyl compounds with a detailed look at **nucleophilic acyl substitution,** a key reaction of carboxylic acids and their derivatives. Substitution at sp^2 hybridized carbon atoms was introduced in Chapter 20 with reactions involving carbon and hydrogen nucleophiles. In Chapter 22, we learn that nucleophilic acyl substitution is a general reaction that occurs with a variety of heteroatomic nucleophiles. This reaction allows the conversion of one carboxylic acid derivative into another. ***Every* reaction in Chapter 22 that begins with a carbonyl compound involves nucleophilic substitution.** Chapter 22 also discusses the properties and chemical reactions of **nitriles,** compounds that contain a carbon–nitrogen triple bond. Nitriles are in the same carbon oxidation state as carboxylic acids, and they undergo reactions that form related products.











22.1 Introduction

Chapter 22 focuses on carbonyl compounds that contain an **acyl group bonded to an electronegative atom.** These include the **carboxylic acids,** as well as carboxylic acid derivatives that can be prepared from them: **acid chlorides, anhydrides, esters, and amides.**




acyl group

Z = an atom more electronegative than carbon


Z = OH	 carboxylic acid	 R = CH_3 acetic acid
Z = Cl	 acid chloride	 R = CH_3 acetyl chloride
Z = OCOR	 anhydride	 R = CH_3 acetic anhydride
Z = OR'	 ester	 R = R' = CH_3 methyl acetate
Z = NR'_2	 R' = H or alkyl amide	 R = CH_3, R' = H acetamide

Anhydrides contain two carbonyl groups joined by a single oxygen atom. **Symmetrical anhydrides** have two identical alkyl groups bonded to the carbonyl carbons, and **mixed anhydrides** have two different alkyl groups. **Cyclic anhydrides** are also known.



symmetrical anhydride | mixed anhydride | cyclic anhydride

As we learned in Section 3.2, **amides** are classified as **1°, 2°,** or **3°** depending on the number of carbon atoms bonded directly to the *nitrogen* atom.



1° amide, 1 C—N bond | **2°** amide, 2 C—N bonds | **3°** amide, 3 C—N bonds


Cyclic esters and amides are called **lactones** and **lactams,** respectively. The ring size of the heterocycle is indicated by a Greek letter. An amide in a four-membered ring is called a **β-lactam,** because the β carbon to the carbonyl is bonded to the heteroatom. An ester in a five-membered ring is called a **γ-lactone.**



γ-lactone | δ-lactone | β-lactam | γ-lactam

Nucleophilic acyl substitution was first discussed in Chapter 20 with R^- and H^- as the nucleophiles. This substitution reaction is general for a variety of nucleophiles, making it possible to form many different substitution products, as discussed in Sections 22.8–22.13.

All of these compounds contain an acyl group bonded to an electronegative atom Z that can serve as a **leaving group.** As a result, these compounds undergo **nucleophilic acyl substitution.** Recall from Chapters 20 and 21 that aldehydes and ketones do *not* undergo nucleophilic substitution because they have no leaving group on the carbonyl carbon.



Z = OH, Cl, OCOR, OR', NR'_2


Nitriles **are compounds that contain a cyano group, C≡N, bonded to an alkyl group.** Nitriles have no carbonyl group, so they are structurally distinct from carboxylic acids and their derivatives. The carbon atom of the cyano group, however, has the same oxidation state as the carbonyl carbon of carboxylic acid derivatives, so there are certain parallels in their chemistry.



nitrile


Each labeled C has three bonds to a more electronegative element.

Problem 22.1 Oxytocin, sold under the trade name Pitocin, is a naturally occurring hormone used to stimulate uterine contractions and induce labor. Classify each amide in oxytocin as 1°, 2°, or 3°.




22.2 Structure and Bonding

The two most important features of any carbonyl group, regardless of the other groups bonded to it, are the following:



- **The carbonyl carbon is sp^2 hybridized and trigonal planar, making it relatively uncrowded.**
- **The electronegative oxygen atom polarizes the carbonyl group, making the carbonyl carbon electrophilic.**

Because carboxylic acid derivatives (RCOZ) all contain an atom Z with a nonbonded electron pair, three resonance structures can be drawn for RCOZ, compared to just two for aldehydes and ketones (Section 20.1). These three resonance structures stabilize RCOZ by delocalizing electron density. In fact, the more resonance structures **2** and **3** contribute to the resonance hybrid, the more stable RCOZ is.



- **The more basic Z is, the more it donates its electron pair, and the more resonance structure 3 contributes to the hybrid.**

To determine the relative basicity of the leaving group Z, we compare the pK_a values of the conjugate acids HZ, given in Table 22.1. The following order of basicity results:





Table 22.1 pK_a Values of the Conjugate Acids (HZ) for Common Z Groups of Acyl Compounds (RCOZ)

	Structure	Leaving group (Z^-)	Conjugate acid (HZ)	pK_a	
Increasing basicity of Z ↓	**RCOCl** acid chloride	Cl^-	HCl	–7	↑ Increasing acidity of HZ
	$(RCO)_2O$ anhydride	RCO_2^-	RCO_2H	3–5	
	RCO_2H carboxylic acid	^-OH	H_2O	15.7	
	RCO_2R' ester	$^-OR'$	R'OH	15.5–18	
	$RCONR'_2$ amide	$^-NR'_2$	R'_2NH	38–40	

Because the basicity of Z determines the relative stability of the carboxylic acid derivatives, the following **order of stability** results:




Thus, an acid chloride is the least stable carboxylic acid derivative because Cl^- is the weakest base. An amide is the most stable carboxylic acid derivative because $^-NR'_2$ is the strongest base.

- **In summary: As the basicity of Z increases, the stability of RCOZ increases because of added resonance stabilization.**

Problem 22.2 Draw the three possible resonance structures for an acid bromide, RCOBr. Then, using the pK_a values in Appendix A, decide if RCOBr is more or less stabilized by resonance than a carboxylic acid (RCOOH).

Problem 22.3 How do the following experimental results support the resonance description of the relative stability of acid chlorides compared to amides? The C – Cl bond lengths in CH_3Cl and CH_3COCl are identical (178 pm), but the C – N bond in $HCONH_2$ is shorter than the C – N bond in CH_3NH_2 (135 pm versus 147 pm).

The structure and bonding in nitriles is very different from the carboxylic acid derivatives, and resembles the carbon–carbon triple bond of alkynes.



- **The carbon atom of the C≡N group is *sp* hybridized, making it linear with a bond angle of 180°.**
- **The triple bond consists of one σ and two π bonds.**

Like the carboxylic acid derivatives, **nitriles contain an electrophilic carbon atom,** making them susceptible to nucleophilic attack.

22.3 Nomenclature

The names of carboxylic acid derivatives are formed from the names of the parent carboxylic acids discussed in Section 19.2. Keep in mind that the common names **formic acid, acetic acid,** and **benzoic acid** are virtually always used for the parent acid, so these common parent names are used for their derivatives as well.

22.3A Naming an Acid Chloride—RCOCl

Acid chlorides are named by naming the acyl group and adding the word ***chloride.*** Two different methods are used.


[1] For acyclic acid chlorides: Change the suffix *-ic acid* of the parent carboxylic acid to the suffix *-yl chloride*; or

[2] When the –COCl group is bonded to a ring: Change the suffix *-carboxylic acid* to *-carbonyl chloride*.




derived from acet*ic acid*

acetyl chloride



derived from cyclohexane*carboxylic acid*

cyclohexanecarbonyl chloride




derived from 2-methylbutan*oic acid*

2-methylbutanoyl chloride


22.3B Naming an Anhydride

The word *anhydride* means "without water." Removing one molecule of water from two molecules of carboxylic acid forms an anhydride.




anhydride

Symmetrical anhydrides are named by changing the ***acid*** ending of the parent carboxylic acid to the word ***anhydride.*** **Mixed anhydrides,** which are derived from two different carboxylic acids, are named by alphabetizing the names for both acids and replacing the word ***acid*** by the word ***anhydride.***



derived from acetic acid

acetic anhydride



derived from acetic acid and benzoic acid

acetic benzoic anhydride

22.3C Naming an Ester—RCOOR'


Esters are often written as RCOOR', where the alkyl group (R') is written *last.* When an ester is named, however, the R' group appears *first* in the name.


An ester has two parts to its structure, each of which must be named: an **acyl group (RCO−)** and an **alkyl group** (designated as **R'**) bonded to an oxygen atom.

- **In the IUPAC system, esters are identified by the suffix *-ate.***

How To Name an Ester (RCO_2R') Using the IUPAC System


Example Give a systematic name for each ester:


a. 

b. 

Step [1] **Name the R' group bonded to the oxygen atom as an alkyl group.**


- The name of the alkyl group, ending in the suffix *-yl,* becomes the *first* part of the ester name.






Step [2] **Name the acyl group (RCO−) by changing the *-ic acid* ending of the parent carboxylic acid to the suffix *-ate.***

- The name of the acyl group becomes the *second* part of the name.



derived from
acet*ic acid* ----→ acet*ate*

Answer: ethyl acetate




derived from
cyclohexanecarboxyl*ic acid* ----→ cyclohexanecarboxyl*ate*

Answer: *tert*-butyl cyclohexanecarboxylate


22.3D Naming an Amide

All 1° amides are named by replacing the ***-ic acid, -oic acid,*** or ***-ylic acid*** ending with the suffix **amide.**




derived from
acet*ic acid*

acetamide



derived from
benz*oic acid*

benzamide




derived from
2-methylcyclopentanecarbox*ylic acid*


2-methylcyclopentanecarboxamide

A 2° or 3° amide has two parts to its structure: an **acyl group** that contains the carbonyl group **(RCO−)** and one or two **alkyl groups** bonded to the nitrogen atom.

How To Name a 2° or 3° Amide

Example Give a systematic name for each amide:


a. 

b. 


How To, continued . . .

Step [1] **Name the alkyl group (or groups) bonded to the N atom of the amide. Use the prefix "*N*-" preceding the name of each alkyl group.**

- The names of the alkyl groups form the first part of each amide name.
- For 3° amides, use the prefix **di-** if the two alkyl groups on N are the same. If the two alkyl groups are different, **alphabetize** their names. One "***N*-**" is needed for each alkyl group, even if both R groups are identical.


a. 

- The compound is a 2° amide with one ethyl group → *N*-ethyl.

b. 


- The compound is a 3° amide with two methyl groups.
- Use the prefix di- and two "*N*-" to begin the name → *N,N*-dimethyl.

Step [2] **Name the acyl group (RCO–) with the suffix *-amide.***

a. 

derived from
form*ic acid* – – – – – → form*amide*

- Change the *-ic acid* or *-oic acid* suffix of the parent carboxylic acid to the suffix *-amide.*
- Put the two parts of the name together.
- **Answer: *N*-ethylformamide**

b. 

derived from
benz*oic acid* – – – → benz*amide*

- Change benz*oic acid* to benz*amide.*
- Put the two parts of the name together.
- **Answer: *N,N*-dimethylbenzamide**

22.3E Naming a Nitrile

In contrast to the carboxylic acid derivatives, **nitriles are named as alkane derivatives.** To name a nitrile using IUPAC rules:

- **Find the longest chain that contains the CN and add the word *nitrile* to the name of the parent alkane. Number the chain to put CN at C1, but omit this number from the name.**

In naming a nitrile, the CN carbon is one carbon atom of the longest chain. **CH_3CH_2CN is propanenitrile, *not* ethanenitrile.**


Common names for nitriles are derived from the names of the carboxylic acid having the same number of carbon atoms by replacing the ***-ic acid*** ending of the carboxylic acid by the suffix ***-onitrile.***

When CN is named as a substituent, it is called a ***cyano*** group. Figure 22.1 illustrates features of nitrile nomenclature.

Table 22.2 summarizes the most important points about the nomenclature of carboxylic acid derivatives and nitriles.


Figure 22.1 Summary of nitrile nomenclature

a. IUPAC name for a nitrile




butane + nitrile
(4 C's)
2-methylbutanenitrile

b. Common name for a nitrile








derived from
acet*ic acid*
acetonitrile

c. CN as a substituent




2-cyanocyclohexanecarboxylic acid


Table 22.2 Summary: Nomenclature of Carboxylic Acid Derivatives and Nitriles

Compound	Name ending	Example	Name
acid chloride	**-yl chloride** or **-carbonyl chloride**		benzoyl chloride
anhydride	**anhydride**		benzoic anhydride
ester	**-ate**		ethyl benzoate
amide	**-amide**		*N*-methylbenzamide
nitrile	**-nitrile** or **-onitrile**		benzonitrile

Sample Problem 22.1

Give the IUPAC name for each compound.


a. 

b. 

Solution


a. The functional group is an acid chloride bonded to a chain of atoms, so the name ends in ***-yl chloride.***

[1] Find and name the longest chain containing the COCl:



hexano*ic acid* ⟶ **hexano*yl chloride***
(6 C's)


[2] Number and name the substituents:



Answer:
2,4-dimethylhexanoyl chloride


b. The functional group is an ester, so the name ends in ***-ate.***

[1] Find and name the longest chain containing the carbonyl group:



pentano*ic acid* ⟶ **pentano*ate***
(5 C's)







[2] Number and name the substituents:



Answer: isopropyl 2-methylpentanoate

The name of the alkyl group on the O atom goes **first** in the name.

Problem 22.4 Give an IUPAC or common name for each compound.

a. 
c. 
e. 
b. 
d. 
f. 


Problem 22.5 Draw the structure corresponding to each name.

a. 5-methylheptanoyl chloride
b. isopropyl propanoate
c. acetic formic anhydride
d. *N*-isobutyl-*N*-methylbutanamide
e. 3-methylpentanenitrile
f. *o*-cyanobenzoic acid
g. *sec*-butyl 2-methylhexanoate
h. *N*-ethylhexanamide

22.4 Physical Properties

Because all carbonyl compounds have a polar carbonyl group, they exhibit **dipole–dipole interactions.** Nitriles also have dipole–dipole interactions because they have a polar C≡N group. Because they contain one or two N−H bonds, 1° and 2° amides are capable of intermolecular hydrogen bonding. The N−H bond of one amide intermolecularly hydrogen bonds to the C=O of another amide, as shown using two acetamide molecules (CH_3CONH_2) in Figure 22.2.


Figure 22.2
Intermolecular hydrogen bonding between two CH_3CONH_2 molecules



Problem 22.6 Explain why the boiling point of CH_3CONH_2 (221 °C) is significantly higher than the boiling point of CH_3CO_2H (118 °C).

How these factors affect the physical properties of carboxylic acid derivatives is summarized in Table 22.3.

Table 22.3 Physical Properties of Carboxylic Acid Derivatives

Property	Observation
Boiling point and melting point	• Primary (1°) and 2° amides have *higher* boiling points and melting points than compounds of comparable molecular weight. • The boiling points and melting points of other carboxylic acid derivatives are similar to those of other polar compounds of comparable size and shape. 
Solubility	• Carboxylic acid derivatives are soluble in organic solvents regardless of size. • Most carboxylic acid derivatives having ≤ 5 C's are H_2O soluble because they can hydrogen bond with H_2O (Section 3.4C). • Carboxylic acid derivatives having > 5 C's are H_2O insoluble because the nonpolar alkyl portion is too large to dissolve in the polar H_2O solvent.

Key: MW = molecular weight

22.5 Spectroscopic Properties

22.5A IR Spectra

The most prominent IR absorptions for carboxylic acid derivatives and nitriles are as follows:

[1] Like all carbonyl compounds, carboxylic acid derivatives have a **strong C=O absorption between 1600 and 1850 cm^{-1}.**

[2] **Primary (1°) and 2° amides** have two additional absorptions due to the N−H bonds:
- one or two N−H stretching peaks at **3200–3400 cm^{-1}**
- an N−H bending absorption at **~1640 cm^{-1}**

[3] **Nitriles** have an absorption at **2250 cm^{-1} for the C≡N.**

The exact location of the carbonyl absorption varies with the identity of Z in the carbonyl compound RCOZ. As detailed in Section 22.2, as the basicity of Z increases, resonance stabilization of RCOZ increases, resulting in the following trend:

- **As the carbonyl π bond becomes more delocalized, the C=O absorption shifts to lower frequency.**




Thus, the carbonyl group of an acid chloride and anhydride, which are least stabilized by resonance, absorb at higher frequency than the carbonyl group of an amide, which is more stabilized by resonance. Table 22.4 lists specific values for the carbonyl absorptions of the carboxylic acid derivatives.

Conjugation and ring size affect the location of these carbonyl absorptions.

The effects of conjugation and ring size on the location of a carbonyl absorption were first discussed in Section 21.4A.

- **Conjugation shifts a carbonyl absorption to lower frequencies.**
- **For cyclic carboxylic acid derivatives, decreasing ring size shifts a carbonyl absorption to higher frequencies.**



Table 22.4 IR Absorptions for the Carbonyl Group of Carboxylic Acid Derivatives



Compound type	Structure (RCOZ)	Carbonyl absorption ($\tilde{\nu}$)
acid chloride		~1800
anhydride		1820 and 1760 (2 peaks)
ester		1735–1745
amide	 R' = H or alkyl	1630–1680



Increasing basicity of Z



Increasing $\tilde{\nu}$ of absorption

Problem 22.7 How would the compounds in each pair differ in their IR spectra?

a.  and 

c.  and 

b.  and 

d.  and 

22.5B NMR Spectra

Carboxylic acid derivatives have two characteristic ^{1}H NMR absorptions.

[1] **Protons on the α carbon to the carbonyl absorb at 2–2.5 ppm.**

[2] **The N–H protons of 1° and 2° amides absorb at 7.5–8.5 ppm.**

In their ^{13}C NMR spectra, carboxylic acid derivatives give a highly deshielded peak at 160–180 ppm due to the carbonyl carbon. This is somewhat upfield from the carbonyl absorption of aldehydes and ketones, which occurs at 190–215 ppm.

Nitriles give a peak at 115–120 ppm in their ^{13}C NMR spectrum due to the *sp* hybridized carbon. This is farther downfield than the signal due to the *sp* hybridized carbon of an alkyne, which occurs at 65–100 ppm.

Problem 22.8 Deduce the structures of compounds **A** and **B,** two of the major components of jasmine oil, from the given data.

Compound **A:** $C_9H_{10}O_2$; IR absorptions at 3091–2895 and 1743 cm^{-1}; ^{1}H NMR signals at 2.06 (singlet, 3 H), 5.08 (singlet, 2 H), and 7.33 (broad singlet, 5 H) ppm.

Compound **B:** $C_{14}H_{12}O_2$; IR absorptions at 3091–2953 and 1718 cm^{-1}; ^{1}H NMR signals at 5.35 (singlet, 2 H) and 7.26–8.15 (multiplets, 10 H) ppm.


22.6 Interesting Esters and Amides

22.6A Esters



The characteristic odor of many fruits is due to low molecular weight esters.

Many low molecular weight esters have pleasant and very characteristic odors.



isoamyl acetate
odor of banana


ethyl butanoate
odor of mango

methyl 2-methylbutanoate
odor of pineapple




The coca plant, *Erythroxylon coca,* is the source of the addictive drug cocaine.

Several esters, including vitamin C and cocaine, have important biological activities.



vitamin C




cocaine

- **Vitamin C** (or **ascorbic acid**) is a water-soluble vitamin containing a five-membered lactone that we first discussed in Section 3.5B. Although vitamin C is synthesized in plants, humans do not have the necessary enzymes to make it, and so they must obtain it from their diet.
- **Cocaine** is an addictive stimulant obtained from the leaves of the coca plant. Chewing coca leaves for pleasure has been practiced by the indigenous peoples of South America for over a thousand years, and coca leaves were a very minor ingredient in Coca-Cola for the first 20 years of its production. Cocaine is a widely abused recreational drug, and the possession and use of cocaine is currently illegal in most countries.

22.6B Amides

An important group of naturally occurring amides consists of ***proteins,*** **polymers of amino acids joined together by amide linkages.** Proteins differ in the length of the polymer chain, as well as in the identity of the R groups bonded to it. The word *protein* is usually reserved for high molecular weight polymers composed of 40 or more amino acid units, while the designation *peptide* is given to polymers of lower molecular weight.




portion of a protein molecule
[Amide bonds are shown in red.]

Peptides and proteins are discussed in detail in Chapter 29.


Proteins and peptides have diverse functions in the cell. They form the structural components of muscle, connective tissue, hair, and nails. They catalyze reactions and transport ions and

molecules across cell membranes. **Met-enkephalin,** for example, a peptide with four amide bonds found predominately in nerve tissue cells, relieves pain and acts as an opiate by producing morphine-like effects.




met-enkephalin
[The four amide bonds are shown in red.]

Several useful drugs are amides. For example, **Gleevec** (generic name imatinib mesylate), an amide sold as a salt with methanesulfonic acid (CH_3SO_3H), is an anticancer drug used for the treatment of chronic myeloid leukemia as well as certain gastrointestinal tumors. Gleevec targets a single molecule to disable the molecular mechanism responsible for a specific type of cancer.




imatinib mesylate
Trade name Gleevec


Penicillins are a group of structurally related antibiotics, known since the pioneering work of Sir Alexander Fleming led to the discovery of penicillin G in the 1920s. All penicillins contain a strained β-lactam fused to a five-membered ring, as well as a second amide located α to the β-lactam carbonyl group. Particular penicillins differ in the identity of the R group in the amide side chain.



penicillin
β-lactam shown in red




penicillin G




amoxicillin

Cephalosporins represent a second group of β-lactam antibiotics that contain a four-membered ring fused to a six-membered ring. Cephalosporins are generally active against a broader range of bacteria than penicillins.




cephalosporin
β-lactam shown in red



cephalexin
Trade name Keflex

22.7 Introduction to Nucleophilic Acyl Substitution

The characteristic reaction of carboxylic acid derivatives is *nucleophilic acyl substitution.* This is a general reaction that occurs with both negatively charged nucleophiles (Nu:$^-$) and neutral nucleophiles (HNu:).



Nu replaces Z.


- Carboxylic acid derivatives (RCOZ) react with nucleophiles because they contain an electrophilic, unhindered carbonyl carbon.
- Substitution occurs, *not* addition, because carboxylic acid derivatives (RCOZ) have a leaving group Z on the carbonyl carbon.

22.7A The Mechanism

The mechanism for nucleophilic acyl substitution was first presented in Section 20.2.

The general mechanism for nucleophilic acyl substitution is a two-step process: **nucleophilic attack** followed by **loss of the leaving group,** as shown in Mechanism 22.1.

Mechanism 22.1 General Mechanism—Nucleophilic Acyl Substitution




Z = OH, Cl, OCOR, OR', NH_2


1. The nucleophile attacks the electrophilic carbonyl group. The π bond is broken, moving an electron pair out on oxygen and forming an sp^3 hybridized carbon.
2. An electron pair on oxygen re-forms the π bond and Z comes off as a leaving group with the electron pair in the C–Z bond.

The overall result of addition of a nucleophile and elimination of a leaving group is substitution of the nucleophile for the leaving group. Recall from Chapter 20 that nucleophilic substitution occurs with carbanions (R^-) and hydride (H^-) as nucleophiles. A variety of oxygen and nitrogen nucleophiles also participate in this reaction.


Oxygen nucleophiles



Nitrogen nucleophiles



Nucleophilic acyl substitution using heteroatomic nucleophiles results in the conversion of one carboxylic acid derivative into another, as shown in two examples.



Each reaction results in the replacement of the leaving group by the nucleophile, regardless of the identity of or charge on the nucleophile. To draw any nucleophilic acyl substitution product:

- Find the sp^2 hybridized carbon with the leaving group.
- Identify the nucleophile.
- Substitute the nucleophile for the leaving group. With a neutral nucleophile a proton must be lost to obtain a neutral substitution product.


22.7B Relative Reactivity of Carboxylic Acids and Their Derivatives

As discussed in Section 20.2B, carboxylic acids and their derivatives differ greatly in reactivity toward nucleophiles. The order of reactivity parallels the leaving group ability of the group Z.

- The better the leaving group, the more reactive RCOZ is in nucleophilic acyl substitution.


Recall that the **best leaving group is the weakest base.** The relative basicity of the common leaving groups, Z, is given in Table 22.1.

Thus, the following trends result:



Based on this order of reactivity, **more reactive acyl compounds (acid chlorides and anhydrides) can be converted to less reactive ones (carboxylic acids, esters, and amides). The reverse is not usually true.**


To see why this is so, recall that nucleophilic addition to a carbonyl group forms a tetrahedral intermediate with two possible leaving groups, Z^- or $:Nu^-$. The group that is subsequently eliminated is the *better* of the two leaving groups. For a reaction to form a substitution product, therefore, Z^- must be the better leaving group, making the starting material RCOZ a more reactive acyl compound.




For a reaction to occur, Z must be a better leaving group than Nu.

To evaluate whether a nucleophilic substitution reaction will occur, **compare the leaving group ability of the incoming nucleophile and the departing leaving group,** as shown in Sample Problem 22.2.

Sample Problem 22.2 Determine whether each nucleophilic acyl substitution is likely to occur.

a. 

b. 


Solution

a. Conversion of CH_3COCl to $CH_3COOCH_2CH_3$ requires the substitution of Cl^- by $^-OCH_2CH_3$. Because Cl^- is a weaker base and therefore a better leaving group than $^-OCH_2CH_3$, **this reaction occurs.**


b. Conversion of $C_6H_5CONH_2$ to $(C_6H_5CO)_2O$ requires the substitution of $^-NH_2$ by $^-OCOC_6H_5$. Because $^-NH_2$ is a stronger base and therefore a poorer leaving group than $^-OCOC_6H_5$, **this reaction does *not* occur.**

Problem 22.9 Without reading ahead in Chapter 22, state whether it should be possible to carry out each of the following nucleophilic substitution reactions.

a. 

c. 

b. 

d. 

Learn the order of reactivity of carboxylic acid derivatives. Keeping this in mind allows you to organize a very large number of reactions.

To summarize:

- **Nucleophilic substitution occurs when the leaving group Z^- is a weaker base and therefore better leaving group than the attacking nucleophile $:Nu^-$.**
- **More reactive acyl compounds can be converted to less reactive acyl compounds by nucleophilic substitution.**

Problem 22.10 Rank the compounds in each group in order of increasing reactivity in nucleophilic acyl substitution.

a. $C_6H_5CO_2CH_3$, C_6H_5COCl, $C_6H_5CONH_2$

b. $CH_3CH_2CO_2H$, $(CH_3CH_2CO)_2O$, $CH_3CH_2CONHCH_3$


Problem 22.11 Explain why trichloroacetic anhydride $[(Cl_3CCO)_2O]$ is more reactive than acetic anhydride $[(CH_3CO)_2O]$ in nucleophilic acyl substitution reactions.

22.7C A Preview of Specific Reactions

Sections 22.8–22.14 are devoted to specific examples of nucleophilic acyl substitution using heteroatoms as nucleophiles. There are a great many reactions, and it is easy to confuse them unless you learn the general order of reactivity of carboxylic acid derivatives. **Keep in mind that every reaction that begins with an acyl starting material involves nucleophilic substitution.**

In this text, all of the nucleophilic substitution reactions are grouped according to the carboxylic acid derivative used as a starting material. We begin with the reactions of acid chlorides, the most reactive acyl compounds, then proceed to less and less reactive carboxylic acid derivatives, ending with amides. Acid chlorides undergo many reactions, because they have the best leaving group of all acyl compounds, whereas amides undergo only one reaction, which must be carried out under harsh reaction conditions, because amides have a poor leaving group.

In general, we will examine nucleophilic acyl substitution with four different nucleophiles, as shown in the following equations.




These reactions are used to make anhydrides, carboxylic acids, esters, and amides, but not acid chlorides, from other acyl compounds. Acid chlorides are the most reactive acyl compounds (they have the best leaving group), so they are not easily formed as a product of nucleophilic substitution reactions. They can only be prepared from carboxylic acids using special reagents, as discussed in Section 22.10A.

22.8 Reactions of Acid Chlorides




The reaction of acid chlorides with water is rapid. Exposure of an acid chloride to moist air on a humid day leads to some hydrolysis, giving the acid chloride a very acrid odor, due to the HCl formed as a by-product.


Acid chlorides readily react with nucleophiles to form nucleophilic substitution products, with HCl usually formed as a reaction by-product. A weak base like pyridine is added to the reaction mixture to remove this strong acid, forming an ammonium salt.



Acid chlorides react with oxygen nucleophiles to form anhydrides, carboxylic acids, and esters.




Acid chlorides also react with ammonia and 1° and 2° amines to form 1°, 2°, and 3° amides, respectively. Two equivalents of NH_3 or amine are used. One equivalent acts as a nucleophile to replace Cl and form the substitution product, while the second equivalent reacts as a base with the HCl by-product to form an ammonium salt.





Insect repellents containing DEET have become particularly popular because of the recent spread of many insect-borne diseases such as West Nile virus and Lyme disease. DEET does not kill insects—it repels them. It is thought that DEET somehow confuses insects so that they can no longer sense the warm moist air that surrounds a human body.


As an example, reaction of an acid chloride with diethylamine forms the 3° amide *N,N*-diethyl-*m*-toluamide, popularly known as **DEET.** DEET, the active ingredient in the most widely used insect repellents, is effective against mosquitoes, fleas, and ticks.



Problem 22.12 Draw the products formed when benzoyl chloride (C_6H_5COCl) is treated with each nucleophile: (a) H_2O, pyridine; (b) CH_3COO^-; (c) NH_3 (excess); (d) $(CH_3)_2NH$ (excess).

With a carboxylate nucleophile the mechanism follows the general, two-step mechanism discussed in Section 22.7: **nucleophilic attack followed by loss of the leaving group,** as shown in Mechanism 22.2.

Mechanism 22.2 Conversion of Acid Chlorides to Anhydrides




1 The nucleophilic carboxylate anion attacks the carbonyl group, forming an sp^3 hybridized carbon.

2 Elimination of the leaving group (Cl^-) forms the substitution product, an anhydride.

Nucleophilic substitution with the neutral nucleophiles (H_2O, R'OH, NH_3, and so forth) requires an additional step for proton transfer. For example, the reaction of an acid chloride with H_2O as nucleophile converts an acid chloride to a carboxylic acid in three steps (Mechanism 22.3).

Mechanism 22.3 Conversion of Acid Chlorides to Carboxylic Acids





1 The nucleophile (H_2O) attacks the carbonyl group, forming an sp^3 hybridized carbon.

2 – 3 Loss of a proton and elimination of the leaving group (Cl^-) form the substitution product, a carboxylic acid.

The exact same three-step process can be written for any neutral nucleophile that reacts with acid chlorides.

Problem 22.13 Draw a stepwise mechanism for the formation of **A** from an alcohol and acid chloride. **A** was converted in one step to blattellaquinone, the sex pheromone of the female German cockroach *Blattella germanica.*






A short laboratory synthesis of blattellaquinone (Problem 22.13), the sex pheromone of the female German cockroach, opens new possibilities for cockroach population control using pheromone-baited traps.

Nucleophilic substitution occurs only when the leaving group is a weaker base and therefore a better leaving group than the attacking nucleophile.


22.9 Reactions of Anhydrides

Although somewhat less reactive than acid chlorides, anhydrides nonetheless readily react with most nucleophiles to form substitution products. Nucleophilic substitution reactions of anhydrides are no different than the reactions of other carboxylic acid derivatives, even though anhydrides contain two carbonyl groups. **Nucleophilic attack occurs at one carbonyl group, while the second carbonyl becomes part of the leaving group.**



Anhydrides can't be used to make acid chlorides, because $RCOO^-$ is a stronger base and therefore a poorer leaving group than Cl^-. Anhydrides can be used to make all other acyl derivatives, however. Reaction with water and alcohols yields **carboxylic acids** and **esters,** respectively.


Reaction with two equivalents of NH_3 or amines forms **1°, 2°,** and **3° amides.** A molecule of carboxylic acid (or a carboxylate salt) is always formed as a by-product.



Problem 22.14 Draw the products formed when benzoic anhydride [$(C_6H_5CO)_2O$] is treated with each nucleophile: (a) H_2O; (b) CH_3OH; (c) NH_3 (excess); (d) $(CH_3)_2NH$ (excess).

The conversion of an anhydride to an amide illustrates the mechanism of nucleophilic acyl substitution with an anhydride as starting material (Mechanism 22.4). Besides the usual steps of **nucleophilic addition** and **elimination of the leaving group,** an additional proton transfer is needed.

Mechanism 22.4 Conversion of an Anhydride to an Amide




1 The nucleophile (NH_3) attacks the carbonyl, forming an sp^3 hybridized carbon.

2 – 3 Loss of a proton and elimination of the leaving group (RCO_2^-) form the substitution product, a 1°amide.


Acetaminophen reduces pain and fever, but it is not anti-inflammatory, so it is ineffective in treating conditions like arthritis, which have a significant inflammatory component. In large doses, acetaminophen causes liver damage, so dosage recommendations must be carefully followed.

Anhydrides react with alcohols and amines with ease, so they are often used in the laboratory to prepare esters and amides. For example, acetic anhydride is used to prepare two analgesics, **acetylsalicylic acid** (aspirin) and **acetaminophen** (the active ingredient in Tylenol).




These are called **acetylation** reactions because they result in the transfer of an acetyl group, CH_3CO-, from one heteroatom to another.

Heroin is prepared by the acetylation of morphine, an analgesic compound isolated from the opium poppy. Both OH groups of morphine are readily acetylated with acetic anhydride to form the diester present in heroin.




22.10 Reactions of Carboxylic Acids

Carboxylic acids are strong organic acids. Because acid–base reactions proceed rapidly, any nucleophile that is also a strong base will react with a carboxylic acid by removing a proton ***first,*** before any nucleophilic substitution reaction can take place.




An acid–base reaction occurs with ^{-}OH, NH_3, and amines, all common nucleophiles used in nucleophilic acyl substitution reactions. Nonetheless, carboxylic acids do undergo nucleophilic acyl substitution and can be converted to a variety of other acyl derivatives using special reagents, with acid catalysis, or sometimes, by using rather forcing reaction conditions. These reactions are summarized in Figure 22.3 and detailed in Sections 22.10A–22.10D.

Figure 22.3
Nucleophilic acyl substitution reactions of carboxylic acids




22.10A Conversion of RCOOH to RCOCl


Carboxylic acids can't be converted to acid chlorides by using Cl^- as a nucleophile, because the attacking nucleophile Cl^- is a weaker base than the departing leaving group, ^-OH. But carboxylic acids *can* be converted to acid chlorides using thionyl chloride, **$SOCl_2$,** a reagent that was introduced in Section 9.12 to convert alcohols to alkyl chlorides.



Treatment of benzoic acid with $SOCl_2$ forms benzoyl chloride. This reaction converts a less reactive acyl derivative (a carboxylic acid) into a more reactive one (an acid chloride). This is possible because thionyl chloride converts the OH group of the acid into a better leaving group, and because it provides the nucleophile (Cl^-) to displace the leaving group. The steps in the process are illustrated in Mechanism 22.5.




Mechanism 22.5 Conversion of Carboxylic Acids to Acid Chlorides



1 – 2 Reaction of the carboxylic acid with $SOCl_2$ and loss of a proton convert the OH group to OSOCl, a good leaving group.


3 – 4 Nucleophilic attack of chloride generates a tetrahedral intermediate, and loss of the leaving group (SO_2 and Cl^-) forms the acid chloride.

Problem 22.15 Draw the products of each reaction.




22.10B Conversion of RCOOH to $(RCO)_2O$

Carboxylic acids cannot be readily converted to anhydrides, but dicarboxylic acids can be converted to cyclic anhydrides by heating to high temperatures. This is a **dehydration** reaction because a water molecule is lost from the diacid.




22.10C Conversion of RCOOH to RCOOR'

Treatment of a carboxylic acid with an alcohol in the presence of an acid catalyst forms an ester. This reaction is called a **Fischer esterification.**



This reaction is an equilibrium. According to Le Châtelier's principle, it is driven to the right by using excess alcohol or by removing the water as it is formed.


Ethyl acetate is a common organic solvent with a characteristic odor. It is used in nail polish remover and model airplane glue.



The mechanism for the Fischer esterification involves the usual two steps of nucleophilic acyl substitution—that is, **addition of a nucleophile followed by elimination of a leaving group.** Because the reaction is acid catalyzed, however, there are additional protonation and deprotonation steps. As always, though, the first step of any mechanism with an oxygen-containing starting material and an acid is to **protonate an oxygen atom** as shown with a general acid HA in Mechanism 22.6.

Mechanism 22.6 Fischer Esterification—Acid-Catalyzed Conversion of Carboxylic Acids to Esters


Part [1] Addition of the nucleophile R'OH



1 Protonation of the carbonyl oxygen makes the carbonyl more electrophilic.

2 – 3 Nucleophilic attack by R'OH forms a tetrahedral intermediate, and deprotonation gives the addition product.


Part [2] Elimination of the leaving group H_2O




4 Protonation of the OH group forms a good leaving group.

5 – 6 Loss of H_2O and deprotonation give the ester.


Esterification of a carboxylic acid occurs in the presence of acid but not in the presence of base. Base removes a proton from the carboxylic acid, forming an electron-rich carboxylate anion, which does not react with an electron-rich nucleophile.



Intramolecular esterification of γ- and δ-hydroxy carboxylic acids forms five- and six-membered lactones.




Problem 22.16 Draw the products of each reaction.




Problem 22.17 Draw the products formed when benzoic acid ($C_6H_5CO_2H$) is treated with CH_3OH having its O atom labeled with ^{18}O ($CH_3{}^{18}OH$). Indicate where the labeled oxygen atom resides in the products.

Problem 22.18 Draw a stepwise mechanism for the following reaction.



22.10D Conversion of RCOOH to RCONR'$_2$

The direct conversion of a carboxylic acid to an amide with NH_3 or an amine is very difficult, even though a more reactive acyl compound is being transformed into a less reactive one. The problem is that carboxylic acids are strong organic acids and NH_3 and amines are bases, so they undergo an **acid–base reaction to form an ammonium salt** before any nucleophilic substitution occurs.




Heating at high temperature (>100 °C) dehydrates the resulting ammonium salt of the carboxylate anion to form an amide, though the yield can be low.

Amides are much more easily prepared from acid chlorides and anhydrides, as discussed in Sections 22.8 and 22.9.


Therefore, the overall conversion of RCOOH to $RCONH_2$ requires two steps:

[1] **Acid–base reaction of RCOOH with NH_3 to form an ammonium salt**


[2] **Dehydration at high temperature (>100 °C)**



A carboxylic acid and an amine readily react to form an amide in the presence of an additional reagent, **dicyclohexylcarbodiimide (DCC),** which is converted to the by-product dicyclohexylurea in the course of the reaction.




DCC is a dehydrating agent. The dicyclohexylurea by-product is formed by adding the elements of H_2O to DCC. DCC promotes amide formation by converting the carboxy OH group into a better leaving group.



The mechanism consists of two parts: [1] conversion of the OH group into a better leaving group, followed by [2] **addition of the nucleophile and loss of the leaving group** to form the product of nucleophilic acyl substitution (Mechanism 22.7).

Mechanism 22.7 Conversion of Carboxylic Acids to Amides with DCC


Part [1] Conversion of OH into a better leaving group



1 Acid–base reaction results in transfer of a proton from the carboxylic acid to DCC.

2 Nucleophilic attack of RCO_2^- on the conjugate acid of DCC forms an addition product. The overall result of Steps [1] and [2] is conversion of OH into a better leaving group.

Part [2] Addition of the nucleophile and loss of the leaving group



3 Nucleophilic attack of the amine on the activated carboxy group forms a tetrahedral intermediate.

4 – 5 Proton transfer and elimination of dicyclohexylurea as the leaving group form the amide.


The reaction of an acid and an amine with DCC is often used in the laboratory to form the amide bond in peptides, as is discussed in Chapter 29.

Problem 22.19 What product is formed when acetic acid is treated with each reagent: (a) CH_3NH_2; (b) CH_3NH_2, then heat; (c) CH_3NH_2 + DCC?


22.11 Reactions of Esters

Esters can be converted into carboxylic acids and amides.

- **Esters are hydrolyzed with water in the presence of either acid or base to form carboxylic acids or carboxylate anions.**




- **Esters react with NH_3 and amines to form 1°, 2°, or 3° amides.**



22.11A Ester Hydrolysis in Aqueous Acid

The first step in acid-catalyzed ester hydrolysis is **protonation on oxygen,** the same first step of any mechanism involving an oxygen-containing starting material and an acid.


The hydrolysis of esters in aqueous acid is a reversible equilibrium reaction that is driven to the right by using a large excess of water.



The mechanism of ester hydrolysis in acid (shown in Mechanism 22.8) is the reverse of the mechanism of ester synthesis from carboxylic acids (Mechanism 22.6). Thus, the mechanism consists of the addition of the nucleophile and the elimination of the leaving group, the two steps common to all nucleophilic acyl substitutions, as well as several proton transfers, because the reaction is acid-catalyzed.

Mechanism 22.8 Acid-Catalyzed Hydrolysis of an Ester to a Carboxylic Acid


Part [1] Addition of the nucleophile H_2O



1 Protonation of the carbonyl oxygen makes the carbonyl more electrophilic.

2 – 3 Nucleophilic attack by H_2O forms a tetrahedral intermediate, and deprotonation gives the addition product.

Part [2] Elimination of the leaving group R'OH




4 Protonation of the OR' group forms a good leaving group.

5 – 6 Loss of R'OH and deprotonation give the carboxylic acid.

22.11B Ester Hydrolysis in Aqueous Base

The word **saponification** comes from the Latin *sapo* meaning **soap.** Soap is prepared by hydrolyzing esters in fats with aqueous base, as explained in Section 22.12B.


Esters are hydrolyzed in aqueous base to form carboxylate anions. Basic hydrolysis of an ester is called **saponification.**



carboxylate anion

The mechanism for this reaction has the usual two steps of the general mechanism for nucleophilic acyl substitution presented in Section 22.7—**addition of the nucleophile** followed by **loss of a leaving group**—plus an additional step involving proton transfer (Mechanism 22.9).

Mechanism 22.9 Base-Promoted Hydrolysis of an Ester to a Carboxylic Acid




1 – 2 Addition of the nucleophile ($^-$OH) followed by elimination of the leaving group ($^-$OR') form a carboxylic acid. These two steps are reversible.

3 Because the carboxylic acid is a strong organic acid and the leaving group ($^-$OR') is a strong base, an acid–base reaction forms the carboxylate anion.


The carboxylate anion is resonance stabilized, and this drives the equilibrium in its favor. Once the reaction is complete and the carboxylate anion is formed, it can be protonated with strong acid to form the neutral carboxylic acid.

Hydrolysis is base promoted, *not* base catalyzed, because the base ($^-$OH) is the nucleophile that adds to the ester and forms part of the product. It participates in the reaction and is not regenerated later.




Where do the oxygen atoms in the product come from? The C–OR' bond in the ester is cleaved, so the OR' group becomes the alcohol by-product (R'OH) and one of the oxygens in the carboxylate anion product comes from $^-$OH (the nucleophile).

Problem 22.20 Fenofibrate is a cholesterol-lowering medication that is converted to fenofibric acid, the active drug, by hydrolysis during metabolism. What is the structure of fenofibric acid?



fenofibrate

Problem 22.21 What product is formed when the esters in ginkgolide B, the chapter-opening molecule, are hydrolyzed in aqueous acid? Indicate the stereochemistry of all stereogenic centers.




ginkgolide B

22.12 Application: Lipid Hydrolysis

22.12A Olestra—A Synthetic Fat

The most prevalent naturally occurring esters are the **triacylglycerols,** which were first discussed in Section 10.6. **Triacylglycerols are the lipids that comprise animal fats and vegetable oils.**


- **Each triacylglycerol is a triester, containing three long hydrocarbon side chains.**
- **Unsaturated triacylglycerols have one or more double bonds in their long hydrocarbon chains, whereas saturated triacylglycerols have none.**



triacylglycerol

R groups have 11–19 C's.
[Three ester groups are labeled in red.]


Figure 22.4
The three-dimensional structure of a saturated triacylglycerol



- This triacylglycerol has no double bonds in the three R groups (each with 11 C's) bonded to the ester carbonyls, making it a saturated fat.

Figure 22.4 contains a ball-and-stick model of a saturated fat.

Animals store energy in the form of triacylglycerols, kept in a layer of fat cells below the surface of the skin. This fat serves to insulate the organism, as well as provide energy for its metabolic needs for long periods. The first step in the metabolism of a triacylglycerol is hydrolysis of the ester bonds to form glycerol and three fatty acids. **This reaction is simply ester hydrolysis.** In cells, this reaction is carried out with enzymes called **lipases.**



triacylglycerol


[The three bonds drawn in red are cleaved in hydrolysis.]

The fatty acids produced on hydrolysis are then oxidized in a stepwise fashion, ultimately yielding CO_2 and H_2O, as well as a great deal of energy. Oxidation of fatty acids yields twice as much energy per gram as oxidation of an equivalent weight of carbohydrate.

Diets high in fat content lead to a large amount of stored fat, ultimately causing an individual to be overweight. One attempt to reduce calories in common snack foods has been to substitute "fake fats" such as **olestra** (trade name **Olean**) for triacylglycerols.




Some snack foods contain the "fake fat" olestra, giving them fewer calories for the calorie-conscious consumer.




olestra

a polyester of sucrose
a synthetic fat
[R groups contain 11–19 C's.]



sucrose



Olestra is a polyester formed from long-chain fatty acids and sucrose, the sweet-tasting carbohydrate in table sugar. Naturally occurring triacylglycerols are also polyesters formed from long-chain fatty acids, but olestra has so many ester units clustered together in close proximity that they are too hindered to be hydrolyzed. As a result, olestra is not metabolized. Instead, it passes through the body unchanged, providing no calories to the consumer.

Thus, olestra's many C−C and C−H bonds make it similar in solubility to naturally occurring triacylglycerols, but its three-dimensional structure makes it inert to hydrolysis because of steric hindrance.

Problem 22.22 How would you synthesize olestra from sucrose?


22.12B The Synthesis of Soap

Soap has been previously discussed in Section 3.6.




All soaps are salts of fatty acids. The main difference between soaps is the addition of other ingredients that do not alter their cleaning properties: dyes for color, scents for a pleasing odor, and oils for lubrication. Soaps that float are aerated, so that they are less dense than water.

Soap is prepared by the basic hydrolysis or saponification of a triacylglycerol. Heating an animal fat or vegetable oil with aqueous base hydrolyzes the three esters to form glycerol and sodium salts of three fatty acids. These carboxylate salts are **soaps,** which clean away dirt because of their two structurally different regions. The nonpolar tail dissolves grease and oil and the polar head makes it soluble in water (Figure 3.5). Most triacylglycerols have two or three different R groups in their hydrocarbon chains, so soaps are usually mixtures of two or three different carboxylate salts.




Soaps are typically made from lard (from hogs), tallow (from cattle or sheep), coconut oil, or palm oil. All soaps work in the same way, but have somewhat different properties depending on the lipid source. The length of the carbon chain in the fatty acids and the number of degrees of unsaturation affect the properties of the soap to some extent.

Problem 22.23 What is the composition of the soap prepared by hydrolysis of the following triacylglycerol?




22.13 Reactions of Amides

Because amides have the poorest leaving group of all the carboxylic acid derivatives, they are the least reactive. Under strenuous reaction conditions, **amides are hydrolyzed in acid or base to form carboxylic acids or carboxylate anions.**



In acid, the amine by-product is protonated as an ammonium ion, whereas in base, a neutral amine is formed.




The relative lack of reactivity of the amide bond is notable in proteins, which are polymers of amino acids connected by amide linkages (Section 22.6B). Proteins are stable in aqueous solution in the absence of acid or base, so they can perform their various functions in the aqueous cellular environment without breaking down. The hydrolysis of the amide bonds in proteins requires a variety of specific enzymes.

The mechanism of amide hydrolysis in acid is exactly the same as the mechanism of ester hydrolysis in acid (Section 22.11A) except that the leaving group is different.

The mechanism of amide hydrolysis in base has the usual two steps of the general mechanism for nucleophilic acyl substitution—**addition of the nucleophile** followed by **loss of a leaving group**—plus an additional proton transfer. The initially formed carboxylic acid reacts further under basic conditions to form the resonance-stabilized carboxylate anion, and this drives the reaction to completion. Mechanism 22.10 is written for a 1° amide.

Mechanism 22.10 Amide Hydrolysis in Base




1 – 2 Addition of the nucleophile ($^-$OH) followed by elimination of the leaving group ($^-NH_2$) form a carboxylic acid. These two steps are reversible.


3 Because the carboxylic acid is a strong organic acid and the leaving group ($^-NH_2$) is a strong base, an acid–base reaction forms the carboxylate anion.

Step [2] of Mechanism 22.10 deserves additional comment. For amide hydrolysis to occur, the tetrahedral intermediate must lose $^-NH_2$, a stronger base and therefore poorer leaving group than $^-$OH. This means that loss of $^-NH_2$ does not often happen. Instead, $^-$OH is lost as the leaving group most of the time, and the starting material is regenerated. But, when $^-NH_2$ is occasionally eliminated, the carboxylic acid product is converted to a lower energy carboxylate anion in Step [3], and this drives the equilibrium to favor its formation.

Problem 22.24 Draw a stepwise mechanism for the following reaction.




Problem 22.25 With reference to the structures of acetylsalicylic acid (aspirin, Chapter 2 opening molecule) and acetaminophen (the active ingredient in Tylenol), explain why acetaminophen tablets can be stored in the medicine cabinet for years, but aspirin slowly decomposes over time.



acetylsalicylic acid acetaminophen

22.14 Application: The Mechanism of Action of β-Lactam Antibiotics


Penicillin and related β-lactams kill bacteria by a nucleophilic acyl substitution reaction. All penicillins have an unreactive amide side chain and a very reactive amide that is part of a β-lactam. The β-lactam is more reactive than other amides because it is part of a strained, four-membered ring that is readily opened with nucleophiles.



penicillin


Unlike mammalian cells, bacterial cells are surrounded by a fairly rigid cell wall, which allows the bacterium to live in many different environments. This protective cell wall is composed of carbohydrates linked together by peptide chains containing amide linkages, formed using the enzyme **glycopeptide transpeptidase.**

Penicillin interferes with the synthesis of the bacterial cell wall. A nucleophilic OH group of the glycopeptide transpeptidase enzyme cleaves the β-lactam ring of penicillin by a nucleophilic acyl substitution reaction. The opened ring of the penicillin molecule remains covalently bonded to the enzyme, thus deactivating the enzyme, halting cell wall construction, and killing the bacterium. Penicillin has no effect on mammalian cells because they are surrounded by a flexible membrane composed of a lipid bilayer (Chapter 3) and not a cell wall.



Thus, penicillin and other β-lactam antibiotics are biologically active precisely because they undergo a nucleophilic acyl substitution reaction with an important bacterial enzyme.

Problem 22.26 Some penicillins cannot be administered orally because their β-lactam is rapidly hydrolyzed by the acidic environment of the stomach. What product is formed in the following hydrolysis reaction?



22.15 Summary of Nucleophilic Acyl Substitution Reactions

To help you organize and remember all of the nucleophilic acyl substitution reactions that can occur at a carbonyl carbon, keep in mind the following two principles:

- **The better the leaving group, the more reactive the carboxylic acid derivative.**
- **More reactive acyl compounds can always be converted to less reactive ones. The reverse is not usually true.**

This results in the following order of reactivity:




Table 22.5 summarizes the specific nucleophilic acyl substitution reactions. Use it as a quick reference to remind you which products can be formed from a given starting material.

Table 22.5 Summary of the Nucleophilic Substitution Reactions of Carboxylic Acids and Their Derivatives

Starting material		Product: RCOCl	$(RCO)_2O$	RCO_2H	RCO_2R'	$RCONR'_2$
[1] RCOCl	→	–	✓	✓	✓	✓
[2] $(RCO)_2O$	→	✗	–	✓	✓	✓
[3] RCO_2H	→	✓	✓	–	✓	✓
[4] RCO_2R'	→	✗	✗	✓	–	✓
[5] $RCONR'_2$	→	✗	✗	✓	✗	–


Table key: ✓ = A reaction occurs.
✗ = No reaction occurs.

22.16 Natural and Synthetic Fibers


All natural and synthetic fibers are high molecular weight polymers. Natural fibers are obtained from either plant or animal sources, and this determines the fundamental nature of their chemical structure. Fibers like **wool and silk obtained from animals are proteins,** so they are formed from amino acids joined together by many amide linkages. **Cotton and linen,** on the other hand, are derived from plants, so they are **carbohydrates having the general structure of cellulose,** formed from glucose monomers. General structures for these polymers are shown in Figure 22.5.

Figure 22.5
The general structure of the common natural fibers

a. Wool and silk—Proteins with many amide bonds



b. Cotton and linen—Carbohydrates like cellulose




An important practical application of organic chemistry has been the synthesis of synthetic fibers, many of which have properties that are different from and sometimes superior to their naturally occurring counterparts. The two most common classes of synthetic polymers are based on polyamides and polyesters.

22.16A Nylon—A Polyamide



DuPont built the first commercial nylon plant in 1938. Although it was initially used by the military to make parachutes, nylon quickly replaced silk in many common products after World War II.


The search for a synthetic fiber led to the discovery of **nylon,** a **polyamide** that is strong and durable and resembles the silk produced by silkworms. There are several different kinds of nylon, but the most well known is called nylon 6,6.



[The amide bonds are labeled in red.]


nylon 6,6

Nylon 6,6 can be synthesized from two six-carbon monomers (hence its name)—adipoyl chloride (**ClOCCH$_2$CH$_2$CH$_2$CH$_2$COCl**) and hexamethylenediamine (**H$_2$NCH$_2$CH$_2$CH$_2$CH$_2$CH$_2$CH$_2$NH$_2$**). This diacid chloride and diamine react together to form new amide bonds, yielding the polymer. Nylon is called a **condensation polymer** because a small molecule, in this case HCl, is eliminated during its synthesis.



nylon 6,6 + 3 HCl

Problem 22.27 What two monomers are needed to prepare nylon 6,10?




nylon 6,10

22.16B Polyesters



As we will learn in Section 30.9, PET is more easily recycled than other common polymers. For example, recycled PET is used to make reusable shopping bags.


Polyesters constitute a second major class of condensation polymers. The most common polyester is polyethylene terephthalate **(PET),** which is sold under a variety of trade names (Dacron, Terylene, and Mylar) depending on its use.



polyethylene terephthalate
PET
(Dacron, Terylene, and Mylar)

Ester bonds (in red) join the carbon skeleton together.

One method of synthesizing a polyester is by acid-catalyzed esterification of a diacid with a diol (Fischer esterification).



Because these polymers are easily and cheaply prepared and form strong and chemically stable materials, they have been used in clothing, films, tires, and many other products.

Problem 22.28 Draw the structure of Kodel, a polyester formed from 1,4-dihydroxymethylcyclohexane and terephthalic acid. Explain why fabrics made from Kodel are stiff and crease resistant.




1,4-dihydroxymethylcyclohexane terephthalic acid


Problem 22.29 Poly(lactic acid) (PLA) has received much recent attention because the lactic acid monomer [$CH_3CH(OH)COOH$] from which it is made can be obtained from carbohydrates rather than petroleum. This makes PLA a more "environmentally friendly" polyester. (A more in-depth discussion of green polymer synthesis is presented in Chapter 30.) Draw the structure of PLA.


22.17 Biological Acylation Reactions

Nucleophilic acyl substitution is a common reaction in biological systems. These acylation reactions are called **acyl transfer reactions** because they result in the transfer of an acyl group from one atom to another (from Z to Nu in this case).




In cells, such acylations occur with the sulfur analogue of an ester, called a **thioester,** having the general structure **RCOSR'.** The most common thioester is called **acetyl coenzyme A,** often referred to merely as **acetyl CoA.**






- A thioester (RCOSR') has a good leaving group (⁻SR'), so, like other acyl compounds, it undergoes substitution reactions with other nucleophiles. With acetyl CoA, an acetyl group is transferred from SCoA to a nucleophile, :Nu⁻.



For example, acetyl CoA undergoes enzyme-catalyzed nucleophilic acyl substitution with choline, forming acetylcholine, a charged compound that transmits nerve impulses between nerve cells.




Many other acyl transfer reactions are important cellular processes. Thioesters of fatty acids react with cholesterol, forming **cholesteryl esters** in an enzyme-catalyzed reaction (Figure 22.6). These

esters are the principal form in which cholesterol is stored and transported in the body. Because cholesterol is a lipid, insoluble in the aqueous environment of the blood, it travels through the bloodstream in particles that also contain proteins and phospholipids. These particles are classified by their density.


- **LDL particles** (low density lipoproteins) transport cholesterol from the liver to the tissues.
- **HDL particles** (high density lipoproteins) transport cholesterol from the tissues back to the liver, where it is metabolized or converted to other steroids.

Atherosclerosis is a disease that results from the buildup of fatty deposits on the walls of arteries, forming deposits called **plaque.** Plaque is composed largely of the cholesterol (esterified as an ester) of LDL particles. LDL is often referred to as "bad cholesterol" for this reason. In contrast, HDL particles are called "good cholesterol" because they reduce the amount of cholesterol in the bloodstream by transporting it back to the liver.

Figure 22.6
Cholesteryl esters and lipoprotein particles



Problem 22.30 Glucosamine is a dietary supplement available in many over-the-counter treatments for osteoarthritis. Reaction of acetyl CoA with glucosamine forms NAG, *N*-acetylglucosamine, the monomer used to form chitin, the carbohydrate that forms the rigid shells of lobsters and crabs. What is the structure of NAG?




glucosamine

22.18 Nitriles


We end Chapter 22 with the chemistry of **nitriles (RC≡N).** Nitriles have a carbon atom in the same oxidation state as in the acyl compounds that are the principal focus of Chapter 22. Moreover, the chemical reactions of nitriles illustrate some of the concepts first discussed earlier in Chapter 22 and in Chapters 20 and 21.

In addition to the cyanohydrins discussed in Section 21.9, two useful biologically active nitriles are **letrozole** and **anastrozole,** new drugs that reduce the recurrence of breast cancer in women whose tumors are estrogen positive.

Letrozole and anastrozole are called **aromatase inhibitors** because they block the activity of the aromatase enzyme, which is responsible for estrogen synthesis. This inhibits tumor growth in those forms of breast cancer that are stimulated by estrogen.




Generic name letrozole
Trade name Femara




Generic name anastrozole
Trade name Arimidex

Nitriles are readily prepared by S_N2 substitution reactions of unhindered methyl and 1° alkyl halides with ^-CN. This reaction adds one carbon to the alkyl halide and **forms a new carbon–carbon bond.**




new C–C bond in red

Because nitriles have no leaving group, they do not undergo nucleophilic substitution reactions like carboxylic acid derivatives. Because the cyano group contains an electrophilic carbon atom that is part of a multiple bond, a nitrile reacts with nucleophiles by a **nucleophilic addition reaction.** The nature of the nucleophile determines the structure of the product.




Nucleophiles attack here.

The reactions of nitriles with water, hydride, and organometallic reagents as nucleophiles are as follows:




22.18A Hydrolysis of Nitriles

Nitriles are hydrolyzed with water in the presence of acid or base to yield **carboxylic acids** or **carboxylate anions.** In this reaction, the three C−N bonds are replaced by three C−O bonds.



The mechanism of this reaction involves the formation of an amide tautomer. Two tautomers can be drawn for any carbonyl compound, and those for a 1° amide are as follows:




Recall from Chapter 11 that tautomers are constitutional isomers that differ in the location of a double bond and a proton.

- **The amide form is the more stable tautomer, having a C=O and an N−H bond.**
- **The imidic acid tautomer is the less stable form, having a C=N and an O−H bond.**

Recall from Chapter 11 that the keto and enol tautomers of a carbonyl compound are in equilibrium, but the keto form is lower in energy, so it is highly favored in most cases.


The imidic acid and amide tautomers are interconverted by treating with acid or base, analogous to the keto–enol tautomers of other carbonyl compounds. In fact, the two amide tautomers are exactly the same as keto–enol tautomers except that a nitrogen atom replaces a carbon atom bonded to the carbonyl group.



The mechanism of nitrile hydrolysis in both acid and base consists of two parts: [1] **nucleophilic addition** to form the imidic acid tautomer followed by **tautomerization** to form the amide, and [2] **hydrolysis of the amide** to form RCO_2H or RCO_2^-. The mechanism is shown for the basic hydrolysis of RCN to RCO_2^- (Mechanism 22.11).

Mechanism 22.11 Hydrolysis of a Nitrile in Base


Part [1] Conversion of a nitrile to a 1° amide



1 – 2 Nucleophilic attack of ⁻OH followed by protonation forms an imidic acid.


3 – 4 Tautomerization occurs by a two-step sequence—deprotonation followed by protonation.

Part [2] Hydrolysis of the 1° amide to a carboxylate anion




Conversion of the amide to the carboxylate occurs by the three-step sequence in Mechanism 22.10.

Problem 22.31 Draw the products of each reaction.




Problem 22.32 Draw a tautomer of each compound.




22.18B Reduction of Nitriles

Nitriles are reduced with metal hydride reagents to form either 1° amines or aldehydes, depending on the reducing agent.

- **Treatment of a nitrile with $LiAlH_4$ followed by H_2O adds two equivalents of H_2 across the triple bond, forming a 1° amine.**




- **Treatment of a nitrile with a milder reducing agent such as DIBAL-H followed by H_2O forms an aldehyde.**



The mechanism of both reactions involves **nucleophilic addition of hydride (H^-) to the polarized C–N triple bond.** Mechanism 22.12 illustrates that reduction of a nitrile to an amine requires addition of two equivalents of $H:^-$ from $LiAlH_4$. It is likely that intermediate nitrogen anions complex with AlH_3 (formed in situ) to facilitate the addition. Protonation of the dianion in Step [4] forms the amine.

Mechanism 22.12 Reduction of a Nitrile with $LiAlH_4$




1 – 2 Addition of one equivalent of $H:^-$ from $LiAlH_4$ forms an intermediate with one new C–H bond, which complexes with AlH_3.

3 – 4 Nucleophilic attack of a second equivalent of $H:^-$ and complexation with AlH_3 form a dianion, which reacts with water to form two new N–H bonds, giving the 1° amine.


With **DIBAL-H,** nucleophilic addition of one equivalent of hydride forms an anion (Step [1]), which is protonated with water to generate an **imine,** as shown in Mechanism 22.13. As described in Section 21.12, imines are hydrolyzed in water to form aldehydes. Mechanism 22.13 is written without complexation of aluminum with the anion formed in Step [1], to emphasize the identity of intermediates formed during reduction.

Mechanism 22.13 Reduction of a Nitrile with DIBAL-H




1. Addition of $H:^-$ from DIBAL-H (drawn as R_2AlH) forms the new C–H bond.
2. Protonation forms an imine, which is hydrolyzed to an aldehyde by the stepwise sequence shown in Mechanism 21.5.

Problem 22.33 Draw the product of each reaction.




22.18C Addition of Grignard and Organolithium Reagents to Nitriles

Both Grignard and organolithium reagents react with nitriles to form ketones with a new carbon–carbon bond.




The reaction occurs by nucleophilic addition of the organometallic reagent to the polarized C−N triple bond to form an anion (Step [1]), which is protonated with water to form an **imine.** Water then hydrolyzes the imine, replacing the C=N by C=O as described in Section 21.12. The final product is a ketone with a new carbon–carbon bond (Mechanism 22.14).

Mechanism 22.14 Addition of Grignard and Organolithium Reagents (R–M) to Nitriles



1. Addition of $R:^-$ from R'M (M = MgX or Li) forms a new C–C bond.
2. Protonation forms an imine, which is hydrolyzed to a ketone by the stepwise sequence shown in Mechanism 21.5.

Problem 22.34 Draw the products of each reaction.



Problem 22.35 What reagents are needed to convert phenylacetonitrile ($C_6H_5CH_2CN$) to each compound: (a) $C_6H_5CH_2COCH_3$; (b) $C_6H_5CH_2COC(CH_3)_3$; (c) $C_6H_5CH_2CHO$; (d) $C_6H_5CH_2COOH$?

Problem 22.36 Outline two different ways that butan-2-one can be prepared from a nitrile and a Grignard reagent.

KEY CONCEPTS

Carboxylic Acids and Their Derivatives—Nucleophilic Acyl Substitution

Summary of Spectroscopic Absorptions of RCOZ (22.5)

IR absorptions

- All RCOZ compounds have a C=O absorption in the region 1600–1850 cm^{-1}.
  - RCOCl: 1800 cm^{-1}
  - $(RCO)_2O$: 1820 and 1760 cm^{-1} (two peaks)
  - RCO_2R': 1735–1745 cm^{-1}
  - $RCONR'_2$: 1630–1680 cm^{-1}
- Additional amide absorptions occur at 3200–3400 cm^{-1} (N–H stretch) and 1640 cm^{-1} (N–H bending).
- Decreasing the ring size of a cyclic lactone, lactam, or anhydride increases the frequency of the C=O absorption.
- Conjugation shifts the C=O to lower wavenumber.

^{1}H NMR absorptions

- C–H α to the C=O absorbs at 2–2.5 ppm.
- N–H of an amide absorbs at 7.5–8.5 ppm.

^{13}C NMR absorption

- C=O absorbs at 160–180 ppm.

Summary of Spectroscopic Absorptions of RCN (22.5)

IR absorption


- C≡N absorbs at ~2250 cm^{-1}.

^{13}C NMR absorption

- C≡N absorbs at 115–120 ppm.

Summary: The Relationship between the Basicity of Z^- and the Properties of RCOZ

- **Increasing basicity of the leaving group** (22.2)
- **Increasing resonance stabilization** (22.2)



- **Increasing leaving group ability** (22.7B)
- **Increasing reactivity** (22.7B)
- **Increasing frequency of the C=O absorption in the IR** (22.5)


General Features of Nucleophilic Acyl Substitution

- The characteristic reaction of compounds having the general structure RCOZ is nucleophilic acyl substitution (22.1).
- The mechanism consists of two steps (22.7A):
  [1] Addition of a nucleophile to form a tetrahedral intermediate
  [2] Elimination of a leaving group
- More reactive acyl compounds can be used to prepare less reactive acyl compounds. The reverse is not necessarily true (22.7B).

Nucleophilic Acyl Substitution Reactions

[1] Reactions that produce acid chlorides (RCOCl)

From RCO_2H (22.10A):




[2] Reactions that produce anhydrides [$(RCO)_2O$]

a. From RCOCl (22.8):




b. From dicarboxylic acids (22.10B):




cyclic anhydride

[3] Reactions that produce carboxylic acids (RCOOH)


a. From RCOCl (22.8):




b. From $(RCO)_2O$ (22.9):



c. From RCO_2R' (22.11):




d. From $RCONR'_2$ (R' = H or alkyl, 22.13):



[4] Reactions that produce esters (RCOOR')


a. From RCOCl (22.8):



b. From $(RCO)_2O$ (22.9):




c. From RCO_2H (22.10C):




[5] Reactions that produce amides ($RCONH_2$) [The reactions are written with NH_3 as nucleophile to form $RCONH_2$. Similar reactions occur with $R'NH_2$ to form RCONHR', and with R'_2NH to form $RCONR'_2$.]


a. From RCOCl (22.8):



b. From $(RCO)_2O$ (22.9):



c. From RCO_2H (22.10D):






d. From RCO_2R' (22.11):




Nitrile Synthesis (22.18)

Nitriles are prepared by S_N2 substitution using unhindered alkyl halides as starting materials.




Reactions of Nitriles


[1] Hydrolysis (22.18A)



[2] Reduction (22.18B)




[3] Reaction with organometallic reagents (22.18C)



PROBLEMS

Problems Using Three-Dimensional Models

22.37 Rank the following compounds in order of increasing reactivity in nucleophilic acyl substitution.




22.38 (a) Give an acceptable name for each compound. (b) Draw the organic products formed when **A** or **B** is treated with each reagent: [1] H_3O^+; [2] ^-OH, H_2O; [3] $CH_3CH_2CH_2MgBr$ (excess), then H_2O; [4] $LiAlH_4$, then H_2O.



A **B**


22.39 Which ester, **C** or **D,** is more reactive in nucleophilic acyl substitution? Explain your reasoning.



C **D**

Nomenclature

22.40 Give the IUPAC or common name for each compound.




22.41 Give the structure corresponding to each name.

a. cyclohexyl propanoate
b. cyclohexanecarboxamide
c. 4-methylheptanenitrile
d. vinyl acetate
e. benzoic propanoic anhydride
f. 3-methylhexanoyl chloride
g. octyl butanoate
h. *N,N*-dibenzylformamide

Properties of Carboxylic Acid Derivatives


22.42 Explain why imidazolides are much more reactive than other amides in nucleophilic acyl substitution.




imidazolide

22.43 Explain why CH_3CONH_2 is a stronger acid and a weaker base than $CH_3CH_2NH_2$.

22.44 (a) Propose an explanation for the difference in the frequency of the carbonyl absorptions of phenyl acetate (1765 cm^{-1}) and cyclohexyl acetate (1738 cm^{-1}). (b) Which carbonyl group is more effectively stabilized by resonance? (c) Which ester reacts faster when treated with aqueous base?



phenyl acetate



cyclohexyl acetate

Reactions


22.45 Draw the product formed when phenylacetic acid ($C_6H_5CH_2COOH$) is treated with each reagent. With some reagents, no reaction occurs.


a. $NaHCO_3$
b. NaOH
c. $SOCl_2$
d. NaCl
e. NH_3 (1 equiv)
f. NH_3, Δ
g. CH_3OH, H_2SO_4
h. CH_3OH, ^-OH
i. [1] NaOH; [2] CH_3COCl
j. CH_3NH_2, DCC
k. [1] $SOCl_2$; [2] $CH_3CH_2CH_2NH_2$ (excess)
l. [1] $SOCl_2$; [2] $(CH_3)_2CHOH$


22.46 Draw the product formed when phenylacetonitrile ($C_6H_5CH_2CN$) is treated with each reagent.


a. H_3O^+
b. H_2O, ^-OH
c. [1] CH_3MgBr; [2] H_2O
d. [1] CH_3CH_2Li; [2] H_2O
e. [1] DIBAL-H; [2] H_2O
f. [1] $LiAlH_4$; [2] H_2O


22.47 Draw the organic products formed in each reaction.


a. 


b. 


c. 

d. 


e. 

f. 


g. 

h. 


22.48 Cinnamoylcocaine, a natural product that occurs in coca leaves, can be converted to cocaine by the following reaction sequence. Identify the structure of cinnamoylcocaine, as well as intermediates **X** and **Y.**




22.49 What products are formed by hydrolysis of each lactone or lactam with acid?



22.50 Identify compounds **A–M** in the following reaction sequence.




22.51 Draw the products of each reaction and indicate the stereochemistry at any stereogenic centers.



22.52 What products are formed when all of the amide and ester bonds are hydrolyzed in each of the following compounds? **Tamiflu** [part (a)] is the trade name of the antiviral agent oseltamivir, thought to be the most effective agent in treating influenza. **Aspartame** [part (b)] is the artificial sweetener used in Equal and many diet beverages. One of the products of this hydrolysis reaction is the amino acid phenylalanine. Infants afflicted with phenylketonuria cannot metabolize this amino acid, so it accumulates, causing mental retardation. When the


affliction is identified early, a diet limiting the consumption of phenylalanine (and compounds like aspartame that are converted to it) can make a normal life possible.

a.




oseltamivir

b.




aspartame

22.53 Identify **F** in the following reaction sequence. **F** was converted in several steps to the antidepressant paroxetine (trade name Paxil; see also Problem 9.9).




Mechanism


22.54 Draw a stepwise mechanism for the following reaction.



22.55 When acetic acid (CH_3COOH) is treated with a trace of acid in water labeled with ^{18}O, the label gradually appears in both oxygen atoms of the carboxylic acid. Draw a mechanism that explains this phenomenon.




22.56 Although γ-butyrolactone (Problem 19.62) is a biologically inactive compound, it is converted in the body to 4-hydroxybutanoic acid (GHB), an addictive and intoxicating recreational drug (Section 19.5). Draw a stepwise mechanism for this conversion in the presence of acid.



γ-butyrolactone


4-hydroxybutanoic acid
GHB

22.57 Aspirin is an anti-inflammatory agent because it inhibits the conversion of arachidonic acid to prostaglandins by the transfer of its acetyl group (CH_3CO-) to an OH group at the active site of an enzyme (Section 19.6). This reaction, called transesterification, results in the conversion of one ester to another by a nucleophilic acyl substitution reaction. Draw a stepwise mechanism for the given transesterification.




aspirin enzyme inactive enzyme salicylic acid


22.58 Draw a stepwise mechanism for the following reaction.



22.59 Draw a stepwise mechanism for the following reaction, one step in the synthesis of the cholesterol-lowering drug ezetimibe (Section 20.6).




22.60 Draw a stepwise mechanism for the conversion of lactone **C** to carboxylic acid **D. C** is a key intermediate in the synthesis of prostaglandins (Section 19.6) by Nobel Laureate E. J. Corey and co-workers at Harvard University.




C **D**

22.61 Draw a stepwise mechanism for the conversion of lactone **A** to ester **B** using HCl in ethanol. **B** is converted in one step to ethyl chrysanthemate, a useful intermediate in the synthesis of a variety of pyrethrins, naturally occurring insecticides with three-membered rings that are isolated from chrysanthemums (Section 26.4).



A **B** ethyl chrysanthemate


22.62 Draw a stepwise mechanism for the following reaction.




22.63 Acid-catalyzed hydrolysis of $HOCH_2CH_2C(CH_3)_2CN$ forms compound **A** ($C_6H_{10}O_2$). **A** shows a strong peak in its IR spectrum at 1770 cm^{-1} and the following signals in its ^1H NMR spectrum: 1.27 (singlet, 6 H), 2.12 (triplet, 2 H), and 4.26 (triplet, 2 H) ppm. Draw the structure for **A** and give a stepwise mechanism that accounts for its formation.


Synthesis

22.64 What carboxylic acid and alcohol are needed to prepare each ester by Fischer esterification?




22.65 Devise a synthesis of each compound using 1-bromobutane ($CH_3CH_2CH_2CH_2Br$) as the only organic starting material. You may use any other inorganic reagents.

a. 



b. 

22.66 Two methods convert an alkyl halide into a carboxylic acid having one more carbon atom.

[1] R—X $\xrightarrow[\text{[2] } H_3O^+]{\text{[1] } {}^-CN}$  (Section 20.18)


[2] R—X $\xrightarrow[\text{[2] } CO_2\\ \text{[3] } H_3O^+]{\text{[1] Mg}}$  (Section 20.14A)


Depending on the structure of the alkyl halide, one or both of these methods may be employed. For each alkyl halide, write out a stepwise sequence that converts it to a carboxylic acid with one more carbon atom. If both methods work, draw both routes. If one method cannot be used, state why it can't.


a. CH_3Cl b.  c. 

22.67 Devise a synthesis of benzocaine, ethyl *p*-aminobenzoate ($H_2NC_6H_4CO_2CH_2CH_3$), from benzene, organic alcohols, and any needed organic or inorganic reagents. Benzocaine is the active ingredient in the topical anesthetic Orajel (Section 18.15C).


22.68 Devise a synthesis of each analgesic compound from phenol (C_6H_5OH) and any other organic or inorganic reagents.


a. 
salicylamide

b. 
acetaminophen

c. 
p-acetophenetidin


22.69 Devise a synthesis of each compound from benzene and organic alcohols containing four or fewer carbons. You may also use any required organic or inorganic reagents.


a. 

b. 


Polymers


22.70 What polyester or polyamide can be prepared from each pair of monomers?

a. 

b. 


22.71 What two monomers are needed to prepare each polymer?

a. 


b. 

General Problem

22.72 Taxotere is the trade name for the synthetic anticancer drug docetaxel, whose structure closely resembles the naturally occurring compound paclitaxel, which is isolated from the Pacific yew tree (Section 5.5).



docetaxel



carbamate

a. Taxol's limited water solubility led to an extensive search for related compounds with increased water solubility. What structural features give docetaxel a better water solubility profile?


b. Docetaxel contains a carbamate (labeled in red). Draw three more resonance structures for a carbamate. Rank all four resonance structures in order of increasing stability.

c. A carbamate with a *tert*-butoxy group [$(CH_3)_3CO-$] is hydrolyzed according to the given equation. Draw a stepwise mechanism for this reaction.


d. Assuming that all ester and carbamate bonds are cleaved, draw all products formed when docetaxel is hydrolyzed with aqueous acid.

Spectroscopy

22.73 How can IR spectroscopy be used to distinguish between each pair of isomers?



22.74 Rank the following compounds in order of increasing frequency of the C=O absorption in their IR spectra.



22.75 Identify the structures of each compound from the given data.

a. Molecular formula	$C_6H_{12}O_2$
IR absorption:	1738 cm^{-1}
^1H NMR:	1.12 (triplet, 3 H), 1.23 (doublet, 6 H), 2.28 (quartet, 2 H), and 5.00 (septet, 1 H) ppm
b. Molecular formula	C_4H_7N
IR absorption:	2250 cm^{-1}
^1H NMR:	1.08 (triplet, 3 H), 1.70 (multiplet, 2 H), and 2.34 (triplet, 2 H) ppm


c. Molecular formula C_8H_9NO
IR absorptions: 3328 and 1639 cm^{-1}
^1H NMR: 2.95 (singlet, 3 H), 6.95 (singlet, 1 H), and 7.3–7.7 (multiplet, 5 H) ppm

d. Molecular formula C_4H_7ClO
IR absorption: 1802 cm^{-1}
^1H NMR: 0.95 (triplet, 3 H), 1.07 (multiplet, 2 H), and 2.90 (triplet, 2 H) ppm


e. Molecular formula $C_{10}H_{12}O_2$
IR absorption: 1740 cm^{-1}
^1H NMR: 1.2 (triplet, 3 H), 2.4 (quartet, 2 H), 5.1 (singlet, 2 H), and 7.1–7.5 (multiplet, 5 H) ppm

22.76 Identify the structures of **A** and **B,** isomers of molecular formula $C_{10}H_{12}O_2$, from their IR data and ^1H NMR spectra.


a. IR absorption for **A** at 1718 cm^{-1}




b. IR absorption for **B** at 1740 cm^{-1}




22.77 Phenacetin is an analgesic compound having molecular formula $C_{10}H_{13}NO_2$. Once a common component in over-the-counter pain relievers such as APC (**a**spirin, **p**henacetin, **c**affeine), phenacetin is no longer used because of its liver toxicity. Deduce the structure of phenacetin from its ^1H NMR and IR spectra.






22.78 Identify the structure of compound **C** (molecular formula $C_{11}H_{15}NO_2$), which has an IR absorption at 1699 cm^{-1} and the ^1H NMR spectrum shown below.




22.79 Identify the structures of **D** and **E,** isomers of molecular formula $C_6H_{12}O_2$, from their IR and ^1H NMR data. Signals at 1.35 and 1.60 ppm in the ^1H NMR spectrum of **D** and 1.90 ppm in the ^1H NMR spectrum of **E** are multiplets.

a. IR absorption for **D** at 1743 cm^{-1}




b. IR absorption for **E** at 1746 cm^{-1}




Challenge Problems

22.80 With reference to amides **A** and **B,** the carbonyl of one amide absorbs at a much higher wavenumber in its IR spectrum than the carbonyl of the other amide. Which absorbs at higher wavenumber and why?



A




B


22.81 The ^1H NMR spectrum of 2-chloroacetamide ($ClCH_2CONH_2$) shows three signals at 4.02, 7.35, and 7.60 ppm. What protons give rise to each signal? Explain why three signals are observed.

22.82 Compelling evidence for the existence of a tetrahedral intermediate in nucleophilic acyl substitution was obtained in a series of elegant experiments carried out by Myron Bender in 1951. The key experiment was the reaction of aqueous ^-OH with ethyl benzoate ($C_6H_5COOCH_2CH_3$) labeled at the carbonyl oxygen with ^{18}O. Bender did not allow the hydrolysis to go to completion, and then examined the presence of a label in the *recovered*


starting material. He found that some of the recovered ethyl benzoate no longer contained a label at the carbonyl oxygen. With reference to the accepted mechanism of nucleophilic acyl substitution, explain how this provides evidence for a tetrahedral intermediate.




22.83 Draw a stepwise mechanism for the following reactions, two steps in R. B. Woodward's classic synthesis of reserpine in 1958. Reserpine, which is isolated from the extracts of the Indian snakeroot *Rauwolfia serpentina Benth,* was used at one time to manage mild hypertension associated with anxiety.



22.84 Draw a stepwise mechanism for the following reaction, the last step in a five-step industrial synthesis of vitamin C that begins with the simple carbohydrate glucose.



22.85 Draw a stepwise mechanism for the following reaction, a key step in the synthesis of linezolid, an antibacterial agent.



23 Substitution Reactions of Carbonyl Compounds at the α Carbon

23.1 Introduction
23.2 Enols
23.3 Enolates
23.4 Enolates of unsymmetrical carbonyl compounds
23.5 Racemization at the α carbon
23.6 A preview of reactions at the α carbon
23.7 Halogenation at the α carbon
23.8 Direct enolate alkylation
23.9 Malonic ester synthesis
23.10 Acetoacetic ester synthesis



Tamoxifen is a potent anticancer drug used widely in the treatment of breast cancer. Tamoxifen binds to estrogen receptors, and in this way inhibits the growth of breast cancers that are estrogen dependent. One method to synthesize tamoxifen forms a new carbon–carbon bond on the α carbon to a carbonyl group using an intermediate enolate. In Chapter 23 we learn about these and other carbon–carbon bond-forming reactions that occur at the α carbon.


Chapters 23 and 24 focus on reactions that occur at the α carbon to a carbonyl group. These reactions are different from the reactions of Chapters 20–22, all of which involved nucleophilic attack at the electrophilic carbonyl carbon. In reactions at the α carbon, the carbonyl compound serves as a *nucleophile* that reacts with a carbon or halogen electrophile to form a new bond to the α carbon.

Chapter 23 concentrates on **substitution reactions at the α carbon,** whereas Chapter 24 concentrates on reactions between two carbonyl compounds, one of which serves as the nucleophile and one of which is the electrophile. Many of the reactions in Chapter 23 form new carbon–carbon bonds, thus adding to your repertoire of reactions that can be used to synthesize more complex organic molecules from simple precursors. As you will see, the reactions introduced in Chapter 23 have been used to prepare a wide variety of interesting and useful compounds.


23.1 Introduction

Up to now, the discussion of carbonyl compounds has centered on their reactions with nucleophiles at the electrophilic carbonyl carbon. **Two general reactions are observed,** depending on the structure of the carbonyl starting material.

- *Nucleophilic addition* occurs when there is no electronegative atom Z on the carbonyl carbon (as with aldehydes and ketones).




- *Nucleophilic acyl substitution* occurs when there is an electronegative atom Z on the carbonyl carbon (as with carboxylic acids and their derivatives).




Reactions can also occur at the α carbon to the carbonyl group. These reactions proceed by way of **enols** or **enolates,** two electron-rich intermediates that react with electrophiles, forming a new bond on the α carbon. This reaction results in the **substitution of the electrophile E for hydrogen.**

Hydrogen atoms on the α carbon are called **α hydrogens.**



23.2 Enols


Recall from Chapter 11 that **enol and keto forms are tautomers of the carbonyl group that differ in the position of a double bond and a proton.** These constitutional isomers are in equilibrium with each other.




keto form enol form

- A keto tautomer has a C=O and an additional C–H bond.
- An enol tautomer has an O–H group bonded to a C=C.

Equilibrium favors the keto form for most carbonyl compounds largely because a C=O is much stronger than a C=C. For simple carbonyl compounds, < 1% of the enol is present at equilibrium. With unsymmetrical ketones, moreover, two different enols are possible, yet they still total < 1%.



With compounds containing two carbonyl groups separated by a single carbon (called β-dicarbonyl compounds or 1,3-dicarbonyl compounds), however, the concentration of the enol form sometimes exceeds the concentration of the keto form.




pentane-2,4-dione
β-dicarbonyl compound
24% keto tautomer

76% enol tautomers

Two factors stabilize the enol of β-dicarbonyl compounds: **conjugation** and **intramolecular hydrogen bonding.** The C=C of the enol is conjugated with the carbonyl group, allowing delocalization of the electron density in the π bonds. Moreover, the OH of the enol can hydrogen bond to the oxygen of the nearby carbonyl group. Such intramolecular hydrogen bonds are especially stabilizing when they form a six-membered ring, as in this case.

Sample Problem 23.1 Convert each compound to its enol or keto tautomer.


a. 

b. 


Solution

a. To convert a carbonyl compound to its enol tautomer, draw a double bond between the carbonyl carbon and the α carbon, and change the C=O to C–OH. In this case, both α carbons are identical, so only one enol is possible.

b. To convert an enol to its keto tautomer, change the C–OH to C=O and add a proton to the other end of the C=C.




Problem 23.1 Draw the enol or keto tautomer(s) of each compound.



23.2A The Mechanism of Tautomerization

Tautomerization, the process of converting one tautomer into another, is catalyzed by both acid and base. Tautomerization always requires two steps **(protonation** and **deprotonation),** but the order of these steps depends on whether the reaction takes place in acid or base. In Mechanisms 23.1 and 23.2 for tautomerization, the keto form is converted to the enol form. All of the steps are reversible, though, so they equally apply to the conversion of the enol form to the keto form.


Mechanism 23.1 Tautomerization in Acid



1 With acid, protonation *precedes* deprotonation. Protonation of the carbonyl forms a resonance-stabilized carbocation.

2 Removal of a proton forms the enol.


Mechanism 23.2 Tautomerization in Base



1 With base, deprotonation *precedes* protonation. Removal of a proton on the α carbon forms a resonance-stabilized enolate.

2 Protonation of the enolate forms the enol.


Problem 23.2 During the metabolism of glucose, glyceraldehyde 3-phosphate is converted to dihydroxyacetone phosphate by a process that involves two keto–enol tautomerizations. Draw a stepwise mechanism for this reaction in the presence of acid.



glyceraldehyde 3-phosphate dihydroxyacetone phosphate


23.2B How Enols React

Like other compounds with carbon–carbon double bonds, **enols are electron rich, so they react as nucleophiles.** Enols are even more electron rich than alkenes, though, because the OH group has a powerful electron-donating resonance effect. A second resonance structure can be drawn for the enol that places a negative charge on one of the carbon atoms. As a result, this carbon atom is especially nucleophilic, and it can react with an electrophile E^+ to form a new bond to carbon. Loss of a proton then forms a neutral product.



two resonance structures for an enol

- **Reaction of an enol with an electrophile E^+ forms a new C–E bond on the α carbon. The net result is substitution of H by E on the α carbon.**




Problem 23.3 When phenylacetaldehyde ($C_6H_5CH_2CHO$) is dissolved in D_2O with added DCl, the hydrogen atoms α to the carbonyl are gradually replaced by deuterium atoms. Write a mechanism for this process that involves enols as intermediates.

23.3 Enolates


Enolates are formed when a base removes a proton on the α carbon to a carbonyl group. A C–H bond on the α carbon is more acidic than many other sp^3 hybridized C–H bonds, because the resulting enolate is resonance stabilized. Moreover, one of the resonance structures is especially stable because it places a negative charge on an electronegative oxygen atom.

Forming enolates from carbonyl compounds was first discussed in Section 21.7.





resonance-stabilized enolate anion

Enolates are always formed by removal of a proton on the **α carbon.**



The pK_a of the α hydrogen in an aldehyde or ketone is **~20.** As shown in Table 23.1, this makes it considerably more acidic than the C–H bonds in CH_3CH_3 and $CH_3CH{=}CH_2$. Although C–H bonds α to a carbonyl are more acidic than many other C–H bonds, they are still less acidic than O–H bonds that always place the negative charge of the conjugate base on an electronegative oxygen atom (c.f. CH_3CH_2OH and CH_3COOH in Table 23.1).

Table 23.1 A Comparison of pK_a Values


Compound	pK_a	Conjugate base	Structural features of the conjugate base
CH_3CH_3	50	$CH_3\ddot{C}H_2$	• The conjugate base has a (–) charge on C, but is not resonance stabilized.
	43		• The conjugate base has a (–) charge on C, and is resonance stabilized.
	19.2		• **The conjugate base has two resonance structures, one of which has a (–) charge on O.**
	16		• The conjugate base has a (–) charge on O, but is not resonance stabilized.
	4.8		• The conjugate base has two resonance structures, both of which have a (–) charge on O.

Increasing acidity
Increasing stability of the conjugate base

- **Resonance stabilization of the conjugate base increases acidity.**
  - $CH_2{=}CHCH_3$ is more acidic than CH_3CH_3.
  - CH_3COOH is more acidic than CH_3CH_2OH.
- **Placing a negative charge on O in the conjugate base increases acidity.**
  - CH_3CH_2OH is more acidic than CH_3CH_3.
  - CH_3COCH_3 is more acidic than $CH_2{=}CHCH_3$.
  - CH_3COOH (with two O atoms) is more acidic than CH_3COCH_3.

The electrostatic potential plots in Figure 23.1 compare the electron density of the acetone enolate, which is resonance stabilized and delocalized, with that of $(CH_3)_2CHO^-$, an alkoxide that is not resonance stabilized.


Figure 23.1
Electron density in an enolate and an alkoxide




- The acetone enolate is resonance stabilized. The negative charge is delocalized on the oxygen atom (pale red) and the carbon atom (pale green).
- The alkoxide anion is *not* resonance stabilized. The negative charge is concentrated on the oxygen atom only (deep red).

23.3A Examples of Enolates and Related Anions

In addition to enolates from aldehydes and ketones, **enolates from esters and 3° amides can be formed as well,** although the α hydrogen is somewhat less acidic. **Nitriles** also have acidic protons on the carbon atom adjacent to the cyano group, because the negative charge of the conjugate base is stabilized by delocalization onto an electronegative nitrogen atom.



The protons on the carbon between the two carbonyl groups of a β-dicarbonyl compound are especially acidic because resonance delocalizes the negative charge on two different oxygen atoms. Table 23.2 lists pK_a values for β-dicarbonyl compounds as well as other carbonyl compounds and nitriles.



Problem 23.4 Draw additional resonance structures for each anion.


















Table 23.2 pK_a Values for Some Carbonyl Compounds and Nitriles

Compound type	Example	pK_a	Compound type	Example	pK_a
[1] Amide		30	[6] 1,3-Diester		13.3
[2] Nitrile		25	[7] 1,3-Dinitrile		11
[3] Ester		25	[8] β-Keto ester		10.7
[4] Ketone		19.2	[9] β-Diketone		9
[5] Aldehyde		17			

Problem 23.5 Which C–H bonds in the following molecules are acidic because the resulting conjugate base is resonance stabilized?


a. 
b. 
c. 
d. 

Problem 23.6 Rank the protons in the labeled CH_2 groups in order of increasing acidity, and explain why you chose this order.



23.3B The Base

The formation of an enolate is an acid–base equilibrium, so the **stronger the base, the more enolate that forms.**



Stronger bases drive the equilibrium to the right.

We have now used the term ***amide*** in two different ways—first as a functional group (e.g., the carboxylic acid derivative $RCONH_2$) and now as a base (e.g., $^-NH_2$, which can be purchased as a sodium or lithium salt, $NaNH_2$ or $LiNH_2$, respectively). In Chapter 23 we will use dialkylamides, $^-NR_2$, in which the two H atoms of $^-NH_2$ have been replaced by R groups.

We can predict the extent of an acid–base reaction by comparing the pK_a of the starting acid (the carbonyl compound in this case) with the pK_a of the conjugate acid formed. **The equilibrium favors the side with the weaker acid (the acid with the higher pK_a value).** The pK_a of many carbonyl compounds is ~20, so a significant amount of enolate will form only if the pK_a of the conjugate acid is > 20.

The common bases used to form enolates are hydroxide (^-OH), various alkoxides (^-OR), hydride (H^-), and dialkylamides ($^-NR_2$). How much enolate is formed using each of these bases is indicated in Table 23.3.

Table 23.3 Enolate Formation with Various Bases: $RCOCH_3$ (p$K_a \simeq 20$) + B: → $RCOCH_2^-$ + HB^+


	Base (B:)	Conjugate acid (HB^+)	pK_a of HB^+	% Enolate
[1]	$Na^+ {}^-OH$	H_2O	15.7	< 1%
[2]	$Na^+ {}^-OCH_2CH_3$	CH_3CH_2OH	16	< 1%
[3]	$K^+ {}^-OC(CH_3)_3$	$(CH_3)_3COH$	18	1–10%
[4]	Na^+H^-	H_2	35	100%
[5]	$Li^+ {}^-N[CH(CH_3)_2]_2$	$HN[CH(CH_3)_2]_2$	40	100%



Enolate formation with LDA is typically carried out at –78 °C, a convenient temperature to maintain in the laboratory because it is the temperature at which dry ice (solid CO_2) sublimes. A low-temperature cooling bath can be made by adding dry ice to acetone until the acetone cools to –78 °C. Immersing a reaction flask in this cooling bath keeps its contents at a constant low temperature.


When the pK_a of the conjugate acid is < 20, as it is for ^-OH and all ^-OR (entries 1–3), only a small amount of enolate is formed at equilibrium. These bases are more useful in forming enolates when more acidic 1,3-dicarbonyl compounds are used as starting materials. They are also used when both the enolate and the carbonyl starting material are involved in the reaction, as is the case for reactions described in Chapter 24.

To form an enolate in essentially 100% yield, a much stronger base such as lithium diisopropylamide, **$Li^+ {}^-N[CH(CH_3)_2]_2$,** abbreviated as **LDA,** is used (entry 5). **LDA is a strong nonnucleophilic base.** Like the other nonnucleophilic bases (Sections 7.8B and 8.1), its bulky isopropyl groups make the nitrogen atom too hindered to serve as a nucleophile. It is still able, though, to remove a proton in an acid–base reaction.



lithium diisopropylamide

LDA




The N atom is too crowded to be a nucleophile.



THF
tetrahydrofuran
a polar aprotic solvent

LDA quickly deprotonates essentially all of the carbonyl starting material, even at –78 °C, to form the enolate product. THF is the typical solvent for these reactions.




pK_a = 20


diisopropylamine
pK_a = 40

Equilibrium greatly favors the products.
Essentially all of the ketone is converted to enolate.

LDA can be prepared by deprotonating diisopropylamine with an organolithium reagent such as butyllithium, and then used immediately in a reaction.




Problem 23.7 Draw the product formed when each starting material is treated with LDA in THF solution at –78 °C.



Problem 23.8 As we learned in Chapter 20, organolithium reagents (RLi) are strong bases that readily react with acidic protons. Why aren't organolithium reagents used to generate enolates?

23.3C General Reactions of Enolates

Enolates are nucleophiles, and as such they react with many electrophiles. Because an enolate is resonance stabilized, however, it has two reactive sites—the carbon and oxygen atoms that bear the negative charge. **A nucleophile with two reactive sites is called an *ambident nucleophile*.** In theory, each of these atoms could react with an electrophile to form two different products, one with a new bond to carbon, and one with a new bond to oxygen.




Because enolates usually react at carbon instead of oxygen, the resonance structure that places the negative charge on oxygen will often be omitted in multistep mechanisms.

An enolate usually reacts at the carbon end, however, because this site is more nucleophilic. Thus, **enolates generally react with electrophiles on the α carbon,** so that many reactions in Chapter 23 follow a two-step path:

[1] Reaction of a carbonyl compound with base forms an enolate.
[2] Reaction of the enolate with an electrophile forms a new bond on the α carbon.

23.4 Enolates of Unsymmetrical Carbonyl Compounds

What happens when an unsymmetrical carbonyl compound like 2-methylcyclohexanone is treated with base? **Two enolates are possible,** one formed by removal of a 2° hydrogen, and one formed by removal of a 3° hydrogen.



Path [1] occurs *faster* than Path [2] because it results in removal of the less hindered 2° hydrogen, forming an enolate on the less substituted α carbon. Path [2] results in removal of a 3° hydrogen, forming the *more stable* enolate with the more substituted double bond. This enolate predominates at equilibrium.

- **The kinetic enolate is formed faster because it is the less substituted enolate.**
- **The thermodynamic enolate is lower in energy because it is the more substituted enolate.**

It is possible to regioselectively form one or the other enolate by the proper use of reaction conditions, because the base, solvent, and reaction temperature all affect the identity of the enolate formed.


Kinetic Enolates

The kinetic enolate forms faster, so mild reaction conditions favor it over slower processes with higher energies of activation. It is the less stable enolate, so it must not be allowed to equilibrate to the more stable thermodynamic enolate. **The kinetic enolate is favored by:**

[1] **A strong nonnucleophilic base.** A strong base assures that the enolate is formed rapidly. A **bulky base like LDA removes the more accessible proton on the less substituted carbon** much faster than a more hindered proton.

[2] **Polar aprotic solvent.** The solvent must be polar to dissolve the polar starting materials and intermediates. It must be aprotic so that it does not protonate any enolate that is formed. **THF** is both polar and aprotic.

[3] **Low temperature.** The temperature must be low **(–78 °C)** to prevent the kinetic enolate from equilibrating to the thermodynamic enolate.



- **A kinetic enolate is formed with a strong, nonnucleophilic base (LDA) in a polar aprotic solvent (THF) at low temperature (–78 °C).**

Thermodynamic Enolates


A thermodynamic enolate is favored by equilibrating conditions. This is often achieved using a **strong base in a protic solvent.** A strong base yields both enolates, but in a protic solvent, enolates can also be protonated to re-form the carbonyl starting material. At equilibrium, the lower energy intermediate always wins out, so that **the more stable, more substituted enolate is present in higher concentration.** Thus, the **thermodynamic enolate is favored by:**

[1] **A strong base. $Na^+ {}^-OCH_2CH_3$, $K^+ {}^-OC(CH_3)_3$,** or other alkoxides are common.

[2] **Protic solvent. CH_3CH_2OH** or other alcohols.


[3] **Room temperature (25 °C).**

To simplify structures, we use abbreviations:
Me = CH_3, so $NaOCH_3$ = NaOMe
Et = CH_2CH_3, so $NaOCH_2CH_3$ = NaOEt
*t*Bu = $C(CH_3)_3$, so $KOC(CH_3)_3$ = KO*t*Bu



- **A thermodynamic enolate is formed with a strong base (RO^-) in a polar protic solvent (ROH) at room temperature.**


Sample Problem 23.2 What is the major enolate formed in each reaction?




Solution

a. LDA is a strong, nonnucleophilic base that removes a proton on the less substituted α carbon to form the kinetic enolate.

b. $NaOCH_2CH_3$ (a strong base) and CH_3CH_2OH (a protic solvent) favor removal of a proton from the more substituted α carbon to form the thermodynamic enolate.



Problem 23.9 What enolate is formed when each ketone is treated with LDA in THF solution? What enolate is formed when these same ketones are treated with $NaOCH_3$ in CH_3OH solution?



23.5 Racemization at the α Carbon


Recall from Section 16.5 that an enolate can be stabilized by the delocalization of electron density only if it possesses the proper geometry and hybridization.

- **The electron pair on the carbon adjacent to the C=O must occupy a *p* orbital that overlaps with the two other *p* orbitals of the C=O, making an enolate conjugated.**

- **Thus, all three atoms of the enolate are sp^2 hybridized and trigonal planar.**


These bonding features are shown in the acetone enolate in Figure 23.2.

Figure 23.2
The hybridization and geometry of the acetone enolate $(CH_3COCH_2)^-$



- The O atom and both C's of the enolate are sp^2 hybridized and lie in a plane.
- Each atom has a *p* orbital extending above and below the plane; these orbitals overlap to delocalize electron density.

When the α carbon to the carbonyl is a stereogenic center, treatment with aqueous base leads to **racemization** by a two-step process: **deprotonation to form an enolate and protonation to re-form the carbonyl compound.** For example, chiral ketone **A** reacts with aqueous ^-OH to form an achiral enolate having an sp^2 hybridized α carbon. Because the enolate is planar, it can be protonated with H_2O with equal probability from both directions, yielding a racemic mixture of two ketones.




Problem 23.10 Explain each observation: (a) When (*R*)-2-methylcyclohexanone is treated with NaOH in H_2O, the optically active solution gradually loses optical activity. (b) When (*R*)-3-methylcyclohexanone is treated with NaOH in H_2O, the solution remains optically active.

23.6 A Preview of Reactions at the α Carbon

Having learned about the synthesis and properties of enolates, we can now turn our attention to their reactions. Like enols, **enolates are nucleophiles,** but because they are negatively charged, enolates are much more nucleophilic than neutral enols. Consequently, they undergo a wider variety of reactions.


Two general types of reactions of enolates—**substitutions** and **reactions with other carbonyl compounds**—will be discussed in the remainder of Chapter 23 and in Chapter 24. Both reactions form new bonds to the carbon α to the carbonyl.

- Enolates react with electrophiles to afford substitution products.



Two different kinds of substitution reactions are examined: **halogenation** with X_2 and **alkylation** with alkyl halides RX. These reactions are detailed in Sections 23.7–23.10.


- Enolates react with other carbonyl groups at the electrophilic carbonyl carbon.



These reactions are more complicated because the initial addition adduct goes on to form different products depending on the structure of the carbonyl group. These reactions form the subject of Chapter 24.

23.7 Halogenation at the α Carbon

The first substitution reaction we examine is **halogenation.** Treatment of a ketone or aldehyde with halogen and either acid or base results in **substitution of X for H on the α carbon,** forming an **α-halo aldehyde or ketone.** Halogenation readily occurs with Cl_2, Br_2, and I_2.




The mechanisms of halogenation in acid and base are somewhat different.

- Reactions done in acid generally involve *enol* intermediates.
- Reactions done in base generally involve *enolate* intermediates.


23.7A Halogenation in Acid

Halogenation is often carried out by treating a carbonyl compound with a halogen in acetic acid. In this way, acetic acid is both the solvent and the acid catalyst for the reaction.



The mechanism of acid-catalyzed halogenation consists of two parts: **tautomerization** of the carbonyl compound to the enol form, and **reaction of the enol with halogen.** Mechanism 23.3 illustrates the reaction of $(CH_3)_2C{=}O$ with Br_2 in CH_3CO_2H.


Mechanism 23.3 Acid-Catalyzed Halogenation at the α Carbon



1 – 2 The ketone is converted to its enol tautomer by the two-step process of protonation followed by deprotonation.

3 – 4 Addition of the halogen to the enol forms a new bond to Br on the α carbon, and deprotonation yields the substitution product.


Problem 23.11 Draw the products of each reaction.



23.7B Halogenation in Base

Halogenation in base is much less useful, because it is often difficult to stop the reaction after addition of just one halogen atom to the α carbon. For example, treatment of propiophenone with Br_2 and aqueous ^-OH yields a dibromo ketone.


Reactions of carbonyl compounds with base invariably involve enolates because the α hydrogens of the carbonyl compound are easily removed.



propiophenone

The mechanism for introduction of each Br atom involves the same two steps: **deprotonation with base followed by reaction with Br_2** to form a new C–Br bond, as shown in Mechanism 23.4.


Mechanism 23.4 Halogenation at the α Carbon in Base




1 Treatment of the ketone with base forms a nucleophilic enolate.

2 Reaction of the enolate with Br_2 forms the substitution product in which one H is replaced by Br on the α carbon.

Only a small amount of the enolate forms at equilibrium using $^-$OH as base, but the enolate is such a strong nucleophile that it readily reacts with Br_2, thus driving the equilibrium to the right. Then, the same two steps introduce the second Br atom on the α carbon: **deprotonation** followed by **nucleophilic attack.**




The electronegative Br stabilizes the negative charge.



Although all ketones with α hydrogens react with base and I_2, only **methyl** ketones form CHI_3 (iodoform), a pale yellow solid that precipitates from the reaction mixture. This reaction is the basis of the **iodoform test,** once a common chemical method to detect methyl ketones. Methyl ketones give a positive iodoform test (appearance of a yellow solid), whereas other ketones give a negative iodoform test (no change in the reaction mixture).

It is difficult to stop this reaction after the addition of one Br atom because the electron-withdrawing inductive effect of Br stabilizes the second enolate. As a result, the α H of α-bromopropiophenone is more acidic than the α H atoms of propiophenone, making it easier to remove with base.

Halogenation of a methyl ketone with excess halogen, called the **haloform reaction,** results in cleavage of a carbon–carbon σ bond and formation of two products, a carboxylate anion and CHX_3 (commonly called **haloform**).




The C–C bond in red is broken.

In the haloform reaction, the three H atoms of the CH_3 group are successively replaced by X to form an intermediate that is oxidatively cleaved with base. Mechanism 23.5 is written with I_2 as halogen, forming CHI_3 (iodoform) as product.


Mechanism 23.5 The Haloform Reaction

Part [1] Halogenation at the α carbon



1 – 2 Removal of a proton forms an enolate that reacts with I_2 to yield an α-iodo ketone. Steps [1] and [2] are repeated twice more to form the triiodo substitution product.

Part [2] Oxidative cleavage with ^-OH



3 Nucleophilic addition of ^-OH forms a tetrahedral intermediate.

4 Elimination of $^-CI_3$ cleaves a carbon–carbon bond, forming the substitution product.

5 Proton transfer generates the carboxylate anion and HCI_3, iodoform.

Steps [3] and [4] result in a **nucleophilic *substitution*** reaction of a *ketone.* Because ketones normally undergo **nucleophilic *addition,*** this two-step sequence makes the haloform reaction unique. Substitution occurs because the three electronegative halogen atoms make CX_3 (CI_3 in the example) a good leaving group.

Figure 23.3 summarizes the three possible outcomes of halogenation at the α carbon, depending on the substrate and chosen reaction conditions.

Problem 23.12 Draw the products of each reaction. Assume excess halogen is present.






Figure 23.3 Summary: Halogenation reactions at the α carbon to a carbonyl group


a. Halogenation in acid—Monosubstitution on the α carbon



b. General halogenation in base—Polysubstitution on the α carbon




c. Halogenation of *methyl* ketones with excess X_2 and base—Oxidative cleavage



23.7C Reactions of α-Halo Carbonyl Compounds

α-Halo carbonyl compounds undergo two useful reactions—**elimination** with base and **substitution** with nucleophiles.

For example, treatment of 2-bromocyclohexanone with the base Li_2CO_3 in the presence of LiBr in the polar aprotic solvent DMF [$HCON(CH_3)_2$] affords cyclohex-2-enone by elimination of the elements of Br and H from the α and β carbons, respectively. Thus, a two-step method can convert a carbonyl compound such as cyclohexanone into an α,β-unsaturated carbonyl compound such as cyclohex-2-enone.




α,β-Unsaturated carbonyl compounds undergo a variety of 1,2- and 1,4-addition reactions as discussed in Section 20.15.

[1] Bromination at the α carbon is accomplished with Br_2 in CH_3CO_2H.

[2] Elimination of Br and H occurs with Li_2CO_3 and LiBr in DMF.

α-Halo carbonyl compounds also react with nucleophiles by S_N2 reactions. For example, reaction of 2-bromocyclohexanone with CH_3NH_2 affords the substitution product **A.** A related intramolecular nucleophilic substitution of an α-halo ketone was a key step in the synthesis of the antimalarial drug quinine, as shown in Figure 23.4.



Problem 23.13 Draw the organic products formed when 2-bromopentan-3-one ($CH_3CH_2COCHBrCH_3$) is treated with each reagent: (a) Li_2CO_3, LiBr, DMF; (b) $CH_3CH_2NH_2$; (c) CH_3SH.

Problem 23.14 Identify the product **M** of the following two-step reaction sequence. **M** was converted to the hallucinogen LSD (Figure 18.4) in several steps.






Figure 23.4
Intramolecular nucleophilic substitution in the synthesis of quinine



- Intramolecular S_N2 reaction of a nitrogen nucleophile with an α-halo ketone affords a compound that can be converted to quinine in a single step. The new C–N bond on the α carbon is labeled in red.


23.8 Direct Enolate Alkylation

Treatment of an aldehyde or ketone with base and an alkyl halide (RX) results in ***alkylation*—the substitution of R for H on the α carbon atom.** Alkylation forms a new carbon–carbon bond on the α carbon.



23.8A General Features

We will begin with the most direct method of alkylation, and then (in Sections 23.9 and 23.10) examine two older, multistep methods that are still used today. Direct alkylation is carried out by a two-step process:




[1] **Deprotonation:** Base removes a proton from the α carbon to generate an enolate. The reaction works best with a strong nonnucleophilic base like LDA in THF solution at low temperature (–78 °C).


[2] **Nucleophilic attack:** The nucleophilic enolate attacks the alkyl halide, displacing the halide (a good leaving group) and forming the alkylation product by an S_N2 reaction.

Because Step [2] is an S_N2 reaction, it works best with unhindered methyl and 1° alkyl halides. Hindered alkyl halides and those with halogens bonded to sp^2 hybridized carbons do not undergo substitution.


R_3CX, $CH_2{=}CHX$, and C_6H_5X do *not* undergo alkylation reactions with enolates, because they are unreactive in S_N2 reactions.




Ester enolates and carbanions derived from nitriles are also alkylated under these conditions.




Problem 23.15 What product is formed when each compound is treated first with LDA in THF solution at low temperature, followed by CH_3CH_2I?




The stereochemistry of enolate alkylation follows the general rule governing the stereochemistry of reactions: **an achiral starting material yields an achiral or racemic product.** For example, when cyclohexanone (an achiral starting material) is converted to 2-ethylcyclohexanone by treatment with base and CH_3CH_2I, a new stereogenic center (labeled in blue) is introduced, and both enantiomers of the product are formed in equal amounts—that is, a **racemic mixture.**



Problem 23.16 Draw the products obtained (including stereochemistry) when each compound is treated with LDA, followed by CH_3I.



Problem 23.17 The analgesic naproxen can be prepared by a stepwise reaction sequence from ester **A.** Using enolate alkylation in one step, what reagents are needed to convert **A** to naproxen? Write the structure of each intermediate. Explain why a racemic product is formed.




A naproxen

23.8B Alkylation of Unsymmetrical Ketones

An unsymmetrical ketone can be regioselectively alkylated to yield one major product. The strategy depends on the use of the appropriate base, solvent, and temperature to form the kinetic or thermodynamic enolate (Section 23.4), which is then treated with an alkyl halide to form the alkylation product.


For example, 2-methylcyclohexanone can be converted to either 2,6-dimethylcyclohexanone (**A**) or 2,2-dimethylcyclohexanone (**B**) by proper choice of reaction conditions.

- **Treatment of 2-methylcyclohexanone with LDA in THF solution at –78 °C gives the less substituted kinetic enolate, which then reacts with CH_3I to form A.**




2-methyl-cyclohexanone **kinetic enolate** 2,6-dimethylcyclohexanone **A**

- **Treatment of 2-methylcyclohexanone with $KOC(CH_3)_3$ in $(CH_3)_3COH$ solution at room temperature forms the more substituted thermodynamic enolate, which then reacts with CH_3I to form B.**



Finally, while enolate alkylation at the less substituted α carbon using LDA is a reliable regioselective reaction, enolate alkylation at the more substituted α carbon with $KOC(CH_3)_3$ may lead to mixtures of products. Regioselectivity depends on the identity of the substrate and the experimental parameters, which sometimes must be carefully monitored to maximize the yield of the desired alkylation product.

Problem 23.18 How can pentan-2-one be converted into each compound?




23.8C Application of Enolate Alkylation: Tamoxifen Synthesis




Tamoxifen has been commercially available since the 1970s, sold under the brand name of Nolvadex.

Tamoxifen, the chapter-opening molecule, is a potent anticancer drug that has been used to treat certain forms of breast cancer for many years. One step in the synthesis of tamoxifen involves the treatment of ketone **A** with NaH as base to form an enolate. Alkylation of this enolate with CH_3CH_2I forms **B** in high yield. **B** is converted to tamoxifen in several steps, some of which are reactions you have already learned.



Only the *Z* isomer of the C=C provides beneficial effects.


Problem 23.19 Identify **A, B,** and **C,** intermediates in the synthesis of the five-membered ring called an α-methylene-γ-butyrolactone. This heterocyclic ring system is present in some antitumor agents.




23.9 Malonic Ester Synthesis

Besides the direct method of enolate alkylation discussed in Section 23.8, a new alkyl group can also be introduced on the α carbon using the malonic ester synthesis and the acetoacetic ester synthesis.

- **The malonic ester synthesis prepares carboxylic acids** having two general structures:




- **The acetoacetic ester synthesis prepares methyl ketones** having two general structures:



23.9A Background for the Malonic Ester Synthesis


- **The malonic ester synthesis is a stepwise method for converting diethyl malonate into a carboxylic acid having one or two alkyl groups on the α carbon.**

To simplify the structures, the CH_3CH_2 groups of the esters are abbreviated as Et.



diethyl malonate
Et = CH_3CH_2


Before writing out the steps in the malonic ester synthesis, recall from Section 22.11 that esters are hydrolyzed by aqueous acid. Thus, heating diethyl malonate with acid and water hydrolyzes both esters to carboxy groups, forming a β-diacid (1,3-diacid).




diethyl malonate

malonic acid
β-diacid


The resulting β-diacids are unstable to heat. They **decarboxylate (lose CO_2),** resulting in cleavage of a carbon–carbon bond and formation of a carboxylic acid. Decarboxylation is not a general reaction of all carboxylic acids. It occurs with β-diacids, however, because CO_2 can be eliminated through a cyclic, six-atom transition state. This forms an enol of a carboxylic acid, which in turn tautomerizes to the more stable keto form.




The net result of decarboxylation is cleavage of a carbon–carbon bond on the α carbon, with loss of CO_2.



Decarboxylation occurs readily whenever a carboxy group (COOH) is bonded to the α carbon of another carbonyl group. For example, β-keto acids also readily lose CO_2 on heating to form ketones.




Problem 23.20 Which of the following compounds will readily lose CO_2 when heated?




23.9B Steps in the Malonic Ester Synthesis

The malonic ester synthesis converts diethyl malonate to a carboxylic acid in three steps.



[1] **Deprotonation.** Treatment of diethyl malonate with $^-$OEt removes the acidic α proton between the two carbonyl groups. Recall from Section 23.3A that these protons are more acidic than other α protons because three resonance structures can be drawn for the enolate, instead of the usual two. Thus, $^-$OEt, rather than the stronger base LDA, can be used for this reaction.




three resonance structures for the conjugate base

[2] **Alkylation.** The nucleophilic enolate reacts with an alkyl halide in an $\mathbf{S_N2}$ reaction to form a substitution product. Because the mechanism is S_N2, the yields are higher when R is CH_3 or a 1° alkyl group.

[3] **Hydrolysis and decarboxylation.** Heating the diester with aqueous acid hydrolyzes the diester to a β-diacid, which loses CO_2 to form a carboxylic acid.


The synthesis of butanoic acid ($CH_3CH_2CH_2COOH$) from diethyl malonate illustrates the basic process:



diethyl malonate

new C–C bond in red

If the first two steps of the reaction sequence are repeated *prior* to hydrolysis and decarboxylation, then a carboxylic acid having *two new alkyl groups* on the α carbon can be synthesized. This is illustrated in the synthesis of 2-benzylbutanoic acid [$CH_3CH_2CH(CH_2C_6H_5)COOH$] from diethyl malonate:



diethyl malonate

new C–C bond in red


new C–C bond in blue

from $CH_2(CO_2Et)_2$


from

from

An intramolecular malonic ester synthesis can be used to form rings having three to six atoms, provided the appropriate dihalide is used as starting material. For example, cyclopentanecarboxylic acid can be prepared from diethyl malonate and 1,4-dibromobutane ($BrCH_2CH_2CH_2CH_2Br$) by the following sequence of reactions:





Problem 23.21 Draw the products of each reaction.

a. $CH_2(CO_2Et)_2$ [1] NaOEt, [2]  H_3O^+, Δ

b. $CH_2(CO_2Et)_2$ [1] NaOEt, [2] CH_3Br [1] NaOEt, [2] CH_3Br H_3O^+, Δ

Problem 23.22 What cyclic product is formed from each dihalide using the malonic ester synthesis?

a. 


b. 

23.9C Retrosynthetic Analysis

To use the malonic ester synthesis you must be able to determine what starting materials are needed to prepare a given compound—that is, you must **work backwards in the retrosynthetic direction.** This involves a two-step process:

[1] Locate the α carbon to the COOH group, and identify all alkyl groups bonded to the α carbon.


[2] Break the molecule into two (or three) components: Each alkyl group bonded to the α carbon comes from an alkyl halide. The remainder of the molecule comes from $CH_2(COOEt)_2$.




Sample Problem 23.3 What starting materials are needed to prepare 2-methylhexanoic acid [$CH_3CH_2CH_2CH_2CH(CH_3)COOH$] using a malonic ester synthesis?

Solution


The target molecule has two different alkyl groups bonded to the α carbon, so three components are needed for the synthesis:




Writing the synthesis in the synthetic direction:



Problem 23.23 What alkyl halides are needed to prepare each carboxylic acid by the malonic ester synthesis?




Problem 23.24 Explain why each of the following carboxylic acids cannot be prepared by a malonic ester synthesis.




23.10 Acetoacetic Ester Synthesis


- **The acetoacetic ester synthesis is a stepwise method for converting ethyl acetoacetate into a ketone having one or two alkyl groups on the α carbon.**



23.10A Steps in the Acetoacetic Ester Synthesis



The steps in the acetoacetic ester synthesis are exactly the same as those in the malonic ester synthesis. Because the starting material, CH_3COCH_2COOEt, is a β-keto ester, the final product is a **ketone,** not a carboxylic acid.




[1] **Deprotonation.** Treatment of ethyl acetoacetate with $^{-}$OEt removes the acidic proton between the two carbonyl groups.

[2] **Alkylation.** The nucleophilic enolate reacts with an alkyl halide (RX) in an S_N2 reaction to form a substitution product. Because the mechanism is S_N2, the yields are higher when R is CH_3 or a 1° alkyl group.


[3] **Hydrolysis and decarboxylation.** Heating the β-keto ester with aqueous acid hydrolyzes the ester to a β-keto acid, which loses CO_2 to form a ketone.

If the first two steps of the reaction sequence are repeated *prior* to hydrolysis and decarboxylation, then a ketone having *two new alkyl groups* on the α carbon can be synthesized.



ethyl acetoacetate — new C–C bond in red — new C–C bond in blue

Problem 23.25 What ketones are prepared by the following reactions?




23.10B Retrosynthetic Analysis

To determine what starting materials are needed to prepare a given ketone using the acetoacetic ester synthesis, you must again work in the **retrosynthetic** direction. This involves a two-step process:


[1] Identify the alkyl groups bonded to the α carbon to the carbonyl group.

[2] Break the molecule into two (or three) components: Each alkyl group bonded to the α carbon comes from an alkyl halide. The remainder of the molecule comes from CH_3COCH_2COOEt.



For a ketone with two R groups on the α carbon, three components are needed.


Sample Problem 23.4 What starting materials are needed to synthesize heptan-2-one using the acetoacetic ester synthesis?




heptan-2-one

Solution


Heptan-2-one has only one alkyl group bonded to the α carbon, so only one alkyl halide is needed in the acetoacetic ester synthesis.




Writing the acetoacetic ester synthesis in the synthetic direction:



Problem 23.26 What alkyl halides are needed to prepare each ketone using the acetoacetic ester synthesis?



Problem 23.27 Treatment of ethyl acetoacetate with NaOEt (2 equiv) and $BrCH_2CH_2Br$ forms compound **X.** This reaction is the first step in the synthesis of illudin-S, an antitumor substance isolated from the jack-o'-lantern, a poisonous, saffron-colored mushroom. What is the structure of **X?**







The jack-o'-lantern, source of the antitumor agent illudin-S

The acetoacetic ester synthesis and direct enolate alkylation are two different methods that prepare similar ketones. Butan-2-one, for example, can be synthesized from acetone by direct enolate alkylation with CH_3I (Method [1]), or by alkylation of ethyl acetoacetate followed by hydrolysis and decarboxylation (Method [2]).

Method [1]
Direct enolate alkylation



Method [2]
Acetoacetic ester synthesis




Why would you ever make butan-2-one from ethyl acetoacetate when you could make it in fewer steps from acetone? There are many factors to consider. First of all, synthetic organic chemists like to have a variety of methods to accomplish a single kind of reaction. Sometimes subtle changes in the structure of a starting material make one reaction work better than another.

In the chemical industry, moreover, cost is an important issue. Any reaction needed to make a large quantity of a useful drug or other consumer product must use cheap starting materials. Direct enolate alkylation usually requires a very strong base like LDA to be successful, whereas the acetoacetic ester synthesis utilizes NaOEt. NaOEt can be prepared from cheaper starting materials, and this makes the acetoacetic ester synthesis an attractive method, even though it involves more steps.

Thus, each method has its own advantages and disadvantages, depending on the starting material, the availability of reagents, the cost, and the occurrence of side reactions.

Problem 23.28 Nabumetone is a pain reliever and anti-inflammatory agent sold under the brand name of Relafen.




nabumetone

a. Write out a synthesis of nabumetone from ethyl acetoacetate.
b. What ketone and alkyl halide are needed to synthesize nabumetone by direct enolate alkylation?

KEY CONCEPTS


Substitution Reactions of Carbonyl Compounds at the α Carbon

Kinetic Versus Thermodynamic Enolates (23.4)



kinetic enolate

Kinetic enolate
- The less substituted enolate
- Favored by strong base, polar aprotic solvent, low temperature: LDA, THF, –78 °C




thermodynamic enolate

Thermodynamic enolate
- The more substituted enolate
- Favored by strong base, protic solvent, higher temperature: $NaOCH_2CH_3$, CH_3CH_2OH, room temperature

Halogenation at the α Carbon

[1] Halogenation in acid (23.7A)




$X_2 = Cl_2$, Br_2, or I_2

α-halo aldehyde or ketone

- The reaction occurs via enol intermediates.
- Monosubstitution of X for H occurs on the α carbon.


[2] Halogenation in base (23.7B)



$X_2 = Cl_2$, Br_2, or I_2

- The reaction occurs via enolate intermediates.
- Polysubstitution of X for H occurs on the α carbon.

[3] Halogenation of *methyl* ketones in base—The haloform reaction (23.7B)




$X_2 = Cl_2$, Br_2, or I_2

- The reaction occurs with methyl ketones and results in cleavage of a carbon–carbon σ bond.


Reactions of α-Halo Carbonyl Compounds (23.7C)

[1] Elimination to form α,β-unsaturated carbonyl compounds



- Elimination of the elements of Br and H forms a new π bond, giving an α,β-unsaturated carbonyl compound.


[2] Nucleophilic substitution



- The reaction follows an S_N2 mechanism, generating an α-substituted carbonyl compound.


Alkylation Reactions at the α Carbon

[1] Direct alkylation at the α carbon (23.8)




- The reaction forms a new C–C bond to the α carbon.
- LDA is a common base used to form an intermediate enolate.
- The alkylation in Step [2] follows an S_N2 mechanism.

[2] Malonic ester synthesis (23.9)



- The reaction is used to prepare carboxylic acids with one or two alkyl groups on the α carbon.
- The alkylation in Step [2] follows an S_N2 mechanism.

[3] Acetoacetic ester synthesis (23.10)





- The reaction is used to prepare ketones with one or two alkyl groups on the α carbon.
- The alkylation in Step [2] follows an S_N2 mechanism.

PROBLEMS


Problems Using Three-Dimensional Models

23.29 Draw enol tautomer(s) for each compound. Ignore stereoisomers.

a. 

b. 

23.30 The cis ketone **A** is isomerized to a trans ketone **B** with aqueous NaOH. A similar isomerization does not occur with ketone **C.** (a) Draw the structure of **B** using a chair cyclohexane. (b) Label the substituents in **C** as cis or trans, and explain the difference in reactivity.




A





C

Enols, Enolates, and Acidic Protons


23.31 Draw enol tautomer(s) for each compound.

a. 


b. 

c.  (mono enol form)

23.32 Both pentane-2,4-dione and ethyl acetoacetate have two carbonyl groups separated by a single carbon atom. Although an equilibrium mixture of pentane-2,4-dione tautomers contains 76% of the enol forms, an equilibrium mixture of ethyl acetoacetate tautomers contains only 8% of the enol forms. Suggest a reason for this difference.







pentane-2,4-dione




ethyl acetoacetate


23.33 Which carbonyl compound in each pair exhibits the higher percentage of the enol tautomer?


a.  or 

b.  or 


23.34 What hydrogen atoms in each compound have a p$K_a \leq 25$?


a. 

b. 


c. 


23.35 Rank the labeled protons in each compound in order of increasing acidity.


a. 


b. 

23.36 What is the major enolate (or carbanion) formed when each compound is treated with LDA?


a. 

b. 

c. 


d. 

23.37 Why is the pK_a of the H_a protons in 1-acetylcyclohexene higher than the pK_a of the H_b protons?




1-acetylcyclohexene

23.38 Acyclovir is an effective antiviral agent used to treat the herpes simplex virus. (a) Draw the enol form of acyclovir, and explain why it is aromatic. (b) Why is acyclovir typically drawn in its keto form, despite the fact that its enol is aromatic?




acyclovir

23.39 Explain why pentane-2,4-dione forms two different alkylation products (**A** or **B**) when the number of equivalents of base is increased from one to two.



A pentane-2,4-dione **B**

23.40 Vitamin C is a stable enediol. Draw the structure of the two keto tautomers in equilibrium with the enediol and explain why the enediol is more stable than the other tautomers.




vitamin C
an enediol

Halogenation


23.41 Acid-catalyzed bromination of pentan-2-one ($CH_3COCH_2CH_2CH_3$) forms two products: $BrCH_2COCH_2CH_2CH_3$ (**A**) and $CH_3COCH(Br)CH_2CH_3$ (**B**). Explain why the major product is **B,** with the Br atom on the more substituted side of the carbonyl group.

23.42 Draw a stepwise mechanism for the following reaction.





Malonic Ester Synthesis

23.43 What alkyl halides are needed to prepare each carboxylic acid using the malonic ester synthesis?




23.44 Use the malonic ester synthesis to prepare each carboxylic acid.


a. 

b. 

23.45 Devise a synthesis of valproic acid [$(CH_3CH_2CH_2)_2CHCO_2H$], a medicine used to treat epileptic seizures, using the malonic ester synthesis.


23.46 Synthesize each compound from diethyl malonate. You may use any other organic or inorganic reagents.

a. 


b. 

23.47 The enolate derived from diethyl malonate reacts with a variety of electrophiles (not just alkyl halides) to form new carbon–carbon bonds. With this in mind, draw the products formed when $Na^+ {}^-CH(CO_2Et)_2$ reacts with each electrophile, followed by treatment with H_2O.

a. 


b. 


c. 


d. 

Acetoacetic Ester Synthesis


23.48 What alkyl halides are needed to prepare each ketone using the acetoacetic ester synthesis?


a. 

b. 

c. 


23.49 Synthesize each compound from ethyl acetoacetate. You may use any other organic or inorganic reagents.


a. 


b. 


Reactions


23.50 Draw the organic products formed in each reaction.


a. 


b. 


c. 

d. 


e. 

f. 


g. 

h. 

23.51 Draw the products formed (including stereoisomers) in each reaction.




23.52 a. Identify intermediates **A–C** in the following stepwise conversion of *p*-isobutylbenzaldehyde to the analgesic ibuprofen.




p-isobutylbenzaldehyde ibuprofen

b. Direct alkylation of **D** by treatment with one equivalent of LDA and CH_3I does not form ibuprofen. Identify the product of this reaction and explain how it is formed.




D

23.53 A key step in the synthesis of the narcotic analgesic meperidine (trade name Demerol) is the conversion of phenylacetonitrile to **X.** (a) What is the structure of **X?** (b) What reactions convert **X** to meperidine?




phenylacetonitrile meperidine

23.54 Clopidogrel is the generic name for Plavix, a drug used to prevent the formation of blood clots in patients that have a history of heart attacks or strokes. A single enantiomer of clopidogrel can be prepared in three steps from the chiral α-hydroxy acid **A.** Identify **B** and **C** in the following reaction sequence, and designate the configuration of the enantiomer formed by this route as *R* or *S*.



A


23.55 What reaction conditions—base, solvent, and temperature—are needed to convert ketone **A** to either **B** or **C** by an intramolecular alkylation reaction?



A **B** **C**

Mechanism


23.56 Although ibuprofen is sold as a racemic mixture, only the *S* enantiomer acts as an analgesic. In the body, however, some of the *R* enantiomer is converted to the *S* isomer by tautomerization to an enol and then protonation to regenerate the carbonyl compound. Write a stepwise mechanism for this isomerization.




R isomer
inactive enantiomer

S isomer
active enantiomer


23.57 Draw a stepwise mechanism showing how two alkylation products are formed in the following reaction.



23.58 Draw a stepwise mechanism for the following reaction.




23.59 Treatment of α,β-unsaturated carbonyl compound **X** with base forms the diastereomer **Y.** Write a stepwise mechanism for this reaction. Explain why one stereogenic center changes configuration but the other does not.




X **Y**

23.60 Draw stepwise mechanisms illustrating how each product is formed.




23.61 A key step in the synthesis of β-vetivone, a major constituent of vetiver, a perennial grass found in tropical and subtropical regions of the world, involved the reaction of compound **A** and dihalide **B** with two equivalents of LDA to form **C.** Draw a stepwise mechanism for this reaction. β-Vetivone contains a spiro ring system—that is, two rings that share a single carbon atom.



A **B** **C** β-vetivone


Synthesis


23.62 (a) Draw two different halo ketones that can form **A** by an intramolecular alkylation reaction. (b) How can **A** be synthesized by an acetoacetic ester synthesis?




A


23.63 Synthesize each compound from cyclohexanone and organic halides having ≤ 4 C's. You may use any other inorganic reagents.

a. 

b. 


c. 

23.64 Bupropion, sold under the trade name of Zyban, is an antidepressant that was approved to aid smoking cessation in 1997. Devise a synthesis of bupropion from benzene, organic compounds that have fewer than five carbons, and any required inorganic reagents.




bupropion

23.65 Devise a synthesis of anastrozole, a drug used to reduce the recurrence of breast cancer (Section 22.18), from the given compounds. You may use any other needed organic compounds or inorganic reagents.




anastrozole

23.66 Synthesize (*Z*)-hept-5-en-2-one from ethyl acetoacetate ($CH_3COCH_2CO_2Et$) and the given starting material. You may also use any other organic compounds or required inorganic reagents.




(*Z*)-hept-5-en-2-one

23.67 Treatment of ketone **A** with LDA followed by CH_3CH_2I did not form the desired alkylation product **B.** What product was formed instead? Devise a multistep method to convert **A** to **B,** a synthetic intermediate used to prepare the anticancer drug tamoxifen (Section 23.8C and the chapter-opening molecule).



A **B**


23.68 Capsaicin, the spicy component of hot peppers, can be prepared from amine **X** and acid chloride **Y.** Devise a synthesis of **Y** from (*E*)-6-methylhept-4-en-1-ol [$(CH_3)_2CHCH{=}CH(CH_2)_3OH$], $CH_2(CO_2Et)_2$, and any required inorganic reagents.




X **Y** capsaicin

Spectroscopy

23.69 Treatment of **W** with CH_3Li, followed by CH_3I, affords compound **Y** ($C_7H_{14}O$) as the major product. **Y** shows a strong absorption in its IR spectrum at 1713 cm^{-1}, and its ^1H NMR spectrum is given below. (a) Propose a structure for **Y.** (b) Draw a stepwise mechanism for the conversion of **W** to **Y.**




W




Challenge Problems


23.70 Explain why H_a is much less acidic than H_b. Then draw a mechanism for the following reaction.



23.71 Devise a stepwise mechanism for the following reaction.




23.72 The last step in the synthesis of β-vetivone (Problem 23.61) involves treatment of **C** with CH_3Li to form an intermediate **X,** which forms β-vetivone with aqueous acid. Identify the structure of **X** and draw a mechanism for converting **X** to β-vetivone.




C β-vetivone


23.73 Keeping in mind the mechanism for the dissolving metal reduction of alkynes to trans alkenes in Chapter 12, write a stepwise mechanism for the following reaction, which involves the conversion of an α,β-unsaturated carbonyl compound to a carbonyl compound with a new alkyl group on the α carbon.



23.74 (–)-Hyoscyamine, an optically active drug used to treat gastrointestinal disorders, is isolated from *Atropa belladonna,* the deadly nightshade plant, by a basic aqueous extraction procedure. If too much base is used during isolation, optically inactive material is isolated. (a) Explain this result by drawing a stepwise mechanism. (b) Explain why littorine, an isomer isolated from the tailflower plant in Australia, can be obtained optically pure regardless of the amount of base used during isolation.



(–)-hyoscyamine



(–)-littorine

24 Carbonyl Condensation Reactions

24.1 The aldol reaction
24.2 Crossed aldol reactions
24.3 Directed aldol reactions
24.4 Intramolecular aldol reactions
24.5 The Claisen reaction
24.6 The crossed Claisen and related reactions
24.7 The Dieckmann reaction
24.8 The Michael reaction
24.9 The Robinson annulation




Ibuprofen is the generic name for the pain reliever known by the trade names of Motrin and Advil. Like aspirin, ibuprofen acts as an anti-inflammatory agent by blocking the synthesis of prostaglandins from arachidonic acid. One step in a commercial synthesis of ibuprofen involves the reaction of a nucleophilic enolate with an electrophilic carbonyl group. In Chapter 24, we learn about the carbon–carbon bond-forming reactions of enolates with carbonyl electrophiles.

In Chapter 24, we examine **carbonyl condensations**—that is, reactions between two carbonyl compounds—a second type of reaction that occurs at the α carbon of a carbonyl group. Much of what is presented in Chapter 24 applies principles you have already learned. Many of the reactions may look more complicated than those in previous chapters, but they are fundamentally the same. Nucleophiles attack electrophilic carbonyl groups to form the products of nucleophilic addition or substitution, depending on the structure of the carbonyl starting material.

Every reaction in Chapter 24 forms a new carbon–carbon bond at the α carbon to a carbonyl group, so these reactions are extremely useful in the synthesis of complex natural products.

24.1 The Aldol Reaction

Chapter 24 concentrates on the second general reaction of enolates—**reaction with other carbonyl compounds.** In these reactions, one carbonyl component serves as the nucleophile and one serves as the electrophile, and a new carbon–carbon bond is formed.




The presence or absence of a leaving group on the electrophilic carbonyl carbon determines the structure of the product. Even though they appear somewhat more complicated, these reactions are often reminiscent of the nucleophilic addition and nucleophilic acyl substitution reactions of Chapters 21 and 22. Four types of reactions are examined:

- **Aldol reaction** (Sections 24.1–24.4)
- **Claisen reaction** (Sections 24.5–24.7)
- **Michael reaction** (Section 24.8)
- **Robinson annulation** (Section 24.9)

24.1A General Features of the Aldol Reaction


In the **aldol reaction,** two molecules of an aldehyde or ketone react with each other in the presence of base to form a **β-hydroxy carbonyl compound.** For example, treatment of acetaldehyde with aqueous $^-$OH forms 3-hydroxybutanal, a **β-hydroxy aldehyde.**

Many aldol products contain an ***ald***ehyde and an alcoh***ol***—hence the name ***aldol.***



The mechanism of the aldol reaction has **three steps,** as shown in Mechanism 24.1. Carbon–carbon bond formation occurs in Step [2], when the nucleophilic enolate reacts with the electrophilic carbonyl carbon.

Mechanism 24.1 The Aldol Reaction



1 The base removes a proton on the α carbon to form a resonance-stabilized enolate.

2 Nucleophilic attack of the enolate on an electrophilic carbonyl in another molecule of aldehyde forms a new C–C bond.


3 Protonation of the alkoxide forms the β-hydroxy aldehyde.

The aldol reaction is a reversible equilibrium, so the position of the equilibrium depends on the base and the carbonyl compound. **$^-$OH is the base** typically used in an aldol reaction. Recall from Section 23.3B that only a small amount of enolate forms with $^-$OH. In this case, that's appropriate because the starting aldehyde is needed to react with the enolate in the second step of the mechanism.

Aldol reactions can be carried out with either aldehydes or ketones. With aldehydes, the equilibrium usually favors the products, but with ketones the equilibrium favors the starting materials. There are ways of driving this equilibrium to the right, however, so we will write aldol products whether the substrate is an aldehyde or a ketone.

- **The characteristic reaction of aldehydes and ketones is nucleophilic addition (Section 21.7). An aldol reaction is a nucleophilic addition in which an enolate is the nucleophile. See the comparison in Figure 24.1.**


Figure 24.1
The aldol reaction—An example of nucleophilic addition




- Aldehydes and ketones react by nucleophilic addition. In an aldol reaction, **an enolate is the nucleophile** that adds to the carbonyl group.

A **second example of an aldol** reaction is shown with propanal as starting material. The two molecules of the aldehyde that participate in the aldol reaction react in opposite ways.

- One molecule of propanal becomes an enolate—an electron-rich *nucleophile.*
- One molecule of propanal serves as the *electrophile* because its carbonyl carbon is electron deficient.




These two examples illustrate the general features of the aldol reaction. **The α carbon of one carbonyl component becomes bonded to the carbonyl carbon of the other component.**




Problem 24.1 Draw the aldol product formed from each compound.

a. b. c. d.



Problem 24.2 Which carbonyl compounds do *not* undergo an aldol reaction when treated with ⁻OH in H_2O?

a. b. c. d.




24.1B Dehydration of the Aldol Product

The β-hydroxy carbonyl compounds formed in the aldol reaction dehydrate more readily than other alcohols. In fact, under the basic reaction conditions, the initial aldol product is often not isolated. Instead, **it loses the elements of H_2O from the α and β carbons to form an α,β-unsaturated carbonyl compound, a conjugated product.**

All alcohols—including β-hydroxy carbonyl compounds—dehydrate in the presence of acid. Only β-hydroxy carbonyl compounds dehydrate in the presence of base. [1] 2

An aldol reaction is often called an **aldol condensation,** because the β-hydroxy carbonyl compound that is initially formed loses H_2O by dehydration. **A *condensation reaction* is one in which a small molecule, in this case H_2O, is eliminated during a reaction.** [2] 2




It may or may not be possible to isolate the β-hydroxy carbonyl compound under the conditions of the aldol reaction. When the α,β-unsaturated carbonyl compound is further conjugated with a

carbon–carbon double bond or a benzene ring, as in the case of Reaction [2], **elimination of H_2O is spontaneous** and the β-hydroxy carbonyl compound cannot be isolated.

The mechanism of dehydration consists of two steps: **deprotonation followed by loss of ^-OH,** as shown in Mechanism 24.2.

Mechanism 24.2 Dehydration of β-Hydroxy Carbonyl Compounds with Base



1 The base removes a proton on the α carbon to form a resonance-stabilized enolate.

2 ^-OH is eliminated as the electron pair of the enolate forms the new π bond.


Like E1 elimination, E1cB requires two steps. Unlike E1, though, the intermediate in E1cB is a *carbanion,* not a carbocation. E1cB stands for **Elimination, unimolecular, conjugate base.**

This elimination mechanism, called the **E1cB mechanism,** differs from the two more general mechanisms of elimination, E1 and E2, which were discussed in Chapter 8. The E1cB mechanism involves two steps, and proceeds by way of an **anionic** intermediate.


Regular alcohols dehydrate only in the presence of acid but not base, because hydroxide is a poor leaving group. When the hydroxy group is β to a carbonyl group, however, loss of H and OH from the α and β carbons forms a **conjugated double bond,** and the stability of the conjugated system makes up for having such a poor leaving group.

Dehydration of the initial β-hydroxy carbonyl compound drives the equilibrium of an aldol reaction to the right, thus favoring product formation. Once the conjugated α,β-unsaturated carbonyl compound forms, it is *not* re-converted to the β-hydroxy carbonyl compound.

Problem 24.3 What unsaturated carbonyl compound is formed by dehydration of each β-hydroxy carbonyl compound?



Problem 24.4 Acid-catalyzed dehydration of β-hydroxy carbonyl compounds occurs by the mechanism discussed in Section 9.8. With this in mind, draw a stepwise mechanism for the following reaction.




24.1C Retrosynthetic Analysis

To utilize the aldol reaction in synthesis, you must be able to determine which aldehyde or ketone is needed to prepare a particular β-hydroxy carbonyl compound or α,β-unsaturated carbonyl compound—that is, you must be able to **work backwards, in the retrosynthetic direction.**

How To Synthesize a Compound Using the Aldol Reaction

Example What starting material is needed to prepare each compound by an aldol reaction?




Step [1] **Locate the α and β carbons of the carbonyl group.**

- When a carbonyl group has two different α carbons, choose the side that contains the OH group (in a β-hydroxy carbonyl compound) or is part of the C=C (in an α,β-unsaturated carbonyl compound).


Step [2] **Break the molecule into two components between the α and β carbons.**

- The α carbon and all remaining atoms bonded to it belong to one carbonyl component. The β carbon and all remaining atoms bonded to it belong to the other carbonyl component. Both components are identical in all aldols we have thus far examined.


a. Break the molecule into two halves at the labeled bond.



b. Break the molecule into two halves at the labeled bond.



Problem 24.5 What aldehyde or ketone is needed to prepare each compound by an aldol reaction?




24.2 Crossed Aldol Reactions

In all of the aldol reactions discussed so far, the electrophilic carbonyl and the nucleophilic enolate have originated from the *same* aldehyde or ketone. Sometimes, though, it is possible to carry out an aldol reaction between two *different* carbonyl compounds.

- **An aldol reaction between two different carbonyl compounds is called a *crossed aldol* or *mixed aldol reaction.***

24.2A A Crossed Aldol Reaction with Two Different Aldehydes, Both Having α H Atoms

When two different aldehydes, both having α H atoms, are combined in an aldol reaction, four different β-hydroxy carbonyl compounds are formed. Four products form, not one, because *both* aldehydes can lose an acidic α hydrogen atom and form an enolate in the presence of base. *Both* enolates can then react with *both* carbonyl compounds, as shown for acetaldehyde and propanal in the following reaction scheme.



- **Conclusion: When two different aldehydes have α hydrogens, a crossed aldol reaction is *not* synthetically useful.**


24.2B Synthetically Useful Crossed Aldol Reactions

Crossed aldols are synthetically useful in two different situations.

- **A crossed aldol occurs when only one carbonyl component has α H atoms.**


When one carbonyl compound has no α hydrogens, a crossed aldol reaction often leads to one product. Two common carbonyl compounds with no α hydrogens used for this purpose are **formaldehyde ($CH_2{=}O$)** and **benzaldehyde (C_6H_5CHO).**

For example, reaction of C_6H_5CHO (as the electrophile) with either acetaldehyde (CH_3CHO) or acetone [$(CH_3)_2C{=}O$] in the presence of base forms a single α,β-unsaturated carbonyl compound after dehydration.




The yield of a single crossed aldol product is increased further if the electrophilic carbonyl component is relatively unhindered (as is the case with most aldehydes), and if it is used in excess.

Problem 24.6 2-Pentylcinnamaldehyde, commonly called flosal, is a perfume ingredient with a jasmine-like odor. Flosal is an α,β-unsaturated aldehyde made by a crossed aldol reaction between benzaldehyde (C_6H_5CHO) and heptanal ($CH_3CH_2CH_2CH_2CH_2CH_2CHO$), followed by dehydration. Draw a stepwise mechanism for the following reaction that prepares flosal.




flosal
(perfume component)

Problem 24.7 Draw the products formed in each crossed aldol reaction.



- **A crossed aldol occurs when one carbonyl component has especially acidic α H atoms.**

A useful crossed aldol reaction takes place between an aldehyde or ketone and a β-dicarbonyl (or similar) compound.




R' = H or alkyl

Y, Z = COOEt, CHO, COR, CN

new C−C σ and π bonds in red


β-dicarbonyl compound
(and related compounds)



benzaldehyde diethyl malonate


As we learned in Section 23.3, the α hydrogens between two carbonyl groups are especially acidic, and so they are more readily removed than other α H atoms. As a result, **the β-dicarbonyl**

Figure 24.2
Crossed aldol reaction between benzaldehyde and $CH_2(COOEt)_2$



compound always becomes the enolate component of the aldol reaction. Figure 24.2 shows the steps for the crossed aldol reaction between diethyl malonate and benzaldehyde. In this type of crossed aldol reaction, the initial β-hydroxy carbonyl compound *always* loses water to form the highly conjugated product.

β-Dicarbonyl compounds are sometimes called **active methylene compounds** because they are more reactive towards base than other carbonyl compounds. **1,3-Dinitriles** and **α-cyano carbonyl compounds** are also active methylene compounds.



Problem 24.8 Draw the products formed in the crossed aldol reaction of phenylacetaldehyde ($C_6H_5CH_2CHO$) with each compound: (a) $CH_2(COOEt)_2$; (b) $CH_2(COCH_3)_2$; (c) CH_3COCH_2CN.

24.2C Useful Transformations of Aldol Products

The aldol reaction is synthetically useful because it forms new carbon–carbon bonds, generating products with two functional groups. Moreover, the β-hydroxy carbonyl compounds formed in aldol reactions are readily transformed into a variety of other compounds. Figure 24.3 illustrates how the crossed aldol product obtained from cyclohexanone and formaldehyde ($CH_2{=}O$) can be converted to other compounds by reactions learned in earlier chapters.

Problem 24.9 What aldol product is formed when two molecules of butanal react together in the presence of base? What reagents are needed to convert this product to each of the following compounds?





Figure 24.3
Conversion of a β-hydroxy carbonyl compound into other compounds



- The β-hydroxy carbonyl compound formed from the crossed aldol reaction can be reduced with $NaBH_4$, CH_3OH (Section 20.4A) to form a 1,3-diol (Reaction [1]) or dehydrated to form an α,β-unsaturated carbonyl compound (Reaction [2]).
- Reduction of the α,β-unsaturated carbonyl compound forms an allylic alcohol with $NaBH_4$ (Reaction [3]), or a ketone with H_2 and Pd-C (Reaction [4]); see Section 20.4C.
- Reaction of the α,β-unsaturated carbonyl compound with an organometallic reagent forms two different products depending on the choice of RM (Reaction [5]); see Section 20.15.

24.3 Directed Aldol Reactions

A **directed aldol reaction** is a variation of the crossed aldol reaction that clearly defines which carbonyl compound becomes the nucleophilic enolate and which reacts at the electrophilic carbonyl carbon. The strategy of a directed aldol reaction is as follows:


[1] Prepare the enolate of one carbonyl component with LDA.

[2] Add the second carbonyl compound (the electrophile) to this enolate.

Because the steps are done sequentially and a strong nonnucleophilic base is used to form the enolate of one carbonyl component only, a variety of carbonyl substrates can be used in the reaction. Both carbonyl components can have α hydrogens because only one enolate is prepared with LDA. Also, when an unsymmetrical ketone is used, LDA selectively forms the **less substituted, kinetic enolate.**

Sample Problem 24.1 illustrates the steps of a directed aldol reaction between a ketone and an aldehyde, both of which have α hydrogens.

Sample Problem 24.1 Draw the product of the following directed aldol reaction.



2-methylcyclohexanone

Solution

2-Methylcyclohexanone forms an enolate on the less substituted carbon, which then reacts with the electrophile, CH_3CHO.





Figure 24.4 illustrates how a directed aldol reaction was used in the synthesis of **periplanone B,** the sex pheromone of the female American cockroach.

Figure 24.4
A directed aldol reaction in the synthesis of periplanone B




To determine the needed carbonyl components for a directed aldol, follow the same strategy used for a regular aldol reaction in Section 24.1C, as shown in Sample Problem 24.2.

Sample Problem 24.2

What starting materials are needed to prepare *ar*-turmerone using a directed aldol reaction? *ar*-Turmerone is a principal component of the essential oil derived from turmeric root.




The dried and ground root of the turmeric plant, a tropical perennial in the ginger family, is an essential ingredient in curry powder.




ar-turmerone

Solution


When the desired product is an α,β-unsaturated carbonyl compound, identify the α and β carbons that are part of the C = C, and break the molecule into two components between these carbons.



Problem 24.10 What carbonyl starting materials are needed to prepare each compound using a directed aldol reaction?




Problem 24.11 Zingerone, a spicy-sweet component of cooked ginger, can be converted to its protected TBDMS ether **A**, as we learned in Chapter 20. How can **A** be converted to gingerol, a compound present in fresh ginger, using a directed aldol reaction as a key step?





Donepezil (trade name Aricept) is a drug used to improve cognitive function in patients suffering from dementia and Alzheimer's disease.


Problem 24.12 A key step in the synthesis of donepezil (trade name Aricept) is a directed aldol reaction that forms α,β-unsaturated carbonyl compound **X.** What carbonyl starting materials are needed to prepare **X** using a directed aldol reaction? What reagents are needed to convert **X** to donepezil?



24.4 Intramolecular Aldol Reactions


Hexane-2,5-dione is called a **1,4-dicarbonyl compound** to emphasize the relative positions of its carbonyl groups. 1,4-Dicarbonyl compounds are starting materials for synthesizing **five-membered rings.**

Aldol reactions with dicarbonyl compounds can be used to make five- and six-membered rings. The enolate formed from one carbonyl group is the nucleophile, and the carbonyl carbon of the other carbonyl group is the electrophile. For example, treatment of hexane-2,5-dione with base forms a five-membered ring.




The steps in this process, shown in Mechanism 24.3, are no different from the general mechanisms of the aldol reaction and dehydration described previously in Section 24.1.

Mechanism 24.3 The Intramolecular Aldol Reaction



1. The base removes a proton on the α carbon to form a resonance-stabilized enolate.
2. Nucleophilic attack of the enolate on the electrophilic carbonyl *in the same molecule* forms a new C–C σ bond, generating the five-membered ring.
3. Protonation of the alkoxide forms the β-hydroxy carbonyl compound.
4 – 5. Dehydration occurs by the two-step E1cB mechanism—loss of a proton to form an enolate and elimination of $^-$OH to form a π bond.

When hexane-2,5-dione is treated with base in Step [1], two different enolates are possible—enolates **A** and **B,** formed by removal of H_a and H_b, respectively. Although enolate **A** goes on to form the five-membered ring, intramolecular cyclization using enolate **B** would lead to a strained three-membered ring.



Because the three-membered ring is much higher in energy than the enolate starting material, equilibrium greatly favors the starting materials and the **three-membered ring does not form.** Under the reaction conditions, enolate **B** is re-protonated to form hexane-2,5-dione, because all steps except dehydration are equilibria. **Thus, equilibrium favors formation of the more stable five-membered ring over the much less stable three-membered ring.**

In a similar fashion, six-membered rings can be formed from the intramolecular aldol reaction of **1,5-dicarbonyl compounds.**





Figure 24.5
The synthesis of progesterone using an intramolecular aldol reaction




- Oxidative cleavage of the alkene with O_3, followed by Zn, H_2O (Section 12.10), gives the 1,5-dicarbonyl compound.
- Intramolecular aldol reaction of the 1,5-dicarbonyl compound with dilute ^-OH in H_2O solution forms progesterone.
- **This two-step reaction sequence converts a five-membered ring into a six-membered ring.** Reactions that synthesize larger rings from smaller ones are called **ring expansion reactions.**


The synthesis of the female sex hormone **progesterone** by W. S. Johnson and co-workers at Stanford University is considered one of the classics in total synthesis. The last six-membered ring needed in the steroid skeleton was prepared by a two-step sequence using an intramolecular aldol reaction, as shown in Figure 24.5.

Problem 24.13 Draw a stepwise mechanism for the conversion of heptane-2,6-dione to 3-methylcyclohex-2-enone with NaOEt, EtOH.

Problem 24.14 What cyclic product is formed when each 1,5-dicarbonyl compound is treated with aqueous ^-OH?




Problem 24.15 Following the two-step reaction sequence depicted in Figure 24.5, write out the steps needed to convert **A** to **B.**



24.5 The Claisen Reaction

The **Claisen reaction** is the second general reaction of enolates with other carbonyl compounds. In the Claisen reaction, two molecules of an ester react with each other in the presence of an alkoxide base to form a **β-keto ester.** For example, treatment of ethyl acetate with NaOEt forms ethyl acetoacetate after protonation with aqueous acid.




Unlike the aldol reaction, which is base-catalyzed, a full equivalent of base is needed to deprotonate the β-keto ester formed in Step [3] of the Claisen reaction.

The mechanism for the Claisen reaction (Mechanism 24.4) resembles the mechanism of an aldol reaction in that it involves nucleophilic addition of an enolate to an electrophilic carbonyl group. Because esters have a leaving group on the carbonyl carbon, however, loss of a leaving group occurs to form the product of **substitution, *not* addition.**


Mechanism 24.4 The Claisen Reaction

Part [1] Formation of the β-keto ester



1. The base removes a proton on the α carbon to form a resonance-stabilized enolate.
2. Nucleophilic attack of the enolate on an electrophilic carbonyl in another molecule of ester forms a new C–C bond.
3. Loss of the leaving group (⁻OEt) forms a β-keto ester.

Part [2] Deprotonation and protonation




4. Because the β-keto ester formed in Step [3] has especially acidic protons between its two carbonyl groups, the base removes a proton to form a resonance-stabilized enolate.
5. Protonation of the enolate with strong acid re-forms the β-keto ester.

Because the generation of a resonance-stabilized enolate from the product β-keto ester drives the Claisen reaction (Step [4] of Mechanism 24.4), **only esters with two or three hydrogens on the α carbon undergo this reaction;** that is, esters must have the general structure CH_3CO_2R' or RCH_2CO_2R'.

- **Keep in mind: The characteristic reaction of esters is nucleophilic substitution. A Claisen reaction is a nucleophilic substitution in which an enolate is the nucleophile.**


Figure 24.6 compares the general reaction for nucleophilic substitution of an ester with the Claisen reaction. Sample Problem 24.3 reinforces the basic features of the Claisen reaction.

Figure 24.6
The Claisen reaction—An example of nucleophilic substitution




- Esters react by **nucleophilic substitution.** In a Claisen reaction, an enolate is the nucleophile that adds to the carbonyl group.

Sample Problem 24.3 Draw the product of the following Claisen reaction.




Solution

To draw the product of any Claisen reaction, form a new carbon–carbon bond between the α carbon of one ester and the carbonyl carbon of another ester, with elimination of the leaving group ($^-OCH_3$ in this case).





new C–C bond in red

Next, write out the steps of the reaction to verify this product.



Problem 24.16 What β-keto ester is formed when each ester is used in a Claisen reaction?

a. 

b. 

24.6 The Crossed Claisen and Related Reactions

Like the aldol reaction, it is sometimes possible to carry out a Claisen reaction with two different carbonyl components as starting materials.


- **A Claisen reaction between two different carbonyl compounds is called a *crossed Claisen reaction.***

24.6A Two Useful Crossed Claisen Reactions

A crossed Claisen reaction is synthetically useful in two different instances.

- **A crossed Claisen occurs between two different esters when only one has α hydrogens.**


When one ester has no α hydrogens, a crossed Claisen reaction often leads to one product. Common esters with no α H atoms include ethyl formate (**HCO_2Et**) and ethyl benzoate (**$C_6H_5CO_2Et$**). For example, the reaction of ethyl benzoate (as the electrophile) with ethyl acetate (which forms the enolate) in the presence of base forms predominately one β-keto ester.



ethyl benzoate — ethyl acetate (Only this ester can form an enolate.) — new C–C bond in red; β-keto ester

- **A crossed Claisen occurs between a ketone and an ester.**


The reaction of a ketone and an ester in the presence of base also forms the product of a crossed Claisen reaction. The enolate is generally formed from the ketone component, and the reaction works best when the ester has no α hydrogens. The product of this crossed Claisen reaction is a **β-dicarbonyl compound,** but ***not*** a β-keto ester.




new C–C bond in red

β-dicarbonyl compound


Problem 24.17 What crossed Claisen product is formed from each pair of compounds?





Sunscreen ingredients


Problem 24.18 Avobenzone is a conjugated compound that absorbs ultraviolet light with wavelengths in the 320–400 nm region, so it is a common ingredient in commercial sunscreens. Write out two different crossed Claisen reactions that form avobenzone.



avobenzone

24.6B Other Useful Variations of the Crossed Claisen Reaction

β-Dicarbonyl compounds are also prepared by reacting an enolate with **ethyl chloroformate** and **diethyl carbonate.**



ethyl chloroformate — diethyl carbonate


These reactions resemble a Claisen reaction because they involve the same three steps:


[1] Formation of an enolate

[2] Nucleophilic addition to a carbonyl group


[3] Elimination of a leaving group

For example, reaction of an ester enolate with diethyl carbonate yields a β-diester (Reaction [1]), whereas reaction of a ketone enolate with ethyl chloroformate forms a β-keto ester (Reaction [2]). New carbon–carbon bonds are shown in red.







Reaction [2] is noteworthy because it provides easy access to **β-keto esters,** which are useful starting materials in the acetoacetic ester synthesis (Section 23.10). In this reaction, Cl^- is eliminated rather than ^-OEt in Step [3], because Cl^- is a better leaving group, as shown in the following steps.



Problem 24.19 Draw the products of each reaction.




Problem 24.20 Two steps in a synthesis of the analgesic ibuprofen, the chapter-opening molecule, include a carbonyl condensation reaction, followed by an alkylation reaction. Identify intermediates **A** and **B** in the synthesis of ibuprofen.




24.7 The Dieckmann Reaction

Intramolecular Claisen reactions of diesters form five- and six-membered rings. The enolate of one ester is the nucleophile, and the carbonyl carbon of the other is the electrophile. An intramolecular Claisen reaction is called a **Dieckmann reaction.** Two types of diesters give good yields of cyclic products.

- **1,6-Diesters yield five-membered rings by the Dieckmann reaction.**




- **1,7-Diesters yield six-membered rings by the Dieckmann reaction.**



The mechanism of the Dieckmann reaction is exactly the same as the mechanism of an intermolecular Claisen reaction. It is illustrated in Mechanism 24.5 for the formation of a six-membered ring.

Mechanism 24.5 The Dieckmann Reaction




1 – 2 The base removes a proton on the α carbon to form an enolate, which attacks the electrophilic carbonyl of the other ester to form a new C–C bond.

3 Elimination of $^-$OEt forms the β-keto ester.


4 – 5 Under the basic reaction conditions, the proton between the two carbonyls is removed with base to form an enolate, which is protonated with acid to re-form the β-keto ester.

Problem 24.21 What two β-keto esters are formed in the Dieckmann reaction of the following diester?




24.8 The Michael Reaction

Like the aldol and Claisen reactions, the **Michael reaction involves two carbonyl components—the enolate of one carbonyl compound and an α,β-unsaturated carbonyl compound.**




Recall from Section 20.15 that α,β-unsaturated carbonyl compounds are resonance stabilized and have **two electrophilic sites—the carbonyl carbon and the β carbon.**




- **The Michael reaction involves the conjugate addition (1,4-addition) of a resonance-stabilized enolate to the β carbon of an α,β-unsaturated carbonyl system.**


All conjugate additions add the **elements of H and Nu across the α and β carbons.**



In the Michael reaction, the **nucleophile is an enolate.** Enolates of active methylene compounds are particularly common. The α,β-unsaturated carbonyl component is often called a **Michael acceptor.**




Problem 24.22 Which of the following compounds can serve as Michael acceptors?



The Michael reaction always forms a new carbon–carbon bond on the β carbon of the Michael acceptor. Reaction [2] is used to illustrate the mechanism of the Michael reaction in Mechanism 24.6. The key step is nucleophilic addition of the enolate to the β carbon of the Michael acceptor in Step [2].


Mechanism 24.6 The Michael Reaction



1. The base removes a proton on the carbon between the two carbonyl groups to form an enolate.
2. Nucleophilic addition of the enolate to the β carbon of the α,β-unsaturated carbonyl compound forms a new carbon–carbon bond and another enolate.
3. Protonation of the enolate forms the 1,4-addition product.


Figure 24.7

Using a Michael reaction in the synthesis of the steroid estrone




When the product of a Michael reaction is also a β-keto ester, it can be hydrolyzed and decarboxylated by heating in aqueous acid, as discussed in Section 23.9. This forms a **1,5-dicarbonyl compound.** Figure 24.7 shows a Michael reaction that was a key step in the synthesis of **estrone,** a female sex hormone.


1,5-Dicarbonyl compounds are starting materials for intramolecular aldol reactions, as described in Section 24.4.



Problem 24.23 What product is formed when each pair of compounds is treated with NaOEt in ethanol?




Problem 24.24 What starting materials are needed to prepare each compound by the Michael reaction?




24.9 The Robinson Annulation

The word **annulation** comes from the Greek word *annulus* for ring. The Robinson annulation is named for English chemist Sir Robert Robinson, who was awarded the 1947 Nobel Prize in Chemistry.

The Robinson annulation is a ring-forming reaction that combines a Michael reaction with an intramolecular aldol reaction. Like the other reactions in Chapter 24, it involves enolates and it forms carbon–carbon bonds. The two starting materials for a Robinson annulation are an α,β-unsaturated carbonyl compound and an enolate.




The Robinson annulation forms a six-membered ring and three new carbon–carbon bonds—two σ bonds and one π bond. The product contains an α,β-unsaturated ketone in a cyclohexane ring—that is, a **cyclohex-2-enone** ring. To generate the enolate component of the Robinson annulation, $^{-}$OH in H_2O and $^{-}$OEt in EtOH are typically used.



The mechanism of the Robinson annulation consists of a **Michael addition** to the α,β-unsaturated carbonyl compound to form a 1,5-dicarbonyl compound, followed by an **intramolecular aldol reaction** to form the six-membered ring. The mechanism is written out in three parts in Mechanism 24.7 for the reaction between methyl vinyl ketone and 2-methylcyclohexane-1,3-dione.

Mechanism 24.7 The Robinson Annulation


Part [1] Michael addition



1 – 2 Base removes the most acidic proton—the proton between the two carbonyl groups—to form an enolate. Conjugate addition of the enolate to the α,β-unsaturated carbonyl compound forms a new carbon–carbon bond, generating an enolate.

3 Protonation of the enolate forms a 1,5-dicarbonyl compound.


Part [2] Intramolecular aldol reaction



4 – 5 The base removes a proton to form an enolate, which attacks a carbonyl group to form a new C–C σ bond, generating the six-membered ring.

6 Protonation of the alkoxide forms the β-hydroxy carbonyl compound.


Part [3] Dehydration of the β-hydroxy carbonyl compound




7 – 8 Dehydration occurs by the two-step E1cB mechanism—loss of a proton to form an enolate and elimination of $^-$OH to form a π bond.

The mechanism begins with the three-step Michael addition that forms the first carbon–carbon σ bond, generating the 1,5-dicarbonyl compound (Part [1]). An intramolecular aldol reaction (Part [2]) forms the second carbon–carbon σ bond, and dehydration of the β-hydroxy ketone (Part [3]) forms the π bond. All of the parts of this mechanism have been discussed in previous sections of Chapter 24. However, the end result of the Robinson annulation—the formation of a cyclohex-2-enone ring—is new.

To draw the product of Robinson annulation without writing out the mechanism each time, **place the α carbon of the compound that becomes the enolate next to the β carbon of the α,β-unsaturated carbonyl compound.** Then, join the appropriate carbons together as shown. If you follow this method of drawing the starting materials, the double bond in the product always ends up in the same position in the six-membered ring.



Sample Problem 24.4 Draw the Robinson annulation product formed from the following starting materials.




Solution


Arrange the starting materials to put the reactive atoms next to each other. For example:

- Place the α,β-unsaturated carbonyl compound *to the left* of the carbonyl compound.
- Determine which α carbon will become the enolate. The most acidic H is always removed with base first, which in this case is the H (in red) on the α carbon between the two carbonyl groups. This α carbon is drawn adjacent to the β carbon of the α,β-unsaturated carbonyl compound.

Then draw the bonds to form the new six-membered ring.




Problem 24.25 Draw the products when each pair of compounds is treated with $CH_3CH_2O^-$, CH_3CH_2OH in a Robinson annulation reaction.



To use the Robinson annulation in synthesis, you must be able to determine what starting materials are needed to prepare a given compound, by working in the retrosynthetic direction.


How To Synthesize a Compound Using the Robinson Annulation

Example **What starting materials are needed to synthesize the following compound using a Robinson annulation?**




Step [1] **Locate the cyclohex-2-enone ring and re-draw the target molecule if necessary.**

- To most easily determine the starting materials, always arrange the α,β-unsaturated carbonyl system in the same location. The target compound may have to be flipped or rotated, and you must be careful not to move any bonds to the wrong location during this process.




Step [2] **Break the cyclohex-2-enone ring into two components.**

- Break the C = C. One half becomes the carbonyl group of the enolate component.
- Break the bond between the β carbon and the carbon to which it is bonded.





Problem 24.26 Which of the following bicyclic ring systems can be prepared by an intermolecular Robinson annulation?




A **B** **C** **D**

Problem 24.27 What starting materials are needed to synthesize each compound by a Robinson annulation?

a. 





b. 

c. 

KEY CONCEPTS

Carbonyl Condensation Reactions


The Four Major Carbonyl Condensation Reactions

Reaction type	Reaction [New C – C bonds are shown in red.]
[1] Aldol reaction (24.1)	 aldehyde (or ketone) → β-hydroxy carbonyl compound → (*E* and *Z*) α,β-unsaturated carbonyl compound
[2] Claisen reaction (24.5)	 ester → β-keto ester
[3] Michael reaction (24.8)	 α,β-unsaturated carbonyl compound + carbonyl compound → 1,5-dicarbonyl compound
[4] Robinson annulation (24.9)	 α,β-unsaturated carbonyl compound + carbonyl compound → cyclohex-2-enone

Useful Variations

[New C – C bonds are shown in red.]


[1] Directed aldol reaction (24.3)




R" = H or alkyl → β-hydroxy carbonyl compound → (*E* and *Z*) α,β-unsaturated carbonyl compound

[2] Intramolecular aldol reaction (24.4)

a. With 1,4-dicarbonyl compounds:




b. With 1,5-dicarbonyl compounds:




[3] Dieckmann reaction (24.7)

a. With 1,6-diesters:





b. With 1,7-diesters:




PROBLEMS

Problems Using Three-Dimensional Models

24.28 Draw the aldol product formed from each pair of starting materials using ^-OH, H_2O.

a.  b. 




24.29 What steps are needed to convert **A** to **B?**




A **B**


The Aldol Reaction

24.30 Draw the product formed from an aldol reaction with the given starting material(s) using ^-OH, H_2O.




a.  only b.  c. 

24.31 Draw the product formed in each directed aldol reaction.


a. 


b. 


24.32 Draw the product formed when each dicarbonyl compound undergoes an intramolecular aldol reaction followed by dehydration.


a.  b.  c. 

24.33 What starting materials are needed to synthesize each compound using an aldol or similar reaction?


a. 

b. 


c. 

d. 

24.34 What product is formed when a solution of **A** and **B** is treated with mild base? This reaction is the first step in the synthesis of rosuvastatin (sold as a calcium salt under the trade name Crestor), a drug used to treat patients with high cholesterol.





A **B**





rosuvastatin

24.35 What dicarbonyl compound is needed to prepare each compound by an intramolecular aldol reaction?


a. 

b. 

c. 


d. 

24.36 Identify the structures of **C** and **D** in the following reaction sequence.



$C_{10}H_{14}O$

24.37 Explain why ketone **K** undergoes aldol reactions but ketone **J** does not.




K




J


The Claisen and Dieckmann Reactions


24.38 Draw the Claisen product formed from each ester.


a. 


b. 

24.39 Draw the product formed from a Claisen reaction with the given starting materials using $^{-}$OEt, EtOH.


a. 


b. 


c. 


d. 

24.40 What starting materials are needed to synthesize each compound by a crossed Claisen reaction?


a. 

c. 

b. 

d. 


24.41 Even though **B** contains three ester groups, a single Dieckmann product results when **B** is treated with $NaOCH_3$ in CH_3OH, followed by H_3O^+. Draw the structure and explain why it is the only product formed.




B


Michael Reaction


24.42 Draw the product formed from a Michael reaction with the given starting materials using ^-OEt, EtOH.


a. 

b. 


24.43 What starting materials are needed to prepare each compound using a Michael reaction?

a. 

b. 

c. 


24.44 β-Vetivone is isolated from vetiver, a perennial grass that yields a variety of compounds used in traditional eastern medicine, pest control, and fragrance. In one synthesis, ketone **A** is converted to β-vetivone by a two-step process: Michael reaction, followed by intramolecular aldol reaction. (a) What Michael acceptor is needed for the conjugate addition? (See Problem 23.61 for another method to form the bicyclic ring system of β-vetivone.) (b) Draw a stepwise mechanism for the aldol reaction, which forms the six-membered ring.





A β-vetivone


Robinson Annulation

24.45 Draw the product of each Robinson annulation from the given starting materials using $^-$OH in H_2O solution.


a. 


c. 


b. 

d. 

24.46 What starting materials are needed to synthesize each compound using a Robinson annulation?


a. 


b. 


c. 


Reactions


24.47 Draw the organic products formed in each reaction.


a. 

d. 


b. 

e. 


c. 

f. 


24.48 Fill in the lettered reagents needed for each reaction.



24.49 What product (including stereochemistry) is formed in each of the following intramolecular reactions?




24.50 Identify compounds **A** and **B,** two synthetic intermediates in the 1979 synthesis of the plant growth hormone gibberellic acid by Corey and Smith. Gibberellic acid induces cell division and elongation, thus making plants tall and leaves large.



gibberellic acid

Mechanisms


24.51 In theory, the intramolecular aldol reaction of 6-oxoheptanal could yield the three compounds shown. It turns out, though, that 1-acetylcyclopentene is by far the major product. Why are the other two compounds formed in only minor amounts? Draw a stepwise mechanism to show how all three products are formed.




6-oxoheptanal

1-acetylcyclopentene
major product

24.52 Draw a stepwise mechanism for the following variation of the aldol reaction, often called a nitro aldol reaction.




24.53 Draw a stepwise mechanism for the following Robinson annulation. This reaction was a key step in a synthesis of the steroid cortisone by R. B. Woodward and co-workers at Harvard University in 1951.




cortisone

24.54 One step in the synthesis of sitagliptin (Problem 17.14, a drug used to treat type 2 diabetes) involves reaction of the mixed anhydride **A** with **B** to form **C.** Draw a stepwise mechanism for this reaction.




A **B** **C**

24.55 Reaction of **X** and phenylacetic acid forms an intermediate **Y,** which undergoes an intramolecular reaction to yield rofecoxib. Rofecoxib is a nonsteroidal anti-inflammatory agent once marketed under the trade name Vioxx, now withdrawn from the market because of increased risk of heart attacks from long-term use in some patients. Identify **Y** and draw a stepwise mechanism for its conversion to rofecoxib.




X phenylacetic acid rofecoxib

24.56 Coumarin, a naturally occurring compound isolated from lavender, sweet clover, and tonka bean, is made in the laboratory from *o*-hydroxybenzaldehyde by the reaction depicted below. Draw a stepwise mechanism for this reaction. Coumarin derivatives are useful synthetic anticoagulants.




o-hydroxybenzaldehyde coumarin

24.57 When **A** is treated with aqueous $^-$OH, the major product is compound **B,** which undergoes ester hydrolysis and decarboxylation to form **C.** Draw a stepwise mechanism for the conversion of **A** to **B.**




A **B** **C**

24.58 One step in a recent short synthesis of a prostaglandin (Section 19.6) involves the conversion of succinaldehyde to the bicyclic hemiacetal **X.** Draw a stepwise mechanism for this process. (Hint: The mechanism begins with an intermolecular aldol reaction.)



succinaldehyde **X**


24.59 (a) Draw a stepwise mechanism for the reaction of ethyl hexa-2,4-dienoate with diethyl oxalate in the presence of base. (b) How does your mechanism explain why a new carbon–carbon bond forms on C6? (c) Why is this reaction an example of a crossed Claisen reaction?




ethyl hexa-2,4-dienoate diethyl oxalate


Synthesis


24.60 Devise a synthesis of each compound from the given starting material. You may use any other organic compounds or required inorganic reagents.


a. 

b. 


24.61 Devise a synthesis of each compound from cyclopentanone, benzene, and organic alcohols having ≤ 3 C's. You may also use any required organic or inorganic reagents.


a. 

b. 

c. 


24.62 Devise a synthesis of each compound from $CH_3CH_2CH_2CO_2Et$, benzene, and alcohols having ≤ 2 C's. You may also use any required organic or inorganic reagents.

a. 

b. 

24.63 Devise a synthesis of 2-methylcyclopentanone from cyclohexene. You may also use any required reagents.


24.64 Octinoxate is an unsaturated ester used as an active ingredient in sunscreens. (a) What carbonyl compounds are needed to synthesize this compound using a condensation reaction? (b) Devise a synthesis of octinoxate from the given organic starting materials and any other needed reagents.



octinoxate

General Problem

24.65 Answer the following questions about 2-acetylcyclopentanone.




2-acetylcyclopentanone


a. What starting materials are needed to form 2-acetylcyclopentanone by a Claisen reaction that forms bond (a)?
b. What starting materials are needed to form 2-acetylcyclopentanone by a Claisen reaction that forms bond (b)?
c. What product is formed when 2-acetylcyclopentanone is treated with $NaOCH_2CH_3$, followed by CH_3I?
d. Draw the Robinson annulation product(s) formed by reaction of 2-acetylcyclopentanone with methyl vinyl ketone ($CH_2=CHCOCH_3$).
e. Draw the structure of the most stable enol tautomer(s).

Challenge Problems


24.66 Draw a stepwise mechanism for the following reaction, which was used in the synthesis of ezetimibe (Section 20.6), a drug used to treat patients with high cholesterol.



24.67 Propose a stepwise mechanism for the following reaction of a β-keto ester. Suggest a reason why this rearrangement reaction occurs.




24.68 Isophorone is formed from three molecules of acetone [$(CH_3)_2C{=}O$] in the presence of base. Draw a mechanism for this process.




isophorone


24.69 Devise a stepwise mechanism for the following reaction. (Hint: The mechanism begins with the conjugate addition of ^-OH.)



24.70 Draw a stepwise mechanism for the following reaction. (Hint: Two Michael reactions are needed.)




24.71 4-Methylpyridine reacts with benzaldehyde (C_6H_5CHO) in the presence of base to form **A.** (a) Draw a stepwise mechanism for this reaction. (b) Would you expect 2-methylpyridine or 3-methylpyridine to undergo a similar type of condensation reaction? Explain why or why not.




4-methylpyridine **A**

24.72 Draw a stepwise mechanism for the following reaction, one step in the synthesis of the cholesterol-lowering drug pitavastatin, marketed in Japan as a calcium salt under the name Livalo.



pitavastatin


24.73 Devise a stepwise mechanism for the following reaction, a key step in the synthesis of the antibiotic abyssomicin C. Abyssomicin C was isolated from sediment collected from almost 1000 ft below the surface in the Sea of Japan. (Hint: The mechanism begins with a Dieckmann reaction.)



abyssomicin C

25 Amines

25.1 Introduction
25.2 Structure and bonding
25.3 Nomenclature
25.4 Physical properties
25.5 Spectroscopic properties
25.6 Interesting and useful amines
25.7 Preparation of amines
25.8 Reactions of amines—General features
25.9 Amines as bases
25.10 Relative basicity of amines and other compounds
25.11 Amines as nucleophiles
25.12 Hofmann elimination
25.13 Reaction of amines with nitrous acid
25.14 Substitution reactions of aryl diazonium salts
25.15 Coupling reactions of aryl diazonium salts
25.16 Application: Synthetic dyes and sulfa drugs




Scopolamine is a complex amine that occurs in angel's trumpets, ornamental plants with large trumpet-shaped flowers native to South America. Transdermal patches that release a minute dose of scopolamine are used to treat the nausea and vomiting associated with motion sickness. Scopolamine is an **alkaloid,** a naturally occurring amine derived from plant sources. In Chapter 25 we learn about the properties and reactions of amines like scopolamine.

We now leave the chemistry of carbonyl compounds to concentrate on **amines,** organic derivatives of ammonia (NH_3), formed by replacing one or more hydrogen atoms by alkyl or aryl groups. **Amines are stronger bases and better nucleophiles than other neutral organic compounds,** so much of Chapter 25 focuses on these properties.


Like that of alcohols, the chemistry of amines does not fit neatly into one reaction class, and this can make learning the reactions of amines challenging. Many interesting natural products and widely used drugs are amines, so you also need to know how to introduce this functional group into organic molecules.

25.1 Introduction

Amines **are organic nitrogen compounds,** formed by replacing one or more hydrogen atoms of ammonia (NH_3) with alkyl groups. As discussed in Section 3.2, amines are classified as 1°, 2°, or 3° by the number of alkyl groups bonded to the *nitrogen* atom.



Like ammonia, **the amine nitrogen atom has a nonbonded electron pair,** making it both a base and a nucleophile. As a result, amines react with electrophiles to form **ammonium salts**—compounds with a positively charged ammonium ion and an anionic counterion.



- The chemistry of amines is dominated by the nonbonded electron pair on the nitrogen atom.

25.2 Structure and Bonding

An amine nitrogen atom is surrounded by three atoms and one nonbonded electron pair, making the N atom ***sp*^{3} hybridized** and **trigonal pyramidal,** with bond angles of approximately 109.5°.





Figure 25.1
Electrostatic potential plots of CH_3NH_2 and $(CH_3)_3N$



- Both amines clearly show the electron-rich region (in red) at the N atom.


Because nitrogen is much more electronegative than carbon or hydrogen, **the C−N and N−H bonds are all polar,** with the N atom electron rich and the C and H atoms electron poor. The electrostatic potential maps in Figure 25.1 show the polar C−N and N−H bonds in CH_3NH_2 (methylamine) and $(CH_3)_3N$ (trimethylamine).

An amine nitrogen atom bonded to an electron pair and three different alkyl groups is technically a stereogenic center, so two nonsuperimposable trigonal pyramids can be drawn.



nonsuperimposable mirror images

This does not mean, however, that such an amine exists as two different enantiomers, because one is rapidly converted to the other at room temperature. The amine flips inside out, passing through a trigonal planar (achiral) transition state. **Because the two enantiomers interconvert, we can ignore the chirality of the amine nitrogen.**



planar transition state


In contrast, the chirality of an ammonium ion with four different groups on N cannot be ignored. Because there is no nonbonded electron pair on the nitrogen atom, interconversion cannot occur, and the N atom is just like a carbon atom with four different groups around it.




chiral ammonium ion

- **The N atom of an ammonium ion is a stereogenic center when N is surrounded by four different groups.**

Problem 25.1 Label the stereogenic centers in each compound.

a. 

b. 

dobutamine
(heart stimulant used in stress tests to measure cardiac fitness)

25.3 Nomenclature

25.3A Primary Amines

Primary amines are named using either systematic or common names.

- To assign the **systematic name,** find the longest continuous carbon chain bonded to the amine nitrogen, and change the ***-e*** ending of the parent alkane to the suffix ***-amine.*** Then use the usual rules of nomenclature to number the chain and name the substituents.
- To assign a **common name,** name the alkyl group bonded to the nitrogen atom and add the suffix ***-amine,*** forming a single word.




Systematic name: **methanamine**
Common name: **methylamine**


Systematic name: **cyclohexanamine**
Common name: **cyclohexylamine**

25.3B Secondary and Tertiary Amines

Secondary and tertiary amines having identical alkyl groups are named by using the prefix **di-** or **tri-** with the name of the primary amine.



triethylamine




diisopropylamine

Secondary and tertiary amines having more than one kind of alkyl group are named as ***N*-substituted primary amines,** using the following procedure.

How To Name 2° and 3° Amines with Different Alkyl Groups


Example Name the following 2° amine: $(CH_3)_2CHNHCH_3$.

Step [1] Designate the longest alkyl chain (or largest ring) bonded to the N atom as the parent amine and assign a systematic name.



3 C's in the longest chain ----→ propan-2-amine


Step [2] Name the other groups on the N atom as alkyl groups, alphabetize the names, and put the prefix *N-* before the name.




methyl substituent

Answer: *N*-methylpropan-2-amine

Sample Problem 25.1 Name each amine.

a. 

b. 


Solution

a. [1] A 1° amine: Find and name the longest chain containing the amine nitrogen.



pentane (5 C's) ----→ **pentan*amine***

[2] Number and name the substituents.




You must use a number to show the location of the NH_2 group.

Answer: 4-methylpentan-1-amine


b. For a 3° amine, one alkyl group on N is the principal R group and the others are substituents.

[1] Name the ring bonded to the N.



cyclopentanamine


[2] Name the substituents.





Two N's are needed, one for each alkyl group.


Answer: *N*-ethyl-*N*-methylcyclopentanamine


Problem 25.2 Name each amine.


a. 

b. 

c. 


d. 

e. 


f. 

25.3C Aromatic Amines


Aromatic amines are named as derivatives of aniline.



aniline



N-ethylaniline



o-bromoaniline

25.3D Miscellaneous Nomenclature Facts


An **NH_2** group named as a substituent is called an **amino group.**

There are many different **nitrogen heterocycles,** and each ring type is named differently depending on the number of N atoms in the ring, the ring size, and whether it is aromatic or not. The structures and names of common nitrogen heterocycles are shown in Figure 25.2.

Problem 25.3 Draw a structure corresponding to each name.

a. 2,4-dimethylhexan-3-amine
b. *N*-methylpentan-1-amine
c. *N*-isopropyl-*p*-nitroaniline
d. *N*-methylpiperidine
e. *N,N*-dimethylethanamine
f. 2-aminocyclohexanone
g. *N*-methylaniline
h. *m*-ethylaniline

Figure 25.2
Common nitrogen heterocycles



- Heterocycles with one N atom are numbered to place the N atom at the "1" position.
- Heterocycles with two N atoms are numbered to place one N atom at the "1" position and give the second N atom the lower number.

25.4 Physical Properties

Amines exhibit dipole–dipole interactions because of the polar C−N and N−H bonds. **Primary and secondary amines are also capable of intermolecular hydrogen bonding,** because they contain N−H bonds. Because nitrogen is less electronegative than oxygen, however, intermolecular hydrogen bonds between N and H are *weaker* than those between O and H. How these factors affect the physical properties of amines is summarized in Table 25.1.




Table 25.1 Physical Properties of Amines

Property	Observation
Boiling point and melting point	• Primary (1°) and 2° amines have higher bp's than similar compounds (like ethers) incapable of hydrogen bonding, but lower bp's than alcohols that have stronger intermolecular hydrogen bonds. MW = 74, bp 38 °C; MW = 73, bp 78 °C; MW = 74, bp 118 °C **Increasing intermolecular forces** **Increasing boiling point** • Tertiary (3°) amines have lower boiling points than 1° and 2° amines of comparable molecular weight, because they have no N−H bonds and are incapable of hydrogen bonding. 3° amine: MW = 73, bp 38 °C, **no N–H bond** 2° amine higher bp: MW = 73, bp 56 °C, **N–H bond**
Solubility	• Amines are soluble in organic solvents regardless of size. • All amines having ≤ 5 C's are H_2O soluble because they can hydrogen bond with H_2O (Section 3.4C). • Amines having > 5 C's are H_2O insoluble because the nonpolar alkyl portion is too large to dissolve in the polar H_2O solvent.

MW = molecular weight

Problem 25.4 Arrange the compounds in order of increasing boiling point.



25.5 Spectroscopic Properties

Amines exhibit characteristic features in their mass spectra, IR spectra, and ^{1}H and ^{13}C NMR spectra.


25.5A Mass Spectra

Amines differ from compounds that contain only C, H, and O atoms, which always have a molecular ion with an *even* mass in their mass spectra. This is apparent in the mass spectrum of butan-1-amine, which is shown in Figure 25.3.

The general molecular formula for an amine with one N atom is $C_nH_{2n+3}N$.

- **Amines with an odd number of N atoms give an odd molecular ion in their mass spectra.**

Figure 25.3
Mass spectrum of butan-1-amine



- The molecular ion for $CH_3CH_2CH_2CH_2NH_2$ occurs at m/z = 73. This odd mass for a molecular ion is characteristic of an amine with an odd number of N atoms.


25.5B IR Spectra

Amines with N−H bonds show characteristic absorptions in their IR spectra.

- **1° Amines show *two* N−H absorptions at 3300–3500 cm^{-1}.**
- **2° Amines show *one* N−H absorption at 3300–3500 cm^{-1}.**

Because 3° amines have no N−H bonds, they do *not* absorb in this region in their IR spectra. The single bond region (> 2500 cm^{-1}) of the IR spectra for 1°, 2°, and 3° amines illustrates these features in Figure 25.4.

Figure 25.4 The single bond region of the IR spectra for a 1°, 2°, and 3° amine



25.5C NMR Spectra


Amines exhibit the following characteristic ^{1}H NMR and ^{13}C NMR absorptions.

- **The NH signal appears between 0.5 and 5.0 ppm.** The exact location depends on the degree of hydrogen bonding and the concentration of the sample.
- The protons on the carbon bonded to the amine nitrogen are deshielded and typically absorb at **2.3–3.0 ppm.**
- In the **^{13}C NMR** spectrum, the carbon bonded to the N atom is deshielded and typically absorbs at **30–50 ppm.**

Like the OH absorption of an alcohol, the **NH absorption is not split by adjacent protons, nor does it cause splitting of adjacent C–H absorptions in a ^{1}H NMR spectrum.** The NH peak of an amine is sometimes somewhat broader than other peaks in the spectrum. The ^{1}H NMR spectrum of *N*-methylaniline is shown in Figure 25.5.

Problem 25.5 What is the structure of an unknown compound with molecular formula $C_6H_{15}N$ that gives the following ^{1}H NMR absorptions: 0.9 (singlet, 1 H), 1.10 (triplet, 3 H), 1.15 (singlet, 9 H), and 2.6 (quartet, 2 H) ppm?

Figure 25.5
The ^{1}H NMR spectrum of *N*-methylaniline



- The CH_3 group appears as a singlet at 2.7 ppm because there is no splitting by the adjacent NH proton.
- The NH proton appears as a broad singlet at 3.6 ppm.
- The five H atoms of the aromatic ring appear as a complex pattern at 6.6–7.2 ppm.

25.6 Interesting and Useful Amines

A great many simple and complex amines occur in nature, and others with biological activity have been synthesized in the lab.


25.6A Simple Amines and Alkaloids

Many low molecular weight amines have *very* foul odors. **Trimethylamine** [$(CH_3)_3N$], formed when enzymes break down certain fish proteins, has the characteristic odor of rotting fish. **Putrescine** ($NH_2CH_2CH_2CH_2CH_2NH_2$) and **cadaverine** ($NH_2CH_2CH_2CH_2CH_2CH_2NH_2$) are both poisonous diamines with putrid odors. They, too, are present in rotting fish, and are partly responsible for the odors of semen, urine, and bad breath.


The word **alkaloid** is derived from the word *alkali,* because aqueous solutions of alkaloids are slightly basic.

Naturally occurring amines derived from plant sources are called **alkaloids.** Alkaloids previously encountered in the text include **quinine** (Problem 17.13), **morphine** (Section 22.9), and **cocaine** (Problem 3.54). Three other common alkaloids are **atropine, nicotine,** and **coniine,** illustrated in Figure 25.6.

Figure 25.6
Three common alkaloids—Atropine, nicotine, and coniine




25.6B Histamine and Antihistamines




Histamine, a rather simple triamine first discussed in Section 17.8, is responsible for a wide variety of physiological effects. Histamine is a vasodilator (it dilates capillaries), so it is released at the site of an injury or infection to increase blood flow. It is also responsible for the symptoms of allergies, including a runny nose and watery eyes. In the stomach, histamine stimulates the secretion of acid.

Understanding the central role of histamine in these biochemical processes has helped chemists design drugs to counteract some of its undesirable effects.



fexofenadine
antihistamine




cimetidine
(Tagamet)
antiulcer drug

Antihistamines bind to the same active site of the enzyme that binds histamine in the cell, but they evoke a different response. An antihistamine like **fexofenadine** (trade name Allegra), for example, inhibits vasodilation, so it is used to treat the symptoms of the common cold and allergies. Unlike many antihistamines, fexofenadine does not cause drowsiness because it binds to histamine receptors but does not cross the blood–brain barrier, so it does not affect the central nervous system. **Cimetidine** (trade name Tagamet) is a histamine mimic that blocks the secretion of hydrochloric acid in the stomach, so it is used to treat individuals with ulcers.


25.6C Derivatives of 2-Phenylethanamine

A large number of physiologically active compounds are derived from **2-phenylethanamine, $C_6H_5CH_2CH_2NH_2$.** Some of these compounds are synthesized in cells and needed to maintain healthy mental function. Others are isolated from plant sources or are synthesized in the laboratory and have a profound effect on the brain because they interfere with normal neurochemistry. These compounds include **adrenaline, noradrenaline, methamphetamine,** and **mescaline.** Each contains a benzene ring bonded to a two-carbon unit with a nitrogen atom (shown in red).




adrenaline
(epinephrine)

a hormone secreted in response to stress
(Figure 7.19)




noradrenaline
(norepinephrine)

a neurotransmitter that increases heart rate
and dilates air passages



methamphetamine

an addictive stimulant sold as
speed, meth, or crystal meth




mescaline

a hallucinogen isolated from peyote, a cactus native
to the southwestern United States and Mexico


Another example, **dopamine,** is a neurotransmitter, a chemical messenger released by one nerve cell (neuron), which then binds to a receptor in a neighboring target cell (Figure 25.7). Dopamine affects brain processes that control movement and emotions, so proper dopamine levels are necessary to maintain an individual's mental and physical health. For example, when dopamine-producing neurons die, the level of dopamine drops, resulting in the loss of motor control symptomatic of Parkinson's disease.

Figure 25.7 Dopamine—A neurotransmitter




Serotonin is a neurotransmitter that plays an important role in mood, sleep, perception, and temperature regulation. A deficiency of serotonin causes depression. Understanding the central role of serotonin in determining one's mood has led to the development of a variety of drugs for the treatment of depression. The most widely used antidepressants today are selective serotonin reuptake inhibitors (SSRIs). These drugs act by inhibiting the reuptake of serotonin by the neurons that produce it, thus effectively increasing its concentration. Fluoxetine (trade name Prozac) is a common antidepressant that acts in this way.

Cocaine, amphetamines, and several other addicting drugs increase the level of dopamine in the brain, which results in a pleasurable "high." With time, the brain adapts to increased dopamine levels, so more drug is required for the same sensation.



serotonin




fluoxetine


Drugs that interfere with the metabolism of serotonin have a profound effect on mental state. For example, bufotenin, isolated from *Bufo* toads from the Amazon jungle, and psilocin, isolated from *Psilocybe* mushrooms, are very similar in structure to serotonin and both cause intense hallucinations.



Bufo toads from the Amazon jungle are the source of the hallucinogen bufotenin.




bufotenin



psilocin


Problem 25.6 LSD (a hallucinogen) and codeine (a narcotic) are structurally more complex derivatives of 2-phenylethanamine. Identify the atoms of 2-phenylethanamine in each of the following compounds.

a.



LSD
lysergic acid diethyl amide

b.



codeine

25.7 Preparation of Amines

In the preparations of a given functional group, many different starting materials form a common product (amines, in this case).

Three types of reactions are used to prepare an amine:

[1] **Nucleophilic substitution** using nitrogen nucleophiles
[2] **Reduction** of other nitrogen-containing functional groups
[3] **Reductive amination** of aldehydes and ketones


25.7A Nucleophilic Substitution Routes to Amines

Nucleophilic substitution is the key step in two different methods for synthesizing amines: direct nucleophilic substitution and the Gabriel synthesis of 1° amines.

Direct Nucleophilic Substitution

Conceptually, the simplest method to synthesize an amine is by **S_N2 reaction of an alkyl halide with NH_3 or an amine.** The method requires two steps:


[1] **Nucleophilic attack** of the nitrogen nucleophile forms an ammonium salt.
[2] **Removal of a proton** on N forms the amine.




The identity of the nitrogen nucleophile determines the type of amine or ammonium salt formed as product. One new carbon–nitrogen bond is formed in each reaction. Because the reaction follows an S_N2 mechanism, the alkyl halide must be unhindered—that is, CH_3X or RCH_2X.

Although this process seems straightforward, polyalkylation of the nitrogen nucleophile limits its usefulness. **Any amine formed by nucleophilic substitution still has a nonbonded electron pair, making it a nucleophile as well.** It will react with remaining alkyl halide to form a more substituted amine. Because of this, a mixture of 1°, 2°, and 3° amines often results. Only the final product—called a **quaternary ammonium salt** because it has four alkyl groups on N—cannot react further, and so the reaction stops.

As a result, this reaction is most useful for preparing 1° amines by using a very large excess of NH_3 (a relatively inexpensive starting material) and for preparing quaternary ammonium salts by alkylating any nitrogen nucleophile with one or more equivalents of alkyl halide.




Problem 25.7 Draw the product of each reaction.



The Gabriel Synthesis of 1° Amines

To avoid polyalkylation, a nitrogen nucleophile can be used that reacts in a single nucleophilic substitution reaction—that is, the reaction forms a product that does not contain a nucleophilic nitrogen atom capable of reacting further.

The **Gabriel synthesis** consists of two steps and uses a resonance-stabilized nitrogen nucleophile to synthesize 1° amines via nucleophilic substitution. The Gabriel synthesis begins with **phthalimide,** one of a group of compounds called **imides.** The **N−H bond of an imide is especially acidic** because the resulting anion is resonance stabilized by the two flanking carbonyl groups.




phthalimide
pK_a = 10

resonance-stabilized anion


An acid–base reaction forms a nucleophilic anion that can react with an unhindered alkyl halide—that is, CH_3X or RCH_2X—in an S_N2 reaction to form a substitution product. This alkylated imide is then hydrolyzed with aqueous base to give a 1° amine and a dicarboxylate. This reaction is similar to the hydrolysis of amides to afford carboxylate anions and amines, as discussed in

Section 22.13. The overall result of this two-step sequence is **nucleophilic substitution of X by NH_2,** so the Gabriel synthesis can be used to prepare 1° amines only.

Steps in the Gabriel synthesis




- **The Gabriel synthesis converts an alkyl halide into a 1° amine by a two-step process: nucleophilic substitution followed by hydrolysis.**




new C–N bond in red

Problem 25.8 What alkyl halide is needed to prepare each 1° amine by a Gabriel synthesis?



Problem 25.9 Which amines cannot be prepared by a Gabriel synthesis? Explain your choices.




25.7B Reduction of Other Functional Groups That Contain Nitrogen

Amines can be prepared by reduction of nitro compounds, nitriles, and amides. Because the details of these reactions have been discussed previously, they are presented here in summary form only.


[1] From nitro compounds (Section 18.15C)

Nitro groups are reduced to 1° amines using a variety of reducing agents.




[2] From nitriles (Section 22.18B)

Nitriles are reduced to 1° amines with $LiAlH_4$.




Because a cyano group is readily introduced by S_N2 substitution of alkyl halides with ^-CN, this provides **a two-step method to convert an alkyl halide to a 1° amine with one more carbon atom.** The conversion of $(CH_3)_2CHCH_2CH_2Br$ to $(CH_3)_2CHCH_2CH_2CH_2NH_2$ illustrates this two-step sequence.




[3] From amides (Section 20.7B)


Primary (1°), 2°, and 3° amides are reduced to 1°, 2°, and 3° amines, respectively, by using $LiAlH_4$.




Problem 25.10 What nitro compound, nitrile, and amide are reduced to each compound?



Problem 25.11 What amine is formed by reduction of each amide?



Problem 25.12 Which amines cannot be prepared by reduction of an amide?




25.7C Reductive Amination of Aldehydes and Ketones

Reductive amination is a two-step method that converts aldehydes and ketones into 1°, 2°, and 3° amines. Let's first examine this method using NH_3 to prepare 1° amines. There are two distinct parts in reductive amination:

[1] **Nucleophilic attack of NH_3 on the carbonyl group forms an imine** (Section 21.11A), which is not isolated; then,

[2] **Reduction of the imine forms an amine** (Section 20.7B).




- Reductive amination replaces a C=O by a C–H and C–N bond.

The most effective reducing agent for this reaction is sodium cyanoborohydride ($NaBH_3CN$). This hydride reagent is a derivative of sodium borohydride ($NaBH_4$), formed by replacing one H atom by CN.

$NaBH_3CN$
sodium cyanoborohydride

Reductive amination combines two reactions we have already learned in a different way. Two examples are shown. The second reaction is noteworthy because the product is **amphetamine,** a potent central nervous system stimulant.



With a 1° or 2° amine as starting material, reductive amination is used to prepare 2° and 3° amines, respectively. Note the result: Reductive amination uses an aldehyde or ketone to replace one H atom on a nitrogen atom by an alkyl group, making a more substituted amine.






Figure 25.8
Synthesis of methamphetamine by reductive amination




- In reductive amination, one of the H atoms bonded to N is replaced by an alkyl group. As a result, a 1° amine is converted to a 2° amine and a 2° amine is converted to a 3° amine. In this reaction, CH_3NH_2 (a 1° amine) is converted to methamphetamine (a 2° amine).

The synthesis of methamphetamine (Section 25.6C) by reductive amination is illustrated in Figure 25.8.

Problem 25.13 Draw the product of each reaction.




Problem 25.14 Enalapril, a drug used to treat hypertension, is prepared from compounds **D** and **E** by reductive amination. What is the structure of enalapril?




To use reductive amination in synthesis, you must be able to determine what aldehyde or ketone and nitrogen compound are needed to prepare a given amine—that is, you must work backwards in the retrosynthetic direction. Keep in mind the following two points:

- **One alkyl group on N comes from the carbonyl compound.**
- **The remainder of the molecule comes from NH_3 or an amine.**




For example, 2-phenylethanamine is a 1° amine, so it has only one alkyl group bonded to N. This alkyl group must come from the carbonyl compound, and the rest of the molecule then comes from the nitrogen component. **For a 1° amine, the nitrogen component must be NH_3.**



There is usually more than one way to use reductive amination to synthesize 2° and 3° amines, as shown in Sample Problem 25.2 for a 2° amine.

Sample Problem 25.2 What aldehyde or ketone and nitrogen component are needed to synthesize *N*-ethylcyclohexanamine by a reductive amination reaction?




N-ethylcyclohexanamine

Solution

Because *N*-ethylcyclohexanamine has two different alkyl groups bonded to the N atom, either R group can come from the carbonyl component and there are two different ways to form a C – N bond by reductive amination.

Possibility [1] Use $CH_3CH_2NH_2$ and cyclohexanone.


Possibility [2] Use cyclohexanamine and an aldehyde.



Because reductive amination adds one R group to a nitrogen atom, both routes to form the 2° amine begin with a 1° amine.


Problem 25.15 What starting materials are needed to prepare each drug using reductive amination? Give all possible pairs of compounds when more than one route is possible.

a.



rimantadine
antiviral used to treat influenza

b.




pseudoephedrine
nasal decongestant

Problem 25.16 (a) Explain why phentermine [$PhCH_2C(CH_3)_2NH_2$] can't be made by a reductive amination reaction. (b) Give a systematic name for phentermine, one of the components of the banned diet drug fen–phen.


25.8 Reactions of Amines—General Features

- The chemistry of amines is dominated by the lone pair of electrons on nitrogen.

Only three elements in the second row of the periodic table have nonbonded electron pairs in neutral organic compounds: nitrogen, oxygen, and fluorine. Because basicity and nucleophilicity decrease across the row, **nitrogen is the most basic and most nucleophilic** of these elements.




- Amines are stronger bases and nucleophiles than other neutral organic compounds.



- Amines react as bases with compounds that contain acidic protons.
- Amines react as nucleophiles with compounds that contain electrophilic carbons.


25.9 Amines as Bases

Amines react as bases with a variety of organic and inorganic acids.



To favor the products, the **pK_a of HA must be < 10.**

What acids can be used to protonate an amine? Equilibrium favors the products of an acid–base reaction when the weaker acid and base are formed. Because the pK_a of many protonated amines is 10–11, the **pK_a of the starting acid must be less than 10** for equilibrium to favor the products. Amines are thus readily protonated by strong inorganic acids like HCl and H_2SO_4, and by carboxylic acids as well.




Equilibrium favors the products.

The principles used in an extraction procedure were detailed in Section 19.12.

Because amines are protonated by aqueous acid, they can be separated from other organic compounds by extraction using a separatory funnel. **Extraction separates compounds based on solubility differences.** When an amine is protonated by aqueous acid, its solubility properties change.

For example, when cyclohexanamine is treated with aqueous HCl, it is protonated, forming an ammonium salt. Because the ammonium salt is ionic, it is soluble in water, but insoluble in organic solvents. A similar acid–base reaction does not occur with other organic compounds like alcohols, which are much less basic.



cyclohexanamine

- insoluble in H_2O
- soluble in CH_2Cl_2

cyclohexanammonium chloride

- soluble in H_2O
- insoluble in CH_2Cl_2


This difference in acid–base chemistry can be used to separate cyclohexanamine and cyclohexanol by the stepwise extraction procedure illustrated in Figure 25.9.

Figure 25.9 Separation of cyclohexanamine and cyclohexanol by an extraction procedure

Step [1] Dissolve cyclohexanamine and cyclohexanol in CH_2Cl_2.

Step [2] Add 10% HCl solution to form two layers.

Step [3] Separate the layers.



- Both compounds dissolve in the organic solvent CH_2Cl_2.


- Adding 10% aqueous HCl solution forms two layers. When the two layers are mixed, the HCl protonates the amine (RNH_2) to form $RNH_3^+Cl^-$, which dissolves in the aqueous layer.
- The cyclohexanol remains in the CH_2Cl_2 layer.


- Draining the lower layer out the bottom stopcock separates the two layers.
- Cyclohexanol (dissolved in CH_2Cl_2) is in one flask. The ammonium salt, $RNH_3^+Cl^-$ (dissolved in water), is in another flask.


- **An amine can be separated from other organic compounds by converting it to a water-soluble ammonium salt by an acid–base reaction.**

Thus, the water-soluble salt $C_6H_{11}NH_3^+Cl^-$ (obtained by protonation of $C_6H_{11}NH_2$) can be separated from water-insoluble cyclohexanol by an aqueous extraction procedure.

Problem 25.17 Draw the products of each acid–base reaction. Indicate whether equilibrium favors the reactants or products.

a. 


c. 

b. 



Many antihistamines and decongestants are sold as their ammonium salts.


Many water-insoluble amines with useful medicinal properties are sold as their water-soluble ammonium salts, which are more easily transported through the body in the aqueous medium of the blood. Benadryl, formed by treating diphenhydramine with HCl, is an over-the-counter antihistamine that is used to relieve the itch and irritation of skin rashes and hives.




diphenhydramine
water insoluble

ammonium salt
Benadryl
(diphenhydramine hydrochloride)
water soluble

Problem 25.18 Write out steps to show how each of the following pairs of compounds can be separated by an extraction procedure.


a. 

b. 

25.10 Relative Basicity of Amines and Other Compounds

The relative acidity of different compounds can be compared using their pK_a values. The relative *basicity* of different compounds (such as amines) can be compared using the pK_a values of their *conjugate acids.*

- **The weaker the conjugate acid, the higher its pK_a and the stronger the base.**




To compare the basicity of two compounds, keep in mind the following:

- **Any factor that increases the electron density on the N atom increases an amine's basicity.**
- **Any factor that decreases the electron density on N decreases an amine's basicity.**

25.10A Comparing an Amine and NH_3

Because alkyl groups are electron donating, they increase the electron density on nitrogen, which makes an amine like $CH_3CH_2NH_2$ more basic than NH_3. In fact, the pK_a of $CH_3CH_2NH_3^+$ is higher than the pK_a of NH_4^+, so **$CH_3CH_2NH_2$ is a stronger base than NH_3.**




The relative basicity of 1°, 2°, and 3° amines depends on additional factors, and will not be considered in this text.

- **Primary (1°), 2°, and 3° alkylamines are more basic than NH_3 because of the electron-donating inductive effect of the R groups.**

Problem 25.19 Which compound in each pair is more basic: (a) $(CH_3)_2NH$ and NH_3; (b) $CH_3CH_2NH_2$ and $ClCH_2CH_2NH_2$?

25.10B Comparing an Alkylamine and an Arylamine

To compare an alkylamine ($CH_3CH_2NH_2$) and an arylamine ($C_6H_5NH_2$, aniline), we must look at the availability of the nonbonded electron pair on N. With $CH_3CH_2NH_2$, the electron pair is localized on the N atom. With an arylamine, however, the electron pair is now delocalized on the benzene ring. This *decreases* the electron density on N, and makes $C_6H_5NH_2$ less basic than $CH_3CH_2NH_2$.



The electron pair is localized on N.

stronger base

The electron pair is delocalized on the benzene ring.

weaker base


The pK_a values support this reasoning. Because the pK_a of $CH_3CH_2NH_3^+$ is higher than the pK_a of $C_6H_5NH_3^+$ (10.8 vs. 4.6), **$CH_3CH_2NH_2$ is a stronger base than $C_6H_5NH_2$.**

- **Arylamines are less basic than alkylamines because the electron pair on N is delocalized.**

Substituted anilines are more or less basic than aniline depending on the nature of the substituent.

- **Electron-donor groups add electron density to the benzene ring, making the arylamine more basic than aniline.**

D = electron-donor group




D makes the amine more basic than aniline.

D
–NH_2
–OH
–OR
–NHCOR
–R

- **Electron-withdrawing groups remove electron density from the benzene ring, making the arylamine less basic than aniline.**

W = electron-withdrawing group



W makes the amine less basic than aniline.


W	
−X	−CN
−CHO	$-SO_3H$
−COR	$-NO_2$
−COOR	$-NR_3^+$
−COOH	

The effect of electron-donating and electron-withdrawing groups on the acidity of substituted benzoic acids was discussed in Section 19.11.


Whether a substituent donates or withdraws electron density depends on the balance of its inductive and resonance effects (Section 18.6 and Figure 18.8).

Sample Problem 25.3


Rank the following compounds in order of increasing basicity.



aniline




p-nitroaniline



p-methylaniline (*p*-toluidine)


Solution

***p*-Nitroaniline: NO_2** is an electron-withdrawing group, making the amine **less basic** than aniline.




The lone pair on N is delocalized on the O atom, decreasing the basicity of the amine.

***p*-Methylaniline:** CH_3 has an electron-donating inductive effect, making the amine **more basic** than aniline.



CH_3 inductively donates electron density, increasing the basicity of the amine.




The electrostatic potential plots in Figure 25.10 demonstrate that the electron density of the nitrogen atoms in these anilines increases in the order shown.

Problem 25.20

Rank the compounds in each group in order of increasing basicity.

a.



b.





Figure 25.10
Electrostatic potential plots of substituted anilines



The NH_2 group gets more electron rich as the para substituent changes from $NO_2 \rightarrow H \rightarrow CH_3$. This is indicated by the color change around NH_2 (from green to yellow to red) in the electrostatic potential plot.

25.10C Comparing an Alkylamine and an Amide


To compare the basicity of an alkylamine (RNH_2) and an amide ($RCONH_2$), we must once again compare the availability of the nonbonded electron pair on nitrogen. With RNH_2, the electron pair is localized on the N atom. With an amide, however, the electron pair is delocalized on the carbonyl oxygen by resonance. This *decreases* the electron density on N, making **an amide much less basic than an alkylamine.**



The electron pair on N is delocalized on O by resonance.


- **Amides are much less basic than amines because the electron pair on N is delocalized.**

In fact, amides are not much more basic than any carbonyl compound. When an amide is treated with acid, **protonation occurs at the carbonyl oxygen, *not* the nitrogen,** because the resulting cation is resonance stabilized.



three resonance structures for the conjugate acid

Problem 25.21 Rank the following compounds in order of increasing basicity.



25.10D Heterocyclic Aromatic Amines

To determine the relative basicity of nitrogen heterocycles that are also aromatic, you must know whether the nitrogen lone pair is part of the aromatic π system.

For example, pyridine and pyrrole are both aromatic, but the nonbonded electron pair on the N atom in these compounds is located in different orbitals. Recall from Section 17.8C that the **lone pair of electrons in pyridine occupies an sp^2 hybridized orbital,** perpendicular to the π bonds of the molecule, so it is *not* part of the aromatic system, whereas that of pyrrole resides in a *p* orbital, making it part of the aromatic system. **The lone pair on pyrrole, therefore, is delocalized on all of the atoms of the five-membered ring,** making pyrrole a much weaker base than pyridine.



pyridine — pyrrole

The lone pair resides in an sp^2 hybrid orbital.

The lone pair resides in a *p* orbital and is delocalized in the ring.

Protonation of pyrrole occurs at a ring carbon, not the N atom, as noted in Problem 17.50.

As a result, the pK_a of the conjugate acid of pyrrole is much less than that for the conjugate acid of pyridine.



higher pK_a weaker acid — pyridine stronger base — lower pK_a stronger acid — pyrrole weaker base

pK_a = 5.3 — pK_a = 0.4

- **Pyrrole is much less basic than pyridine because its lone pair of electrons is part of the aromatic π system.**

25.10E Hybridization Effects

The effect of hybridization on the acidity of an H–A bond was first discussed in Section 2.5D.

The hybridization of the orbital that contains an amine's lone pair also affects its basicity. This is illustrated by comparing the basicity of **piperidine** and **pyridine,** two nitrogen heterocycles. The lone pair in piperidine resides in an sp^3 hybrid orbital that has 25% *s*-character. The lone pair in pyridine resides in an sp^2 hybrid orbital that has 33% *s*-character.



piperidine — pyridine

The lone pair is in an sp^3 hybrid orbital.

The lone pair is in an sp^2 hybrid orbital.

- **The higher the percent *s*-character of the orbital containing the lone pair, the more tightly the lone pair is held, and the weaker the base.**

Pyridine is a weaker base than piperidine because its nonbonded pair of electrons resides in an sp^2 hybrid orbital. Although pyridine is an aromatic amine, its lone pair is *not* part of the delocalized π system, so its basicity is determined by the hybridization of its N atom. As a result, the

pK_a value for the conjugate acid of pyridine is much lower than that for the conjugate acid of piperidine, making pyridine the weaker base.



lower pK_a
stronger acid

pK_a = 5.3

weaker base


higher pK_a
weaker acid

pK_a = 11.1

stronger base

Problem 25.22 Which nitrogen atom in each compound is more basic?

a.



DMAP
4-(*N*,*N*-dimethylamino)pyridine

b.



nicotine


25.10F Summary

Acid–base chemistry is central to many processes in organic chemistry, so it has been a constant theme throughout this text. Tables 25.2 and 25.3 organize and summarize the acid–base principles discussed in Section 25.10. The principles in these tables can be used to determine the most basic site in a molecule that has more than one nitrogen atom, as shown in Sample Problem 25.4.

Table 25.2 Factors That Determine Amine Basicity

Factor	Example
[1] **Inductive effects:** Electron-donating groups bonded to N increase basicity.	• RNH_2, R_2NH, and R_3N are more basic than NH_3.
[2] **Resonance effects:** Delocalizing the lone pair on N decreases basicity.	• Arylamines ($C_6H_5NH_2$) are less basic than alkylamines (RNH_2). • Amides ($RCONH_2$) are much less basic than amines (RNH_2).
[3] **Aromaticity:** Having the lone pair on N as part of the aromatic π system decreases basicity.	• Pyrrole is less basic than pyridine.  less basic more basic
[4] **Hybridization effects:** Increasing the percent *s*-character in the orbital with the lone pair decreases basicity.	• Pyridine is less basic than piperidine.  less basic more basic

Table 25.3 Table of pK_a Values of Some Representative Organic Nitrogen Compounds

	Compound	pK_a of the conjugate acid
Ammonia	NH_3	9.3
Alkylamines[a]		11.1
	$(CH_3CH_2)_2NH$	11.1
	$(CH_3CH_2)_3N$	11.0
	$CH_3CH_2NH_2$	10.8
Arylamines[b]	*p*-$CH_3OC_6H_4NH_2$	5.3
	p-$CH_3C_6H_4NH_2$	5.1
	$C_6H_5NH_2$	4.6
	p-$NO_2C_6H_4NH_2$	1.0
Heterocyclic aromatic amines[c]		5.3
		0.4
Amides	$RCONH_2$	−1


[a] Alkylamines have pK_a values of ~10—11.
[b] The pK_a *decreases* as the electron density of the benzene ring *decreases*.
[c] The pK_a depends on whether the lone pair of N is localized or delocalized.

Sample Problem 25.4

Which N atom in chloroquine is the strongest base?




Since 1945 chloroquine has been used to treat malaria, an infectious disease caused by a protozoan parasite that is spread by the *Anopheles* mosquito.



chloroquine

Solution

Examine the nitrogen atoms in chloroquine, labeled in red, blue, and green, and recall that decreasing the electron density on N decreases basicity.



- **N** is bonded to an aromatic ring, so its lone pair is delocalized in the ring like aniline, decreasing basicity.
- The lone pair is localized on **N**, but **N** is sp^2 hybridized. Increasing percent *s*-character decreases basicity.
- **N** has a localized lone pair and is sp^3 hybridized, making it the most basic site in the molecule.


Problem 25.23

Which N atom in each compound is more basic? What product is formed when each compound is treated with HCl?




Lupin (Problems 25.23 and 25.78) is a genus of flowering plant that is abundant in the Andes of Peru and along the roadside in parts of Alaska.

a.



matrine
alkaloid in lupin

b.



quinine
antimalarial drug

25.11 Amines as Nucleophiles


Amines react as nucleophiles with electrophilic carbon atoms. The details of these reactions have been described in Chapters 21 and 22, so they are summarized here only to emphasize the similar role that the amine nitrogen plays.

- **Amines attack carbonyl groups to form products of nucleophilic addition or substitution.**

The nature of the product depends on the carbonyl electrophile. These reactions are limited to 1° and 2° amines, because only these compounds yield neutral organic products.


[1] Reaction of 1° and 2° amines with aldehydes and ketones (Sections 21.11–21.12)

Aldehydes and ketones react with 1° amines to form **imines** and with 2° amines to form **enamines.** Both reactions involve nucleophilic addition of the amine to the carbonyl group to form a carbinolamine, which then loses water to form the final product.




[2] Reaction of NH_3 and 1° and 2° amines with acid chlorides and anhydrides (Sections 22.8–22.9)


Acid chlorides and anhydrides react with NH_3, 1° amines, and 2° amines to form **1°, 2°, and 3° amides,** respectively. These reactions involve attack of the nitrogen nucleophile on the carbonyl group followed by elimination of a leaving group (Cl^- or RCO_2^-). The overall result of this reaction is substitution of the leaving group by the nitrogen nucleophile.



Problem 25.24 Draw the products formed when each carbonyl compound reacts with the following amines: [1] $CH_3CH_2CH_2NH_2$; [2] $(CH_3CH_2)_2NH$.



The conversion of amines to amides is useful in the synthesis of substituted anilines. For example, aniline itself does not undergo Friedel–Crafts reactions (Section 18.10B). Instead, its basic lone pair on N reacts with the Lewis acid ($AlCl_3$) to form a deactivated complex that does not undergo further reaction.



The N atom of an amide, however, is much less basic than the N atom of an amine, so it does not undergo a similar Lewis acid–base reaction with $AlCl_3$. A three-step reaction sequence involving an intermediate amide can thus be used to form the products of the Friedel–Crafts reaction.


[1] **Convert the amine (aniline) into an amide (acetanilide).**

[2] **Carry out the Friedel–Crafts reaction.**

[3] **Hydrolyze the amide** to generate the free amino group.

This three-step procedure is illustrated in Figure 25.11. In this way, **the amide serves as a protecting group for the NH_2 group,** in much the same way that *tert*-butyldimethylsilyl ethers and acetals are used to protect alcohols and carbonyls, respectively (Sections 20.12 and 21.15).

Problem 25.25 Devise a synthesis of each compound from aniline ($C_6H_5NH_2$).

a. 



b. 

Figure 25.11
An amide as a protecting group for an amine



A three-step sequence uses an amide as a protecting group.

[1] Treatment of aniline with acetyl chloride (CH_3COCl) forms an amide (acetanilide).

[2] Acetanilide, having a much less basic N atom compared to aniline, undergoes electrophilic aromatic substitution under Friedel–Crafts conditions, forming a mixture of ortho and para products.

[3] Hydrolysis of the amide forms the Friedel–Crafts substitution products.

25.12 Hofmann Elimination


Amines, like alcohols, contain a poor leaving group. To undergo a β elimination reaction, for example, a 1° amine would need to lose the elements of NH_3 across two adjacent atoms. The leaving group, $^-NH_2$, is such a strong base, however, that this reaction does not occur.




The only way around this obstacle is to convert $^-NH_2$ into a better leaving group. The most common method to accomplish this is called a **Hofmann elimination,** which converts an amine into a quaternary ammonium salt prior to β elimination.

25.12A Details of the Hofmann Elimination

The **Hofmann elimination** converts an amine into an alkene.



The Hofmann elimination consists of three steps, as shown for the conversion of propan-1-amine to propene.




- In Step [1], the amine reacts as a nucleophile in an S_N2 reaction with excess CH_3I to form a quaternary ammonium salt. **The $N(CH_3)_3$ group thus formed is a much better leaving group than $^-NH_2$.**
- Step [2] converts one ammonium salt into another one with a different anion. The silver(I) oxide, Ag_2O, replaces the I^- anion with ^-OH, a strong base.
- When the ammonium salt is heated in Step [3], **^-OH removes a proton from the β carbon atom,** forming the new π bond of the alkene. The mechanism of elimination is **E2,** so:

- **All bonds are broken and formed in a single step.**
- **Elimination occurs through an anti periplanar geometry—that is, H and $N(CH_3)_3$ are oriented on opposite sides of the molecule.**


The general E2 mechanism for the Hofmann elimination is shown in Mechanism 25.1.

Mechanism 25.1 The E2 Mechanism for the Hofmann Elimination



Elimination occurs with an anti periplanar arrangement of H and $N(CH_3)_3$. Base removes a proton on the β carbon, the electron pair in the C–H bond forms the π bond, and $N(CH_3)_3$ comes off as the leaving group.


All Hofmann elimination reactions result in the formation of a new π bond between the α and β carbon atoms, as shown for cyclohexanamine and 2-phenylethanamine.




To help remember the reagents needed for the steps of the Hofmann elimination, keep in mind what happens in each step.

- **Step [1]** makes a good leaving group by forming a quaternary ammonium salt.
- **Step [2]** provides the strong base, $^-$OH, needed for elimination.
- **Step [3]** is the E2 elimination that forms the new π bond.

Problem 25.26 Draw the product formed by treating each compound with excess CH_3I, followed by Ag_2O, and then heat.

a. 

b. 


c. 

25.12B Regioselectivity of the Hofmann Elimination

There is one major difference between a Hofmann elimination and other E2 eliminations.

- **When constitutional isomers are possible, the major alkene has the *less* substituted double bond in a Hofmann elimination.**

For example, Hofmann elimination of the elements of H and $N(CH_3)_3$ from 2-methylcyclopentanamine, which has two different β carbons (labeled β_1 and β_2), yields two constitutional isomers: the disubstituted alkene **A** (the major product) and the trisubstituted alkene **B** (the minor product).




2-methylcyclopentanamine

A
major product
disubstituted alkene

B
minor product
trisubstituted alkene


This regioselectivity distinguishes a Hofmann elimination from other E2 eliminations, which form the more substituted double bond by the Zaitsev rule (Section 8.5). This result is sometimes explained by the size of the leaving group, $N(CH_3)_3$. **In a Hofmann elimination, the base removes a proton from the less substituted, more accessible β carbon atom, because of the bulky leaving group on the nearby α carbon.**

Sample Problem 25.5 Draw the major product formed from Hofmann elimination of the following amine.



Solution

The amine has three β carbons but two of them are identical, so two alkenes are possible. Draw elimination products by forming alkenes having a C=C between the α and β carbons. The major product has the **less substituted double bond**—that is, the alkene with the C=C between the α and β_1 carbons in this example.



major product
disubstituted alkene

minor product
trisubstituted alkene

Figure 25.12
A comparison of E2 elimination reactions using alkyl halides and amines






Figure 25.12 contrasts the products formed by E2 elimination reactions using an alkyl halide and an amine as starting materials. Treatment of the alkyl halide (2-bromopentane) with base forms the more substituted alkene as the major product, following the **Zaitsev rule.** In contrast, the three-step Hofmann sequence of an amine (pentan-2-amine) forms the less substituted alkene as major product.

Problem 25.27 Draw the major product formed by treating each amine with excess CH_3I, followed by Ag_2O, and then heat.




Problem 25.28 Draw the major product formed in each reaction.




25.13 Reaction of Amines with Nitrous Acid

Nitrous acid, HNO_2, is a weak, unstable acid formed from $NaNO_2$ and a strong acid like HCl.




In the presence of acid, nitrous acid decomposes to **$^+$NO,** the **nitrosonium ion.** This electrophile then goes on to react with the nucleophilic nitrogen atom of amines to form **diazonium salts ($RN_2^+Cl^-$)** from 1° amines and ***N*-nitrosamines ($R_2NN{=}O$)** from 2° amines.



25.13A Reaction of ^{+}NO with 1° Amines


Nitrous acid reacts with 1° alkylamines and arylamines to form **diazonium salts.** This reaction is called **diazotization.**



The mechanism for this reaction consists of many steps. It begins with nucleophilic attack of the amine on the nitrosonium ion, and it can conceptually be divided into two parts: formation of an ***N*-nitrosamine,** followed by loss of H_2O, as shown in Mechanism 25.2.


Mechanism 25.2 Formation of a Diazonium Salt from a 1° Amine

Part [1] Formation of an *N*-nitrosamine



1 – 2 Nucleophilic attack of the amine on the nitrosonium ion (^{+}NO) followed by loss of a proton form an *N*-nitrosamine.


Part [2] Formation of a diazonium salt



3 – 5 Three proton transfers form an intermediate with a good leaving group (H_2O).

6 Loss of water forms a diazonium ion (RN_2^{+}). The diazonium salt formed in this reaction consists of the diazonium ion (RN_2^{+}) and a chloride anion (Cl^{-}).

Alkyl diazonium salts are generally not useful compounds. They readily decompose below room temperature to form carbocations with loss of N_2, a very good leaving group. These carbocations usually form a complex mixture of substitution, elimination, and rearrangement products.




Care must be exercised in handling diazonium salts, because they can explode if allowed to dry.

On the other hand, **aryl diazonium salts are very useful synthetic intermediates.** Although they are rarely isolated and are generally unstable above 0 °C, they are useful starting materials in two general kinds of reactions described in Section 25.14.


25.13B Reaction of ^{+}NO with 2° Amines

Secondary alkylamines and arylamines react with nitrous acid to form *N*-nitrosamines.




Many *N*-nitrosamines are potent carcinogens found in some food and tobacco smoke. Nitrosamines in food can be formed in the same way they are formed in the laboratory: **reaction of a 2° amine with the nitrosonium ion,** formed from nitrous acid (HNO_2). Mechanism 25.3 is shown for the conversion of dimethylamine [$(CH_3)_2NH$] to *N*-nitrosodimethylamine [$(CH_3)_2NN{=}O$].

Mechanism 25.3 Formation of an *N*-Nitrosamine from a 2° Amine



1 – 2 Nucleophilic attack of the amine on ^{+}NO, followed by loss of a proton, forms the *N*-nitrosamine.


Problem 25.29 Draw the product formed when each compound is treated with $NaNO_2$ and HCl.




25.14 Substitution Reactions of Aryl Diazonium Salts

Aryl diazonium salts undergo two general reactions.

- **Substitution of N_2 by an atom or a group of atoms Z forms a variety of substituted benzenes.**




- **Coupling of a diazonium salt with another benzene derivative forms an azo compound, a compound containing a nitrogen–nitrogen double bond.**




Y = NH_2, NHR, NR_2, OH (a strong electron-donor group)

25.14A Specific Substitution Reactions

Aryl diazonium salts react with a variety of reagents to form products in which Z (an atom or group of atoms) replaces N_2, a very good leaving group. The mechanism of these reactions varies with the identity of Z, so we will concentrate on the products of the reactions, not the mechanisms.




[1] Substitution by OH—Synthesis of phenols




A diazonium salt reacts with H_2O to form a **phenol.**

[2] Substitution by Cl or Br—Synthesis of aryl chlorides and bromides




A diazonium salt reacts with copper(I) chloride or copper(I) bromide to form an **aryl chloride** or **aryl bromide,** respectively. This is called the **Sandmeyer reaction.** It provides an alternative to direct chlorination and bromination of an aromatic ring using Cl_2 or Br_2 and a Lewis acid catalyst.

[3] Substitution by F—Synthesis of aryl fluorides




A diazonium salt reacts with fluoroboric acid (HBF_4) to form an **aryl fluoride.** This is a useful reaction because aryl fluorides cannot be produced by direct fluorination with F_2 and a Lewis acid catalyst, because F_2 reacts too violently (Section 18.3).

[4] Substitution by I—Synthesis of aryl iodides




A diazonium salt reacts with sodium or potassium iodide to form an aryl iodide. This, too, is a useful reaction because aryl iodides cannot be produced by direct iodination with I_2 and a Lewis acid catalyst, because I_2 reacts too slowly (Section 18.3).

[5] Substitution by CN—Synthesis of benzonitriles




A diazonium salt reacts with copper(I) cyanide to form a benzonitrile. Because a cyano group can be hydrolyzed to a carboxylic acid, reduced to an amine or aldehyde, or converted to a ketone with organometallic reagents, this reaction provides easy access to a wide variety of benzene derivatives using chemistry described in Section 22.18.

[6] Substitution by H—Synthesis of benzene




A diazonium salt reacts with hypophosphorus acid (H_3PO_2) to form benzene. This reaction has limited utility because it reduces the functionality of the benzene ring by replacing N_2 with a hydrogen atom. Nonetheless, this reaction *is* useful in synthesizing compounds that have substitution patterns that are not available by other means.

For example, it is not possible to synthesize 1,3,5-tribromobenzene from benzene by direct bromination. Because Br is an ortho, para director, bromination with Br_2 and $FeBr_3$ will not add Br substituents meta to each other on the ring.




It is possible, however, to add three Br atoms meta to each other when aniline is the starting material. Because an NH_2 group is a very powerful ortho, para director, three Br atoms are introduced in a single step on halogenation (Section 18.10A). Then, the NH_2 group can be removed by diazotization and reaction with H_3PO_2.



The complete synthesis of 1,3,5-tribromobenzene from benzene is outlined in Figure 25.13.


Figure 25.13
The synthesis of 1,3,5-tribromobenzene from benzene



- Nitration followed by reduction forms aniline ($C_6H_5NH_2$) from benzene (Steps [1] and [2]).
- Bromination of aniline yields the tribromo derivative in Step [3].
- The NH_2 group is removed by a two-step process: diazotization with $NaNO_2$ and HCl (Step [4]), followed by substitution of the diazonium ion by H with H_3PO_2.


25.14B Using Diazonium Salts in Synthesis

Diazonium salts provide easy access to many different benzene derivatives. Keep in mind the following four-step sequence, because it will be used to synthesize many substituted benzenes.



Sample Problems 25.6 and 25.7 apply these principles to two different multistep syntheses.


Sample Problem 25.6 Synthesize *m*-chlorophenol from benzene.



Solution

Both OH and Cl are ortho, para directors, but they are located meta to each other. The OH group must be formed from a diazonium salt, which can be made from an NO_2 group by a stepwise method.


Retrosynthetic Analysis



Working backwards:


- [1] Form the OH group from NO_2 by a three-step procedure using a diazonium salt.
- [2] Introduce Cl meta to NO_2 by halogenation.
- [3] Add the NO_2 group by nitration.

Synthesis



- Nitration followed by chlorination meta to the NO_2 group forms the meta disubstituted benzene (Steps [1]–[2]).
- Reduction of the nitro group (Step [3]) followed by diazotization forms the diazonium salt in Step [4], which is then converted to the desired phenol by treatment with H_2O (Step [5]).


Sample Problem 25.7 Synthesize *p*-bromobenzaldehyde from benzene.



Solution

Because the two groups are located para to each other and Br is an ortho, para director, Br should be added to the ring first. To add the CHO group, recall that it can be formed from CN by reduction.


Retrosynthetic Analysis



Working backwards:


- [1] Form the CHO group by reduction of CN.
- [2] Prepare the CN group from an NO_2 group by a three-step sequence using a diazonium salt.
- [3] Introduce the NO_2 group by nitration, para to the Br atom.
- [4] Introduce Br by bromination with Br_2 and $FeBr_3$.


Synthesis





- Bromination followed by nitration forms a disubstituted benzene with two para substituents (Steps [1]–[2]), which can be separated from its undesired ortho isomer.
- Reduction of the NO_2 group (Step [3]) followed by diazotization forms the diazonium salt in Step [4], which is converted to a nitrile by reaction with CuCN (Step [5]).
- Reduction of the CN group with DIBAL-H (a mild reducing agent) forms the CHO group and completes the synthesis.

Problem 25.30 Draw the product formed in each reaction.


a. 


b. 


c. 


d. 

Problem 25.31 Devise a synthesis of each compound from benzene.

a. 


b. 

c. 

d. 


25.15 Coupling Reactions of Aryl Diazonium Salts

The second general reaction of diazonium salts is **coupling.** When a diazonium salt is treated with an aromatic compound that contains a strong electron-donor group, the two rings join together to form an **azo compound,** a compound with a nitrogen–nitrogen double bond.




Synthetic dyes are described in more detail in Section 25.16.

Azo compounds are highly conjugated, rendering them colored (Section 16.15). Many of these compounds, such as the azo compound "butter yellow," are synthetic dyes. Butter yellow was once used to color margarine.



This reaction is another example of **electrophilic aromatic substitution,** with the **diazonium salt acting as the electrophile.** Like all electrophilic substitutions (Section 18.2), the mechanism has two steps: addition of the electrophile (the diazonium ion) to form a resonance-stabilized carbocation, followed by deprotonation, as shown in Mechanism 25.4.


Mechanism 25.4 Azo Coupling




1 The diazonium ion reacts with the benzene ring to form a resonance-stabilized carbocation.
2 Loss of a proton regenerates the aromatic ring.

Because a diazonium salt is weakly electrophilic, the reaction occurs only when the benzene ring has **a strong electron-donor group Y, where Y = NH_2, NHR, NR_2, or OH.** Although these groups activate both the ortho and para positions, para substitution occurs unless the para position already has another substituent present.

To determine what starting materials are needed to synthesize a particular azo compound, always divide the molecule into two components: **one has a benzene ring with a diazonium ion, and one has a benzene ring with a very strong electron-donor group.**




Sample Problem 25.8 What starting materials are needed to synthesize the following azo compound?




Solution


Both benzene rings in methyl orange have a substituent, but only one group, $N(CH_3)_2$, is a strong electron donor. In determining the two starting materials, the **diazonium ion must be bonded to the ring that is *not* bonded to $N(CH_3)_2$.**



Problem 25.32 Draw the product formed when $C_6H_5N_2^+Cl^-$ reacts with each compound.



Problem 25.33 What starting materials are needed to synthesize each azo compound?




25.16 Application: Synthetic Dyes and Sulfa Drugs

Azo compounds have two important applications: as dyes and as sulfa drugs, the first synthetic antibiotics.

25.16A Natural and Synthetic Dyes


Until 1856, all dyes were natural in origin, obtained from plants, animals, or minerals. Three natural dyes known for centuries are **indigo, tyrian purple,** and **alizarin.**




The blue dye **indigo,** derived from the plant *Indigofera tinctoria,* has been used in India for thousands of years. Traders introduced it to the Mediterranean area and then to Europe. **Tyrian purple,** a natural dark purple dye obtained from the mucous gland of a Mediterranean snail of the genus *Murex,* was a symbol of royalty before the collapse of the Roman empire. **Alizarin,** a bright red dye obtained from madder root (*Rubia tinctorum*), a plant native to India and northeastern Asia, has been found in cloth entombed with Egyptian mummies.


A purple shawl dyed with Perkin's mauveine


Because all three of these dyes were derived from natural sources, they were difficult to obtain, making them expensive and available only to the privileged. This all changed when William Henry Perkin, an 18-year-old student with a makeshift home laboratory, serendipitously prepared a purple dye, which would later be called mauveine, during his failed attempt to synthesize the antimalarial drug quinine. Mauveine is a mixture of two compounds that differ in the presence of only one methyl group on one of the aromatic rings.






Perkin's discovery marked the beginning of the chemical industry. He patented the dye and went on to build a factory to commercially produce it on a large scale. This event began the surge of research in organic chemistry, not just in the synthesis of dyes, but in the production of perfumes, anesthetics, inks, and drugs as well. Perkin was a wealthy man when he retired at the age of 36 to devote the rest of his life to basic chemical research. The most prestigious award given by the American Chemical Society is named the Perkin Medal in his honor.


Many common synthetic dyes, such as alizarine yellow R, para red, and Congo red, are **azo compounds,** prepared by the diazonium coupling reaction described in Section 25.15.



alizarine yellow R



para red



Congo red


Although natural and synthetic dyes are quite varied in structure, **all of them are colored because they are highly conjugated.** A molecule with eight or more π bonds in conjugation absorbs light in the visible region of the electromagnetic spectrum (Section 16.15A), taking on the color from the visible spectrum that it does *not* absorb.

Problem 25.34 (a) What two components are needed to prepare para red by azo coupling? (b) What two components are needed to prepare alizarine yellow R?

25.16B Sulfa Drugs


Although they may seem quite unrelated, the synthesis of colored dyes led to the development of the first synthetic antibiotics. Much of the early effort in this field was done by the German chemist Paul Ehrlich, who worked with synthetic dyes and used them to stain tissues. This led him on a search for dyes that were lethal to bacteria without affecting other tissue cells, hoping that these dyes could treat bacterial infections. For many years this effort was unsuccessful.

Then, in 1935, Gerhard Domagk, a German physician working for a dye manufacturer, first used a synthetic dye as a drug to kill bacteria. His daughter had contracted a streptococcal infection, and as she neared death, he gave her **prontosil,** an azo dye that inhibited the growth of certain bacteria in mice. His daughter recovered, and the modern era of synthetic antibiotics was initiated. For his pioneering work, Domagk was awarded the Nobel Prize in Physiology or Medicine in 1939.




prontosil — sulfanilamide, active antibacterial agent

Prontosil and other sulfur-containing antibiotics are collectively called **sulfa drugs.** Prontosil is not the active agent itself. In cells, it is metabolized to **sulfanilamide,** the active drug. To understand how sulfanilamide functions as an antibacterial agent we must examine **folic acid,** which microorganisms synthesize from *p*-aminobenzoic acid.




p-aminobenzoic acid (PABA) — folic acid

Sulfanilamide and *p*-aminobenzoic acid are similar in size and shape and have related functional groups. Thus, when sulfanilamide is administered, bacteria attempt to use it in place of *p*-aminobenzoic acid to synthesize folic acid. Derailing folic acid synthesis means that the bacteria cannot grow and reproduce. Sulfanilamide only affects bacterial cells, though, because humans do not synthesize folic acid, and must obtain it from their diets.




sulfanilamide




p-aminobenzoic acid

Many other compounds of similar structure have been prepared and are still widely used as antibiotics. The structures of two other sulfa drugs are shown in Figure 25.14.

Figure 25.14
Two common sulfa drugs



sulfamethoxazole



sulfisoxazole

- Sulfamethoxazole is the sulfa drug in Bactrim, and sulfisoxazole is sold as Gantrisin. Both drugs are commonly used in the treatment of ear and urinary tract infections.

KEY CONCEPTS

Amines

General Facts

- Amines are organic nitrogen compounds having the general structure RNH_2, R_2NH, or R_3N, with a lone pair of electrons on N (25.1).
- Amines are named using the suffix *-amine* (25.3).
- All amines have polar C – N bonds. Primary (1°) and 2° amines have polar N – H bonds and are capable of intermolecular hydrogen bonding (25.4).
- The lone pair on N makes amines strong organic bases and nucleophiles (25.8).

Summary of Spectroscopic Absorptions (25.5)


Mass spectra	Molecular ion	Amines with an odd number of N atoms give an odd molecular ion.
IR absorptions	N – H	3300–3500 cm^{-1} (two peaks for RNH_2, one peak for R_2NH)
^{1}H NMR absorptions	NH	0.5–5 ppm (no splitting with adjacent protons)
	CH – N	2.3–3.0 ppm (deshielded C_{sp^3} – H)
^{13}C NMR absorption	C – N	30–50 ppm

Comparing the Basicity of Amines and Other Compounds (25.10)

- Alkylamines (RNH_2, R_2NH, and R_3N) are more basic than NH_3 because of the electron-donating R groups (25.10A).
- Alkylamines (RNH_2) are more basic than arylamines ($C_6H_5NH_2$), which have a delocalized lone pair from the N atom (25.10B).
- Arylamines with electron-donor groups are more basic than arylamines with electron-withdrawing groups (25.10B).
- Alkylamines (RNH_2) are more basic than amides ($RCONH_2$), which have a delocalized lone pair from the N atom (25.10C).
- Aromatic heterocycles with a localized electron pair on N are more basic than those with a delocalized lone pair from the N atom (25.10D).
- Alkylamines with a lone pair in an sp^3 hybrid orbital are more basic than those with a lone pair in an sp^2 hybrid orbital (25.10E).


Preparation of Amines (25.7)

[1] Direct nucleophilic substitution with NH_3 and amines (25.7A)



- The mechanism is S_N2.
- The reaction works best for CH_3X or RCH_2X.
- The reaction works best to prepare 1° amines and quaternary ammonium salts.


[2] Gabriel synthesis (25.7A)




- The mechanism is S_N2.
- The reaction works best for CH_3X or RCH_2X.
- Only 1° amines can be prepared.

[3] Reduction methods (25.7B)


a. From nitro compounds




b. From nitriles



c. From amides




[4] Reductive amination (25.7C)



- Reductive amination adds one alkyl group (from an aldehyde or ketone) to a nitrogen nucleophile.
- Primary (1°), 2°, and 3° amines can be prepared.

Reactions of Amines

[1] Reaction as a base (25.9)




[2] Nucleophilic addition to aldehydes and ketones (25.11)

With 1° amines: With 2° amines:



R = H or alkyl — imine — R = H or alkyl — enamine


[3] Nucleophilic substitution with acid chlorides and anhydrides (25.11)



Z = Cl or OCOR
R' = H or alkyl

1°, 2°, and 3° amides

[4] Hofmann elimination (25.12)




alkene

- The less substituted alkene is the major product.

[5] Reaction with nitrous acid (25.13)

With 1° amines: With 2° amines:




alkyl diazonium salt


N-nitrosamine

Reactions of Diazonium Salts

[1] Substitution reactions (25.14)




phenol — aryl chloride or aryl bromide (X = Cl or Br) — aryl fluoride



aryl iodide — benzonitrile — benzene

[2] Coupling to form azo compounds (25.15)




Y = NH_2, NHR, NR_2, OH (a strong electron-donor group)


azo compound

PROBLEMS


Problems Using Three-Dimensional Models

25.35 Give a systematic or common name for each compound.


a. 

b. 

25.36 Which compound is the stronger base?




A



B


25.37 Varenicline (trade name Chantix) is a drug used to help smokers quit their habit. (a) Which N atom in varenicline is most basic? Explain your choice. (b) What product is formed when varenicline is treated with HCl?





varenicline


Nomenclature


25.38 Give a systematic or common name for each compound.


a. 


b. 


c. 

d. 

e. 

f. 

g. 


h. 

25.39 Draw the structure that corresponds to each name.

a. *N*-isobutylcyclopentanamine
b. tri-*tert*-butylamine
c. *N*,*N*-diisopropylaniline
d. *N*-methylpyrrole
e. *N*-methylcyclopentanamine
f. 3-methylhexan-2-amine
g. 2-*sec*-butylpiperidine
h. (*S*)-heptan-2-amine


Chiral Compounds


25.40 How many stereogenic centers are present in the following tetraalkylammonium salt? Draw all possible stereoisomers.




Basicity


25.41 Rank the compounds in each group in order of increasing basicity.

a. 

b. 

25.42 Decide which N atom in each molecule is most basic and draw the product formed when each compound is treated with CH_3CO_2H. Zolpidem (trade name Ambien) is used to treat insomnia, whereas aripiprazole (trade name Abilify) is used to treat depression, schizophrenia, and bipolar disorders.

a. 
zolpidem


b. 
aripiprazole


25.43 Explain why pyrimidine is less basic than pyridine.


pyridine


pyrimidine


25.44 Rank the nitrogen atoms in each compound in order of increasing basicity. Isoniazid is a drug used to treat tuberculosis, whereas histamine (Section 25.6B) causes the runny nose and watery eyes associated with allergies.


a. 
isoniazid

b. 
histamine

25.45 Explain why *m*-nitroaniline is a stronger base than *p*-nitroaniline.

25.46 Explain the observed difference in the pK_a values of the conjugate acids of amines **A** and **B.**


A
pK_a = 5.2


B
pK_a = 7.29


25.47 Why is pyrrole more acidic than pyrrolidine?



pyrrole
pK_a = 23



pyrrolidine
pK_a = 44

Preparation of Amines


25.48 What amide(s) can be used to prepare each amine by reduction?


a. 


b. 

c. 


25.49 What carbonyl and nitrogen compounds are needed to make each compound by reductive amination? When more than one set of starting materials is possible, give all possible methods.


a. 


b. 


c. 

25.50 Draw the product of each reductive amination reaction.

a. 

c. 

b. 

d. 

Extraction


25.51 How would you separate toluene ($C_6H_5CH_3$), benzoic acid ($C_6H_5CO_2H$), and aniline ($C_6H_5NH_2$) by an extraction procedure?


Reactions


25.52 Draw the products formed when *p*-methylaniline (*p*-$CH_3C_6H_4NH_2$) is treated with each reagent.


a. HCl
b. CH_3COCl
c. $(CH_3CO)_2O$
d. excess CH_3I
e. $(CH_3)_2C{=}O$
f. CH_3COCl, $AlCl_3$
g. CH_3CO_2H
h. $NaNO_2$, HCl
i. Part (b), then CH_3COCl, $AlCl_3$
j. CH_3CHO, $NaBH_3CN$

25.53 Draw the products formed when each amine is treated with [1] CH_3I (excess); [2] Ag_2O; [3] Δ. Indicate the major product when a mixture results.


a. 

b. 

c. 

d. 


25.54 Answer the following questions about benzphetamine, a habit-forming diet pill sold under the trade name Didrex.




benzphetamine

a. Label the stereogenic center(s).
b. What amide(s) can be reduced to form benzphetamine?
c. What carbonyl compound and amine can be used to make benzphetamine by reductive amination? Draw all possible methods.
d. What products are formed by Hofmann elimination from benzphetamine? Label the major product.


25.55 Draw the organic products formed in each reaction.




25.56 What is the major Hofmann elimination product formed from each amine?




25.57 Identify **A, B,** and **C,** three intermediates in the synthesis of the pain reliever and anesthetic fentanyl.



25.58 Identify the intermediates (**A**–**C**) in the following reaction sequence, which was used to prepare racemic ofloxacin. One enantiomer of the product, levofloxacin, is an antibiotic used to treat severe bacterial infections that have not responded to other drugs.



25.59 Draw the product formed when **A** is treated with each series of reagents.




A


a. [1] H_2O; [2] NaH; [3] CH_3Br
b. [1] CuCN; [2] DIBAL-H; [3] H_2O
c. [1] $C_6H_5NH_2$; [2] CH_3COCl

25.60 A chiral amine **A** having the *R* configuration undergoes Hofmann elimination to form an alkene **B** as the major product. **B** is oxidatively cleaved with ozone, followed by CH_3SCH_3, to form $CH_2=O$ and $CH_3CH_2CH_2CHO$. What are the structures of **A** and **B?**


Mechanism

25.61 Draw a stepwise mechanism for each reaction.


a. 

b. 


25.62 Draw a stepwise mechanism for the following reaction.



25.63 Alkyl diazonium salts decompose to form carbocations, which go on to form products of substitution, elimination, and (sometimes) rearrangement. Keeping this in mind, draw a stepwise mechanism that forms all of the following products.





25.64 Tertiary (3°) aromatic amines react with $NaNO_2$ and HCl to afford products of electrophilic aromatic substitution. Draw a stepwise mechanism for this nitrosation reaction and explain why it occurs only on benzene rings with strong ortho, para activating groups.





Synthesis


25.65 Devise a synthesis of each compound from benzene. You may use any other organic or inorganic reagents.


a. 

b. 


c. 


d. 


e. 

f. 

25.66 Devise a synthesis of each compound from aniline ($C_6H_5NH_2$) as starting material.

a. 

b. 

c. 


25.67 Devise at least three different methods to prepare *N*-methylbenzylamine ($PhCH_2NHCH_3$) from benzene, any one-carbon organic compounds, and any required reagents.


25.68 Safrole, which is isolated from sassafras (Problem 21.33), can be converted to the illegal stimulant MDMA (3,4-methylenedioxymethamphetamine, "Ecstasy") by a variety of methods. (a) Devise a synthesis that begins with safrole and uses a nucleophilic substitution reaction to introduce the amine. (b) Devise a synthesis that begins with safrole and uses reductive amination to introduce the amine.




MDMA safrole


25.69 Synthesize each compound from benzene. Use a diazonium salt as one of the synthetic intermediates.

a. 


b. 

c. 

25.70 Devise a synthesis of each biologically active compound from benzene.


a. 


acetaminophen
(analgesic)


b. 

pseudoephedrine
(nasal decongestant)

25.71 Devise a synthesis of each compound from benzene, any organic alcohols having four carbons or fewer, and any required reagents.

a. 


b. 

c. 


Spectroscopy

25.72 Three isomeric compounds, **A, B,** and **C,** all have molecular formula $C_8H_{11}N$. The ^1H NMR and IR spectral data of **A, B,** and **C** are given below. What are their structures?


Compound **A:** IR peak at 3400 cm^{-1}




Compound **B:** IR peak at 3310 cm^{-1}




Compound **C:** IR peaks at 3430 and 3350 cm^{-1}




25.73 Treatment of compound **D** with $LiAlH_4$ followed by H_2O forms compound **E. D** shows a molecular ion in its mass spectrum at *m/z* = 71 and IR absorptions at 3600–3200 and 2263 cm^{-1}. **E** shows a molecular ion in its mass spectrum at *m/z* = 75 and IR absorptions at 3636 and 3600–3200 cm^{-1}. Propose structures for **D** and **E** from these data and the given ^1H NMR spectra.






Challenge Problems

25.74 The pK_a of the conjugate acid of guanidine is 13.6, making it one of the strongest neutral organic bases. Offer an explanation.




guanidine $pK_a = 13.6$

25.75 Rank the following compounds in order of increasing basicity and explain the order you chose.




pyrrole imidazole thiazole

25.76 Draw the product **Y** of the following reaction sequence. **Y** was an intermediate in the remarkable synthesis of cyclooctatetraene by Wilstatter in 1911.




25.77 Devise a synthesis of each compound from the given starting material(s). Albuterol is a bronchodilator and proparacaine is a local anesthetic.

a.




albuterol

b.



proparacaine

25.78 Heating compound **X** with aqueous formaldehyde forms **Y** ($C_{17}H_{23}NO$), which has been converted to a mixture of lupinine and epilupinine, alkaloids isolated from lupin, a perennial ornamental plant commonly seen on the roadside in parts of Alaska (Section 25.10). Identify **Y** and explain how it is formed.



lupinine epilupinine

Carbon–Carbon Bond-Forming Reactions in Organic Synthesis

26

26.1 Coupling reactions of organocuprate reagents
26.2 Suzuki reaction
26.3 Heck reaction
26.4 Carbenes and cyclopropane synthesis
26.5 Simmons–Smith reaction
26.6 Metathesis



Ingenol mebutate is an ester derived from ingenol, a natural product obtained from the sap of *Euphorbia peplus*, a type of milkweed native to Europe, northern Africa, and western Asia. Because ingenol derivatives exhibited useful biological activity and isolation from the natural source did not provide easy access to the material, scientists developed an efficient laboratory synthesis. A gel formulation of ingenol mebutate (trade name Picato) has been approved for the treatment of actinic keratosis, a skin condition resulting from over-exposure to the sun that may result in squamous cell carcinoma, a form of skin cancer. In Chapter 26 we learn about carbon–carbon bond-forming reactions that prepare complex compounds like ingenol.

To form the carbon skeletons of complex molecules, organic chemists need an extensive repertoire of carbon–carbon bond-forming reactions. In Chapter 20, for example, we learned about the reactions of organometallic reagents—organolithium reagents, Grignard reagents, and organocuprates—with carbonyl substrates. In Chapters 23 and 24, we studied the reactions of nucleophilic enolates that form new carbon–carbon bonds.

Chapter 26 presents more carbon–carbon bond-forming reactions that are especially useful tools in organic synthesis. While previous chapters have concentrated on the reactions of one or two functional groups, the reactions in this chapter utilize a variety of starting materials and conceptually different reactions that form many types of products. All follow one central theme—they form new carbon–carbon bonds under mild conditions, making them versatile synthetic methods.


26.1 Coupling Reactions of Organocuprate Reagents

Several carbon–carbon bond-forming reactions involve the coupling of an organic halide (R'X) with an organometallic reagent or alkene. Three useful reactions are discussed in Sections 26.1–26.3:

[1] Reaction of an organic halide with an organocuprate reagent (Section 26.1)




[2] Suzuki reaction: Reaction of an organic halide with an organoboron reagent in the presence of a palladium catalyst (Section 26.2)



[3] Heck reaction: Reaction of an organic halide with an alkene in the presence of a palladium catalyst (Section 26.3)

A complete list of reactions that form C−C bonds appears in Appendix D.




26.1A General Features of Organocuprate Coupling Reactions

In addition to their reactions with acid chlorides, epoxides, and α,β-unsaturated carbonyl compounds (Sections 20.13–20.15), organocuprate reagents (R_2CuLi) also react with organic halides R'−X to form coupling products R−R' that contain a new C−C bond. Only one R group of the organocuprate is transferred to form the product, while the other becomes part of RCu, a reaction by-product.




A variety of organic halides can be used, including methyl and 1° alkyl halides, as well as vinyl and aryl halides that contain X bonded to an sp^2 hybridized carbon. Some cyclic 2° alkyl halides

give reasonable yields of product, but 3° alkyl halides are too sterically hindered. The halogen X in R'X may be Cl, Br, or I.




Coupling reactions with vinyl halides are **stereospecific.** For example, reaction of *trans*-1-bromohex-1-ene with $(CH_3)_2CuLi$ forms *trans*-hept-2-ene as the only stereoisomer (Equation [3]).

Problem 26.1 Draw the product of each coupling reaction.




Problem 26.2 Identify reagents **A** and **B** in the following reaction scheme. This synthetic sequence was used to prepare the C_{18} juvenile hormone (Figure 20.6).




26.1B Using Organocuprate Couplings to Synthesize Hydrocarbons

Since organocuprate reagents (R_2CuLi) are prepared in two steps from alkyl halides (RX), this method ultimately converts two organic halides (RX and R'X) into a hydrocarbon R−R' with a new carbon–carbon bond. A hydrocarbon can often be made by two different routes, as shown in Sample Problem 26.1.



Two organic halides are needed as starting materials.

Sample Problem 26.1 Devise a synthesis of 1-methylcyclohexene from 1-bromocyclohexene and CH_3I.




1-methylcyclohexene 1-bromocyclohexene


Solution

In this example, either halide can be used to form an organocuprate, which can then be coupled with the second halide.

Possibility [1]




Possibility [2]




Problem 26.3 Synthesize each product from the given starting materials using an organocuprate coupling reaction.


a.



b.



c.



The mechanism of this reaction may vary with the identity of R' in R'−X. Since coupling occurs with organic halides having the halogen X on either an sp^3 or sp^2 hybridized carbon, an S_N2 mechanism cannot explain all the observed results.


26.2 Suzuki Reaction

The **Suzuki reaction** is the first of two reactions that utilize a palladium catalyst and proceed by way of an intermediate organopalladium compound. The second is the Heck reaction (Section 26.3).

26.2A General Features of Reactions with Pd Catalysts

Reactions with palladium compounds share many common features with reactions involving other transition metals. During a reaction, palladium is coordinated to a variety of groups called **ligands,** which donate electron density to (or sometimes withdraw electron density from)

the metal. A common electron-donating ligand is a **phosphine,** such as triphenylphosphine, tri(*o*-tolyl)phosphine, or tricyclohexylphosphine.



PPh_3
triphenylphosphine

$P(o\text{-tolyl})_3$
tri(*o*-tolyl)phosphine
abbreviated as **PAr_3**
Ar = an aryl group

PCy_3
tricyclohexylphosphine


A general ligand bonded to a metal is often designated as **L.** Pd bonded to four ligands is denoted as PdL_4.

Organopalladium compounds—compounds that contain a carbon–palladium bond—are generally prepared in situ during the course of a reaction, from another palladium reagent such as $Pd(OAc)_2$ or $Pd(PPh_3)_4$. In most useful reactions only a catalytic amount of palladium reagent is utilized.


Ac is the abbreviation for an acetyl group, **$CH_3C{=}O$,** so OAc (or $^-$OAc) is the abbreviation for acetate, **$CH_3CO_2^-$.**

Two common processes, called **oxidative addition** and **reductive elimination,** dominate many reactions of palladium compounds.

- ***Oxidative addition*** **is the addition of a reagent (such as RX) to a metal, often increasing the number of groups around the metal by two.**




- ***Reductive elimination*** **is the elimination of two groups that surround the metal, often forming new C–H or C–C bonds.**




Reaction mechanisms with palladium compounds are often multistep. During the course of a reaction, the identity of some groups bonded to Pd will be known with certainty, while the identity of other ligands might not be known. Consequently, only the crucial reacting groups around a metal are usually drawn and the other ligands are not specified.


26.2B Details of the Suzuki Reaction


The **Suzuki reaction** is a palladium-catalyzed coupling of an organic halide (R'X) with an organoborane (RBY_2) to form a product (R–R') with a new C–C bond. $Pd(PPh_3)_4$ is the typical palladium catalyst, and the reaction is carried out in the presence of a base such as NaOH or $NaOCH_2CH_3$.



Vinyl halides and aryl halides, both of which contain a halogen X bonded directly to an sp^2 hybridized carbon, are most often used, and the halogen is usually Br or I. The Suzuki reaction is completely **stereospecific,** as shown in Example [3]; a cis vinyl halide and a trans vinylborane form a *cis,trans*-1,3-diene.

[1] 

[2] 


[3] 

cis vinyl bromide + trans vinylborane

new C–C bonds in red

The organoboranes used in the Suzuki reaction are prepared from two sources.


- **Vinylboranes,** which have a boron atom bonded to a carbon–carbon double bond, are prepared by hydroboration of an alkyne using catecholborane, a commercially available reagent. **Hydroboration adds the elements of H and B in a syn fashion to form a trans vinylborane.** With terminal alkynes, hydroboration always places the boron atom on the *less substituted* terminal carbon.



catecholborane — trans vinylborane


syn addition of H and B


- **Arylboranes,** which have a boron atom bonded to a benzene ring, are prepared from organolithium reagents by reaction with trimethyl borate [$B(OCH_3)_3$].





trimethyl borate — arylborane

Problem 26.4 Draw the product of each reaction.

a. 

b. 


c. 

d. 



The structure of bombykol, the sex pheromone of the female silkworm moth *Bombyx mori,* was elucidated in 1959 using 6.4 mg of material obtained from 500,000 silkworm moths.


Problem 26.5 One step in the synthesis of the nonsteroidal anti-inflammatory drug rofecoxib (trade name Vioxx) involves Suzuki coupling of **A** and **B.** What product is formed in this reaction?



The mechanism of the Suzuki reaction consists of oxidative addition of R'−X to the palladium catalyst, transfer of an alkyl group from the organoborane to palladium, and reductive elimination of R−R', forming a new carbon–carbon bond. A general halide R'−X and organoborane $R-BY_2$ are used to illustrate this process in Mechanism 26.1. Since the palladium reagent is regenerated during reductive elimination, only a catalytic amount of palladium is needed.

The Suzuki reaction was a key step in the synthesis of **bombykol,** the sex pheromone of the female silkworm moth, and **humulene,** a lipid isolated from hops, as shown in Figure 26.1. The synthesis of humulene illustrates that an intramolecular Suzuki reaction can form a ring. Sample Problem 26.2 shows how a conjugated diene can be prepared from an alkyne and vinyl halide using a Suzuki reaction.

Mechanism 26.1 Suzuki Reaction




1 – 2 Loss of two triphenylphosphine ligands followed by oxidative addition of R'X forms an organopalladium reagent.


3 Reaction of the organoborane RBY_2 with ^-OH forms a nucleophilic boron intermediate that transfers an alkyl group from boron to palladium.

4 Reductive elimination of R'–R forms a new carbon-carbon bond, and the palladium catalyst $Pd(PPh_3)_2$ is regenerated.

Figure 26.1 Synthesis of two natural products using the Suzuki reaction




Sample Problem 26.2 Devise a synthesis of (1*Z*,3*E*)-1-phenylocta-1,3-diene from hex-1-yne and (*Z*)-2-bromostyrene using a Suzuki coupling.



(1*Z*,3*E*)-1-phenylocta-1,3-diene (*Z*)-2-bromostyrene

Solution

This synthesis can be accomplished in two steps. Hydroboration of hex-1-yne with catecholborane forms a vinylborane. Coupling of this vinylborane with (*Z*)-2-bromostyrene gives the desired 1,3-diene. The *E* configuration of the vinylborane and the *Z* configuration of the vinyl bromide are both retained in the product.




syn addition of H and B

(1*Z*,3*E*)-1-phenylocta-1,3-diene


new C–C bond in red

Problem 26.6 Synthesize each compound from the given starting materials.


a.



b.




c.



26.3 Heck Reaction


Richard Heck and Akira Suzuki won the 2010 Nobel Prize in Chemistry for the discovery of the carbon–carbon bond-forming reactions detailed in Sections 26.2 and 26.3.

The **Heck reaction is a palladium-catalyzed coupling of a vinyl or aryl halide with an alkene to form a more highly substituted alkene with a new C−C bond.** Palladium(II) acetate [$Pd(OAc)_2$] in the presence of a triarylphosphine [P(*o*-tolyl)$_3$] is the typical catalyst, and the reaction is carried out in the presence of a base such as triethylamine (Et_3N). The Heck reaction is a **substitution reaction** in which one H atom of the alkene starting material is replaced by the R' group of the vinyl or aryl halide.




R' = vinyl or aryl
X = Br or I


The alkene component is typically ethylene or a monosubstituted alkene (CH_2=CHZ), and the halogen X is usually Br or I. When Z = Ph, COOR, or CN in a monosubstituted alkene, **the new C−C bond is formed on the *less* substituted carbon to afford a trans alkene.** When a vinyl halide is used as the organic halide, the reaction is **stereospecific,** as shown in Example [3]; the trans stereochemistry of the vinyl iodide is retained in the product.




Problem 26.7 Draw the coupling product formed when each pair of compounds is treated with $Pd(OAc)_2$, $P(o\text{-tolyl})_3$, and Et_3N.



To use the Heck reaction in synthesis, you must determine what alkene and what organic halide are needed to prepare a given compound. To work backwards, locate the double bond with the aryl, COOR, or CN substituent, and break the molecule into two components at the end of the C=C not bonded to one of these substituents. Sample Problem 26.3 illustrates this retrosynthetic analysis.




Sample Problem 26.3 What starting materials are needed to prepare each alkene using a Heck reaction?




Solution

To prepare an alkene of general formula R'CH=CHZ by the Heck reaction, two starting materials are needed—an alkene (CH_2=CHZ) and a vinyl or aryl halide (R'X).




Problem 26.8 What starting materials are needed to prepare each compound using a Heck reaction?



The actual palladium catalyst in the Heck reaction is thought to contain a palladium atom bonded to two tri(*o*-tolyl)phosphine ligands, abbreviated as $Pd(PAr_3)_2$. In this way it resembles the divalent palladium catalyst used in the Suzuki reaction. The mechanism of the Heck reaction consists of oxidative addition of the halide R'X to the palladium catalyst, addition of the resulting organopalladium reagent to the alkene, and two successive eliminations. A general organic halide R'X and alkene CH_2=CHZ are used to illustrate the process in Mechanism 26.2.


Mechanism 26.2 Heck Reaction



1. Oxidative addition of R'X forms an organopalladium reagent.
2. Addition of R' and Pd to the π bond of CH_2=CHZ places the Pd on the carbon with the Z substituent.
3. Elimination of H and Pd forms the π bond in the reaction product and transfers a hydrogen to Pd.
4. Reductive elimination of HX regenerates the palladium catalyst $Pd(PAr_3)_2$.


26.4 Carbenes and Cyclopropane Synthesis

Another method of carbon–carbon bond formation involves the conversion of alkenes to cyclopropane rings using **carbene** intermediates.






Pyrethrin I and **decamethrin** both contain cyclopropane rings. Pyrethrin I is a naturally occurring biodegradable insecticide obtained from chrysanthemums, whereas **decamethrin** is a more potent synthetic analogue that is widely used as an insecticide in agriculture.




pyrethrin I



decamethrin

26.4A Carbenes

A *carbene*, $R_2C{:}$, is a neutral reactive intermediate that contains a divalent carbon surrounded by six electrons—the lone pair and two each from the two R groups. These three groups make the carbene carbon ***sp*² hybridized,** with a vacant *p* orbital extending above and below the plane containing the C and the two R groups. The lone pair of electrons occupies an sp^2 hybrid orbital.




The carbene carbon is sp^2 hybridized.

Carbenes share two features in common with carbocations and carbon radicals.

- A carbene is highly reactive because carbon does not have an octet of electrons.
- A carbene is electron deficient, so it behaves as an electrophile.

26.4B Preparation and Reactions of Dihalocarbenes


Dihalocarbenes, $:CX_2$, are especially useful reactive intermediates because they are readily prepared from trihalomethanes (CHX_3) by reaction with a strong base. Treatment of chloroform, $CHCl_3$, with $KOC(CH_3)_3$ forms dichlorocarbene, $:CCl_2$.



chloroform — dichlorocarbene

Dichlorocarbene is formed by a two-step process that results in the elimination of the elements of H and Cl from the *same* carbon, as shown in Mechanism 26.3. Loss of two elements from the same carbon is called **α elimination,** to distinguish it from the β eliminations discussed in Chapter 8, in which two elements are lost from *adjacent* carbons.


Mechanism 26.3 Formation of Dichlorocarbene



1 Three electronegative Cl atoms acidify the C–H of $CHCl_3$, so it can be removed by strong base to form a carbanion.


2 Elimination of Cl^- forms the carbene.

Since dihalocarbenes are electrophiles, they readily react with double bonds to afford cyclopropanes, forming two new carbon–carbon bonds.



Cyclopropanation is a concerted reaction, so both C−C bonds are formed in a single step, as shown in Mechanism 26.4.

Mechanism 26.4 Addition of Dichlorocarbene to an Alkene




Carbene addition occurs in a **syn** fashion from either side of the planar double bond. The relative position of substituents in the alkene reactant is retained in the cyclopropane product. **Carbene addition is thus a stereospecific reaction,** because cis and trans alkenes yield different stereoisomers as products, as illustrated in Sample Problem 26.4.

Sample Problem 26.4 Draw the products formed when *cis*- and *trans*-but-2-ene are treated with $CHCl_3$ and $KOC(CH_3)_3$.


Solution

To draw each product, add the carbene carbon from either side of the alkene, and keep all substituents in their original orientations. The **cis** methyl groups in *cis*-but-2-ene become **cis** substituents in the cyclopropane. Addition from either side of the alkene yields the same compound—**an achiral meso compound that contains two stereogenic centers**—labeled in blue.




Products are identical, an achiral meso compound.

The **trans** methyl groups in *trans*-but-2-ene become **trans** substituents in the cyclopropane. Addition from either side of the alkene yields an equal amount of two enantiomers—**a racemic mixture.**




Products are enantiomers.

Problem 26.9 Draw all stereoisomers formed when each alkene is treated with $CHCl_3$ and $KOC(CH_3)_3$.

a.  b.  c. 

Finally, dihalo cyclopropanes can be converted to dialkyl cyclopropanes by reaction with organocuprates (Section 26.1). For example, cyclohexene can be converted to a bicyclic product having four new C−C bonds by the following two-step sequence: cyclopropanation with dibromocarbene ($:CBr_2$) and reaction with lithium dimethylcuprate, $LiCu(CH_3)_2$.




two new C–C bonds in red two new C–C bonds in blue

Problem 26.10 What reagents are needed to convert 2-methylpropene [$(CH_3)_2C{=}CH_2$] to each compound? More than one step may be required.

a.  b.  c. 


26.5 Simmons–Smith Reaction


Although the reaction of dihalocarbenes with alkenes gives good yields of halogenated cyclopropanes, this is not usually the case with **methylene, $:CH_2$,** the simplest carbene. Methylene is readily formed by heating diazomethane, CH_2N_2, which decomposes and loses N_2, but the reaction of $:CH_2$ with alkenes often affords a complex mixture of products. Thus, this reaction cannot be reliably used for cyclopropane synthesis.



diazomethane methylene


Nonhalogenated cyclopropanes can be prepared by the reaction of an alkene with diiodomethane, CH_2I_2, in the presence of a copper-activated zinc reagent called zinc–copper couple [Zn(Cu)]. This process, the **Simmons–Smith reaction,** is named for H. E. Simmons and R. D. Smith, DuPont chemists who discovered the reaction in 1959.






The Simmons–Smith reaction does not involve a free carbene. Rather, the reaction of CH_2I_2 with Zn(Cu) forms (iodomethyl)zinc iodide, which transfers a CH_2 group to an alkene, as shown in Mechanism 26.5.

Mechanism 26.5 Simmons–Smith Reaction




1. Reaction of CH_2I_2 with zinc–copper couple forms ICH_2ZnI [(iodomethyl)zinc iodide], the **Simmons–Smith reagent.** This intermediate is called a *carbenoid,* because the CH_2 does not exist as a free carbene.
2. The Simmons–Smith reagent transfers a CH_2 to an alkene, forming two new C–C bonds.

The Simmons–Smith reaction is stereospecific. The relative position of substituents in the alkene reactant is *retained* in the cyclopropane product, as shown for the conversion of *cis*-hex-3-ene to *cis*-1,2-diethylcyclopropane.



cis-hex-3-ene → *cis*-1,2-diethylcyclopropane

Problem 26.11 What product is formed when each alkene is treated with CH_2I_2 and Zn(Cu)?




Problem 26.12 What stereoisomers are formed when *trans*-hex-3-ene is treated with CH_2I_2 and Zn(Cu)?

26.6 Metathesis


Recall from Section 10.1 that **olefin** is another name for an **alkene.**

Alkene metathesis, more commonly called **olefin metathesis,** is a reaction between two alkene molecules that results in the interchange of the carbons of their double bonds. Two σ and two π bonds are broken and two new σ and two new π bonds are formed.



The word *metathesis* is derived from the Greek words *meta* (change) and *thesis* (position). The 2005 Nobel Prize in Chemistry was awarded to Robert Grubbs of the California Institute of Technology, Yves Chauvin of the Institut Français du Pétrole, and Richard Schrock of the Massachusetts Institute of Technology for their work on olefin metathesis.


Olefin metathesis occurs in the presence of a complex transition metal catalyst that contains a carbon–metal double bond. The metal is typically ruthenium (Ru), tungsten (W), or molybdenum (Mo). In a widely used catalyst, called **Grubbs catalyst,** the metal is Ru.



Grubbs catalyst

Olefin metathesis is an equilibrium process and, with many alkene substrates, a mixture of starting material and two or more alkene products is present at equilibrium, making the reaction useless for preparative purposes. With terminal alkenes, however, one metathesis product is $CH_2{=}CH_2$ (a gas), which escapes from the reaction mixture and drives the equilibrium to the right.

As a result, monosubstituted alkenes ($RCH{=}CH_2$) and 2,2-disubstituted alkenes ($R_2C{=}CH_2$) are excellent metathesis substrates because high yields of a single alkene product are obtained, as shown in Equations [1] and [2].



To draw the products of any metathesis reaction:

[1] Arrange two molecules of the starting alkene adjacent to each other as in Figure 26.2 where styrene ($PhCH{=}CH_2$) is used as the starting material.

[2] Then, break the double bonds in the starting material and form two new double bonds using carbon atoms that were *not* previously bonded to each other in the starting alkenes.

There are always two ways to arrange the starting alkenes (Pathways [1] and [2] in Figure 26.2). In this example, the two products of the reaction, $PhCH{=}CHPh$ and $CH_2{=}CH_2$ are formed in the first reaction pathway (Pathway [1]), while starting material is re-formed in the second pathway (Pathway [2]). Whenever the starting alkene is regenerated, it can go on to form product when the catalytic cycle is repeated.

Problem 26.13 Draw the products formed when each alkene is treated with Grubbs catalyst.






Figure 26.2
Drawing the products of olefin metathesis using styrene ($PhCH{=}CH_2$) as starting material



- Overall reaction: $2\ PhCH{=}CH_2 \rightarrow PhCH{=}CHPh + CH_2{=}CH_2$.
- There are always two ways to join the C's of a single alkene together to form metathesis products (Pathways [1] and [2]).
- When like C's of the alkene substrate are joined in the first reaction (Pathway [1]), $PhCH{=}CHPh$ (in a cis and trans mixture) and $CH_2{=}CH_2$ are formed. Because $CH_2{=}CH_2$ escapes as a gas from the reaction mixture, only $PhCH{=}CHPh$ is isolated as product.
- When unlike C's of $PhCH{=}CH_2$ are joined in the second reaction (Pathway [2]), starting material is formed, which can re-enter the catalytic cycle to form product by the first pathway.
- In this way, a single constitutional isomer, $PhCH{=}CHPh$, is isolated.


Problem 26.14 What products are formed when *cis*-pent-2-ene undergoes metathesis? Use this reaction to explain why metathesis of a 1,2-disubstituted alkene (RCH=CHR') is generally not a practical method for alkene synthesis.

The mechanism for olefin metathesis is complex, and involves **metal–carbene intermediates—intermediates that contain a metal–carbon double bond.** The mechanism is drawn for the reaction of a terminal alkene ($RCH{=}CH_2$) with Grubbs catalyst, abbreviated as **Ru=CHPh,** to form $RCH{=}CHR$ and $CH_2{=}CH_2$. To begin metathesis, Grubbs catalyst reacts with the alkene substrate to form two new metal–carbenes **A** and **B** by a two-step process: addition of Ru=CHPh to the alkene to yield two different metallocyclobutanes (Step [1]), followed by elimination to form **A** and **B** (Steps [2a] and [2b]). The alkene by-products formed in this process (RCH=CHPh and $PhCH{=}CH_2$) are present in only a small amount since Grubbs reagent is used catalytically.



Each of these metal–carbene intermediates **A** and **B** then reacts with more starting alkene to form metathesis products, as shown in Mechanism 26.6. This mechanism is often written in a circle to emphasize the catalytic cycle. The mechanism demonstrates how two molecules of $RCH{=}CH_2$ are converted to $RCH{=}CHR$ and $CH_2{=}CH_2$. The mechanism can be written beginning with reagent **A** or **B,** and all steps are equilibria.


Mechanism 26.6 Olefin Metathesis: 2 $RCH{=}CH_2 \rightarrow RCH{=}CHR + CH_2{=}CH_2$




1. Reaction of $Ru{=}CH_2$ (**A**) with $RCH{=}CH_2$ forms a metallocyclobutane. Ru can bond to either the more or less substituted end of the alkene, but product is formed only when Ru bonds to the *more* substituted end, as shown.
2. Elimination forms one metathesis product, $CH_2{=}CH_2$, and metal–carbene complex **B.**
3. Reaction of **B** with $RCH{=}CH_2$ forms a metallocyclobutane. Ru can bond to either the more or less substituted end of the alkene, but product is formed only when Ru bonds to the *less* substituted end, as shown.
4. Elimination forms the other metathesis product, RCH=CHR, and metal–carbene complex **A.** The catalyst is regenerated and the cycle begins again.


A metathesis reaction that forms a ring is called **ring-closing metathesis (RCM).**

When a diene is used as starting material, ring closure occurs.




These reactions are typically run in very dilute solution, so that the two reactive ends of the *same* molecule have a higher probability of finding each other for reaction than two functional groups in *different* molecules. These high-dilution conditions thus favor ***intra*molecular** rather than *inter*molecular metathesis. Two examples are shown.





new C=C in red




new C=C in red

Because metathesis catalysts are compatible with the presence of many functional groups (such as OH, OR, and C=O) and because virtually any ring size can be prepared, metathesis has been used to prepare many complex natural products (Figure 26.3).


Figure 26.3

Ring-closing metathesis in the synthesis of epothilone A and Sch38516



(*E* and *Z* isomers formed)


epothilone A
anticancer drug




Sch38516
antiviral agent

- **Epothilone A,** a promising anticancer agent, was first isolated from soil bacteria collected from the banks of the Zambezi River in South Africa.
- **Sch38516** is an antiviral agent active against influenza A.
- The new C–C bonds formed during metathesis are indicated in red. In both metathesis reactions, $CH_2=CH_2$ is also formed.


Problem 26.15 Draw the product formed from ring-closing metathesis of each compound.



Problem 26.16 What product is formed by ring-closing metathesis of compound **V,** a key intermediate in the synthesis of ingenol, a natural product mentioned in the chapter opener?



V




ingenol




Ingenol is isolated from the milky liquid obtained from *Euphorbia ingens,* a large cactus commonly called the candelabra tree, which is native to dry areas in southern Africa.

Sample Problem 26.5 What starting material is needed to synthesize each compound by a ring-closing metathesis reaction?





Solution

To work in the retrosynthetic direction, cleave the C=C in the product, and bond each carbon of the original alkene to a CH_2 group using a double bond.





The resulting compound has a carbon chain with two terminal alkenes.






Problem 26.17 What starting material is needed to synthesize each compound by a ring-closing metathesis reaction?

a. 

b. 

c. 

KEY CONCEPTS

Carbon–Carbon Bond-Forming Reactions in Organic Synthesis


Coupling Reactions

[1] Coupling reactions of organocuprate reagents (26.1)




- R'X can be CH_3X, RCH_2X, 2° cyclic halides, vinyl halides, and aryl halides.
- X may be Cl, Br, or I.
- With vinyl halides, coupling is stereospecific.

[2] Suzuki reaction (26.2)



- R'X is most often a vinyl halide or aryl halide.
- With vinyl halides, coupling is stereospecific.


[3] Heck reaction (26.3)



- R'X is a vinyl halide or aryl halide.
- Z = H, Ph, COOR, or CN.
- With vinyl halides, coupling is stereospecific.
- The reaction forms trans alkenes.


Cyclopropane Synthesis

[1] Addition of dihalocarbenes to alkenes (26.4)



- The reaction occurs with syn addition.
- The position of substituents in the alkene is retained in the cyclopropane.


[2] Simmons–Smith reaction (26.5)



- The reaction occurs with syn addition.
- The position of substituents in the alkene is retained in the cyclopropane.


Metathesis (26.6)

[1] Intermolecular reaction



- Metathesis works best when $CH_2{=}CH_2$, a gas that escapes from the reaction mixture, is one of the products formed.

[2] Intramolecular reaction





- Ring-closing metathesis forms rings of any size from diene starting materials.

PROBLEMS


Problems Using Three-Dimensional Models


26.18 In addition to organic halides, alkyl tosylates (R'OTs, Section 9.13) also react with organocuprates (R_2CuLi) to form coupling products R–R'. When 2° alkyl tosylates are used as starting materials (R_2CHOTs), inversion of the configuration at a stereogenic center results. Keeping this in mind, draw the product formed when each compound is treated with $(CH_3)_2CuLi$.

a. 

b. 


26.19 What product is formed by ring-closing metathesis of each compound?


a. 


b. 


Coupling Reactions


26.20 Draw the products formed in each reaction.


a. 


b. 


c. 

d. 


e. 


g. 


f. 

h. 


26.21 What organic halide is needed to convert lithium divinylcuprate [$(CH_2{=}CH)_2CuLi$] to each compound?


a. 


b. 


c. 


26.22 How can you convert ethynylcyclohexane to dienes **A–C** using a Suzuki reaction? You may use any other organic compounds and inorganic reagents. Is it possible to synthesize diene **D** using a Suzuki reaction? Explain why or why not.


ethynylcyclohexane



A



C



B


D

26.23 What compound is needed to convert styrene ($C_6H_5CH{=}CH_2$) to each product using a Heck reaction?


a. 

b. 


c. 

26.24 What steps are needed to convert but-1-ene ($CH_3CH_2CH{=}CH_2$) to octane [$CH_3(CH_2)_6CH_3$] using a coupling reaction with an organocuprate reagent? All carbon atoms in octane must come from but-1-ene.


26.25 What product is formed in the Suzuki coupling of **A** and **B?** This reaction was a key step in the synthesis of losartan, a drug used to treat hypertension.



A




B




losartan

26.26 Identify **X,** an intermediate that was converted to eletriptan (trade name Relpax), a drug used to treat migraines.



Cyclopropanes


26.27 Draw the products (including stereoisomers) formed in each reaction.




26.28 Treatment of cyclohexene with $C_6H_5CHI_2$ and Zn(Cu) forms two stereoisomers of molecular formula $C_{13}H_{16}$. Draw their structures and explain why two compounds are formed.

Metathesis


26.29 What ring-closing metathesis product is formed when each substrate is treated with Grubbs catalyst under high-dilution conditions?




26.30 What starting material is needed to prepare each compound by a ring-closing metathesis reaction?



26.31 Metathesis reactions can be carried out with two *different* alkene substrates in one reaction mixture. Depending on the substitution pattern around the C=C, the reaction may lead to one major product or a mixture of many products. For each pair of alkene substrates, draw all metathesis products formed. (Disregard any starting materials that may also be present at equilibrium.) With reference to the three examples, discuss when alkene metathesis with two different alkenes is a synthetically useful reaction.




26.32 Draw the structure of the two products of molecular formula $C_{15}H_{26}O_2$ formed when **M** is treated with Grubbs catalyst under high-dilution conditions.



M

26.33 When certain cycloalkenes are used in metathesis reactions, **ring-opening metathesis polymerization (ROMP)** occurs to form a high molecular weight polymer, as shown with cyclopentene as the starting material. The reaction is driven to completion by relief of strain in the cycloalkene.





What products are formed by ring-opening metathesis polymerization of each alkene?


a. 
b. 
c. 


General Reactions


26.34 Draw the products formed in each reaction.


a. 


b. 


c. 

d. 

e. 


f. 

g. 

h. 


Mechanisms

26.35 In addition to using CHX_3 and base to synthesize dihalocarbenes (Section 26.4), dichlorocarbene ($:CCl_2$) can be prepared by heating sodium trichloroacetate. Draw a stepwise mechanism for this reaction.




sodium trichloroacetate


26.36 Draw a stepwise mechanism for the following reaction.




26.37 Sulfur ylides, like the phosphorus ylides of Chapter 21, are useful intermediates in organic synthesis. Methyl *trans*-chrysanthemate, an intermediate in the synthesis of the insecticide pyrethrin I (Section 26.4), can be prepared from diene **A** and a sulfur ylide. Draw a stepwise mechanism for this reaction.



26.38 Although diazomethane (CH_2N_2) is often not a useful reagent for preparing cyclopropanes, other diazo compounds give good yields of more complex cyclopropanes. Draw a stepwise mechanism for the conversion of diazo compound **A** to **B,** an intermediate in the synthesis of sirenin, the sperm attractant produced by the female gametes of the water mold *Allomyces.*




26.39 The reaction of cyclohexene with iodobenzene under Heck conditions forms **E,** a coupling product with the new phenyl group on the allylic carbon, but none of the "expected" coupling product **F** with the phenyl group bonded directly to the carbon–carbon double bond.



a. Draw a stepwise mechanism that illustrates how **E** is formed.
b. Step [2] in Mechanism 26.2 proceeds with syn addition of Pd and R' to the double bond. What does the formation of **E** suggest about the stereochemistry of the elimination reaction depicted in Step [3] of Mechanism 26.2?


Synthesis

26.40 Devise a synthesis of diene **A** from (*Z*)-2-bromostyrene as the only organic starting material. Use a Suzuki reaction in one step of the synthesis.





26.41 Devise a synthesis of (*E*)-1-phenylhex-1-ene ($CH_3CH_2CH_2CH_2CH{=}CHPh$) using hydrocarbons having ≤ 6 C's and a Suzuki reaction as one of the steps.

26.42 Devise a synthesis of the given trans vinylborane, which can be used for bombykol synthesis (Figure 26.1). All of the carbon atoms in the vinylborane must come from acetylene, nonane-1,9-diol, and catecholborane.





26.43 Devise a synthesis of each compound using a Heck reaction as one step. You may use benzene, $CH_2=CHCO_2Et$, organic alcohols having two carbons or fewer, and any required inorganic reagents.

a. 


b. 


26.44 Devise a synthesis of each compound from cyclohexene and any required organic or inorganic reagents.

a. 

b. 

26.45 Devise a synthesis of each compound from benzene. You may also use any organic compounds having four carbons or fewer, and any required inorganic reagents.


a. 

b. 


26.46 Devise a synthesis of each substituted cyclopropane. Use acetylene ($HC\equiv CH$) as a starting material in part (a), and cyclohexanone as a starting material in part (b). You may use any other organic compounds and any needed reagents.


a. 


+ enantiomer

b. 


26.47 Biaryls, compounds containing two aromatic rings joined by a C–C bond, can often be efficiently made by two different Suzuki couplings; that is, either aromatic ring can be used to form the organoborane needed for coupling. In some cases, however, only one route is possible. With this in mind, synthesize each of the following biaryls using benzene as the starting material for each aromatic ring. When more than one route is possible, draw both of them. You may use any required organic or inorganic reagents.


a. 

b. 


c. 


26.48 Draw the product formed from the ring-closing metathesis of each compound. Then, devise a synthesis of each metathesis starting material using any of the following compounds: $CH_2(CO_2Et)_2$, alcohols with less than five carbons, and any needed organic and inorganic reagents.

a. 


b. 

26.49 Draw the product formed from the ring-closing metathesis of each compound. Then, devise a synthesis of each metathesis starting material from benzene, alcohols with less than five carbons, and any needed organic and inorganic reagents.


a. 

b. 


26.50 What reagents are needed to carry out transformations [1]–[3] in the synthesis of aldehyde **A**? **A** can be converted to the antitumor agent maytansine in several steps.




26.51 Devise a synthesis of each of the following compounds. Besides inorganic reagents, you may use hydrocarbons and halides having ≤ 6 C's, and $CH_2=CHCOOCH_3$ as starting materials. Each synthesis must use at least one of the carbon–carbon bond-forming reactions in this chapter.




(two enantiomers)



(two enantiomers)








(+ enantiomer)

Challenge Problems

26.52 Many variations of ring-closing metathesis have now been reported. Tandem ring-opening–ring-closing metathesis can occur with cyclic alkenes that contain two additional carbon–carbon double bonds. In this reaction, the cycloalkene is cleaved, and two new rings are formed. [1] What compounds are formed in this tandem reaction with the following substrates? [2] Devise a synthesis of the substrate in part (b) that uses a Diels–Alder reaction with diethyl maleate as the dienophile.









diethyl maleate

26.53 The following conversion, carried out in the presence of Grubbs catalyst and ethylene gas, involves a cascade of metathesis reactions. Draw a reaction sequence that illustrates how the reactant is converted to the product **Z,** and indicate where each labeled atom in the reactant ends up in **Z.**




Z

26.54 Suzuki coupling of aryl iodide **A** and vinylborane **B** affords compound **C,** which is converted to **D** in the presence of aqueous acid. Identify compounds **C** and **D** and draw a stepwise mechanism for the conversion of **C** to **D.**




A **B**

26.55 Dimethyl cyclopropanes can be prepared by the reaction of an α,β-unsaturated carbonyl compound **X** with two equivalents of a Wittig reagent **Y.** Draw a stepwise mechanism for this reaction.



X **Y**


26.56 Dienynes undergo metathesis to afford fused bicyclic ring systems. (a) Explain how **A** is converted to **B.** (b) Keeping this reaction in mind, draw the two products formed by dienyne metathesis of **C.**



A **B** **C**

27 Pericyclic Reactions

27.1 Types of pericyclic reactions
27.2 Molecular orbitals
27.3 Electrocyclic reactions
27.4 Cycloaddition reactions
27.5 Sigmatropic rearrangements
27.6 Summary of rules for pericyclic reactions



Periplanone B, an unusual diepoxide with a 10-membered ring, is a potent sex pheromone of the female American cockroach. Although periplanone B was isolated in 1952, its structure was not determined until 1976 using 200 μg of material obtained from more than 75,000 female cockroaches. This structure was confirmed by synthesis in 1979, and several subsequent syntheses have been reported. Key steps in an elegant 1984 synthesis of periplanone B involve pericyclic reactions, a group of powerful, stereospecific reactions discussed in Chapter 27.

Many of the reactions thus far encountered in our study of organic chemistry occur by way of reactive intermediates—cations, anions, and radicals. For example, the S_N1 reaction in Chapter 7 and electrophilic aromatic substitutions in Chapter 18 involve carbocations, while the aldol and Claisen reactions in Chapter 24 occur via enolate anions. Other reactions, such as the halogenation of alkanes and the polymerization of alkenes discussed in Chapter 15, take place via radical intermediates.

In Chapter 27, we learn about a small but versatile group of reactions, **pericyclic reactions,** which occurs in a concerted process—all bonds are broken and formed in a single step—with a cyclic transition state. The Diels–Alder reaction in Chapter 16 is an example of one type of pericyclic reaction. Pericyclic reactions involve π bonds, and they are governed by a set of rules that allows us to predict the identity and stereochemistry of the products formed. Consequently, pericyclic reactions are valuable tools for synthesizing organic molecules.

27.1 Types of Pericyclic Reactions

Although most organic reactions take place by way of ionic or radical intermediates, a number of useful reactions occur in one-step processes that do *not* form reactive intermediates.


- **A pericyclic reaction is a concerted reaction that proceeds through a cyclic transition state.**

Stereospecific reactions were first discussed in Chapter 10.

Pericyclic reactions require light or heat and are completely stereospecific; that is, a particular stereoisomer of the reactant forms a particular stereoisomer of the product. There are three categories of pericyclic reactions: **electrocyclic reactions, cycloadditions,** and **sigmatropic rearrangements.**

An **electrocyclic reaction** is a reversible reaction that can involve ring closure or ring opening of one molecule of reactant to form one molecule of product.


- **An electrocyclic ring closure is an intramolecular reaction that forms a cyclic product containing one more σ bond and one fewer π bond than the reactant.**



hexa-1,3,5-triene
3 π bonds

cyclohexa-1,3-diene
2 π bonds

- **An electrocyclic ring opening is a reaction in which a σ bond of a cyclic reactant is cleaved to form a conjugated product with one more π bond.**




cyclobutene
1 π bond

buta-1,3-diene
2 π bonds


Cycloaddition reactions form a ring. The Diels–Alder reaction in Chapter 16 is one example of a cycloaddition.

- **A cycloaddition is a reaction between two compounds with π bonds to form a cyclic product with two new σ bonds.**



In contrast to electrocyclic reactions and cycloadditions, in which the number of π bonds differs in the reactants and products, the number of π bonds does *not* change in a **sigmatropic rearrangement.**

- **A sigmatropic rearrangement is a reaction in which a σ bond is broken in the reactant, the π bonds rearrange, and a σ bond is formed in the product.**




Hoffmann and Fukui received the 1981 Nobel Prize in Chemistry for developing theories that explain the course of pericyclic reactions.

Two features determine the course of the reactions: the number of π bonds involved and whether the reaction occurs in the presence of heat (thermal conditions) or light (photochemical conditions). These reactions follow a set of rules based on orbitals and symmetry first proposed by R. B. Woodward and Roald Hoffmann in 1965, and derived from theory described by Kenichi Fukui in 1954.

To understand pericyclic reactions we must review and expand upon what we learned about the molecular orbitals of systems with π bonds in Chapter 17.

Problem 27.1 Classify each reaction as an electrocyclic reaction, a cycloaddition, or a sigmatropic rearrangement. Label the σ bonds that are broken or formed in each reaction.



27.2 Molecular Orbitals


In Section 17.9 we learned that molecular orbital (MO) theory describes bonds as the mathematical combination of atomic orbitals that forms a new set of orbitals called **molecular orbitals (MOs). The number of atomic orbitals used *equals* the number of molecular orbitals formed.**

Because pericyclic reactions involve π bonds, let's examine the molecular orbitals that result from *p* orbital overlap in ethylene, buta-1,3-diene, and hexa-1,3,5-triene, molecules that contain one, two, and three π bonds, respectively. Keep in mind that the two lobes of a *p* orbital are opposite in phase, with a node of electron density at the nucleus.

27.2A Ethylene

The π bond in ethylene ($CH_2{=}CH_2$) is formed by side-by-side overlap of two *p* orbitals on adjacent carbons. Two *p* orbitals can combine in two different ways. As shown in Figure 27.1, when two *p* orbitals of similar phase overlap, a π bonding molecular orbital (designated as ψ_1) results. Two electrons occupy this lower-energy bonding molecular orbital. When two *p* orbitals of opposite phase combine, a π* antibonding molecular orbital (designated as ψ_2*) results. A destabilizing node between the orbitals occurs when two orbitals of opposite phase combine.


Figure 27.1
The π and π* molecular orbitals of ethylene



27.2B Buta-1,3-diene

The two π bonds of buta-1,3-diene ($CH_2=CH-CH=CH_2$) are formed by overlap of four *p* orbitals on four adjacent carbons. As shown in Figure 27.2, four *p* orbitals can combine in four different ways to form four molecular orbitals designated as ψ_1–ψ_4. Two are bonding molecular orbitals (ψ_1 and ψ_2), and two are antibonding molecular orbitals (ψ_3* and ψ_4*). The two bonding MOs are lower in energy than the *p* orbitals from which they are formed, whereas the two antibonding MOs are higher in energy than the *p* orbitals from which they are formed. **As the number of bonding interactions decreases and the number of nodes increases, the energy of the molecular orbital increases.**

Figure 27.2
The four π molecular orbitals of buta-1,3-diene



- The two lowest energy molecular orbitals, ψ_1 and ψ_2, are bonding MOs.
- The two highest energy molecular orbitals, ψ_3* and ψ_4*, are antibonding MOs.

- In the ground-state electronic arrangement, the four π electrons occupy the two bonding molecular orbitals.

Also recall from Section 17.9:

- The highest energy orbital that contains electrons is called the highest occupied molecular orbital (HOMO). In the ground state of buta-1,3-diene, ψ_2 is the HOMO.
- The lowest energy orbital that contains no electrons is called the lowest unoccupied molecular orbital (LUMO). In the ground state of buta-1,3-diene, ψ_3^* is the LUMO.

The thermal reactions discussed in Section 27.3 utilize reactants in their ground state electronic configuration.

When buta-1,3-diene absorbs light of appropriate energy, an electron is promoted from ψ_2 (the HOMO) to ψ_3^* (the LUMO) to form a higher energy electronic configuration, the **excited state.** In the excited state, the HOMO is now ψ_3^*. **In the photochemical reactions in Section 27.3, the reactant is in its excited state.** As a result, the HOMO is ψ_3^* and the LUMO is ψ_4^* for buta-1,3-diene.

All conjugated dienes can be described by a set of molecular orbitals that are similar to those drawn in Figure 27.2 for buta-1,3-diene.

Problem 27.2 For each molecular orbital in Figure 27.2, count the number of bonding interactions (interactions between adjacent orbitals of similar phase) and the number of nodes. (a) How do these two values compare for a bonding molecular orbital? (b) How do these two values compare for an antibonding molecular orbital?

27.2C Hexa-1,3,5-triene


The three π bonds of hexa-1,3,5-triene ($CH_2=CH-CH=CH-CH=CH_2$) are formed by overlap of six *p* orbitals on six adjacent carbons. As shown in Figure 27.3, six *p* orbitals can combine in six different ways to form six molecular orbitals designated as ψ_1–ψ_6. Three are bonding molecular orbitals (ψ_1–ψ_3), and three are antibonding molecular orbitals (ψ_4^*–ψ_6^*).

In the ground state electronic configuration, the six π electrons occupy the three bonding MOs, ψ_3 is the HOMO, and ψ_4^* is the LUMO. In the excited state, which results from promotion of an electron from ψ_3 to ψ_4^*, ψ_4^* is the HOMO and ψ_5^* is the LUMO.

Problem 27.3 (a) Using Figure 27.2 as a guide, draw the molecular orbitals for hexa-2,4-diene. (b) Label the HOMO and the LUMO in the ground state. (c) Label the HOMO and the LUMO in the excited state.


Problem 27.4 (a) How many π molecular orbitals are present in deca-1,3,5,7,9-pentaene ($CH_2=CH-CH=CH-CH=CH-CH=CH-CH=CH_2$)? (b) How many are bonding MOs and how many are antibonding MOs? (c) How many nodes are present in ψ_1? (d) How many nodes are present in ψ_{10}^*?

Figure 27.3 The six π molecular orbitals of hexa-1,3,5-triene



27.3 Electrocyclic Reactions

An electrocyclic reaction is a reversible reaction that involves ring closure of a conjugated polyene to a cycloalkene, or ring opening of a cycloalkene to a conjugated polyene. For example, ring closure of hexa-1,3,5-triene forms cyclohexa-1,3-diene, a product with one more σ bond and one fewer π bond than the reactant. Ring opening of cyclobutene forms buta-1,3-diene, a product with one fewer σ bond and one more π bond than the reactant.




Arrows may be drawn in a clockwise or counterclockwise direction to show the flow of electrons.

- **To draw the product in each reaction, use curved arrows and begin at a π bond. Move the π electrons to an adjacent carbon–carbon bond and continue in a cyclic fashion.**

In a ring-closing reaction, this process forms a new σ bond that now joins the ends of the conjugated polyene. In a ring-opening reaction, this process breaks a σ bond to form a conjugated polyene with one more π bond.


Whether the reactant or product predominates at equilibrium depends on the ring size of the cyclic compound. Generally, a six-membered ring is favored over an acyclic triene at equilibrium. In contrast, an acyclic diene is favored over a strained four-membered ring.

Problem 27.5 Use curved arrows and draw the product of each electrocyclic reaction.




27.3A Stereochemistry and Orbital Symmetry

Electrocyclic reactions are completely stereospecific. For example, ring closure of (2*E*,4*Z*,6*E*)-octa-2,4,6-triene yields a single product with cis methyl groups on the ring. Ring opening of *cis*-3,4-dimethylcyclobutene forms a single conjugated diene with one *Z* alkene and one *E* alkene.



Moreover, the stereochemistry of the product of an electrocyclic reaction depends on whether the reaction is carried out under thermal or photochemical reaction conditions—that is, with heat or light, respectively. Cyclization of (2*E*,4*E*)-hexa-2,4-diene with heat forms a cyclobutene with trans methyl groups, whereas cyclization with light forms a cyclobutene with cis methyl groups.

Electrocyclic ring closure generally forms either an achiral meso compound or a mixture of chiral enantiomers. When enantiomers form, only one enantiomer is drawn in these reactions.




To understand these results, we must focus on the HOMO of the acyclic conjugated polyene that is either the reactant or product in an electrocyclic reaction. In particular, we must examine the *p* orbitals on the terminal carbons of the HOMO, and determine whether like phases of the orbitals are on the same side or on opposite sides of the molecule.




- **An electrocyclic reaction occurs only when like phases of orbitals can overlap to form a bond. Such a reaction is *symmetry allowed.***
- **An electrocyclic reaction cannot occur between lobes of opposite phase. Such a reaction is *symmetry forbidden.***

To form a bond, the *p* orbitals on the terminal carbons must rotate so that like phases can interact to form the new σ bond. Two modes of rotation are possible.

- **When like phases of the *p* orbitals are on the same side of the molecule, the two orbitals must rotate in opposite directions—one clockwise and one counterclockwise. Rotation in opposite directions is said to be *disrotatory.***



- **When like phases of the *p* orbitals are on opposite sides of the molecule, the two orbitals must rotate in the same direction—both clockwise or both counterclockwise. Rotation in the same direction is said to be *conrotatory.***



27.3B Thermal Electrocyclic Reactions


To explain the stereochemistry observed in electrocyclic reactions, we must examine the symmetry of the molecular orbital that contains the most loosely held π electrons. In a thermal reaction, we consider the **HOMO of the ground state electronic configuration.** Rotation occurs in a disrotatory or conrotatory fashion so that like phases of the *p* orbitals on the terminal carbons of this molecular orbital combine.

- **The number of double bonds in the conjugated polyene determines whether rotation is conrotatory or disrotatory.**

Two examples illustrate different outcomes.

Thermal electrocyclic ring closure of (2*E*,4*Z*,6*E*)-octa-2,4,6-triene yields a single product with cis methyl groups on the ring.

Only the *p* orbitals on the terminal carbons of the HOMO are drawn for clarity.



counterclockwise clockwise

ground state HOMO of (2*E*,4*Z*,6*E*)-octa-2,4,6-triene

cis-5,6-dimethylcyclohexa-1,3-diene

cis product


Cyclization occurs in a disrotatory fashion because the HOMO of a conjugated triene has like phases of the outermost *p* orbitals on the *same* side of the molecule (Figure 27.3). A disrotatory ring closure is symmetry allowed because like phases of the *p* orbitals overlap to form the new σ bond of the ring. In the disrotatory ring closure, both methyl groups are pushed *down* (or *up*), making them *cis* in the product.

This is a specific example of the general process observed for conjugated polyenes with an *odd* number of π bonds. The HOMO of a conjugated polyene with an odd number of π bonds has like phases of the outermost *p* orbitals on the *same* side of the molecule. As a result:

- **Thermal electrocyclic reactions occur in a *disrotatory* fashion for a conjugated polyene with an *odd* number of π bonds.**

In contrast, thermal electrocyclic ring closure of (2*E*,4*E*)-hexa-2,4-diene forms a cyclobutene with trans methyl groups.

The conrotatory ring closure of (2*E*,4*E*)-hexa-2,4-diene is drawn with two clockwise rotations. The conrotatory ring closure could also be drawn with two counterclockwise rotations, leading to the enantiomer of the trans product drawn. Both enantiomers are formed in equal amounts.




Cyclization occurs in a conrotatory fashion because the HOMO of a conjugated diene has like phases of the outermost *p* orbitals on *opposite* sides of the molecule (Figure 27.2). A conrotatory ring closure is symmetry allowed because like phases of the *p* orbitals overlap to form the new σ bond of the ring. In the conrotatory ring closure, one methyl group is pushed *down* and one methyl group is pushed *up,* making them *trans* in the product.


This is a specific example of the general process observed for conjugated polyenes with an *even* number of π bonds. The HOMO of a conjugated polyene with an even number of π bonds has like phases of the outermost *p* orbitals on *opposite* sides of the molecule. As a result:

- **Thermal electrocyclic reactions occur in a *conrotatory* fashion for a conjugated polyene with an *even* number of π bonds.**

Since electrocyclic reactions are reversible, **electrocyclic ring-opening reactions follow the same rules** as electrocyclic ring closures. Thus, thermal ring opening of *cis*-3,4-dimethylcyclobutene—which ring opens to a diene with an *even* number of π bonds—occurs in a conrotatory fashion to form (2*E*,4*Z*)-hexa-2,4-diene as the only product.



Sample Problem 27.1 Draw the product of each thermal electrocyclic ring closure.




Solution

Count the number of π bonds in the conjugated polyene to determine the mode of ring closure in a thermal electrocyclic reaction.

- A conjugated polyene with an **odd** number of π bonds undergoes **disrotatory** cyclization.
- A conjugated polyene with an **even** number of π bonds undergoes **conrotatory** cyclization.


a. (2*E*,4*Z*,6*Z*)-Octa-2,4,6-triene contains three π bonds. The HOMO of a conjugated polyene with an odd number of π bonds has like phases of the outermost *p* orbitals on the *same* side of the molecule, and this results in disrotatory cyclization.



counterclockwise clockwise
(2*E*,4*Z*,6*Z*)-octa-2,4,6-triene

trans-5,6-dimethylcyclohexa-1,3-diene
trans product


b. Diene **B** contains two π bonds. The HOMO of a conjugated polyene with an even number of π bonds has like phases of the outermost *p* orbitals on *opposite* sides of the molecule, and this results in conrotatory cyclization.




clockwise clockwise
B

trans product

Problem 27.6 What product is formed when each compound undergoes thermal electrocyclic ring opening or ring closure? Label each process as conrotatory or disrotatory and clearly indicate the stereochemistry around tetrahedral stereogenic centers and double bonds.




Problem 27.7 What cyclic product is formed when each decatetraene undergoes thermal electrocyclic ring closure?



27.3C Photochemical Electrocyclic Reactions

Photochemical electrocyclic reactions follow similar principles as those detailed in thermal reactions with one important difference. **In photochemical reactions, we must consider the orbitals of the HOMO of the *excited* state to determine the course of the reaction.** As a photon is absorbed, an electron in the ground state HOMO is excited to the ground state LUMO. As a result, the excited state HOMO is one energy level higher than before (see Figures 27.2 and 27.3). The excited state HOMO has the *opposite* orientation of the outermost *p* orbitals compared to the HOMO of the ground state. As a result, **the method of ring closure of a photochemical electrocyclic reaction is *opposite* to that of a thermal electrocyclic reaction for the same number of π bonds.**

Photochemical electrocyclic ring closure of (2*E*,4*Z*,6*E*)-octa-2,4,6-triene yields a cyclic product with trans methyl groups on the ring.




clockwise clockwise
excited state HOMO of
(2*E*,4*Z*,6*E*)-octa-2,4,6-triene

trans-5,6-dimethylcyclohexa-1,3-diene
trans product

Cyclization occurs in a conrotatory fashion because the excited state HOMO of a conjugated triene has like phases of the outermost *p* orbitals on the *opposite* sides of the molecule (Figure 27.3). In the conrotatory ring closure, one methyl group is pushed *down* and one methyl group is pushed *up,* making them *trans* in the product. This is a specific example of the general process observed for conjugated polyenes with an *odd* number of π bonds.

- **Photochemical electrocyclic reactions occur in a *conrotatory* fashion for a conjugated polyene with an *odd* number of π bonds.**

Photochemical electrocyclic ring closure of (2*E*,4*E*)-hexa-2,4-diene forms a cyclobutene with cis methyl groups.



clockwise counterclockwise

excited state HOMO of (2*E*,4*E*)-hexa-2,4-diene

cis-3,4-dimethylcyclobutene

cis product

Cyclization occurs in a disrotatory fashion because the excited state HOMO of a conjugated diene has like phases of the outermost *p* orbitals on the *same* side of the molecule (Figure 27.3). In the disrotatory ring closure, both methyl groups are pushed *down* (or *up*), making them *cis* in the product. This is a specific example of the general process observed for conjugated polyenes with an *even* number of π bonds.


- **Photochemical electrocyclic reactions occur in a *disrotatory* fashion for a conjugated polyene with an *even* number of π bonds.**

Problem 27.8 What product is formed when each compound in Problem 27.6 undergoes photochemical electrocyclic ring opening or ring closure? Label each process as conrotatory or disrotatory and clearly indicate the stereochemistry around tetrahedral stereogenic centers and double bonds.



Vitamin D (Problem 27.9) regulates calcium absorption, so adequate vitamin D levels are needed for proper bone growth. Vitamin D-fortified milk sold in the United States is produced by exposing milk to ultraviolet light.

Problem 27.9 Vitamin D_3, the most abundant of the D vitamins, is synthesized from 7-dehydrocholesterol, a compound found in milk and fatty fish such as salmon and mackerel. When the skin is exposed to sunlight, a photochemical electrocyclic ring opening forms provitamin D_3, which is then converted to vitamin D_3 by a sigmatropic rearrangement (Section 27.5). Draw the structure of provitamin D_3.



7-dehydrocholesterol


27.3D Summary of Electrocyclic Reactions

Table 27.1 summarizes the rules, often called the **Woodward–Hoffmann rules,** for electrocyclic reactions under thermal or photochemical reaction conditions. The number of π bonds refers to the acyclic conjugated polyene that is either the reactant or product of an electrocyclic reaction.

Table 27.1 Woodward–Hoffmann Rules for Electrocyclic Reactions


Number of π bonds	Thermal reaction	Photochemical reaction
Even	Conrotatory	Disrotatory
Odd	Disrotatory	Conrotatory

Sample Problem 27.2 Identify **A** and **B** in the following reaction sequence. Label each process as conrotatory or disrotatory.





Solution

Ring opening of a cyclohexadiene forms a hexatriene with **three** π bonds. A conjugated polyene with an odd number of π bonds undergoes a thermal electrocyclic reaction in a disrotatory fashion (Table 27.1). The resulting hexatriene (**A**) then undergoes a photochemical electrocyclic reaction in a conrotatory fashion to form a cyclohexadiene with cis methyl groups (**B**).




Problem 27.10 Draw the product formed when each triene undergoes electrocyclic reaction under [1] thermal conditions; [2] photochemical conditions.

a. 

b. 

Problem 27.11 What product would be formed by the disrotatory cyclization of the given triene? Would this reaction occur under photochemical or thermal conditions?




27.4 Cycloaddition Reactions


A cycloaddition is a reaction between two compounds with π bonds to form a cyclic product with two new σ bonds. Like electrocyclic reactions, cycloadditions are concerted, stereospecific reactions, and the course of the reaction is determined by the symmetry of the molecular orbitals of the reactants.

Cycloadditions can be initiated by heat (thermal conditions) or light (photochemical conditions). Cycloadditions are identified by the number of π electrons in the two reactants.


The Diels–Alder reaction is a thermal [4 + 2] cycloaddition that occurs between a diene with four π electrons and an alkene (dienophile) with two π electrons (Sections 16.12–16.14).



A photochemical [2 + 2] cycloaddition occurs between two alkenes, each with two π electrons, to form a cyclobutane. Thermal [2 + 2] cycloadditions do *not* take place.




Sample Problem 27.3 What type of cycloaddition is shown in each equation?




Solution

Count the number of π electrons *involved in each reactant* to classify the cycloaddition.


a. [2 + 2] Cycloaddition



b. [4 + 2] Cycloaddition



Problem 27.12 Consider cycloheptatrienone and ethylene, and draw a possible product formed from each type of cycloaddition: (a) [2 + 2]; (b) [4 + 2]; (c) [6 + 2].




cycloheptatrienone


27.4A Orbital Symmetry and Cycloadditions

To understand cycloaddition reactions, we examine the *p* orbitals of the terminal carbons of both reactants. Bonding can take place only when like phases of both sets of *p* orbitals can combine. Two modes of reaction are possible.

- A suprafacial cycloaddition occurs when like phases of the *p* orbitals of both reactants are on the *same* side of the π system, so that two bonding interactions result.



- An antarafacial cycloaddition occurs when one π system must *twist* to align like phases of the *p* orbitals of the terminal carbons of the reactants.




Because of the geometrical constraints of small rings, **cycloadditions that form four- or six-membered rings must take place by suprafacial pathways.**

Since cycloaddition involves the donation of electron density from one reactant to another, one reactant donates its most loosely held electrons—those occupying its HOMO—to a vacant orbital that can accept electrons—the LUMO—of the second reactant. The HOMO of either reactant can be used for analysis.

- In a cycloaddition we examine the bonding interactions of the HOMO of one component with the LUMO of the second component.

27.4B [4 + 2] Cycloadditions

To examine the course of a [4 + 2] cycloaddition, let's arbitrarily choose the HOMO of the diene and the LUMO of the alkene, and look at the symmetry of the *p* orbitals on the terminal carbons of both components. Since two bonding interactions result from overlap of the like phases of both sets of *p* orbitals, **a [4 + 2] cycloaddition occurs readily by suprafacial reaction under thermal conditions.**




This is a specific example of a general cycloaddition involving an odd number of π bonds (three π bonds total, two from the diene and one from the alkene).

- Thermal cycloadditions involving an *odd* number of π bonds proceed by a *suprafacial* pathway.

In Section 16.13, we learned that the stereochemistry of the dienophile is retained in the Diels–Alder product.

Because a Diels–Alder reaction follows a concerted, suprafacial pathway, the **stereochemistry of the diene is retained in the Diels–Alder product.** As a result, reaction of (2*E*,4*E*)-hexa-2,4-diene with ethylene forms a cyclohexene with cis substituents (Reaction [1]), whereas reaction of (2*E*,4*Z*)-hexa-2,4-diene with ethylene forms a cyclohexene with trans substituents (Reaction [2]).







Reserpine (Problem 27.15) is isolated from the roots of *Rauwolfia serpentina,* a plant used for centuries in traditional Indian medicine for the treatment of mental disorders and venomous bites.

Problem 27.13 Show that a thermal suprafacial addition is symmetry allowed in a [4 + 2] cycloaddition by using the HOMO of the alkene and the LUMO of the diene.

Problem 27.14 Draw the product (including stereochemistry) formed from each pair of reactants in a thermal [4 + 2] cycloaddition reaction.




Problem 27.15 (a) Draw the product of the following [4 + 2] cycloaddition, which was carried out in the early stages of the synthesis of the alkaloid reserpine (Problem 22.83). Indicate the stereochemistry at any newly formed stereogenic centers. (b) Draw the *p* orbitals of the alkene and the terminal carbons of the conjugated diene, and show how the orientation of the reactants and orbital overlap lead to the observed stereochemistry.




27.4C [2 + 2] Cycloadditions

In contrast to a [4 + 2] cycloaddition, a [2 + 2] cycloaddition does *not* occur under thermal conditions, but *does* take place photochemically. This result is explained by examining the symmetry of the HOMO and LUMO of the alkene reactants.

In a thermal [2 + 2] cycloaddition, like phases of the *p* orbitals on only one set of terminal carbons can overlap. For like phases to overlap on the other terminal carbon, the molecule must twist to allow for an antarafacial pathway. This process *cannot* occur to form small rings.



In a photochemical [2 + 2] cycloaddition, light energy promotes an electron from the ground state HOMO to form the excited state HOMO (designated as ψ_2^* in Figure 27.1). Interaction of this excited state HOMO with the LUMO of the second alkene then allows for overlap of the like phases of both sets of *p* orbitals. Two bonding interactions result and the reaction occurs by a suprafacial pathway.



This is a specific example of a general cycloaddition involving an even number of π bonds (two π bonds total, one from each alkene).

- **Photochemical cycloadditions involving an *even* number of π bonds proceed by a *suprafacial* pathway.**

Problem 27.16 Draw the product formed in each cycloaddition.



27.4D Summary of Cycloaddition Reactions

Table 27.2 summarizes the Woodward–Hoffmann rules that govern cycloaddition reactions. The number of π bonds refers to the total number of π bonds from both components of the cycloaddition. For a given number of π bonds, the mode of cycloaddition is always *opposite* in thermal and photochemical reactions.

Table 27.2 Woodward–Hoffmann Rules for Cycloaddition Reactions

Number of π bonds	Thermal reaction	Photochemical reaction
Even	Antarafacial	Suprafacial
Odd	Suprafacial	Antarafacial


Problem 27.17 Using the Woodward–Hoffmann rules, predict the stereochemical pathway for each cycloaddition: (a) a [6 + 4] photochemical reaction; (b) an [8 + 2] thermal reaction.

Problem 27.18 Using orbital symmetry, explain why a Diels–Alder reaction does not take place under photochemical reaction conditions.

27.5 Sigmatropic Rearrangements

A sigmatropic rearrangement is an intramolecular pericyclic reaction in which a σ bond is broken in a reactant, the π bonds rearrange, and a new σ bond is formed in the product. In a sigmatropic rearrangement, the number of π bonds in the reactant and product is constant, and


the σ bonds broken and formed are **allylic** C−H, C−C, or C−Z bonds (Z = N, O, or S). A sigmatropic rearrangement that results in cleavage and formation of a C−H bond is shown.




Sigmatropic rearrangements are characterized by a set of numbers in brackets, [*n*,*m*], to indicate the location of the new σ bond relative to the broken σ bond. To designate a sigmatropic rearrangement:

- **Locate the σ bond broken in the reactant and label both atoms in the bond with "1's."**
- **Locate the new σ bond in the product, and count the number of atoms from the broken σ bond to the new σ bond for each fragment.**
- **Place both numbers in brackets, with the lower number first. In a rearrangement involving a C−H bond, the first number is always "1."**

For example, a [3,3] sigmatropic rearrangement converts diene **A** into diene **B** when an allylic C−C bond in **A** is broken and a new allylic C−C bond is formed in **B.**




Sample Problem 27.4 What type of sigmatropic rearrangement is illustrated in each equation?




Solution

Locate the atoms in the broken σ bond and label them with 1's. Locate the atoms in the new σ bond, and count the number of atoms from the bond broken to the bond formed. When a C−H bond is broken, the first number in the [*n*,*m*] designation must be 1, because the H atom is bonded to no other atom.


a. A C−H bond is broken on the allylic C and a new C−H bond is formed on C5, so the reaction is a **[1,5] sigmatropic rearrangement.**



b. The reaction is a **[3,3] sigmatropic rearrangement,** because a C−O σ bond is broken and a new allylic C−C σ bond is formed between carbons that are three atoms removed from the broken bond.




Problem 27.19 What type of sigmatropic rearrangement is illustrated in each equation?




27.5A Sigmatropic Rearrangements and Orbital Symmetry

The stereochemistry of a sigmatropic rearrangement, like that of other pericyclic reactions, is determined by the symmetry of the orbitals involved in the reaction. In sigmatropic rearrangements, we consider the orbitals of the σ bond that is broken and the terminal *p* orbital of the π bond at which the new σ bond forms. Two modes of rearrangement are possible: **suprafacial** and **antarafacial.**

- **In a suprafacial rearrangement, the new σ bond forms on the *same* side of the π system as the broken σ bond.**



- **In an antarafacial rearrangement, the new σ bond forms on the *opposite* side of the π system as the broken σ bond.**




Sigmatropic rearrangements can occur under thermal or photochemical conditions, and follow the same rules observed in cycloaddition reactions. With sigmatropic rearrangements we count the total number of electron pairs in the σ bond that is broken and the π bonds that rearrange (Table 27.3). Because sigmatropic rearrangements involve cyclic transition states and small rings have geometrical constraints, **reactions involving six atoms or fewer must take place by suprafacial pathways.**

Table 27.3 Woodward–Hoffmann Rules for Sigmatropic Rearrangements

Number of electron pairs	Thermal reaction	Photochemical reaction
Even	Antarafacial	Suprafacial
Odd	Suprafacial	Antarafacial

For example, a [1,5] sigmatropic rearrangement of **X** to **Y** involves three electron pairs, one from the σ bond that is broken and two from the π bonds that rearrange.




According to Table 27.3, this reaction must occur in a suprafacial mode under thermal conditions and in an antarafacial mode under photochemical conditions. Since this reaction involves only six atoms (including the H atom that migrates), it must take place under thermal conditions in a suprafacial fashion.

Sample Problem 27.5 Classify the following sigmatropic rearrangement and determine whether it takes place readily under thermal or photochemical reaction conditions.




Solution

First, classify the rearrangement as in Sample Problem 27.4: Label the atoms in the broken σ bond with 1's, locate the new σ bond, and count the number of atoms from the bond broken to the bond formed. Then, count the number of electron pairs involved in the reaction, and use Table 27.3 to determine the stereochemical pathway of the reaction. Keep in mind that reactions involving six atoms or fewer must take place by suprafacial pathways.




This reaction is a [1,3] sigmatropic rearrangement, involving two electron pairs: the C–H σ bond broken and one π bond. Since the reaction involves four atoms, it must take place via a suprafacial pathway, which occurs under photochemical conditions.

Problem 27.20 (a) What product is formed from the [1,7] sigmatropic rearrangement of a deuterium in the following triene? (b) Does this reaction proceed in a suprafacial or antarafacial manner under thermal conditions? (c) Does this reaction proceed in a suprafacial or antarafacial manner under photochemical conditions?



27.5B [3,3] Sigmatropic Rearrangements


Two widely used [3,3] sigmatropic rearrangements in organic synthesis are the **Cope rearrangement** of a 1,5-diene to an isomeric 1,5-diene, and the **Claisen rearrangement** of an unsaturated ether to a γ,δ-unsaturated carbonyl compound.



Both reactions involve three electron pairs—two π bonds and one σ bond—and six atoms, and take place readily in a **suprafacial pathway under thermal conditions.**


Cope Rearrangement

Since a Cope rearrangement involves isomeric 1,5-dienes as reactant and product, the more stable diene is favored at equilibrium. Useful Cope rearrangements occur when the reactant 1,5-diene is considerably less stable than the product, as in the case of *cis*-1,2-divinylcyclobutane, which rearranges to cycloocta-1,5-diene with loss of strain from the cyclobutane ring.




cis-1,2-divinylcyclobutane cycloocta-1,5-diene


The **oxy-Cope rearrangement** is an especially powerful variation of a Cope rearrangement using an unsaturated alcohol. [3,3] Sigmatropic rearrangement forms an enol initially, which then tautomerizes to form a carbonyl group.




Moreover, ***anionic* oxy-Cope rearrangements** often give high yields of rearranged product under very mild reaction conditions. In an anionic oxy-Cope rearrangement, the unsaturated alcohol reactant is first treated with strong base, usually KH in the presence of 18-crown-6 (Section 9.5B), to form an alkoxide. [3,3] Sigmatropic rearrangement then yields a resonance-stabilized enolate, which is protonated to form a carbonyl product.




Problem 27.21 What product is formed from the Cope or oxy-Cope rearrangement of each starting material?

a. 


b. 

c. 

Problem 27.22 One step in the synthesis of periplanone B, the chapter-opening molecule, involved anionic oxy-Cope rearrangement of the following unsaturated alcohol. Draw the product that results after protonation of the intermediate enolate.




Problem 27.23 What compound forms geranial (Figure 21.6) by a Cope rearrangement?




geranial

Claisen Rearrangement

A Claisen rearrangement is a [3,3] sigmatropic rearrangement of an unsaturated ether, either an allyl vinyl ether or an allyl aryl ether. With an allyl vinyl ether, a γ,δ-unsaturated carbonyl compound is formed directly by the concerted rearrangement. With an allyl aryl ether, Claisen rearrangement initially generates a cyclohexadienone intermediate, which tautomerizes to a phenol that contains an allyl group ortho to the OH group.





allyl vinyl ether γ,δ-unsaturated carbonyl compound




allyl aryl ether cyclohexadienone phenol

Problem 27.24 What product is formed from the Claisen rearrangement of each starting material?

a. 


b. 

c. 


Problem 27.25 (a) What product is formed by the Claisen rearrangement of compound **Z?** (b) Using what you have learned about ring-closing metathesis in Chapter 26, draw the product formed when the product in part (a) is treated with Grubbs catalyst. These two reactions are key steps in the synthesis of garsubellin A, a biologically active natural product that stimulates the synthesis of the neurotransmitter acetylcholine. Compounds of this sort may prove to be useful drugs for the treatment of neurodegenerative diseases such as Alzheimer's disease.



Garsubellin A (Problem 27.25) is isolated from the wood of *Garcinia subelliptica,* a tree grown in Okinawa, Japan.



Z



garsubellin A

27.6 Summary of Rules for Pericyclic Reactions

Table 27.4 summarizes the rules that govern pericyclic reactions, and in truth, this table holds a great deal of information. To keep track of this information, it may be helpful to **learn one row in the table only,** and then note the result when one or more conditions change. For example,

- A *thermal* reaction involving an *even* number of electron pairs is *conrotatory* or *antarafacial.*
- If *one* of the reaction conditions changes—either from thermal to photochemical or from an even to an odd number of electron pairs—the stereochemistry of the reaction changes to disrotatory or suprafacial.
- If *both* reaction conditions change—that is, a photochemical reaction with an odd number of electron pairs—the stereochemistry does *not* change.

Table 27.4 Summary of the Stereochemical Rules for Pericyclic Reactions

Reaction conditions	Number of electron pairs	Stereochemistry
Thermal	Even	Conrotatory or antarafacial
	Odd	Disrotatory or suprafacial
Photochemical	Even	Disrotatory or suprafacial
	Odd	Conrotatory or antarafacial

Problem 27.26 Using the Woodward–Hoffmann rules in Table 27.4, predict the stereochemistry of each reaction.

a. a [6 + 4] thermal cycloaddition
b. photochemical electrocyclic ring closure of deca-1,3,5,7,9-pentaene
c. a [4 + 4] photochemical cycloaddition
d. a thermal [5,5] sigmatropic rearrangement

KEY CONCEPTS

Pericyclic Reactions


Electrocyclic Reactions (27.3)

Woodward–Hoffmann rules for electrocyclic reactions

Number of π bonds	Thermal reaction	Photochemical reaction
Even	Conrotatory	Disrotatory
Odd	Disrotatory	Conrotatory

Examples

The stereochemistry of a thermal electrocyclic reaction is opposite to that of a photochemical electrocyclic reaction.



(2*E*,4*E*)-hexa-2,4-diene
2 π bonds

- A thermal electrocyclic reaction with an even number of π bonds occurs in a conrotatory fashion.
- A photochemical electrocyclic reaction with an even number of π bonds occurs in a disrotatory fashion.


Cycloaddition Reactions (27.4)

Woodward–Hoffmann rules for cycloaddition reactions

Number of π bonds	Thermal reaction	Photochemical reaction
Even	Antarafacial	Suprafacial
Odd	Suprafacial	Antarafacial


Examples

[1] A thermal [4 + 2] cycloaddition takes place in a suprafacial fashion with an odd number of π bonds. An antarafacial photochemical [4 + 2] cycloaddition to form a six-membered ring cannot occur, because of the geometrical constraints of forming a six-membered ring.



cis product only

[2] A photochemical [2 + 2] cycloaddition takes place in a suprafacial fashion with an even number of π bonds. An antarafacial thermal [2 + 2] cycloaddition to form a four-membered ring cannot occur, because of the geometrical constraints of forming a four-membered ring.



cis product only

Sigmatropic Rearrangements (27.5)

Woodward–Hoffmann rules for sigmatropic rearrangements

Number of electron pairs	Thermal reaction	Photochemical reaction
Even	Antarafacial	Suprafacial
Odd	Suprafacial	Antarafacial


Examples

[1] A **Cope rearrangement** is a thermal [3,3] sigmatropic rearrangement that converts a 1,5-diene into an isomeric 1,5-diene.




1,5-diene isomeric 1,5-diene

[2] An **oxy-Cope rearrangement** is a thermal [3,3] sigmatropic rearrangement that converts a 1,5-dien-3-ol into a δ,ε-unsaturated carbonyl compound, after tautomerization of an intermediate enol.



1,5-dien-3-ol δ,ε-unsaturated carbonyl compound

[3] A **Claisen rearrangement** is a thermal [3,3] sigmatropic rearrangement that converts an unsaturated ether into a γ,δ-unsaturated carbonyl compound.




unsaturated ether γ,δ-unsaturated carbonyl compound

PROBLEMS


Problems Using Three-Dimensional Models


27.27 (a) What product is formed when each compound undergoes a thermal electrocyclic ring opening? (b) What product is formed when each compound undergoes a photochemical electrocyclic ring opening?



A **B**


27.28 What product is formed by the [3,3] sigmatropic rearrangement of each compound?


a. 


b. 

Types of Pericyclic Reactions

27.29 Classify each pericyclic reaction as an electrocyclic reaction, cycloaddition, or sigmatropic rearrangement. Indicate whether the stereochemistry is conrotatory, disrotatory, suprafacial, or antarafacial.


a. 


b. 

c. 

Electrocyclic Reactions


27.30 What product is formed when each compound undergoes thermal electrocyclic ring opening or ring closure? Label each process as conrotatory or disrotatory and clearly indicate the stereochemistry around tetrahedral stereogenic centers and double bonds.


a. 

b. 

27.31 What product is formed when each compound in Problem 27.30 undergoes photochemical electrocyclic reaction? Label each process as conrotatory or disrotatory and clearly indicate the stereochemistry around tetrahedral stereogenic centers and double bonds.

27.32 What cyclic product is formed when each decatetraene undergoes photochemical electrocyclic ring closure?


a. 

b. 


27.33 Draw the product of each electrocyclic reaction.

a. the thermal electrocyclic ring closure of (2*E*,4*Z*,6*Z*)-nona-2,4,6-triene
b. the photochemical electrocyclic ring closure of (2*E*,4*Z*,6*Z*)-nona-2,4,6-triene
c. the thermal electrocyclic ring opening of *cis*-5-ethyl-6-methylcyclohexa-1,3-diene
d. the photochemical electrocyclic ring opening of *trans*-5-ethyl-6-methylcyclohexa-1,3-diene

27.34 Consider the following electrocyclic ring closure. Does the product form by a conrotatory or disrotatory process? Would this reaction occur under photochemical or thermal conditions?




27.35 Draw the product formed when diene **M** undergoes disrotatory cyclization. Indicate the stereochemistry at new sp^3 hybridized carbons. Will the reaction occur under thermal or photochemical conditions?




M

27.36 (a) What product is formed when triene **N** undergoes thermal electrocyclic ring closure? (b) What product is formed when triene **N** undergoes photochemical ring closure? (c) Label each process as conrotatory or disrotatory.



N


27.37 The bicyclic alkene **P** can be prepared by thermal electrocyclic ring closure from cyclodecadiene **Q** or by photochemical electrocyclic ring closure from cyclodecadiene **R.** Draw the structures of **Q** and **R,** and indicate the stereochemistry of the process by which each reaction occurs.



P

Cycloaddition Reactions

27.38 What type of cycloaddition occurs in Reaction [1]? Draw the product of a similar process in Reaction [2]. Would you predict that these reactions occur under thermal or photochemical conditions?




27.39 Draw the product of each Diels–Alder reaction and indicate the stereochemistry at all stereogenic centers.



27.40 What cycloaddition products are formed in each reaction? Indicate the stereochemistry of each product.




27.41 What starting materials are needed to synthesize each compound by a thermal [4 + 2] cycloaddition?




27.42 Explain why heating buta-1,3-diene forms 4-vinylcyclohexene but not cycloocta-1,5-diene.

27.43 How can **X** be prepared from a constitutional isomer by a series of [2 + 2] cycloaddition reactions? Interest in molecules that contain several cyclobutane rings fused together has been fueled by the discovery of pentacycloanammoxic acid methyl ester, a lipid isolated from the membrane of organelles in the bacterium *Candidatus Brocadia anammoxidans.* The role of this unusual natural product is as yet unknown.




X




pentacycloanammoxic acid methyl ester

Sigmatropic Rearrangements

27.44 What type of sigmatropic rearrangement is illustrated in each reaction?




27.45 Draw the product of the [3,3] sigmatropic rearrangement of each compound.




27.46 A solution of 5-methylcyclopenta-1,3-diene rearranges at room temperature to a mixture containing 1-methyl-, 2-methyl-, and 5-methylcyclopenta-1,3-diene. (a) Show how both isomeric products are formed from the starting material by a sigmatropic rearrangement involving a C–H bond. (b) Explain why 2-methylcyclopenta-1,3-diene is not formed directly from 5-methylcyclopenta-1,3-diene by a [1,3] rearrangement.

27.47 What product is formed from the [5,5] sigmatropic rearrangement of the following unsaturated ether?




27.48 Heating **A** results in two successive [3,3] sigmatropic rearrangements—Claisen reaction followed by Cope reaction—to afford β-sinensal, a component of mandarin orange oil. What is the structure of β-sinensal?




A

27.49 Draw structures for **A, B,** and **C** in the following reaction sequence and identify the process that converts **B** to **C.**




General Pericyclic Reactions


27.50 What type of pericyclic reaction is illustrated in each reaction?



27.51 Draw the product formed (including stereochemistry) in each pericyclic reaction.




27.52 Draw the products of each reaction.




Mechanisms


27.53 When both carbons ortho to the aryl oxygen are not bonded to hydrogen, an allyl aryl ether rearranges to a para-substituted phenol. Draw a stepwise mechanism for the following reaction, which contains two [3,3] sigmatropic rearrangements.




27.54 Draw a stepwise, detailed mechanism for the following reaction.



27.55 Show how the following starting material is converted to the given product by a series of two pericyclic reactions. Account for the observed stereochemistry.




27.56 Use curved arrows to show how **E** is converted to **F** by a two-step reaction sequence consisting of a [1,5] sigmatropic rearrangement followed by a [4 + 2] cycloaddition.




E **F**

27.57 Show how the following starting material is converted to the given product by a series of two pericyclic reactions. Account for the observed stereochemistry.




27.58 Draw a stepwise, detailed mechanism for the following reaction.




Challenge Problems

27.59 What product is formed by [3,3] sigmatropic rearrangement of the following compound? Clearly indicate the stereochemistry around all tetrahedral stereogenic centers.




27.60 (a) What is the structure of **C,** which is formed by oxy-Cope rearrangement of **B** with NaOEt? (b) Draw a stepwise mechanism for the conversion of **C** to the bicyclic alcohol **D.**




B **D**

27.61 Draw a stepwise mechanism for the Carroll rearrangement, a reaction that prepares a γ,δ-unsaturated carbonyl compound from a β-keto ester and allylic alcohol in the presence of base.




27.62 The endiandric acids comprise a group of unsaturated carboxylic acids isolated from a tree that grows in the rain forests of eastern Australia. The methyl esters of endiandric acids D and E have been prepared from polyene **Y** by a series of two successive electrocyclic reactions: thermal ring closure of the conjugated tetraene followed by ring closure of the resulting conjugated triene. (a) Draw the structures (including stereochemistry) of the methyl esters of endiandric acids D and E. (b) The methyl ester of endiandric acid E undergoes an intramolecular [4 + 2] cycloaddition to form the methyl ester of endiandric acid A. Propose a possible structure for endiandric acid A.



Y


27.63 [4 + 2] Cycloadditions with *o*-quinones such as **B** are often complex because a variety of products are possible.

a. Draw arrows to illustrate how each product is formed when **B** reacts with styrene, and label the "diene" and "dienophile" components.




o-quinone **B** styrene

b. Draw arrows to illustrate how each product is formed when **B** reacts with buta-1,3-diene, and label the "diene" and "dienophile" components.



o-quinone **B** buta-1,3-diene


c. *o*-Quinone **C** reacts with diene **D** to form heterocycle **E** by a process that involves a cycloaddition followed by a [3,3] sigmatropic rearrangement. Use curved arrows to illustrate how this two-step sequence occurs. **E** is *not* formed directly from **C** by a Diels–Alder reaction.



C

28 Carbohydrates

28.1 Introduction
28.2 Monosaccharides
28.3 The family of D-aldoses
28.4 The family of D-ketoses
28.5 Physical properties of monosaccharides
28.6 The cyclic forms of monosaccharides
28.7 Glycosides
28.8 Reactions of monosaccharides at the OH groups
28.9 Reactions at the carbonyl group—Oxidation and reduction
28.10 Reactions at the carbonyl group—Adding or removing one carbon atom
28.11 Disaccharides
28.12 Polysaccharides
28.13 Other important sugars and their derivatives



The leaves, stems, and green spots on potatoes contain the toxin **solanine,** which is produced by the plant as a defense against insects and predators. Solanine is a carbohydrate derivative, formed from a complex amine and three monosaccharide rings, which are joined together by acetals called glycosides. The many hydroxy groups of the carbohydrate portion of solanine increase its water solubility, a useful feature in aqueous biological systems. In Chapter 28, we learn about carbohydrates and carbohydrate derivatives like solanine.

Chapters 28, 29, and 31 discuss ***biomolecules,*** **organic compounds found in biological systems.** You have already learned many facts about these compounds in previous chapters while you studied other organic compounds having similar properties. In Chapter 10 (Alkenes), for example, you learned that the presence of double bonds determines whether a fatty acid is part of a fat or an oil. In Chapter 19 (Carboxylic Acids and the Acidity of the O−H Bond), you learned that amino acids are the building blocks of proteins.

Chapter 28 focuses on carbohydrates, the largest group of biomolecules in nature, comprising ~50% of the earth's biomass. Chapter 29 concentrates on proteins (and the amino acids that compose them), whereas Chapter 31 explores lipids. These compounds are all organic molecules, so many of the same principles and chemical reactions that you have already studied will be examined once again. But, as you will see, each class of compound has its own unique features that we must learn as well.

28.1 Introduction


Carbohydrates were given their name because molecular formulas of simple carbohydrates could be written as $C_n(H_2O)_n$, making them **hydrates of carbon.**

Carbohydrates such as glucose and cellulose were discussed in Sections 5.1, 6.4, and 21.17.

Carbohydrates, commonly referred to as sugars and starches, are polyhydroxy aldehydes and ketones, or compounds that can be hydrolyzed to them. The cellulose in plant stems and tree trunks and the chitin in the exoskeletons of arthropods and mollusks are both complex carbohydrates. Four examples are shown in Figure 28.1. They include not only glucose and cellulose, but also doxorubicin (an anticancer drug) and 2'-deoxyadenosine 5'-monophosphate (a nucleotide base from DNA), both of which have a carbohydrate moiety as part of a larger molecule.

Carbohydrates are storehouses of chemical energy. They are synthesized in green plants and algae by **photosynthesis,** a process that uses the energy from the sun to convert carbon dioxide and water


Figure 28.1
Some examples of carbohydrates



These compounds illustrate the structural diversity of carbohydrates and their derivatives. **Glucose** is the most common simple sugar, whereas **cellulose,** which comprises wood, plant stems, and grass, is the most common carbohydrate in the plant world. **Doxorubicin,** an anticancer drug that has a carbohydrate ring as part of its structure, has been used in the treatment of leukemia, Hodgkin's disease, and cancers of the breast, bladder, and ovaries. **2'-Deoxyadenosine 5'-monophosphate** is one of the four nucleotides that form DNA.


Although the metabolism of lipids provides more energy per gram than the metabolism of carbohydrates, glucose is the preferred source when a burst of energy is needed during exercise. Glucose is water soluble, so it can be quickly and easily transported through the bloodstream to the tissues.

into glucose and oxygen. This energy is released when glucose is metabolized. The oxidation of glucose is a multistep process that forms carbon dioxide, water, and a great deal of energy (Section 6.4).



28.2 Monosaccharides

The simplest carbohydrates are called **monosaccharides** or **simple sugars. Monosaccharides have three to seven carbon atoms** in a chain, with a **carbonyl group** at either the terminal carbon (C1) or the carbon adjacent to it (C2). In most carbohydrates, each of the remaining carbon atoms has a **hydroxy group.** Monosaccharides are often drawn vertically, with the carbonyl group at the top. When this convention is used, monosaccharides look different from molecules encountered in prior chapters.




- **Monosaccharides with an aldehyde carbonyl group at C1 are called aldoses.**
- **Monosaccharides with a ketone carbonyl group at C2 are called ketoses.**



D-Fructose is almost twice as sweet as normal table sugar (sucrose) with about the same number of calories per gram. "Lite" food products use only half as much fructose as sucrose for the same level of sweetness, and so they have fewer calories.

Several examples of simple carbohydrates are shown. D-Glyceraldehyde and dihydroxyacetone have the same molecular formula, so they are **constitutional isomers,** as are D-glucose and D-fructose.





Dihydroxyacetone is the active ingredient in many artificial tanning agents.

All carbohydrates have common names. The simplest aldehyde, glyceraldehyde, and the simplest ketone, dihydroxyacetone, are the only monosaccharides whose names do not end in the suffix ***-ose.*** (The prefix "D-" is explained in Section 28.2C.)

A monosaccharide is called:

- **a triose if it has 3 C's;**
- **a tetrose if it has 4 C's;**
- **a pentose if it has 5 C's;**
- **a hexose if it has 6 C's, and so forth.**


These terms are then combined with the words *aldose* and *ketose* to indicate both the number of carbon atoms in the monosaccharide and whether it contains an aldehyde or ketone. Thus, glyceraldehyde is an aldotriose (three C atoms and an aldehyde), glucose is an aldohexose (six C atoms and an aldehyde), and fructose is a ketohexose (six C atoms and a ketone).

Problem 28.1 Draw the structure of (a) a ketotetrose; (b) an aldopentose; (c) an aldotetrose.

28.2A Fischer Projection Formulas

A striking feature of carbohydrate structure is the presence of stereogenic centers. **All carbohydrates except for dihydroxyacetone contain one or more stereogenic centers.**

The simplest aldehyde, glyceraldehyde, has one stereogenic center, so there are two possible **enantiomers.** Only the enantiomer with the *R* configuration occurs naturally.




(*R*)-glyceraldehyde
naturally occurring enantiomer

(*S*)-glyceraldehyde

The stereogenic centers in sugars are often depicted following a different convention than is usually seen for other stereogenic centers. Instead of drawing a tetrahedron with two bonds in the plane, one in front of the plane, and one behind it, the **tetrahedron is tipped so that horizontal bonds come forward (drawn on wedges) and vertical bonds go behind (on dashed wedges).** This structure is then abbreviated by a **cross formula,** also called a **Fischer projection formula.** In a Fischer projection formula:

- A carbon atom is located at the intersection of the two lines of the cross.
- The horizontal bonds come forward, on wedges.
- The vertical bonds go back, on dashed wedges.
- In a carbohydrate, the aldehyde or ketone carbonyl is put at or near the top.

Carbon atoms that are not stereogenic centers are generally drawn in. Using a Fischer projection formula, (*R*)-glyceraldehyde becomes:



Tip red bonds in the plane forward.


(*R*)-glyceraldehyde

- Horizontal bonds come forward.
- Vertical bonds go back.

Fischer projection formula
(*R*)-glyceraldehyde


Do not rotate a Fischer projection formula in the plane of the page, because you might inadvertently convert a compound into its enantiomer. When using Fischer projections it is usually best to convert them to structures with wedges and dashed wedges, and then manipulate them. Although a Fischer projection formula can be used for the stereogenic center in any compound, it is most commonly used for monosaccharides.

Sample Problem 28.1 Convert each compound to a Fischer projection formula.




Solution

Rotate and re-draw each molecule to place the horizontal bonds in front of the plane and the vertical bonds behind the plane. Then use a cross to represent the stereogenic center.



Problem 28.2 Draw each stereogenic center using a Fischer projection formula.




R,S designations can be assigned to any stereogenic center drawn as a Fischer projection formula in the following manner:

[1] Assign priorities ($1 \rightarrow 4$) to the four groups bonded to the stereogenic center using the rules detailed in Section 5.6.

[2] When the lowest priority group occupies a vertical bond—that is, it projects *behind* the plane on a dashed wedge—tracing a circle in the clockwise direction (from priority group $1 \rightarrow 2 \rightarrow 3$) gives the *R* configuration. Tracing a circle in the counterclockwise direction gives the *S* configuration.

[3] When the lowest priority group occupies a horizontal bond—that is, it projects *in front* of the plane on a wedge—reverse the answer obtained in Step [2] to designate the configuration.

Sample Problem 28.2 Re-draw each Fischer projection formula using wedges and dashed wedges for the stereogenic center, and label the center as *R* or *S*.




Solution


For each molecule:

[1] Convert the Fischer projection formula to a representation with wedges and dashed wedges.


[2] Assign priorities (Section 5.6).

[3] Determine *R* or *S* in the usual manner. Reverse the answer if priority group [4] is oriented forward (on a wedge).






Problem 28.3 Label each stereogenic center as *R* or *S*.




28.2B Monosaccharides with More Than One Stereogenic Center

The number of possible stereoisomers of a monosaccharide increases exponentially with the number of stereogenic centers present. **An aldohexose has four stereogenic centers, and so it has $2^4 = 16$ possible stereoisomers,** or eight pairs of enantiomers.




Fischer projection formulas are also used for compounds like aldohexoses that contain several stereogenic centers. In this case, the molecule is drawn with a vertical carbon skeleton and the stereogenic centers are stacked one above another. Using this convention, **all horizontal bonds project forward (on wedges).**




Although Fischer projections are commonly used to depict monosaccharides with many stereogenic centers, care must be exercised in using them, because they do not give a true picture of the three-dimensional structures they represent. Each stereogenic center is drawn in the less stable

Figure 28.2
A Fischer projection and the 3-D structure of glucose




eclipsed conformation, so the Fischer projection of glucose really represents the molecule in a cylindrical conformation, as shown in Figure 28.2.

Sample Problem 28.3 Convert the ball-and-stick model to a Fischer projection.




Solution

Since the ball-and-stick model is shown in the more stable staggered conformation, it must be converted to the less stable eclipsed conformation used in a Fischer projection. First, re-draw the model as a skeletal structure (**A**), and rotate it to place the carbonyl group at the top (**B**). To convert the all-staggered form to the all-eclipsed form, rotate around the bonds in **B** to swing two carbons (labeled in red) 180°, forming **C.** Re-draw **C** so that all bonds to H and OH on the four stereogenic centers are drawn on wedges, forming **D.** Groups that are on wedges in **C** (in red) are on the left side of the carbon skeleton in **D,** and groups on dashed wedges in **C** (in blue) are on the right side of the carbon skeleton in **D.** Finally, replace the wedges with crosses to form the Fischer projection.




Problem 28.4 Convert the ball-and-stick model to a Fischer projection.



Problem 28.5 Assign *R,S* designations to each stereogenic center in glucose.

28.2C D and L Monosaccharides

Although the prefixes *R* and *S* can be used to designate the configuration of stereogenic centers in monosaccharides, an older system of nomenclature uses the prefixes D- and L-, instead. Naturally occurring glyceraldehyde with the *R* configuration is called the **D-isomer.** Its enantiomer, (*S*)-glyceraldehyde, is called the **L-isomer.**




The letters **D** and **L** are used to label all monosaccharides, even those with multiple stereogenic centers. **The configuration of the stereogenic center *farthest* from the carbonyl group determines whether a monosaccharide is D- or L-.**

- **A D-sugar has the OH group on the stereogenic center farthest from the carbonyl on the right in a Fischer projection (like D-glyceraldehyde).**
- **An L-sugar has the OH group on the stereogenic center farthest from the carbonyl on the left in a Fischer projection (like L-glyceraldehyde).**


The two designations, D and *d,* refer to very different phenomena. The "D" designates the configuration around a stereogenic center in a monosaccharide. The "*d,*" on the other hand, is an abbreviation for "dextrorotatory"; that is, a *d*-compound rotates the plane of polarized light in the clockwise direction. A D-sugar may be dextrorotatory or it may be levorotatory. **There is no direct correlation between D and *d* or L and *l*.**



Glucose and all other naturally occurring sugars are D-sugars. L-Glucose, a compound that does not occur in nature, is the enantiomer of D-glucose. L-Glucose has the opposite configuration at *every* stereogenic center.



Problem 28.6 (a) Label compounds **A, B,** and **C** as D- or L-sugars. (b) How are compounds **A** and **B** related? **A** and **C? B** and **C?** Choose from enantiomers, diastereomers, or constitutional isomers.




28.3 The Family of D-Aldoses

Beginning with D-glyceraldehyde, one may formulate other D-aldoses having four, five, or six carbon atoms by adding carbon atoms (each bonded to H and OH), one at a time, between C1 and C2. **Two D-aldotetroses can be formed from D-glyceraldehyde,** one with the new OH


The common name of each monosaccharide indicates both the number of atoms it contains and the configuration at each of the stereogenic centers. Because the common names are firmly entrenched in the chemical literature, no systematic method has ever been established to name these compounds.

group on the right and one with the new OH group on the left. Their names are D-erythrose and D-threose. They are two diastereomers, each with two stereogenic centers, labeled in blue.



Because each aldotetrose has two stereogenic centers, there are 2^2 or four possible stereoisomers. D-Erythrose and D-threose are two of them. The other two are their enantiomers, called L-erythrose and L-threose, respectively. The configuration around each stereogenic center is exactly the opposite in its enantiomer. All four stereoisomers of the aldotetroses are shown in Figure 28.3.

Figure 28.3
The four stereoisomeric aldotetroses



D-Ribose, D-arabinose, and D-xylose are all common aldopentoses in nature. D-Ribose is the carbohydrate component of RNA, the polymer that translates the genetic information of DNA for protein synthesis.

To continue forming the family of D-aldoses, we must add another carbon atom (bonded to H and OH) just below the carbonyl of either tetrose. Because there are two D-aldotetroses to begin with, and there are two ways to place the new OH (right or left), there are now four D-aldopentoses: D-ribose, D-arabinose, D-xylose, and D-lyxose. Each aldopentose now has three stereogenic centers, so there are $2^3 = 8$ possible stereoisomers, or four pairs of enantiomers. The D-enantiomer of each pair is shown in Figure 28.4.

Finally, to form the D-aldohexoses, we must add another carbon atom (bonded to H and OH) just below the carbonyl of all the aldopentoses. Because there are four D-aldopentoses to begin with, and there are two ways to place the new OH (right or left), there are now eight D-aldohexoses. Each aldohexose now has four stereogenic centers, so there are $2^4 = 16$ possible stereoisomers, or eight pairs of enantiomers. Only the D-enantiomer of each pair is shown in Figure 28.4.

The tree of D-aldoses (Figure 28.4) is arranged in pairs of compounds that are bracketed together. Each pair of compounds, such as D-glucose and D-mannose, has the same configuration around all of its stereogenic centers except for one.

Of the D-aldohexoses, only D-glucose and D-galactose are common in nature. **D-Glucose is by far the most abundant of all D-aldoses.** D-Glucose comes from the hydrolysis of starch and cellulose, and D-galactose comes from the hydrolysis of fruit pectins.

- **Two diastereomers that differ in the configuration around one stereogenic center only are called *epimers*.**





Figure 28.4
The family of D-aldoses having three to six carbon atoms



Problem 28.7 How many different aldoheptoses are there? How many are D-sugars? Draw all D-aldoheptoses having the *R* configuration at C2 and C3.

Problem 28.8 Draw two possible epimers of D-erythrose. Name each of these compounds using Figure 28.4.

28.4 The Family of D-Ketoses

The family of D-ketoses, shown in Figure 28.5, is formed from dihydroxyacetone by adding a new carbon (bonded to H and OH) between C2 and C3. Having a carbonyl group at C2 decreases the number of stereogenic centers in these monosaccharides, so that there are only four D-ketohexoses. The most common naturally occurring ketose is D-fructose.


Problem 28.9 Referring to the structures in Figures 28.4 and 28.5, classify each pair of compounds as enantiomers, epimers, diastereomers but not epimers, or constitutional isomers of each other.

a. D-allose and L-allose
b. D-altrose and D-gulose
c. D-galactose and D-talose
d. D-mannose and D-fructose
e. D-fructose and D-sorbose
f. L-sorbose and L-tagatose

Problem 28.10
a. Draw the enantiomer of D-fructose.
b. Draw an epimer of D-fructose at C4. What is the name of this compound?
c. Draw an epimer of D-fructose at C5. What is the name of this compound?

Problem 28.11 Referring to Figure 28.5, which D-ketohexoses have the *S* configuration at C3?

Figure 28.5
The family of D-ketoses having three to six carbon atoms




28.5 Physical Properties of Monosaccharides

Monosaccharides have the following physical properties:

- They are all **sweet tasting,** but their relative sweetness varies a great deal.
- They are polar compounds with **high melting points.**
- The presence of so many polar functional groups capable of hydrogen bonding makes them **water soluble.**
- Unlike most other organic compounds, monosaccharides are so polar that they are **insoluble in organic solvents like diethyl ether.**

28.6 The Cyclic Forms of Monosaccharides


Although the monosaccharides in Figures 28.4 and 28.5 are drawn as acyclic carbonyl compounds containing several hydroxy groups, the hydroxy and carbonyl groups of monosaccharides can undergo intramolecular cyclization reactions to form **hemiacetals** having either five or six atoms in the ring. This process was first discussed in Section 21.16.



- A six-membered ring containing an O atom is called a *pyranose* ring.
- A five-membered ring containing an O atom is called a *furanose* ring.

Cyclization of a hydroxy carbonyl compound always forms a stereogenic center at the hemiacetal carbon, called the **anomeric carbon.** The two hemiacetals are called **anomers.**


- Anomers are stereoisomers of a cyclic monosaccharide that differ in the position of the OH group at the hemiacetal carbon.



Cyclization forms the more stable ring size in a given molecule. **The most common monosaccharides, the aldohexoses like glucose, typically form a pyranose ring,** so our discussion begins with forming a cyclic hemiacetal from D-glucose.


28.6A Drawing Glucose as a Cyclic Hemiacetal

Which of the five OH groups in glucose is at the right distance from the carbonyl group to form a six-membered ring? The **O atom on the stereogenic center farthest from the carbonyl** (C5) is six atoms from the carbonyl carbon, placing it in the proper position for cyclization to form a pyranose ring.




To translate the acyclic form of glucose into a cyclic hemiacetal, we must draw the hydroxy aldehyde in a way that suggests the position of the atoms in the new ring, and then draw the ring. **By convention the O atom in the new pyranose ring is drawn in the upper right-hand corner of the six-membered ring.**

Rotating the groups on the bottom stereogenic center in **A** places all six atoms needed for the ring (including the OH) in a vertical line (**B**). Re-drawing this representation as a Fischer projection makes the structure appear less cluttered (**C**). Twisting this structure and rotating it 90° forms **D.** Structures **A–D** are four different ways of drawing the same acyclic structure of D-glucose.



We are now set to draw the cyclic hemiacetal formed by nucleophilic attack of the OH group on C5 on the aldehyde carbonyl. Because cyclization creates a new stereogenic center, there are **two cyclic forms of D-glucose,** an **α anomer** and a **β anomer.** All the original stereogenic centers maintain their configuration in both of the products formed.

- **The α anomer of a D monosaccharide has the OH group drawn *down,* trans to the CH_2OH group at C5. The α anomer of D-glucose is called α-D-glucose, or α-D-glucopyranose (to emphasize the six-membered ring).**
- **The β anomer of a D monosaccharide has the OH group drawn *up,* cis to the CH_2OH group at C5. The β anomer is called β-D-glucose, or β-D-glucopyranose (to emphasize the six-membered ring).**




The **α anomer** in any monosaccharide has the anomeric OH group and the CH_2OH group **trans.** The **β anomer** has the anomeric OH group and the CH_2OH group **cis.**

These flat, six-membered rings used to represent the cyclic hemiacetals of glucose and other sugars are called **Haworth projections.** The cyclic forms of glucose now have **five stereogenic centers, the four from the starting hydroxy aldehyde and the new anomeric carbon.** α-D-Glucose and β-D-glucose are **diastereomers,** because only the anomeric carbon has a different configuration.

The mechanism for this transformation is exactly the same as the mechanism that converts a hydroxy aldehyde to a cyclic hemiacetal (Mechanism 21.10). The acyclic aldehyde and two cyclic hemiacetals are all in equilibrium. Each cyclic hemiacetal can be isolated and crystallized separately, but when any one compound is placed in solution, an equilibrium mixture of all three forms results. This process is called **mutarotation.** At equilibrium, the mixture has 37% of the α anomer, 63% of the β anomer, and only trace amounts of the acyclic hydroxy aldehyde (Figure 28.6). Also shown are representations of the three forms of glucose using wedges and dashed wedges.

Figure 28.6
The three forms of glucose



- Bonds above the ring in a Haworth projection are drawn as wedged bonds.
- Bonds below the ring in a Haworth projection are drawn as dashed wedges.

Problem 28.12 Label each Haworth projection as an α or β anomer and convert the Haworth projection to a six-membered ring with wedges and dashed wedges.




28.6B Haworth Projections

To convert an acyclic monosaccharide to a Haworth projection, follow a stepwise procedure.

How To Draw a Haworth Projection from an Acyclic Aldohexose

Example Convert D-mannose to a Haworth projection.




—Continued

How To, continued . . .


Step [1] **Place the O atom in the upper right corner of a hexagon, and add the CH_2OH group on the first carbon counterclockwise from the O atom.**

- For **D-sugars,** the CH_2OH group is drawn **up.** For **L-sugars,** the CH_2OH group is drawn **down.**



D-sugar


D-mannose




CH_2OH is drawn up.

Step [2] **Place the anomeric carbon on the first carbon clockwise from the O atom.**

- For an α **anomer,** the **OH** is drawn **down** in a D-sugar.
- For a β **anomer,** the **OH** is drawn **up** in a D-sugar.






α anomer β anomer

The anomeric carbon is a new stereogenic center at C1.

- Remember: The carbonyl carbon becomes the anomeric carbon (a new stereogenic center).

Step [3] **Add the substituents at the three remaining stereogenic centers clockwise around the ring.**


- The substituents on the **right side** of the Fischer projection are drawn **down.**
- The substituents on the **left** are drawn **up.**




α anomer β anomer

Problem 28.13 Convert each aldohexose to the indicated anomer using a Haworth projection.


a. Draw the α anomer of:



b. Draw the α anomer of:




c. Draw the β anomer of:



Sample Problem 28.4 shows how to convert a Haworth projection back to the acyclic form of a monosaccharide. It doesn't matter whether the hemiacetal is the α or β anomer, because both anomers give the *same* hydroxy aldehyde.


Sample Problem 28.4 Convert the following Haworth projection to the acyclic form of the aldohexose.



Solution


To convert the substituents to the acyclic form, start at the pyranose O atom, and work in a ***counterclockwise*** fashion around the ring, and from bottom-to-top along the chain.

[1] Draw the carbon skeleton, placing the CHO on the top and the CH_2OH on the bottom.



[2] Classify the sugar as D- or L-.


- The CH_2OH is drawn **up,** so it is a **D-sugar.**
- A D-sugar has the OH group on the bottom stereogenic center on the **right.**



[3] Add the three other stereogenic centers.


- **Up** substituents go on the **left.**
- **Down** substituents go on the **right.**

Answer:



- The anomeric carbon becomes the C=O at C1.

Problem 28.14 Convert each Haworth projection to its acyclic form.




28.6C Three-Dimensional Representations for D-Glucose

Because the chair form of a six-membered ring gives the truest picture of its three-dimensional shape, we must learn to convert Haworth projections into chair forms.

To convert a Haworth projection to a chair form:

- Draw the pyranose ring with the O atom as an "up" atom.
- The "up" substituents in a Haworth projection become the "up" bonds (either axial or equatorial) on a given carbon atom on a puckered six-membered ring.
- The "down" substituents in a Haworth projection become the "down" bonds (either axial or equatorial) on a given carbon atom on a puckered six-membered ring.

As a result, the three-dimensional chair form of β-D-glucose is drawn in the following manner:




chair form for β-D-glucose

- "Up" substituents are labeled in red.
- "Down" substituents are labeled in blue.

Glucose has all substituents larger than a hydrogen atom in the more roomy equatorial positions, making it the most stable and thus most prevalent monosaccharide. The β anomer is the major isomer at equilibrium, moreover, because the hemiacetal OH group is in the equatorial position, too. Figure 28.7 shows both anomers of D-glucose drawn as chair conformations.

Problem 28.15 Convert each Haworth projection in Problem 28.14 to a three-dimensional representation using a chair pyranose ring.

Figure 28.7 Three-dimensional representations for both anomers of D-glucose



α anomer β anomer

28.6D Furanoses


Certain monosaccharides—notably aldopentoses and ketohexoses—predominantly form furanose rings, rather than pyranose rings, in solution. The same principles apply to drawing these structures as for drawing pyranose rings, except the ring size is one atom smaller.

- Cyclization always forms a new stereogenic center at the anomeric carbon, so two different anomers are possible. For a D-sugar, the OH group is drawn down in the α anomer and up in the β anomer.
- Use the same drawing conventions for adding substituents to the five-membered ring. With D-sugars, the CH_2OH group is drawn up.




Honey was the first and most popular sweetening agent until it was replaced by sugar (from sugarcane) in modern times. Honey is a mixture consisting largely of D-fructose and D-glucose.

With D-ribose, the OH group used to form the five-membered furanose ring is located on C4. Cyclization yields two anomers at the new stereogenic center, which are called **α-D-ribofuranose** and **β-D-ribofuranose.**



The same procedure can be used to draw the furanose form of D-fructose, the most common ketohexose. Because the carbonyl group is at C2 (instead of C1, as in the aldoses), the OH group at C5 reacts to form the hemiacetal in the five-membered ring. Two anomers are formed.



Problem 28.16 Aldotetroses exist in the furanose form. Draw both anomers of D-erythrose.

28.7 Glycosides

Keep in mind the difference between a hemiacetal and an acetal:



hemiacetal
- one OH group
- one OR group



acetal
- two OR groups

Because monosaccharides exist in solution in an equilibrium between acyclic and cyclic forms, they undergo three types of reactions:


- **Reaction of the hemiacetal**
- **Reaction of the hydroxy groups**
- **Reaction of the carbonyl group**

Even though the acyclic form of a monosaccharide may be present in only trace amounts, the equilibrium can be tipped in its favor by Le Châtelier's principle (Section 9.8). Suppose, for example, that the carbonyl group of the acyclic form reacts with a reagent, thus depleting its equilibrium concentration. The equilibrium will then shift to compensate for the loss, thus producing more of the acyclic form, which can react further.

Note, too, that monosaccharides have two different types of OH groups. Most are "regular" alcohols, and as such, undergo reactions characteristic of alcohols. The anomeric OH group, on the other hand, is part of a hemiacetal, giving it added reactivity.

28.7A Glycoside Formation

Treatment of a monosaccharide with an alcohol and HCl converts the hemiacetal into an acetal called a **glycoside.** For example, treatment of α-D-glucose with CH_3OH and HCl forms two glycosides that are diastereomers at the acetal carbon. The α and β labels are assigned in the same way as anomers: with a D-sugar, an α glycoside has the new OR group (OCH_3 group in this example) down, and a β glycoside has the new OR group up.



α-D-glucose — α glycoside — β glycoside

Only the hemiacetal OH reacts.

Mechanism 28.1 explains why a single anomer forms two glycosides. The reaction proceeds by way of a **planar carbocation,** which undergoes nucleophilic attack from two different directions to give a mixture of diastereomers. Because both α- and β-D-glucose form the same planar carbocation, each yields the same mixture of two glycosides.

The mechanism also explains why only the hemiacetal OH group reacts. Protonation of the hemiacetal OH, followed by loss of H_2O, forms a resonance-stabilized carbocation in Step [2]. A resonance-stabilized carbocation is not formed by loss of H_2O from any other OH group.


Unlike cyclic hemiacetals, **glycosides are acetals, and so they do not undergo mutarotation.** When a single glycoside is dissolved in H_2O, it is *not* converted to an equilibrium mixture of α and β glycosides.

- **Glycosides are acetals with an alkoxy group (OR) bonded to the anomeric carbon.**

Problem 28.17 What glycosides are formed when each monosaccharide is treated with CH_3CH_2OH, HCl:
(a) β-D-mannose; (b) α-D-gulose; (c) β-D-fructose?


Mechanism 28.1 Glycoside Formation

Part [1] Loss of H_2O from the hemiacetal



1 – 2 Protonation of the hemiacetal OH followed by loss of H_2O forms a resonance-stabilized carbocation.


Part [2] Formation of the glycosides



3 – 4 Nucleophilic attack by CH_3OH occurs from both sides of the planar carbocation to yield α and β glycosides after loss of a proton.

28.7B Glycoside Hydrolysis


Because glycosides are acetals, **they are hydrolyzed with acid and water to cyclic hemiacetals and a molecule of alcohol.** A mixture of two anomers is formed from a single glycoside. For example, treatment of methyl α-D-glucopyranoside with aqueous acid forms a mixture of α- and β-D-glucose and methanol.



methyl α-D-glucopyranoside α-D-glucose β-D-glucose


The mechanism for glycoside hydrolysis is just the reverse of glycoside formation. It involves two parts: **formation of a planar carbocation,** followed by **nucleophilic attack of H_2O** to form anomeric hemiacetals, as shown in Mechanism 28.2.

Problem 28.18 Draw a stepwise mechanism for the following reaction.




Mechanism 28.2 Glycoside Hydrolysis

Part [1] Loss of CH_3OH from the glycoside



1 – 2 Protonation of the acetal OCH_3 followed by loss of CH_3OH forms a resonance-stabilized carbocation.

Part [2] Formation of the hemiacetals




3 – 4 Nucleophilic attack by H_2O occurs from both sides of the planar carbocation to yield α and β anomers after loss of a proton.

28.7C Naturally Occurring Glycosides




The berries of the black nightshade plant (*Solanum nigrum*) are a source of the poisonous alkaloid solanine.

Salicin and **solanine** are two naturally occurring compounds that contain glycoside bonds as part of their structure. Salicin is an analgesic isolated from willow bark, and solanine, the chapter-opening molecule, is a poisonous compound produced in the leaves, stem, and green spots on the skin of potatoes. Solanine is also isolated from the berries of the deadly nightshade plant. It is believed that the role of the sugar rings in both salicin and solanine is to increase their water solubility.



salicin

[The O atoms that are part of the glycoside bonds are drawn in red.]




solanine

Glycosides are common in nature. All disaccharides and polysaccharides are formed by joining monosaccharides together with glycosidic linkages. These compounds are discussed in detail beginning in Section 28.11.

Problem 28.19 (a) Label all the O atoms that are part of a glycoside in rebaudioside A. Rebaudioside A, marketed under the trade name Truvia, is a sweet glycoside obtained from the stevia plant, which has been used for centuries in Paraguay to sweeten foods. (b) The alcohol or phenol formed from the hydrolysis of a glycoside is called an **aglycon.** What aglycon and monosaccharides are formed by the hydrolysis of rebaudioside A?



Rebaudioside A, a naturally occurring glycoside about 400 times sweeter than table sugar, is obtained from the leaves of the stevia plant, a shrub native to Central and South America.




rebaudioside A
Trade name Truvia

28.8 Reactions of Monosaccharides at the OH Groups

Because monosaccharides contain OH groups, they undergo reactions typical of alcohols—that is, they are converted to **ethers** and **esters.** Because the cyclic hemiacetal form of a monosaccharide contains an OH group, this form of a monosaccharide must be drawn as the starting material for any reaction that occurs at an OH group.

All OH groups of a cyclic monosaccharide are converted to ethers by treatment with base and an alkyl halide. For example, α-D-glucose reacts with silver(I) oxide (Ag_2O, a base) and excess CH_3I to form a pentamethyl ether.



hemiacetal OH in red


pentamethyl ether

- The acetal OCH_3 group is in red.
- Ether OCH_3 groups are in blue.


Ag_2O removes a proton from each alcohol, forming an alkoxide (RO^-), which then reacts with CH_3I in an S_N2 reaction. Because no C−O bonds are broken, the configuration of all substituents in the starting material is **retained,** forming a single product.

The product contains two different types of ether bonds. There are four "regular" ethers formed from the "regular" hydroxyls. The new ether from the hemiacetal is now part of an **acetal**—that is, a **glycoside.**

The four ether bonds that are *not* part of the acetal do not react with any reagents except strong acids like HBr and HI (Section 9.14). **The acetal ether, on the other hand, is hydrolyzed with aqueous acid** (Section 28.7B). Aqueous hydrolysis of a single glycoside (like the pentamethyl ether of α-D-glucose) yields both anomers of the product monosaccharide.





The OH groups of monosaccharides can also be converted to esters. For example, treatment of β-D-glucose with either acetic anhydride or acetyl chloride in the presence of pyridine (a base) converts all OH groups into acetate esters.






Since it is cumbersome and tedious to draw in all the atoms of the esters, the abbreviation **Ac** is used for the acetyl group, **$CH_3C{=}O$.** The esterification of β-D-glucose can then be written as follows:







Monosaccharides are so polar that they are insoluble in common organic solvents, making them difficult to isolate and use in organic reactions. Monosaccharide derivatives that have five ether or ester groups in place of the OH groups, however, are readily soluble in organic solvents.

Problem 28.20 Draw the products formed when β-D-galactose is treated with each reagent.

a. $Ag_2O + CH_3I$
b. $NaH + C_6H_5CH_2Cl$
c. The product in (b), then H_3O^+
d. Ac_2O + pyridine
e. C_6H_5COCl + pyridine
f. The product in (c), then C_6H_5COCl + pyridine


28.9 Reactions at the Carbonyl Group—Oxidation and Reduction

Oxidation and reduction reactions occur at the carbonyl group of monosaccharides, so they all begin with the monosaccharide drawn in the acyclic form. We will confine our discussion to aldoses as starting materials.

28.9A Reduction of the Carbonyl Group

Glucitol occurs naturally in some fruits and berries. It is sometimes used as a substitute for sucrose (table sugar). With six polar OH groups capable of hydrogen bonding, glucitol is readily hydrated. It is used as an additive to prevent certain foods from drying out.

Like other aldehydes, the **carbonyl group of an aldose is reduced to a 1° alcohol using $NaBH_4$.** This alcohol is called an **alditol.** For example, reduction of D-glucose with $NaBH_4$ in CH_3OH yields glucitol (also called sorbitol).




D-glucose glucitol (sorbitol)

Problem 28.21 A 2-ketohexose is reduced with $NaBH_4$ in CH_3OH to form a mixture of D-galactitol and D-talitol. What is the structure of the 2-ketohexose?

28.9B Oxidation of Aldoses

Aldoses contain 1° and 2° alcohols and an aldehyde, all of which are oxidizable functional groups. Two different types of oxidation reactions are particularly useful—oxidation of the aldehyde to a carboxylic acid (an **aldonic acid**) and oxidation of both the aldehyde and the 1° alcohol to a diacid (an **aldaric acid**).




aldose aldonic acid aldaric acid

[1] Oxidation of the aldehyde to a carboxylic acid

The aldehyde carbonyl is the most easily oxidized functional group in an aldose, and so a variety of reagents oxidize it to a carboxy group, forming an **aldonic acid.**

Three reagents used for this process produce a characteristic color change because the oxidizing agent is reduced to a colored product that is easily visible. As described in Section 20.8, **Tollens reagent** oxidizes aldehydes to carboxylic acids using Ag_2O in NH_4OH, and forms a mirror of Ag as a by-product. **Benedict's** and **Fehling's reagents** use a blue Cu^{2+} salt as an oxidizing agent, which is reduced to Cu_2O, a brick-red solid. Unfortunately, none of these reagents gives a high yield of aldonic acid. When the aldonic acid is needed to carry on to other reactions, **Br_2 + H_2O** is used as the oxidizing agent.




D-glucose D-gluconic acid

- Any carbohydrate that exists as a *hemiacetal* is in equilibrium with a small amount of acyclic aldehyde, so it is oxidized to an aldonic acid.
- Glycosides are acetals, not hemiacetals, so they are *not* oxidized to aldonic acids.


Carbohydrates that can be oxidized with Tollens, Benedict's, or Fehling's reagent are called **reducing sugars.** Those that do not react with these reagents are called **nonreducing sugars.** Figure 28.8 shows examples of reducing and nonreducing sugars.

Figure 28.8
Examples of reducing and nonreducing sugars




- Carbohydrates containing a hemiacetal are in equilibrium with an acyclic aldehyde, making them reducing sugars.
- Glycosides are acetals, so they are *not* in equilibrium with any acyclic aldehyde, making them nonreducing sugars.

Problem 28.22 Classify each compound as a reducing or nonreducing sugar.



[2] Oxidation of both the aldehyde and 1° alcohol to a diacid


Both the aldehyde and 1° alcohol of an aldose are oxidized to carboxy groups by treatment with warm nitric acid, forming an **aldaric acid.** Under these conditions, D-glucose is converted to D-glucaric acid.



D-glucose

D-glucaric acid
an aldaric acid

Because aldaric acids have identical functional groups on both terminal carbons, some aldaric acids contain a plane of symmetry, making them achiral molecules. For example, oxidation of D-allose forms an achiral, optically inactive aldaric acid. This contrasts with D-glucaric acid formed from glucose, which has no plane of symmetry, and is thus still optically active.




Problem 28.23 Draw the products formed when D-arabinose is treated with each reagent: (a) Ag_2O, NH_4OH; (b) Br_2, H_2O; (c) HNO_3, H_2O.

Problem 28.24 Which aldoses are oxidized to optically inactive aldaric acids: (a) D-erythrose; (b) D-lyxose; (c) D-galactose?

28.10 Reactions at the Carbonyl Group—Adding or Removing One Carbon Atom


Two common procedures in carbohydrate chemistry result in adding or removing one carbon atom from the skeleton of an aldose. The **Wohl degradation** shortens an aldose chain by one carbon, whereas the **Kiliani–Fischer synthesis** lengthens it by one. Both reactions involve cyanohydrins as intermediates. Recall from Section 21.9 that cyanohydrins are formed from aldehydes by addition of the elements of HCN. Cyanohydrins can also be re-converted to carbonyl compounds by treatment with base.




- Forming a cyanohydrin adds one carbon to a carbonyl group.
- Re-converting a cyanohydrin to a carbonyl compound removes one carbon.

28.10A The Wohl Degradation

The Wohl degradation is a stepwise procedure that shortens the length of an aldose chain by cleavage of the C1−C2 bond. As a result, an aldohexose is converted to an aldopentose having the same configuration at its bottom three stereogenic centers (C3−C5). For example, the Wohl degradation converts D-glucose into D-arabinose.



The Wohl degradation consists of three steps, illustrated here beginning with D-glucose.




[1] Treatment of D-glucose with hydroxylamine (NH_2OH) forms an **oxime** by nucleophilic addition. This reaction is analogous to the formation of imines discussed in Section 21.11.

[2] Dehydration of the oxime to a nitrile occurs with acetic anhydride (Ac_2O) and sodium acetate (NaOAc). The nitrile product is a cyanohydrin.

[3] **Treatment of the cyanohydrin with base results in loss of the elements of HCN to form an aldehyde having one fewer carbon.**


The Wohl degradation converts a stereogenic center at C2 in the original aldose to an sp^2 hybridized C=O. As a result, a pair of aldoses that are epimeric at C2, such as D-galactose and D-talose, yield the same aldose (D-lyxose, in this case) upon Wohl degradation.




Problem 28.25 What two aldoses yield D-xylose on Wohl degradation?

28.10B The Kiliani–Fischer Synthesis

The Kiliani–Fischer synthesis lengthens a carbohydrate chain by adding one carbon to the aldehyde end of an aldose, thus forming a new stereogenic center at C2 of the product. The product consists of epimers that differ only in their configuration about the one new stereogenic center. For example, the Kiliani–Fischer synthesis converts D-arabinose into a mixture of D-glucose and D-mannose.



The Kiliani–Fischer synthesis, shown here beginning with D-arabinose, consists of three steps. "Squiggly" lines are meant to indicate that two different stereoisomers are formed at the new stereogenic center. As with the Wohl degradation, the key intermediate is a cyanohydrin.



[1] Treating an aldose with NaCN and HCl adds the elements of HCN to the carbonyl group, forming a **cyanohydrin** and a new carbon–carbon bond. Because the sp^2 hybridized carbonyl carbon is converted to an sp^3 hybridized carbon with four different groups, a new stereogenic center is formed in this step.

[2] Reduction of the nitrile with H_2 and Pd-$BaSO_4$, a poisoned Pd catalyst, forms an **imine.**

[3] Hydrolysis of the imine with aqueous acid forms an aldehyde that has one more carbon than the aldose that began the sequence.

Note that the **Wohl degradation and the Kiliani–Fischer synthesis are conceptually opposite transformations.**

- **The Wohl degradation *removes* a carbon atom from the aldehyde end of an aldose. Two aldoses that are epimers at C2 form the same product.**
- **The Kiliani–Fischer synthesis *adds* a carbon to the aldehyde end of an aldose, forming two epimers at C2.**

Problem 28.26 What aldoses are formed when the following aldoses are subjected to the Kiliani–Fischer synthesis: (a) D-threose; (b) D-ribose; (c) D-galactose?

28.10C Determining the Structure of an Unknown Monosaccharide

The reactions in Sections 28.9–28.10 can be used to determine the structure of an unknown monosaccharide, as shown in Sample Problem 28.5.

Sample Problem 28.5 A D-aldopentose **A** is oxidized to an optically inactive aldaric acid with HNO_3. **A** is formed by the Kiliani–Fischer synthesis of a D-aldotetrose **B,** which is also oxidized to an optically inactive aldaric acid with HNO_3. What are the structures of **A** and **B?**


Solution

Use each fact to determine the relative orientation of the OH groups in the D-aldopentose.

Fact [1] **A D-aldopentose A is oxidized to an optically *inactive* aldaric acid with HNO_3.**


An optically inactive aldaric acid must contain a **plane of symmetry.** Because the OH group on C4 must be on the right for the D-sugar, there are only two ways to arrange the OH groups in a five-carbon D-aldaric acid. Thus, only two structures are possible for **A,** labeled **A'** and **A''.**

Possible optically inactive D-aldaric acids:




Fact [2] **A is formed by the Kiliani–Fischer synthesis from a D-aldotetrose B.**

A' and **A''** are each prepared from a D-aldotetrose (**B'** and **B''**) that has the same configuration at the bottom two stereogenic centers.




Fact [3] **The D-aldotetrose is oxidized to an optically *inactive* aldaric acid upon treatment with HNO_3.**

Only the aldaric acid from **B'** has a plane of symmetry, making it optically inactive. Thus, **B'** is the correct structure for the D-aldotetrose **B,** and therefore **A'** is the structure of the D-aldopentose **A.**



Answer:




Problem 28.27 D-Aldopentose **A** is oxidized to an optically inactive aldaric acid. On Wohl degradation, **A** forms an aldotetrose **B** that is oxidized to an optically active aldaric acid. What are the structures of **A** and **B?**


Problem 28.28 A D-aldohexose **A** is formed from an aldopentose **B** by the Kiliani–Fischer synthesis. Reduction of **A** with $NaBH_4$ forms an optically inactive alditol. Oxidation of **B** forms an optically active aldaric acid. What are the structures of **A** and **B?**

28.11 Disaccharides

Disaccharides contain two monosaccharides joined together by a glycosidic linkage. The **general features of a disaccharide** include the following:



disaccharide
glycosidic linkage in red
acetal carbon labeled in blue



1⟶4-β-glycosidic linkage

[1] Two monosaccharide rings may be five- or six-membered, but six-membered rings are much more common. The two rings are connected by an O atom that is part of an acetal, called a **glycosidic linkage,** which may be oriented α or β.

[2] The **glycoside is formed from the anomeric carbon of one monosaccharide and any OH group on the other monosaccharide.** All disaccharides have **one acetal,** together with either a hemiacetal or another acetal.

[3] With pyranose rings, the carbon atoms in each ring are numbered beginning with the anomeric carbon. The most common disaccharides contain two monosaccharides in which the hemiacetal carbon of one ring (Cl) is joined to C4 of the other ring.


The three most abundant disaccharides are **maltose, lactose,** and **sucrose.**

28.11A Maltose




Maltose gets its name from malt, the liquid obtained from barley and other cereal grains.

Maltose, a disaccharide formed by the hydrolysis of starch, is found in germinated grains such as barley. Maltose contains two glucose units joined together by a 1→4-α-glycoside bond. Maltose contains one acetal carbon (in red) and one hemiacetal carbon (in blue).



Because one glucose ring of maltose still contains a hemiacetal, it exists as a mixture of α and β anomers. Only the β anomer is shown. Maltose exhibits two properties of all carbohydrates that contain a hemiacetal: it undergoes **mutarotation,** and it reacts with oxidizing agents, making it a **reducing sugar.**

Hydrolysis of maltose forms two molecules of glucose. The **C1−O** bond is cleaved in this process, and a mixture of glucose anomers forms. The mechanism for this hydrolysis is exactly the same as the mechanism for glycoside hydrolysis in Section 28.7B.




Problem 28.29 Draw the α anomer of maltose. What products are formed on hydrolysis of this form of maltose?

28.11B Lactose



Milk contains the disaccharide lactose.

Lactose is the principal disaccharide found in milk from both humans and cows. Unlike many mono- and disaccharides, lactose is not appreciably sweet. Lactose consists of **one galactose** and **one glucose unit,** joined by a **1→4-β-glycoside bond** from the anomeric carbon of galactose to C4 of glucose.



Like maltose, lactose also contains a hemiacetal, so it exists as a mixture of α and β anomers. The β anomer is drawn. Lactose undergoes **mutarotation,** and it reacts with oxidizing agents, making it a **reducing sugar.**


Lactose is digested in the body by first cleaving the 1→4-β-glycoside bond using the enzyme *lactase.* Many individuals, mainly of Asian and African descent, lack adequate amounts of lactase, so they are unable to digest and absorb lactose. This condition, lactose intolerance, is associated with abdominal cramping and recurrent diarrhea when milk and dairy products are ingested.

Problem 28.30 Cellobiose, a disaccharide obtained by the hydrolysis of cellulose, is composed of two glucose units joined together in a 1→4-β-glycoside bond. What is the structure of cellobiose?

28.11C Sucrose

Sucrose, the disaccharide found in sugarcane and used as table sugar (Figure 28.9), is the most common disaccharide in nature. It contains **one glucose unit** and **one fructose unit.**

Figure 28.9 Sucrose




The structure of sucrose has several features that make it different from maltose and lactose. First of all, sucrose contains one six-membered ring (glucose) and one five-membered ring (fructose), whereas both maltose and lactose contain two six-membered rings. In sucrose the six-membered glucose ring is joined by an α-glycosidic bond to C2 of a fructofuranose ring. The numbering in a fructofuranose is different from the numbering in a pyranose ring. The anomeric carbon is now designated as C2, so the anomeric carbons of the glucose and fructose rings are both used to form the glycosidic linkage.

As a result, **sucrose contains two acetals but no hemiacetal.** Sucrose, therefore, is a **nonreducing sugar** and **it does not undergo mutarotation.**

Sucrose's pleasant sweetness has made it a widely used ingredient in baked goods, cereals, bread, and many other products. It is estimated that the average American ingests 100 lb of sucrose annually. Like other carbohydrates, however, sucrose contains many calories. To reduce caloric intake while maintaining sweetness, a variety of artificial sweeteners have been developed. These include sucralose, aspartame, and saccharin (Figure 28.10). These compounds are much sweeter than sucrose so only a small amount of each compound is needed to achieve the same level of perceived sweetness.

Figure 28.10
Artificial sweeteners



The sweetness of these three artificial sweeteners was discovered accidentally. The sweetness of sucralose was discovered in 1976 when a chemist misunderstood his superior, and so he *tasted* rather than *tested* his compound. Aspartame was discovered in 1965 when a chemist licked his dirty fingers in the lab and tasted its sweetness. Saccharin, the oldest known artificial sweetener, was discovered in 1879 by a chemist who failed to wash his hands after working in the lab. Saccharin was not used extensively until sugar shortages occurred during World War I. Although there were concerns in the 1970s that saccharin causes cancer, there is no proven link between cancer occurrence and saccharin intake at normal levels.


28.12 Polysaccharides

Polysaccharides contain three or more monosaccharides joined together. Three prevalent polysaccharides in nature are **cellulose, starch,** and **glycogen,** each of which consists of repeating glucose units joined by different glycosidic bonds.

28.12A Cellulose

The structure of cellulose was discussed previously in Section 5.1.

Cellulose is found in the cell walls of nearly all plants, where it gives support and rigidity to wood and plant stems. Cotton is essentially pure cellulose.




cellulose
1→4-β-glycosidic linkages in red

Ball-and-stick models showing the three-dimensional structures of cellulose and starch were given in Figure 5.2.

Cellulose is an unbranched polymer composed of repeating glucose units joined in a 1→4-β-glycosidic linkage. The β-glycosidic linkage forms long linear chains of cellulose molecules that stack in sheets, creating an extensive three-dimensional array. A network of intermolecular hydrogen bonds between the chains and sheets means that only the few OH groups on the surface are available to hydrogen bond to water, making this very polar compound water insoluble.

Cellulose acetate, a cellulose derivative, is made by treating cellulose with acetic anhydride and sulfuric acid. The resulting product has acetate esters in place of every OH group. Cellulose acetate is spun into fibers that are used for fabrics called *acetates,* which have a deep luster and satin appearance.




cellulose

Ac_2O, H_2SO_4

cellulose acetate

Cellulose can be hydrolyzed to glucose by cleaving all of the β-glycosidic bonds, yielding both anomers of glucose.



β-D-glucose α-D-glucose


A **β-glycosidase** is the general name of an enzyme that hydrolyzes a β-glycoside linkage.

In cells, the hydrolysis of cellulose is accomplished by an enzyme called a **β-glucosidase,** which cleaves all the β-glycoside bonds formed from glucose. Humans do not possess this enzyme, and therefore cannot digest cellulose. Ruminant animals, on the other hand, such as cattle, deer, and camels, have bacteria containing a β-glucosidase in their digestive systems, so they can derive nutritional benefit from eating grass and leaves.

28.12B Starch


Starch is the main carbohydrate found in the seeds and roots of plants. Corn, rice, wheat, and potatoes are common foods that contain a great deal of starch.

Starch is a polymer composed of repeating glucose units joined in α-glycosidic linkages. Both starch and cellulose are polymers of glucose, but starch contains α glycoside bonds, whereas cellulose contains β glycoside bonds. The two common forms of starch are **amylose** and **amylopectin.**



amylose
(the linear form of starch)

1→4-α-glycosidic linkages in red



amylopectin
(the branched form of starch)

1→4-α-glycosidic linkages in red
1→6-α-glycosidic linkage in blue

Amylose, which comprises about 20% of starch molecules, has an unbranched skeleton of glucose molecules with **1→4-α-glycoside bonds.** Because of this linkage, an amylose chain adopts a helical arrangement, giving it a very different three-dimensional shape from the linear chains of cellulose. Amylose was first described in Section 5.1.

Amylopectin, which comprises about 80% of starch molecules, likewise consists of a backbone of glucose units joined in **α-glycosidic bonds,** but it also contains considerable branching along the chain. The linear linkages of amylopectin are formed by **1→4-α-glycoside bonds,** similar to amylose. The branches are linked to the chain with **1→6-α-glycosidic linkages.**

Both forms of starch are water soluble. Because the OH groups in these starch molecules are not buried in a three-dimensional network, they are more available for hydrogen bonding with water molecules, leading to greater water solubility than cellulose has.

The ability of amylopectin to form branched polymers is a unique feature of carbohydrates. Other types of polymers in the cell, such as the proteins discussed in Chapter 29, occur in nature only as linear molecules.

α-Glycosidase is the general name of an enzyme that hydrolyzes an α-glycoside linkage.

Both amylose and amylopectin are hydrolyzed to glucose with cleavage of the glycosidic bonds. The human digestive system has the necessary **α-glucosidase** enzymes needed to catalyze this process. Bread and pasta made from wheat flour, rice, and corn tortillas are all sources of starch that are readily digested.

28.12C Glycogen

Glycogen is the major form in which polysaccharides are stored in animals. Glycogen, a polymer of glucose containing **α-glycosidic bonds,** has a branched structure similar to amylopectin, but the branching is much more extensive.

Glycogen is stored principally in the liver and muscle. When glucose is needed for energy in the cell, glucose units are hydrolyzed from the ends of the glycogen polymer, and then further metabolized with the release of energy. Because glycogen has a highly branched structure, there are many glucose units at the ends of the branches that can be cleaved whenever the body needs them.

Problem 28.31 Draw the structure of: (a) a polysaccharide formed by joining D-mannose units in 1→4-β-glycosidic linkages; (b) a polysaccharide formed by joining D-glucose units in 1→6-α-glycosidic linkages. The polysaccharide in (b) is dextran, a component of dental plaque.

28.13 Other Important Sugars and Their Derivatives


Many other examples of simple and complex carbohydrates with useful properties exist in the biological world. In Section 28.13, we examine some carbohydrates that contain nitrogen atoms.

28.13A Amino Sugars and Related Compounds




Dietary supplements containing glucosamine are used by individuals suffering from osteoarthritis.

Amino sugars contain an NH_2 group instead of an OH group at a non-anomeric carbon. The most common amino sugar in nature, **D-glucosamine,** is formally derived from D-glucose by replacing the OH at C2 with NH_2. Although it is not classified as a drug, and therefore not regulated by the Food and Drug Administration, glucosamine is available in many over-the-counter treatments for osteoarthritis.



D-glucosamine — *N*-acetyl-D-glucosamine **NAG**

Acetylation of glucosamine with acetyl CoA (Section 22.17) forms ***N*-acetyl-D-glucosamine,** abbreviated as **NAG. Chitin,** the second most abundant carbohydrate polymer, is a polysaccharide formed from NAG units joined together in **1→4-β-glycosidic linkages.** Chitin is identical in structure to cellulose, except that each OH group at C2 is now replaced by $NHCOCH_3$. The exoskeletons of lobsters, crabs, and shrimp are composed of chitin. Like those of cellulose, chitin chains are held together by an extensive network of hydrogen bonds, forming water-insoluble sheets.




1→4-β-glycosidic linkages in red

chitin




The rigidity of a crab shell is due to chitin, a high molecular weight carbohydrate molecule. Chitin-based coatings have found several commercial applications, such as extending the shelf life of fruits. Processing plants now convert the shells of crabs, lobsters, and shrimp to chitin and various derivatives for use in many consumer products.

Several trisaccharides containing amino sugars are potent antibiotics used in the treatment of certain severe and recurrent bacterial infections. These compounds, such as tobramycin and amikacin, are called **aminoglycoside antibiotics.**



tobramycin




amikacin

Problem 28.32 Treating chitin with H_2O, ^-OH hydrolyzes its amide linkages, forming a compound called chitosan. What is the structure of chitosan? Chitosan has been used in shampoos, fibers for sutures, and wound dressings.


28.13B *N*-Glycosides

***N*-Glycosides** are formed when a monosaccharide is reacted with an amine in the presence of mild acid (Reactions [1] and [2]).



The mechanism of *N*-glycoside formation is analogous to the mechanism for glycoside formation, and both anomers of the *N*-glycoside are formed as products.

Problem 28.33 Draw the products of each reaction.



Problem 28.34 Draw a stepwise mechanism for the conversion of β-D-glucose to both anomers of *N*-ethyl glucopyranoside, the equation written in Reaction [1].


The prefix ***deoxy*** means "without oxygen."

The *N*-glycosides of two sugars, **D-ribose** and **2-deoxy-D-ribose,** are especially noteworthy, because they form the building blocks of RNA and DNA, respectively. 2-Deoxyribose is so named because it lacks an OH group at C2 of ribose.




- **Reaction of D-ribose with certain amine heterocycles forms *N*-glycosides called ribonucleosides.**
- **This same reaction of 2-deoxy-D-ribose forms deoxyribonucleosides.**

An example of a **ribonucleoside** and a **deoxyribonucleoside** are drawn. These *N*-glycosides have the β orientation. Numbering in the sugar ring begins at the anomeric carbon (1'), and proceeds in a clockwise fashion around the ring.



cytidine

a ribonucleoside




2-deoxyadenosine


a deoxyribonucleoside

Only five common nitrogen heterocycles are used to form these nucleosides. Three compounds have one ring, and are derived from a nitrogen heterocycle called **pyrimidine.** Two are bicyclic, and are derived from a nitrogen heterocycle called **purine.** These five amines are referred to as ***bases.*** Each base is designated by a one-letter abbreviation, as shown in the names and structures drawn. Uracil (U) occurs only in ribonucleosides and thymine (T) occurs only in deoxyribonucleosides.

- Each nucleoside has two parts, a sugar and a base, joined together by a β *N*-glycosidic linkage.




When one OH group of the sugar nucleus is bonded to a phosphate, the derivatives are called **ribonucleotides** and **deoxyribonucleotides.**



cytidine monophosphate

a ribonucleotide




deoxyadenosine monophosphate

a deoxyribonucleotide

- Ribonucleotides are the building blocks of the polymer ribonucleic acid, or RNA, the messenger molecules that convert genetic information into proteins.
- Deoxyribonucleotides are the building blocks of the polymer deoxyribonucleic acid, or DNA, the molecules that are responsible for the storage of all genetic information.

Figure 28.11 Short segments of RNA and DNA



Short segments of both RNA and DNA are shown in Figure 28.11. Note the central role of the sugar moiety in both RNA and DNA. The sugar residues are bonded to two phosphate groups, thus connecting the chain of RNA or DNA together. The sugar residues are also bonded to the nitrogen base via the anomeric carbon.





Figure 28.12 DNA—A double helix



Individual base pairs that hydrogen bond to hold two strands of DNA together

- Two polynucleotide strands form the double helix of DNA. The backbone of each polymer strand is composed of sugar–phosphate residues. Hydrogen bonding of base pairs (A–T and C–G) holds the two strands of DNA together.

DNA is composed of two polynucleotide strands that wind around each other to form a double helix, resembling a spiral ladder. The sides of the ladder are composed of the sugar–phosphate backbone of the polymer and the rungs are composed of the bases, as shown in Figure 28.12.

The nitrogen bases on one strand of DNA hydrogen bond to nitrogen bases on the other strand. A purine base on one strand hydrogen bonds with a pyrimidine base on the other strand. Two types of bases, called **base pairs,** hydrogen bond to each other: adenine hydrogen bonds with thymine (A–T) and cytosine hydrogen bonds with guanine (C–G).

Problem 28.35 Draw the structures of the nucleosides formed from each of the following components: (a) ribose + uracil; (b) 2-deoxyribose + guanine.

Problem 28.36 (a) Why can't two purine bases (A and G) form a base pair and hydrogen bond to each other on two strands of DNA in the double helix? (b) Why is hydrogen bonding between guanine and cytosine more favorable than hydrogen bonding between guanine and thymine?


KEY CONCEPTS

Carbohydrates

Important Terms


- **Aldose** A monosaccharide containing an aldehyde (28.2)
- **Ketose** A monosaccharide containing a ketone (28.2)
- **D-Sugar** A monosaccharide with the OH bonded to the stereogenic center farthest from the carbonyl group drawn on the right in the Fischer projection (28.2C)
- **Epimers** Two diastereomers that differ in configuration around one stereogenic center only (28.3)
- **Anomers** Monosaccharides that differ in configuration at the hemiacetal OH group (28.6)
- **Glycoside** An acetal derived from a monosaccharide hemiacetal (28.7)

Acyclic, Haworth, and 3-D Representations for D-Glucose (28.6)




Reactions of Monosaccharides Involving the Hemiacetal

[1] Glycoside formation (28.7A)



- Only the hemiacetal OH reacts.
- A mixture of α and β glycosides forms.


[2] Glycoside hydrolysis (28.7B)



- A mixture of α and β anomers forms.


Reactions of Monosaccharides at the OH Groups

[1] Ether formation (28.8)



- All OH groups react.
- The stereochemistry at all stereogenic centers is retained.


[2] Ester formation (28.8)



- All OH groups react.
- The stereochemistry at all stereogenic centers is retained.


Reactions of Monosaccharides at the Carbonyl Group

[1] Oxidation of aldoses (28.9B)




- Aldonic acids are formed using:
  - Ag_2O, NH_4OH
  - Cu^{2+}
  - Br_2, H_2O
- Aldaric acids are formed with HNO_3, H_2O.

[2] Reduction of aldoses to alditols (28.9A)




[3] Wohl degradation (28.10A)



- The C1 – C2 bond is cleaved to shorten an aldose chain by one carbon.
- The stereochemistry at all other stereogenic centers is retained.
- Two epimers at C2 form the same product.


[4] Kiliani–Fischer synthesis (28.10B)



- One carbon is added to the aldehyde end of an aldose.
- Two epimers at C2 are formed.


Other Reactions

[1] Hydrolysis of disaccharides (28.11)



A mixture of anomers is formed.

[2] Formation of *N*-glycosides (28.13B)





- Two anomers are formed.

PROBLEMS


Problems Using Three-Dimensional Models

28.37 Convert each ball-and-stick model to a Fischer projection.


a. 

b. 

28.38 (a) Convert each cyclic monosaccharide into a Fischer projection of its acyclic form. (b) Name each monosaccharide. (c) Label the anomer as α or β.




A




B


Fischer Projections


28.39 Classify each compound as identical to **A** or its enantiomer.




A


a. 


b. 


c. 


28.40 Convert each compound to a Fischer projection and label each stereogenic center as *R* or *S*.


a. 

c. 

e. 

b. 

d. 


f. 


Monosaccharide Structure and Stereochemistry


28.41 For D-arabinose:


a. Draw its enantiomer.
b. Draw an epimer at C3.
c. Draw a diastereomer that is not an epimer.
d. Draw a constitutional isomer that still contains a carbonyl group.


28.42 Consider the following six compounds (**A–F**).



A


B


C


D


E


F

How are the two compounds in each pair related? Choose from enantiomers, epimers, diastereomers but not epimers, constitutional isomers, and identical compounds.


a. **A** and **B** b. **A** and **C** c. **B** and **C** d. **A** and **D** e. **E** and **F**

28.43 Draw a Haworth projection for each compound using the structures in Figures 28.4 and 28.5.


a. β-D-talopyranose b. α-D-galactopyranose c. α-D-tagatofuranose

28.44 Draw the β anomer of a monosaccharide epimeric with D-glucose at C2 using a Haworth projection.

28.45 Draw both pyranose anomers of each aldohexose using a three-dimensional representation with a chair pyranose. Label each anomer as α or β.



28.46 Convert each cyclic monosaccharide into its acyclic form.



28.47 D-Arabinose can exist in both pyranose and furanose forms.

a. Draw the α and β anomers of D-arabinofuranose.

b. Draw the α and β anomers of D-arabinopyranose.

28.48 The most stable conformation of the pyranose ring of most D-aldohexoses places the largest group, CH_2OH, in the equatorial position. An exception to this is the aldohexose D-idose. Draw the two possible chair conformations of either the α or β anomer of D-idose. Explain why the more stable conformation has the CH_2OH group in the axial position.

Monosaccharide Reactions

28.49 Draw the products formed when α-D-gulose is treated with each reagent.


a. CH_3I, Ag_2O
b. CH_3OH, HCl
c. Ac_2O, pyridine
d. The product in (a), then H_3O^+
e. The product in (b), then Ac_2O, pyridine
f. The product in (d), then $C_6H_5CH_2Cl$, Ag_2O


28.50 Draw the products formed when D-altrose is treated with each reagent.

a. $(CH_3)_2CHOH$, HCl
b. $NaBH_4$, CH_3OH
c. Br_2, H_2O
d. HNO_3, H_2O
e. [1] NH_2OH; [2] $(CH_3CO)_2O$, $NaOCOCH_3$; [3] $NaOCH_3$
f. [1] NaCN, HCl; [2] H_2, Pd-$BaSO_4$; [3] H_3O^+
g. CH_3I, Ag_2O
h. $C_6H_5CH_2NH_2$, mild H^+

28.51 What aglycon and monosaccharides are formed when salicin and solanine (Section 28.7C) are each hydrolyzed with aqueous acid?


28.52 Draw a Fischer projection of the monosaccharide from which each of the following glycosides was prepared.


a. 

b. 


28.53 What two aldohexoses yield D-arabinose upon Wohl degradation?


28.54 What products are formed when each compound is subjected to a Kiliani–Fischer synthesis?


a. 

b. 

28.55 How would you convert D-glucose into each compound? More than one step is required.


a. 


b. 


c. 

28.56 Which D-aldopentoses are reduced to optically inactive alditols using $NaBH_4$, CH_3OH?

28.57 What products are formed when each compound is treated with aqueous acid?


a. 

b. 


c. 

Mechanisms


28.58 Draw a stepwise mechanism for the following reaction.



28.59 Draw a stepwise mechanism for the following hydrolysis.



28.60 The following isomerization reaction, drawn using D-glucose as starting material, occurs with all aldohexoses in the presence of base. Draw a stepwise mechanism that illustrates how each compound is formed.



Identifying Monosaccharides

28.61 Which D-aldopentose is oxidized to an optically active aldaric acid and undergoes the Wohl degradation to yield a D-aldotetrose that is oxidized to an optically active aldaric acid?

28.62 Identify compounds **A–D.** A D-aldopentose **A** is oxidized with HNO_3 to an optically inactive aldaric acid **B. A** undergoes the Kiliani–Fischer synthesis to yield **C** and **D. C** is oxidized to an optically active aldaric acid. **D** is oxidized to an optically inactive aldaric acid.


28.63 A D-aldopentose **A** is reduced to an optically active alditol. Upon Kiliani–Fischer synthesis, **A** is converted to two D-aldohexoses, **B** and **C. B** is oxidized to an optically inactive aldaric acid. **C** is oxidized to an optically active aldaric acid. What are the structures of **A–C?**

Disaccharides and Polysaccharides


28.64 Draw the structure of a disaccharide formed from two mannose units joined by a 1→4-α-glycosidic linkage.

28.65 a. Identify the glycosidic linkage in disaccharide **C,** classify the glycosidic bond as α or β, and use numbers to designate its location.

b. Identify the lettered compounds in the following reaction.



28.66 Consider the tetrasaccharide stachyose drawn below. Stachyose is found in white jasmine, soybeans, and lentils. Because humans cannot digest it, its consumption causes flatulence.



a. Label all glycoside bonds.
b. Classify each glycosidic linkage as α or β and use numbers to designate its location between two rings (e.g., 1→4-β).
c. What products are formed when stachyose is hydrolyzed with H_3O^+?
d. Is stachyose a reducing sugar?
e. What product is formed when stachyose is treated with excess CH_3I, Ag_2O?
f. What products are formed when the product in (e) is treated with H_3O^+?


28.67 Deduce the structure of the disaccharide isomaltose from the following data.

[1] Hydrolysis yields D-glucose exclusively.

[2] Isomaltose is cleaved with α-glycosidase enzymes.

[3] Isomaltose is a reducing sugar.

[4] Methylation with excess CH_3I, Ag_2O and then hydrolysis with H_3O^+ forms two products:



(Both anomers are present.)

28.68 Draw the structure of each of the following compounds:

a. a polysaccharide formed by joining D-glucosamine in 1→6-α-glycosidic linkages


b. a disaccharide formed by joining D-mannose and D-glucose in a 1→4-β-glycosidic linkage using mannose's anomeric carbon

c. an α-*N*-glycoside formed from D-arabinose and $C_6H_5CH_2NH_2$

d. a ribonucleoside formed from D-ribose and thymine


Challenge Problems

28.69 (a) Draw the more stable chair form of fucose, an essential monosaccharide needed in the diet and a component of carbohydrates on mammalian and plant cell surfaces. (b) Classify fucose as a D- or L-monosaccharide. (c) What two structural features are unusual in fucose?




fucose

28.70 As we have seen in Chapter 28, monosaccharides can be drawn in a variety of ways, and in truth, often a mixture of cyclic compounds is present in a solution. Identify each monosaccharide, including its proper D,L designation, drawn in a less-than-typical fashion.




28.71 Draw a stepwise mechanism for the following reaction.



29 Amino Acids and Proteins

29.1 Amino acids
29.2 Synthesis of amino acids
29.3 Separation of amino acids
29.4 Enantioselective synthesis of amino acids
29.5 Peptides
29.6 Peptide sequencing
29.7 Peptide synthesis
29.8 Automated peptide synthesis
29.9 Protein structure
29.10 Important proteins



Myoglobin is a globular protein that contains 153 amino acids joined together, as well as a non-protein portion called a heme unit. The heme group consists of a large nitrogen heterocycle complexed with the Fe^{2+} cation. The Fe^{2+} cation binds oxygen in the blood and stores it in tissues. Whales have a particularly high myoglobin concentration in their muscles. It serves as an oxygen reservoir for the whale while it is submerged for long periods of time. In Chapter 29, we discuss the properties of proteins and the amino acids from which they are synthesized.

Of the four major groups of biomolecules—lipids, carbohydrates, nucleic acids, and proteins—proteins have the widest array of functions. **Keratin** and **collagen,** for example, are part of a large group of structural proteins that form long insoluble fibers, giving strength and support to tissues. Hair, horns, hooves, and fingernails are all made up of keratin. **Collagen** is found in bone, connective tissue, tendons, and cartilage. **Enzymes** are proteins that catalyze and regulate all aspects of cellular function. **Membrane proteins** transport small organic molecules and ions across cell membranes. **Insulin,** the hormone that regulates blood glucose levels, **fibrinogen** and **thrombin,** which form blood clots, and **hemoglobin,** which transports oxygen from the lungs to tissues, are all proteins.


In Chapter 29 we discuss proteins and their primary components, the amino acids.

29.1 Amino Acids

Amino acids were first discussed in Section 19.14.

Naturally occurring amino acids have an amino group (NH_2) bonded to the α carbon of a carboxy group (COOH), and so they are called **α-amino acids.**

- **All proteins are polyamides formed by joining amino acids together.**



α-amino acid **portion of a protein molecule**

29.1A General Features of α-Amino Acids

The 20 amino acids that occur naturally in proteins differ in the identity of the R group bonded to the α carbon. The R group is called the **side chain** of the amino acid.

The simplest amino acid, called glycine, has R = H. **All other amino acids (R ≠ H) have a stereogenic center on the α carbon.** As is true for monosaccharides, the prefixes **D** and **L** are used to designate the configuration at the stereogenic center of amino acids. Common, naturally occurring amino acids are called **L-amino acids.** Their enantiomers, D-amino acids, are rarely found in nature. These general structures are shown in Figure 29.1. According to *R,S* designations, all L-amino acids except cysteine have the ***S* configuration.**

All amino acids have common names. These names can be represented by either a one-letter or a three-letter abbreviation. Figure 29.2 is a listing of the 20 naturally occurring amino acids, together with their abbreviations. Note the variability in the R groups. A side chain can be a simple alkyl group, or it can have additional functional groups such as OH, SH, COOH, or NH_2.

- **Amino acids with an additional COOH group in the side chain are called acidic amino acids.**
- **Those with an additional basic N atom in the side chain are called basic amino acids.**
- **All others are neutral amino acids.**

Figure 29.1
The general features of an α-amino acid





















Figure 29.2 The 20 naturally occurring amino acids

Neutral amino acids




Name	Structure	Abbreviations
Alanine		Ala A
Asparagine		Asn N
Cysteine		Cys C
Glutamine		Gln Q
Glycine		Gly G
Isoleucine*		Ile I
Leucine*		Leu L
Methionine*		Met M

Name	Structure	Abbreviations
Phenylalanine*		Phe F
Proline		Pro P
Serine		Ser S
Threonine*		Thr T
Tryptophan*		Trp W
Tyrosine		Tyr Y
Valine*		Val V

Acidic amino acids

Name	Structure	Abbreviations
Aspartic acid		Asp D
Glutamic acid		Glu E

Basic amino acids

Name	Structure	Abbreviations
Arginine*		Arg R
Histidine*		His H
Lysine*		Lys K


Essential amino acids are labeled with an asterisk (*).



The essential amino acid leucine is sold as a dietary supplement that is used by body builders to help prevent muscle loss and heal muscle tissue after injury.

Look closely at the structures of proline, isoleucine, and threonine.

- **All amino acids are 1° amines except for proline,** which has its N atom in a five-membered ring, making it a **2° amine.**
- **Isoleucine** and **threonine** contain an additional stereogenic center at the β carbon, so there are four possible stereoisomers, only one of which is naturally occurring.



L-proline
2° amine

L-isoleucine


L-threonine

Humans can synthesize only 10 of these 20 amino acids. The remaining 10 are called **essential amino acids** because they must be obtained from the diet. These are labeled with an asterisk in Figure 29.2.

Problem 29.1 Draw the other three stereoisomers of L-isoleucine, and label the stereogenic centers as *R* or *S*.

29.1B Acid–Base Behavior

Recall from Section 19.14B that an amino acid has both an acidic and a basic functional group, so proton transfer forms a salt called a **zwitterion.**



This neutral form of an amino acid does ***not*** really exist.

The zwitterion is neutral.

This salt is the neutral form of an amino acid.

This form exists at **pH ≈ 6.**

The structures in Figure 29.2 show the charged form of the amino acids at the physiological pH of the blood.


- **Amino acids do not exist to any appreciable extent as uncharged neutral compounds. They exist as salts, giving them high melting points and making them water soluble.**

Amino acids exist in different charged forms, as shown in Figure 29.3, depending on the pH of the aqueous solution in which they are dissolved. For neutral amino acids, the overall charge is +1, 0, or –1. Only at pH ~6 does the zwitterionic form exist.

The $-COOH$ and $-NH_3^+$ groups of an amino acid are ionizable, because they can lose a proton in aqueous solution. As a result, they have different pK_a values. The pK_a of the $-COOH$ group is typically ~2, whereas that of the $-NH_3^+$ group is ~9, as shown in Table 29.1.

Some amino acids, such as aspartic acid and lysine, have acidic or basic side chains. These additional ionizable groups complicate somewhat the acid–base behavior of these amino acids. Table 29.1 lists the pK_a values for these acidic and basic side chains as well.

Figure 29.3
How the charge of a neutral amino acid depends on the pH



overall (+1) charge
pH ≈ 2

neutral
pH ≈ 6

overall (–1) charge
pH ≈ 10

Table 29.1 pK_a Values for the Ionizable Functional Groups of an α-Amino Acid


Amino acid	α-COOH	α-NH_3^+	Side chain	p*I*
Alanine	2.35	9.87	—	6.11
Arginine	2.01	9.04	12.48	10.76
Asparagine	2.02	8.80	—	5.41
Aspartic acid	2.10	9.82	3.86	2.98
Cysteine	2.05	10.25	8.00	5.02
Glutamic acid	2.10	9.47	4.07	3.08
Glutamine	2.17	9.13	—	5.65
Glycine	2.35	9.78	—	6.06
Histidine	1.77	9.18	6.10	7.64
Isoleucine	2.32	9.76	—	6.04
Leucine	2.33	9.74	—	6.04
Lysine	2.18	8.95	10.53	9.74
Methionine	2.28	9.21	—	5.74
Phenylalanine	2.58	9.24	—	5.91
Proline	2.00	10.60	—	6.30
Serine	2.21	9.15	—	5.68
Threonine	2.09	9.10	—	5.60
Tryptophan	2.38	9.39	—	5.88
Tyrosine	2.20	9.11	10.07	5.63
Valine	2.29	9.72	—	6.00

Table 29.1 also lists the isoelectric points (p*I*) for all of the amino acids. Recall from Section 19.14C that the **isoelectric point is the pH at which an amino acid exists primarily in its neutral form,** and that it can be calculated from the average of the pK_a values of the α-COOH and α-NH_3^+ groups (for neutral amino acids only).

Problem 29.2 What form exists at the isoelectric point of each of the following amino acids: (a) valine; (b) leucine; (c) proline; (d) glutamic acid?

Problem 29.3 Explain why the pK_a of the $-NH_3^+$ group of an α-amino acid is lower than the pK_a of the ammonium ion derived from a 1° amine (RNH_3^+). For example the pK_a of the $-NH_3^+$ group of alanine is 9.87 but the pK_a of $CH_3NH_3^+$ is 10.63.

Problem 29.4 L-Thyroxine, a thyroid hormone and oral medication used to treat thyroid hormone deficiency, is an amino acid that does not exist in proteins. Draw the zwitterionic form of L-thyroxine.




L-thyroxine

29.2 Synthesis of Amino Acids

Amino acids can be prepared in a variety of ways in the laboratory. Three methods are described, each of which is based on reactions learned in previous chapters.

29.2A S_N2 Reaction of α-Halo Acids with NH_3

The most direct way to synthesize an α-amino acid is by **S_N2 reaction of an α-halo carboxylic acid with a large excess of NH_3.**



Although the alkylation of ammonia with simple alkyl halides does not generally afford high yields of 1° amines (Section 25.7A), this reaction using α-halo carboxylic acids does form the desired amino acids in good yields. In this case, the amino group in the product is both less basic and more sterically crowded than other 1° amines, so that a single alkylation occurs and the desired amino acid is obtained.

Problem 29.5 What α-halo carbonyl compound is needed to synthesize each amino acid: (a) glycine; (b) isoleucine; (c) phenylalanine?

29.2B Alkylation of a Diethyl Malonate Derivative

The second method for preparing amino acids is based on the malonic ester synthesis. Recall from Section 23.9 that this synthesis converts diethyl malonate to a carboxylic acid with a new alkyl group on its α carbon atom.




diethyl malonate

This reaction can be adapted to the synthesis of α-amino acids by using a commercially available derivative of diethyl malonate as starting material. This compound, **diethyl acetamidomalonate,** has a nitrogen atom on the α carbon, which ultimately becomes the NH_2 group on the α carbon of the amino acid.



diethyl acetamidomalonate

The malonic ester synthesis consists of three steps, and so does this variation to prepare an amino acid.




diethyl acetamidomalonate

[1] **Deprotonation** of diethyl acetamidomalonate with NaOEt forms an enolate by removal of the acidic proton between the two carbonyl groups.

[2] **Alkylation** of the enolate with an unhindered alkyl halide (usually CH_3X or RCH_2X) forms a substitution product with a new R group on the α carbon.

[3] Heating the alkylation product with aqueous acid results in **hydrolysis** of both esters and the amide, followed by **decarboxylation** to form the amino acid.

Phenylalanine, for example, can be synthesized as follows:




The charge on the amino acid product (+1, –1, or 0) depends on the reaction conditions. Phenylalanine bears a net positive charge because the last step in its synthesis uses strong acid.

Problem 29.6 The enolate derived from diethyl acetamidomalonate is treated with each of the following alkyl halides. After hydrolysis and decarboxylation, what amino acid is formed: (a) CH_3I; (b) $(CH_3)_2CHCH_2Cl$; (c) $CH_3CH_2CH(CH_3)Br$?


Problem 29.7 What amino acid is formed when $CH_3CONHCH(CO_2Et)_2$ is treated with the following series of reagents: [1] NaOEt; [2] $CH_2=O$; [3] H_3O^+, Δ?

29.2C Strecker Synthesis

The third method, the **Strecker amino acid synthesis,** converts an aldehyde into an amino acid by a two-step sequence that adds one carbon atom to the aldehyde carbonyl. Treating an aldehyde with NH_4Cl and NaCN first forms an **α-amino nitrile,** which can then be hydrolyzed in aqueous acid to an amino acid.



The Strecker synthesis of alanine, for example, is as follows:




Mechanism 29.1 for the formation of the α-amino nitrile from an aldehyde (the first step in the Strecker synthesis) consists of **nucleophilic addition of NH_3** to form an imine, followed by **addition of cyanide** to the C=N bond. Both parts are related to earlier mechanisms involving imines (Section 21.11) and cyanohydrins (Section 21.9).

The details of the second step of the Strecker synthesis, the hydrolysis of a nitrile (RCN) to a carboxylic acid (RCOOH), have already been presented in Section 22.18A.

Figure 29.4 shows how the amino acid methionine can be prepared by all three methods in Section 29.2.

Problem 29.8 What aldehyde is needed to synthesize each amino acid by the Strecker synthesis: (a) valine; (b) leucine; (c) phenylalanine?


Mechanism 29.1 Formation of an α-Amino Nitrile



1 – 3 Nucleophilic attack of NH_3 followed by proton transfer and loss of H_2O forms an imine. Loss of H_2O occurs by the same three-step process outlined in Mechanism 21.5.

4 – 5 Protonation of the imine followed by nucleophilic attack of ^-CN gives the α-amino nitrile.


Figure 29.4
The synthesis of methionine by three different methods



Three methods of amino acid synthesis:

[1] S_N2 reaction using an α-halo carboxylic acid
[2] Alkylation of diethyl acetamidomalonate
[3] Strecker synthesis

Problem 29.9 Draw the products of each reaction.



29.3 Separation of Amino Acids

No matter which of the preceding methods is used to synthesize an amino acid, all three yield a racemic mixture. Naturally occurring amino acids exist as a single enantiomer, however, so the two enantiomers obtained must be separated if they are to be used in biological applications. This is not an easy task. Two enantiomers have the same physical properties, so they cannot be separated by common physical methods, such as distillation or chromatography. Moreover, they react in the same way with achiral reagents, so they cannot be separated by chemical reactions either.

Nonetheless, strategies have been devised to separate two enantiomers using physical separation techniques and chemical reactions. We examine two different strategies in Section 29.3. Then,

in Section 29.4, we will discuss a method that affords optically active amino acids without the need for separation.

- The separation of a racemic mixture into its component enantiomers is called *resolution.* Thus, a racemic mixture is *resolved* into its component enantiomers.

29.3A Resolution of Amino Acids

The oldest, and perhaps still the most widely used method to separate enantiomers exploits the following fact: **enantiomers have the *same* physical properties, but diastereomers have *different* physical properties.** Thus, a racemic mixture can be resolved using the following general strategy.


[1] **Convert a pair of enantiomers into a pair of diastereomers,** which are now separable because they have different melting points and boiling points.

[2] **Separate the diastereomers.**

[3] **Re-convert each diastereomer into the original enantiomer,** now separated from the other.

This general three-step process is illustrated in Figure 29.5.


Figure 29.5
Resolution of a racemic mixture by converting it to a mixture of diastereomers




Enantiomers A and B can be separated by reaction with a single enantiomer of a chiral reagent, Y. The process of resolution requires three steps:

[1] Reaction of enantiomers **A** and **B** with **Y** forms two diastereomers, **AY** and **BY.**
[2] Diastereomers **AY** and **BY** have different physical properties, so they can be separated by physical methods such as fractional distillation or crystallization.
[3] **AY** and **BY** are then re-converted to **A** and **B** by a chemical reaction. The two enantiomers **A** and **B** are now separated from each other, and resolution is complete.


To resolve a racemic mixture of amino acids such as (*R*)- and (*S*)-alanine, the racemate is first treated with acetic anhydride to form ***N*-acetyl amino acids.** Each of these amides contains one stereogenic center and they are still enantiomers, so they are *still inseparable.*





(*R*)-α-methylbenzylamine

a resolving agent




Both enantiomers of *N*-acetyl alanine have a free carboxy group that can react with an amine in an acid–base reaction. **If a chiral amine is used, such as (*R*)-α-methylbenzylamine, the two**

salts formed are diastereomers, *not* enantiomers. Diastereomers can be physically separated from each other, so the compound that converts enantiomers into diastereomers is called a **resolving agent.** Either enantiomer of the resolving agent can be used.


How To Use (*R*)-α-Methylbenzylamine to Resolve a Racemic Mixture of Amino Acids

Step [1] React both enantiomers of an *N*-acetyl amino acid with the *R* isomer of the chiral amine.




These salts have the *same* configuration around one stereogenic center, but the *opposite* configuration about the other stereogenic center.

Step [2] Separate the diastereomers.



Step [3] Regenerate the amino acid by hydrolysis of the amide.





Step [1] is just an acid–base reaction in which the racemic mixture of *N*-acetyl alanines reacts with the same enantiomer of the resolving agent, in this case (*R*)-α-methylbenzylamine. The salts that form are **diastereomers, *not* enantiomers,** because they have the same configuration about one stereogenic center, but the opposite configuration about the other stereogenic center.


In **Step [2],** the diastereomers are separated by some physical technique, such as crystallization or distillation.


In **Step [3],** the amides can be hydrolyzed with aqueous base to regenerate the amino acids. The amino acids are now separated from each other. The optical activity of the amino acids can be measured and compared to their known rotations to determine the purity of each enantiomer.

Problem 29.10 Which of the following amines can be used to resolve a racemic mixture of amino acids?

a. 

b. 

c. 

d. 


strychnine
(a powerful poison)

Problem 29.11 Write out a stepwise sequence that shows how a racemic mixture of leucine enantiomers can be resolved into optically active amino acids using (*R*)-α-methylbenzylamine.

29.3B Kinetic Resolution of Amino Acids Using Enzymes


A second strategy used to separate amino acids is based on the fact that two enantiomers react differently with chiral reagents. An **enzyme** is typically used as the chiral reagent.

To illustrate this strategy, we begin again with the two enantiomers of *N*-acetyl alanine, which were prepared by treating a racemic mixture of (*R*)- and (*S*)-alanine with acetic anhydride (Section 29.3A). Enzymes called **acylases** hydrolyze amide bonds, such as those found in *N*-acetyl alanine, but only for amides of L-amino acids. Thus, when a racemic mixture of *N*-acetyl alanines is treated with an acylase, only the amide of L-alanine (the *S* stereoisomer) is hydrolyzed to generate L-alanine, whereas the amide of D-alanine (the *R* stereoisomer) is untouched. The reaction mixture now consists of one amino acid and one *N*-acetyl amino acid. Because they have different functional groups with different physical properties, they can be physically separated.



- **Separation of two enantiomers by a chemical reaction that selectively occurs for only one of the enantiomers is called *kinetic resolution.***

Problem 29.12 Draw the organic products formed in the following reaction.




(mixture of enantiomers)

29.4 Enantioselective Synthesis of Amino Acids

Although the two methods introduced in Section 29.3 for resolving racemic mixtures of amino acids make enantiomerically pure amino acids available for further research, half of the reaction product is useless because it has the undesired configuration. Moreover, each of these procedures is costly and time-consuming.


If we use a chiral reagent to synthesize an amino acid, however, it is possible to favor the formation of the desired enantiomer over the other, without having to resort to a resolution. For example, single enantiomers of amino acids have been prepared by using **enantioselective (or asymmetric) hydrogenation reactions.** The success of this approach depends on finding a chiral catalyst, in much the same way that a chiral catalyst is used for the Sharpless asymmetric epoxidation (Section 12.15).

The necessary starting material is an alkene. Addition of H_2 to the double bond forms an *N*-acetyl amino acid with a new stereogenic center on the α carbon to the carboxy group. With proper choice of a chiral catalyst, the naturally occurring *S* configuration can be obtained as product.




achiral alkene — new stereogenic center on the α carbon — With proper choice of catalyst, the naturally occurring *S* isomer is formed.

Several chiral catalysts with complex structures have now been developed for this purpose. Many contain **rhodium** as the metal, complexed to a chiral molecule containing one or more phosphorus atoms. One example, abbreviated simply as **Rh*,** is drawn below.




chiral hydrogenation catalyst

Figure 29.6
The structure of BINAP



2,2'-bis(diphenylphosphino)-1,1'-binaphthyl

BINAP



3-D model of BINAP

- The two naphthalene rings are oriented at right angles to each other, creating a rigid shape that makes the molecule chiral.


Ryoji Noyori shared the 2001 Nobel Prize in Chemistry for developing methods for asymmetric hydrogenation reactions using the chiral BINAP catalyst.

Twistoflex and helicene (Section 17.5) are two more aromatic compounds whose shape makes them chiral.

This catalyst is synthesized from a rhodium salt and a phosphorus compound, 2,2'-bis(diphenylphosphino)-1,1'-binaphthyl **(BINAP).** It is the BINAP moiety (Figure 29.6) that makes the catalyst chiral.

BINAP is one of a small number of molecules that is chiral even though it has no tetrahedral stereogenic centers. Its shape makes it a chiral molecule. The two naphthalene rings of the BINAP molecule are oriented at almost 90° to each other to minimize steric interactions between the hydrogen atoms on adjacent rings. This rigid three-dimensional shape makes BINAP nonsuperimposable on its mirror image, and thus it is a chiral compound.

Enantioselective hydrogenation can be used to synthesize a single stereoisomer of phenylalanine. Treating achiral alkene **A** with H_2 and the chiral rhodium catalyst Rh* forms the *S* isomer of *N*-acetyl phenylalanine in 100% *ee*. Hydrolysis of the acetyl group on nitrogen then yields a single enantiomer of phenylalanine.




Problem 29.13 What alkene is needed to synthesize each amino acid by an enantioselective hydrogenation reaction using H_2 and Rh*: (a) alanine; (b) leucine; (c) glutamine?

29.5 Peptides

When amino acids are joined together by amide bonds, they form larger molecules called **peptides** and **proteins.**

- A *dipeptide* has two amino acids joined together by *one* amide bond.
- A *tripeptide* has three amino acids joined together by *two* amide bonds.



Two amino acids joined together. Three amino acids joined together.

[Amide bonds are drawn in red.]


Polypeptides and **proteins** both have many amino acids joined together in long linear chains, but the term **protein** is usually reserved for polymers of more than 40 amino acids.

- The amide bonds in peptides and proteins are called *peptide bonds.*
- The individual amino acids are called *amino acid residues.*


29.5A Simple Peptides

To form a dipeptide, the amino group of one amino acid forms an amide bond with the carboxy group of another amino acid. Because each amino acid has both an amino group and a carboxy group, **two different dipeptides can be formed.** This is illustrated with alanine and cysteine.

[1] **The COO^- group of alanine can combine with the NH_3^+ group of cysteine.**



[2] **The COO^- group of cysteine can combine with the NH_3^+ group of alanine.**



These compounds are **constitutional isomers** of each other. Both have a free amino group (protonated as NH_3^+) at one end of their chains and a free carboxy group (deprotonated as a carboxylate anion, COO^-) at the other.

- The amino acid with the free amino group is called the *N-terminal amino acid.*
- The amino acid with the free carboxy group is called the *C-terminal amino acid.*


By convention, **the N-terminal amino acid is always written at the left end of the chain and the C-terminal amino acid at the right.** The peptide can be abbreviated by writing the one- or three-letter symbols for the amino acids in the chain from the N-terminal to the C-terminal end. Thus, Ala–Cys has alanine at the N-terminal end and cysteine at the C-terminal end, whereas Cys–Ala has cysteine at the N-terminal end and alanine at the C-terminal end. Sample Problem 29.1 shows how this convention applies to a tripeptide.

Sample Problem 29.1 Draw the structure of the following tripeptide, and label its N-terminal and C-terminal amino acids: Ala–Gly–Ser.

Solution

Draw the structures of the amino acids in order from **left to right,** placing the COO^- of one amino acid *next* to the NH_3^+ group of the adjacent amino acid. Always draw the **NH_3^+ group on the *left***

and the **COO^- group on the *right*.** Then, join adjacent COO^- and NH_3^+ groups together in amide bonds to form the tripeptide.



tripeptide **Ala–Gly–Ser**
[The new peptide bonds are drawn in red.]

The N-terminal amino acid is **alanine,** and the C-terminal amino acid is **serine.**


The tripeptide in Sample Problem 29.1 has one N-terminal amino acid, one C-terminal amino acid, and two peptide bonds.

- **No matter how many amino acid residues are present, there is only *one* N-terminal amino acid and *one* C-terminal amino acid.**
- **For *n* amino acids in the chain, the number of amide bonds is *n* – 1.**

Problem 29.14 Draw the structure of each peptide. Label the N-terminal and C-terminal amino acids and all amide bonds.

a. Val–Glu b. Gly–His–Leu c. M–A–T–T


Problem 29.15 Name each peptide using both the one-letter and the three-letter abbreviations for the names of the component amino acids.



Problem 29.16 How many different tripeptides can be formed from three different amino acids?

29.5B The Peptide Bond


The carbonyl carbon of an amide is ***sp^2* hybridized** and has **trigonal planar** geometry. A second resonance structure can be drawn that delocalizes the nonbonded electron pair on the N atom. Amides are more resonance stabilized than other acyl compounds, so the resonance structure having the C=N makes a significant contribution to the hybrid.



two resonance structures for the peptide bond

Recall from Section 16.6 that buta-1,3-diene can also exist as *s*-cis and *s*-trans conformations. In buta-1,3-diene, the *s*-cis conformation has the two double bonds on the same side of the single bond (dihedral angle = 0°), whereas the *s*-trans conformation has them on opposite sides (dihedral angle = 180°).

Resonance stabilization has important consequences. Rotation about the C–N bond is restricted because it has partial double bond character. As a result, there are two possible conformations.




s-trans *s*-cis

- The *s*-trans conformation has the two R groups oriented on *opposite* sides of the C – N bond.
- The *s*-cis conformation has the two R groups oriented on the *same* side of the C – N bond.
- The *s*-trans conformation of a peptide bond is typically more stable than the *s*-cis, because the *s*-trans has the two bulky R groups located farther from each other.

The planar geometry of the peptide bond is analogous to the planar geometry of ethylene (or any other alkene), where the double bond between sp^2 hybridized carbon atoms makes all of the bond angles ~120° and puts all six atoms in the same plane.


A second consequence of resonance stabilization is that **all six atoms involved in the peptide bond lie in the same plane.** All bond angles are ~120° and the C=O and N–H bonds are oriented 180° from each other.



These six atoms lie in a plane.

The structure of a tetrapeptide illustrates the results of these effects in a long peptide chain.

- The *s*-trans arrangement makes a long chain with a zigzag arrangement.
- In each peptide bond, the N – H and C = O bonds lie parallel and at 180° with respect to each other.



tetrapeptide


29.5C Interesting Peptides

Even relatively simple peptides can have important biological functions. **Bradykinin,** for example, is a peptide hormone composed of nine amino acids. It stimulates smooth muscle contraction, dilates blood vessels, and causes pain. Bradykinin is a component of bee venom.


Arg–Pro–Pro–Gly–Phe–Ser–Pro–Phe–Arg

bradykinin

Oxytocin and **vasopressin** are nonapeptide hormones, too. Their sequences are identical except for two amino acids, yet this is enough to give them very different biological activities. Oxytocin induces labor by stimulating the contraction of uterine muscles, and it stimulates the flow of milk in nursing mothers. Vasopressin, on the other hand, controls blood pressure by regulating smooth muscle contraction. The N-terminal amino acid in both hormones is a cysteine residue, and the C-terminal residue is glycine. Instead of a free carboxy group, both peptides have an NH_2 group in place of OH, so this is indicated with the additional NH_2 group drawn at the end of the chain.




oxytocin



vasopressin


- The N-terminal amino acid is labeled in red.
- The amino acids that differ are labeled in blue.

Figure 29.7
Three-dimensional structures of vasopressin and oxytocin




The oxidation of thiols to disulfides was discussed in Section 9.15.

The structure of both peptides includes a **disulfide bond,** a form of covalent bonding in which the −SH groups from two cysteine residues are oxidized to form a sulfur–sulfur bond. In oxytocin and vasopressin, the disulfide bonds make the peptides cyclic. Three-dimensional structures of oxytocin and vasopressin are shown in Figure 29.7.



thiol disulfide


The artificial sweetener **aspartame** (Figure 28.10) is the methyl ester of the dipeptide Asp–Phe. This synthetic peptide is 180 times sweeter (on a gram-for-gram basis) than sucrose (common table sugar). Both of the amino acids in aspartame have the naturally occurring L-configuration. If the D-amino acid is substituted for either Asp or Phe, the resulting compound tastes bitter.



aspartame
the methyl ester of Asp–Phe
a synthetic artificial sweetener

Problem 29.17 Draw the structure of leu-enkephalin, a pentapeptide that acts as an analgesic and opiate, and has the following sequence: Tyr–Gly–Gly–Phe–Leu. (The structure of a related peptide, met-enkephalin, appeared in Section 22.6B.)

Problem 29.18 Glutathione, a powerful antioxidant that destroys harmful oxidizing agents in cells, is composed of glutamic acid, cysteine, and glycine, and has the following structure:



glutathione

a. What product is formed when glutathione reacts with an oxidizing agent?
b. What is unusual about the peptide bond between glutamic acid and cysteine?

29.6 Peptide Sequencing

To determine the structure of a peptide, we must know not only what amino acids comprise it, but also the sequence of the amino acids in the peptide chain. Although mass spectrometry has become an increasingly powerful method for the analysis of high molecular weight proteins (Section 13.4), chemical methods to determine peptide structure are still widely used and presented in this section.

29.6A Amino Acid Analysis


The structure determination of a peptide begins by analyzing the total amino acid composition. The amide bonds are first hydrolyzed by heating with hydrochloric acid for 24 h to form the individual amino acids. The resulting mixture is then separated using high-performance liquid chromatography (HPLC), a technique in which a solution of amino acids is placed on a column and individual amino acids move through the column at characteristic rates, often dependent on polarity.

This process determines both the identity of the individual amino acids and the amount of each present, but it tells nothing about the order of the amino acids in the peptide. For example, complete hydrolysis and HPLC analysis of the tetrapeptide Gly–Gly–Phe–Tyr would indicate the presence of three amino acids—glycine, phenylalanine, and tyrosine—and show that there are twice as many glycine residues as phenylalanine or tyrosine residues. The exact order of the amino acids in the peptide chain must then be determined by additional methods

29.6B Identifying the N-Terminal Amino Acid—The Edman Degradation

To determine the sequence of amino acids in a peptide chain, a variety of procedures are often combined. One especially useful technique is to identify the N-terminal amino acid using the **Edman degradation.** In the Edman degradation, amino acids are cleaved one at a time from the N-terminal end, the identity of the amino acid determined, and the process repeated until the entire sequence is known. Automated sequencers using this methodology are now available to sequence peptides containing up to about 50 amino acids.

The Edman degradation is based on the reaction of the nucleophilic NH_2 group of the N-terminal amino acid with the electrophilic carbon of phenyl isothiocyanate, $C_6H_5N{=}C{=}S$. When the N-terminal amino acid is removed from the peptide chain, two products are formed: **an *N*-phenylthiohydantoin (PTH) and a new peptide with one *fewer* amino acid.**




The *N*-phenylthiohydantoin derivative contains the atoms of the N-terminal amino acid. This product identifies the N-terminal amino acid in the peptide because the PTH derivatives of all 20 naturally occurring amino acids are known and characterized. The new peptide formed in the Edman degradation has one amino acid fewer than the original peptide. Moreover, it contains a new N-terminal amino acid, so the process can be repeated.

Mechanism 29.2 illustrates some of the key steps of the Edman degradation. The nucleophilic N-terminal NH_2 group adds to the electrophilic carbon of phenyl isothiocyanate to form an *N*-phenylthiourea, the product of nucleophilic addition (Part [1]). Intramolecular cyclization followed by elimination results in cleavage of the terminal amide bond in Part [2] to form a new peptide with one fewer amino acid. A sulfur heterocycle, called a thiazolinone, is also formed, which rearranges by a multistep pathway to form an *N*-phenylthiohydantoin. The R group in this product identifies the amino acid located at the N-terminal end.


Mechanism 29.2 Edman Degradation

Part [1] Formation of an *N*-phenylthiourea



1 – 2 Addition of the amino group of the N-terminal amino acid to phenyl isothiocyanate followed by proton transfer forms an *N*-phenylthiourea.

Part [2] Formation of the N-terminal amino acid and *N*-phenylthiohydantoin (PTH)



3 Nucleophilic addition of the S atom to the amide carbonyl forms a five-membered ring.

4 Loss of the amino group forms two products—a thiazolinone ring and a peptide chain that contains one fewer amino acid than the original peptide.

5 The thiazolinone rearranges by a multistep pathway to form an *N*-phenylthiohydantoin (PTH) that contains the original amino acid.

In theory a protein of any length can be sequenced using the Edman degradation, but in practice, the accumulation of small quantities of unwanted by-products limits sequencing to proteins having fewer than approximately 50 amino acids.

Problem 29.19 Draw the structure of the *N*-phenylthiohydantoin formed by initial Edman degradation of each peptide: (a) Ala–Gly–Phe–Phe; (b) Val–Ile–Tyr.

29.6C Partial Hydrolysis of a Peptide

Additional structural information can be obtained by cleaving some, but not all, of the amide bonds in a peptide. Partial hydrolysis of a peptide with acid forms smaller fragments in a random fashion. Sequencing these peptides and identifying sites of overlap can be used to determine the sequence of the complete peptide, as shown in Sample Problem 29.2.

Sample Problem 29.2 Give the amino acid sequence of a hexapeptide that contains the amino acids Ala, Val, Ser, Ile, Gly, Tyr, and forms the following fragments when partially hydrolyzed with HCl: Gly–Ile–Val, Ala–Ser–Gly, and Tyr–Ala.

Solution

Looking for points of overlap in the sequences of the smaller fragments shows how the fragments should be pieced together. In this example, the fragment Ala–Ser–Gly contains amino acids common to the two other fragments, thus showing how the three fragments can be joined together.



Problem 29.20 Give the amino acid sequence of an octapeptide that contains the amino acids Tyr, Ala, Leu (2 equiv), Cys, Gly, Glu, and Val, and forms the following fragments when partially hydrolyzed with HCl: Val–Cys–Gly–Glu, Ala–Leu–Tyr, and Tyr–Leu–Val–Cys.

Peptides can also be hydrolyzed at specific sites using enzymes. The enzyme carboxypeptidase catalyzes the hydrolysis of the amide bond nearest the C-terminal end, forming the C-terminal amino acid and a peptide with one fewer amino acid. In this way, carboxypeptidase is used to identify the C-terminal amino acid.

Other enzymes catalyze the hydrolysis of amide bonds formed with specific amino acids. For example:

- **Trypsin catalyzes the hydrolysis of amides with a carbonyl group that is part of the basic amino acids arginine and lysine.**
- **Chymotrypsin hydrolyzes amides with carbonyl groups that are part of the aromatic amino acids phenylalanine, tyrosine, and tryptophan.**




Table 29.2 summarizes these enzyme specificities used in peptide sequencing.

Table 29.2 Cleavage Sites of Specific Enzymes in Peptide Sequencing

Enzyme	Site of cleavage
Carboxypeptidase	Amide bond nearest to the C-terminal amino acid
Chymotrypsin	Amide bond with a carbonyl group from Phe, Tyr, or Trp
Trypsin	Amide bond with a carbonyl group from Arg or Lys

Problem 29.21 (a) What products are formed when each peptide is treated with trypsin? (b) What products are formed when each peptide is treated with chymotrypsin?

[1] Gly–Ala–Phe–Leu–Lys–Ala
[2] Phe–Tyr–Gly–Cys–Arg–Ser
[3] Thr–Pro–Lys–Glu–His–Gly–Phe–Cys–Trp–Val–Val–Phe

Sample Problem 29.3 Deduce the sequence of a pentapeptide that contains the amino acids Ala, Glu, Gly, Ser, and Tyr, from the following experimental data. Edman degradation cleaves Gly from the pentapeptide, and carboxypeptidase forms Ala and a tetrapeptide. Treatment of the pentapeptide with chymotrypsin forms a dipeptide and a tripeptide. Partial hydrolysis forms Gly, Ser, and the tripeptide Tyr–Glu–Ala.

Solution

Use each result to determine the location of an amino acid in the pentapeptide.


Experiment		Result
• Edman degradation identifies the N-terminal amino acid—in this case, Gly.	→	Gly– __ – __ – __ – __
• Carboxypeptidase identifies the C-terminal amino acid (Ala) when it is cleaved from the end of the chain.	→	Gly– __ – __ – __ –Ala
• Chymotrypsin cleaves amide bonds that contain a carbonyl group from an aromatic amino acid—Tyr in this case. Because a dipeptide and tripeptide are obtained after treatment with chymotrypsin, Tyr must be the C-terminal amino acid of either the di- or tripeptide. As a result, Tyr must be either the second or third amino acid in the pentapeptide chain.	→	Gly–Tyr– __ – __ –Ala or Gly– __ –Tyr– __ –Ala
• Partial hydrolysis forms the tripeptide Tyr–Glu–Ala. Because Ala is the C-terminal amino acid, this result identifies the last three amino acids in the chain.	→	Gly– __ –Tyr–Glu–Ala
• The last amino acid, Ser, must be located at the only remaining position, the second amino acid in the pentapeptide, and the complete sequence is determined.	→	Gly–Ser–Tyr–Glu–Ala

Problem 29.22 Deduce the sequence of a heptapeptide that contains the amino acids Ala, Arg, Glu, Gly, Leu, Phe, and Ser, from the following experimental data. Edman degradation cleaves Leu from the heptapeptide, and carboxypeptidase forms Glu and a hexapeptide. Treatment of the heptapeptide with chymotrypsin forms a hexapeptide and a single amino acid. Treatment of the heptapeptide with trypsin forms a pentapeptide and a dipeptide. Partial hydrolysis forms Glu, Leu, Phe, and the tripeptides Gly–Ala–Ser and Ala–Ser–Arg.

29.7 Peptide Synthesis

The synthesis of a specific dipeptide, such as Ala–Gly from alanine and glycine, is complicated because both amino acids have two functional groups. As a result, four products—namely, Ala–Ala, Ala–Gly, Gly–Gly, and Gly–Ala—are possible.

From two amino acids… **…there are four possible dipeptides.**




How do we selectively join the COOH group of alanine with the NH_2 group of glycine?


- **Protect the functional groups that we don't want to react, and then form the amide bond.**

How To Synthesize a Dipeptide from Two Amino Acids

Example




Step [1] **Protect the NH_2 group of alanine.**




- In the neutral amino acid, the NH_2 group exists largely as an ammonium ion, $-NH_3^+$.

Step [2] **Protect the COOH group of glycine.**




- In the neutral amino acid, the COOH group exists largely as a carboxylate anion, $-COO^-$.


Step [3] **Form the amide bond with DCC.**




Dicyclohexylcarbodiimide (**DCC**) is a reagent commonly used to form amide bonds (see Section 22.10D). DCC makes the OH group of the carboxylic acid a better leaving group, thus **activating the carboxy group toward nucleophilic attack.**




Step [4] **Remove one or both protecting groups.**




Two widely used amino protecting groups convert an amine into a **carbamate,** a functional group having a carbonyl bonded to both an oxygen and a nitrogen atom. Since the N atom of the carbamate is bonded to a carbonyl group, the protected amino group is no longer nucleophilic.




For example, the ***tert*-butoxycarbonyl protecting group,** abbreviated as **Boc,** is formed by reacting the amino acid with di-*tert*-butyl dicarbonate in a nucleophilic acyl substitution reaction.







tert-butoxycarbonyl
Boc

To be a useful protecting group, the Boc group must be removed under reaction conditions that do not affect other functional groups in the molecule. It can be removed with an acid such as **trifluoroacetic acid, HCl,** or **HBr.**




$(Boc)_2O$






9-fluorenylmethoxycarbonyl
Fmoc

A second amino protecting group, the **9-fluorenylmethoxycarbonyl protecting group,** abbreviated as **Fmoc,** is formed by reacting the amino acid with 9-fluorenylmethyl chloroformate in a nucleophilic acyl substitution reaction.




9-fluorenylmethyl
chloroformate
Fmoc – Cl


While the Fmoc protecting group is stable to most acids, it can be removed by treatment with base (NH_3 or an amine).




The carboxy group is usually protected as a **methyl** or **benzyl ester** by reaction with an alcohol and an acid.



These esters are usually removed by hydrolysis with aqueous base.




One advantage of using a benzyl ester for protection is that it can also be removed with H_2 in the presence of a Pd catalyst. This process is called **hydrogenolysis.** These conditions are especially mild, because they avoid the use of either acid or base. Benzyl esters can also be removed with HBr in acetic acid.




The specific reactions needed to synthesize the dipeptide Ala–Gly are illustrated in Sample Problem 29.4.

Sample Problem 29.4 Draw out the steps in the synthesis of the dipeptide Ala–Gly.




Solution


Step [1] **Protect the NH_2 group of alanine using a Boc group.**



Step [2] **Protect the COOH group of glycine as a benzyl ester.**




Step [3] **Form the amide bond with DCC.**




Step [4] **Remove one or both protecting groups.**

The protecting groups can be removed in a stepwise fashion, or in a single reaction.




This method can be applied to the synthesis of tripeptides and even larger polypeptides. After the protected dipeptide is prepared in Step [3], only one of the protecting groups is removed, and this dipeptide is coupled to a third amino acid with one of its functional groups protected, as illustrated in the following equations.



Problem 29.23 Devise a synthesis of each peptide from amino acid starting materials: (a) Leu–Val; (b) Ala–Ile–Gly.

Problem 29.24 Devise a synthesis of the following dipeptide from amino acid starting materials.




29.8 Automated Peptide Synthesis

Development of the solid phase technique earned Merrifield the 1984 Nobel Prize in Chemistry and has made possible the synthesis of many polypeptides and proteins.

The method described in Section 29.7 works well for the synthesis of small peptides. It is extremely time-consuming to synthesize larger proteins by this strategy, however, because each step requires isolation and purification of the product. The synthesis of larger polypeptides is usually accomplished by using the **solid phase technique** originally developed by R. Bruce Merrifield of Rockefeller University.


In the **Merrifield method** an amino acid is attached to an **insoluble polymer.** Amino acids are sequentially added, one at a time, thereby forming successive peptide bonds. Because impurities and by-products are not attached to the polymer chain, they are removed simply by washing them away with a solvent at each stage of the synthesis.

A commonly used polymer is a **polystyrene derivative** that contains $-CH_2Cl$ groups bonded to some of the benzene rings in the polymer chain. The Cl atoms serve as handles that allow attachment of amino acids to the chain.



polystyrene derivative with Cl leaving groups

An Fmoc-protected amino acid is attached to the polymer at its carboxy group by an S_N2 reaction.




The amino acid is now bound to the insoluble polymer.

Once the first amino acid is bound to the polymer, additional amino acids can be added sequentially. The steps of the solid phase peptide synthesis technique are illustrated in the accompanying scheme. In the last step, HF cleaves the polypeptide chain from the polymer.


How To Synthesize a Peptide Using the Merrifield Solid Phase Technique

Step [1]
Attach an Fmoc-protected amino acid to the polymer.



new bond to the polymer in red


Step [2]
Remove the protecting group.



free amino group


How To, continued . . .

Step [3]
Form the amide bond with DCC.




new amide bond in red

Step [4]
Repeat Steps [2] and [3].



new amide bond in red

Step [5]
Remove the protecting group and detach the peptide from the polymer.



tripeptide

The Merrifield method has now been completely automated, so it is possible to purchase peptide synthesizers that automatically carry out all of the above operations and form polypeptides in high yield in a matter of hours, days, or weeks, depending on the length of the chain of the desired product. For example, the protein ribonuclease, which contains 128 amino acids, has been prepared by this technique in an overall yield of 17%. This remarkable synthesis involved 369 separate reactions, and thus the yield of each individual reaction was > 99%.

Problem 29.25 Outline the steps needed to synthesize the tetrapeptide Ala–Leu–Ile–Gly using the Merrifield technique.


29.9 Protein Structure

Now that you have learned some of the chemistry of amino acids, it's time to study proteins, the large polymers of amino acids that are responsible for so much of the structure and function of all living cells. We begin with a discussion of the **primary, secondary, tertiary, and quaternary structure** of proteins.

29.9A Primary Structure

The *primary structure* of proteins is the particular sequence of amino acids that is joined together by peptide bonds. The most important element of this primary structure is the **amide bond.**


- Rotation around the amide C–N bond is *restricted* because of electron delocalization, and the *s*-trans conformation is the more stable arrangement.
- In each peptide bond, the N–H and C=O bonds are directed 180° from each other.



Although rotation about the amide bonds is restricted, **rotation about the other σ bonds in the protein backbone is not.** As a result, the peptide chain can twist and bend into a variety of different arrangements that constitute the secondary structure of the protein.

29.9B Secondary Structure

The three-dimensional conformations of localized regions of a protein are called its secondary structure. These regions arise due to hydrogen bonding between the N–H proton of one amide and C=O oxygen of another. Two arrangements that are particularly stable are called the **α-helix** and the **β-pleated sheet.**



α-Helix

The **α-helix** forms when a peptide chain twists into a right-handed or clockwise spiral, as shown in Figure 29.8. Four important features of the α-helix are as follows:

[1] **Each turn of the helix has 3.6 amino acids.**

[2] **The N–H and C=O bonds point along the axis of the helix.** All C=O bonds point in one direction, and all N–H bonds point in the opposite direction.

[3] **The C=O group of one amino acid is hydrogen bonded to an N–H group four amino acid residues farther along the chain.** Thus, hydrogen bonding occurs between two amino acids ***in the same chain.*** Note, too, that the hydrogen bonds are parallel to the axis of the helix.


[4] The **R groups of the amino acids extend outward** from the core of the helix.

An α-helix can form only if there is rotation about the bonds at the α carbon of the amide carbonyl group, and not all amino acids can do this. For example, proline, the amino acid whose nitrogen atom forms part of a five-membered ring, is more rigid than other amino acids, and its C_α–N bond cannot rotate the necessary amount. Additionally, it has no N–H proton with which to form an intramolecular hydrogen bond to stabilize the helix. Thus, proline cannot be part of an α-helix.

Both the myosin in muscle and α-keratin in hair are proteins composed almost entirely of α-helices.


Figure 29.8
Two different illustrations of the α-helix

a. The right-handed α-helix



All atoms of the α-helix are drawn in this representation. All C=O bonds are pointing up and all N−H bonds are pointing down.

b. The backbone of the α-helix



Only the peptide backbone is drawn in this representation. The hydrogen bonds between the C=O and N−H of amino acids four residues away from each other are shown.

β-Pleated Sheet


The **β-pleated sheet** secondary structure forms when two or more peptide chains, called **strands,** line up side-by-side, as shown in Figure 29.9. All β-pleated sheets have the following characteristics:

[1] **The C=O and N−H bonds lie in the plane of the sheet.**

[2] **Hydrogen bonding often occurs between the N−H and C=O groups of nearby amino acid residues.**

[3] The **R groups are oriented above and below the plane** of the sheet, and alternate from one side to the other along a given strand.

Figure 29.9
Three-dimensional structure of the β-pleated sheet



- The β-pleated sheet consists of extended strands of the peptide chains held together by hydrogen bonding. The C=O and N−H bonds lie in the plane of the sheet, and the R groups (shown as orange balls) alternate above and below the plane.


The β-pleated sheet arrangement most commonly occurs with amino acids with small R groups, like alanine and glycine. With larger R groups steric interactions prevent the chains from getting close together and so the sheet cannot be stabilized by hydrogen bonding.

The peptide strands of β-pleated sheets can actually be oriented in two different ways, as shown in Figure 29.10.

- In a *parallel* β-pleated sheet, the strands run in the *same* direction from the N- to C-terminal amino acid.
- In an *antiparallel* β-pleated sheet, the strands run in the *opposite* direction.

Figure 29.10 The parallel and antiparallel forms of the β-pleated sheet


Parallel β-pleated sheet



The two peptide chains are arranged in the same direction. Hydrogen bonds occur between N–H and C=O bonds in adjacent chains.

[Note: R groups on the carbon chain are omitted for clarity.]

Antiparallel β-pleated sheet




The two peptide chains are arranged in opposite directions. Hydrogen bonding between the N–H and C=O groups still holds the two chains together.

Most proteins have regions of α-helix and β-pleated sheet, in addition to other regions that cannot be characterized by either of these arrangements. Shorthand symbols are often used to indicate regions of a protein that have α-helix or β-pleated sheet. A **flat helical ribbon** is used for the α-helix, and a **flat wide arrow** is used for the β-pleated sheet. These representations are often used in **ribbon diagrams** to illustrate protein structure.



α-helix shorthand




β-pleated sheet shorthand

Proteins are drawn in a variety of ways to illustrate different aspects of their structure. Figure 29.11 illustrates three different representations of the protein lysozyme, an enzyme found in both plants and animals. Lysozyme catalyzes the hydrolysis of bonds in bacterial cell walls, weakening them, often causing the bacteria to burst.


Spider dragline silk is a strong yet elastic protein because it has regions of β-pleated sheet and regions of α-helix (Figure 29.12). α-Helical regions impart elasticity to the silk because the peptide chain is twisted (not fully extended), so it can stretch. β-Pleated sheet regions are almost fully extended, so they can't be stretched further, but their highly ordered three-dimensional structure imparts strength to the silk. Thus, spider silk suits the spider by comprising both types of secondary structure with beneficial properties.

Figure 29.11 Lysozyme

a. Ball-and-stick model



b. Space-filling model




c. Ribbon diagram



(a) The ball-and-stick model of lysozyme shows the protein backbone with color-coded C, N, O, and S atoms. Individual amino acids are most clearly located using this representation. (b) The space-filling model uses color-coded balls for each atom in the backbone of the enzyme and illustrates how the atoms fill the space they occupy. (c) The ribbon diagram shows regions of α-helix and β-pleated sheet that are not clearly in evidence in the other two representations.

Figure 29.12
Different regions of secondary structure in spider silk



Spider silk has regions of α-helix and β-pleated sheet that make it both strong and elastic. The green coils represent the α-helical regions, and the purple arrows represent the β-pleated sheet regions. The yellow lines represent other areas of the protein that are neither α-helix nor β-pleated sheet.


Problem 29.26 Consider two molecules of a tetrapeptide composed of only alanine residues. Draw the hydrogen bonding interactions that result when these two peptides adopt a parallel β-pleated sheet arrangement. Answer this same question for the antiparallel β-pleated sheet arrangement.

29.9C Tertiary and Quaternary Structure

The three-dimensional shape adopted by the entire peptide chain is called its tertiary structure. A peptide generally folds into a conformation that maximizes its stability. In the aqueous environment of the cell, proteins often fold in such a way as to place a large number of polar and charged groups on their outer surface, to maximize the dipole–dipole and hydrogen bonding interactions with water. This generally places most of the nonpolar side chains in the interior of the protein, where van der Waals interactions between these hydrophobic groups help stabilize the molecule, too.

In addition, polar functional groups hydrogen bond with each other (not just water), and amino acids with charged side chains like $-COO^-$ and $-NH_3^+$ can stabilize tertiary structure by electrostatic interactions.

Finally, **disulfide bonds are the only covalent bonds that stabilize tertiary structure.** As previously mentioned, these strong bonds form by oxidation of two cysteine residues either on the same polypeptide chain or another polypeptide chain of the same protein.




disulfide bond between two SH groups on the same chain


disulfide bond between two SH groups on different chains

The nonapeptides **oxytocin** and **vasopressin** (Section 29.5C) contain intramolecular disulfide bonds. **Insulin,** on the other hand, consists of two separate polypeptide chains (**A** and **B**) that are covalently linked by two intermolecular disulfide bonds, as shown in Figure 29.13. The **A** chain, which also has an intramolecular disulfide bond, has 21 amino acid residues, whereas the **B** chain has 30.

Figure 29.13
Insulin



Insulin is a small protein consisting of two polypeptide chains (designated as the **A** and **B** chains) held together by two disulfide bonds. An additional disulfide bond joins two cysteine residues within the **A** chain.



Synthesized by groups of cells in the pancreas called the islets of Langerhans, insulin is the protein that regulates the levels of glucose in the blood. Insufficiency of insulin results in diabetes. Many of the abnormalities associated with this disease can be controlled by the injection of insulin. Until the availability of human insulin through genetic engineering techniques, all insulin used by diabetics was obtained from pigs and cattle. The amino acid sequences of these insulin proteins is slightly different from that of human insulin. Pig insulin differs in one amino acid only, whereas bovine insulin has three different amino acids. This is shown in the accompanying table.

	Chain A			Chain B
Position of residue →	**8**	**9**	**10**	**30**
Human insulin	Thr	Ser	Ile	Thr
Pig insulin	Thr	Ser	Ile	Ala
Bovine insulin	Ala	Ser	Val	Ala

Figure 29.14
The stabilizing interactions in secondary and tertiary protein structure





Figure 29.14 schematically illustrates the many different kinds of intramolecular forces that stabilize the secondary and tertiary structures of polypeptide chains.

The shape adopted when two or more folded polypeptide chains aggregate into one protein complex is called the **quaternary structure** of the protein. Each individual polypeptide chain is called a **subunit** of the overall protein. **Hemoglobin,** for example, consists of two α and two β subunits held together by intermolecular forces in a compact three-dimensional shape. The unique function of hemoglobin is possible only when all four subunits are together.

The four levels of protein structure are summarized in Figure 29.15.

Figure 29.15 The primary, secondary, tertiary, and quaternary structure of proteins



Problem 29.27 What types of stabilizing interactions exist between each of the following pairs of amino acids?

a. Ser and Tyr b. Val and Leu c. two Phe residues

Problem 29.28 The fibroin proteins found in silk fibers consist of large regions of β-pleated sheets stacked one on top of another. (a) Explain why having a glycine at every other residue allows the β-pleated sheets to stack on top of each other. (b) Why are silk fibers insoluble in water?

29.10 Important Proteins

Proteins are generally classified according to their three-dimensional shapes.

- **Fibrous proteins** are composed of long linear polypeptide chains that are bundled together to form rods or sheets. These proteins are insoluble in water and serve structural roles, giving strength and protection to tissues and cells.
- **Globular proteins** are coiled into compact shapes with hydrophilic outer surfaces that make them water soluble. Enzymes and transport proteins are globular to make them soluble in the blood and other aqueous environments in cells.

29.10A α-Keratins

α-Keratins are the proteins found in hair, hooves, nails, skin, and wool. They are composed almost exclusively of long sections of α-helix units, having large numbers of alanine and leucine residues. Because these nonpolar amino acids extend outward from the α-helix, these proteins are very water insoluble. Two α-keratin helices coil around each other, forming a structure called a **supercoil** or **superhelix.** These, in turn, form larger and larger bundles of fibers, ultimately forming a strand of hair, as shown schematically in Figure 29.16.





Figure 29.16 Anatomy of a hair—It begins with α-keratin.

α-Keratins also have a number of cysteine residues, and because of this, disulfide bonds are formed between adjacent helices. The number of disulfide bridges determines the strength of the material. Claws, horns, and fingernails have extensive networks of disulfide bonds, making them extremely hard.

Straight hair can be made curly by cleaving the disulfide bonds in α-keratin, and then rearranging and re-forming them, as shown schematically in Figure 29.17. First, the disulfide bonds in the straight hair are reduced to thiol groups, so the bundles of α-keratin chains are no longer held in their specific "straight" orientation. Then, the hair is wrapped around curlers and treated with an oxidizing agent that converts the thiol groups back to disulfide bonds, now with twists and turns in the keratin backbone. This makes the hair look curly and is the chemical basis for a "permanent."

Figure 29.17
The chemistry of a "permanent"—Making straight hair curly




To make straight hair curly, the disulfide bonds holding the α-helical chains together are cleaved by reduction. This forms free thiol groups (–SH). The hair is turned around curlers and then an oxidizing agent is applied. This re-forms the disulfide bonds in the hair, but between different thiol groups, now giving it a curly appearance.

29.10B Collagen

Collagen, the most abundant protein in vertebrates, is found in connective tissues such as bone, cartilage, tendons, teeth, and blood vessels. Glycine and proline account for a large fraction of its amino acid residues, whereas cysteine accounts for very little. Because of the high proline content, it cannot form a right-handed α-helix. Instead, it forms an elongated left-handed helix, and then three of these helices wind around each other to form a right-handed **superhelix** or **triple helix.** The side chain of glycine is only a hydrogen atom, so the high glycine content allows the collagen superhelices to lie compactly next to each other, thus stabilizing the superhelices via hydrogen bonding. Two views of the collagen superhelix are shown in Figure 29.18.

Figure 29.18
Two different representations for the triple helix of collagen




- In collagen, three polypeptide chains having an unusual left-handed helix wind around each other in a right-handed triple helix. The high content of small glycine residues allows the chains to lie close to each other, permitting hydrogen bonding between the chains.

29.10C Hemoglobin and Myoglobin

Hemoglobin and **myoglobin,** two globular proteins, are called **conjugated proteins** because they are composed of a protein unit and a nonprotein molecule called a **prosthetic group.** The prosthetic group in hemoglobin and myoglobin is **heme,** a complex organic compound containing the


Figure 29.19
Protein ribbon diagrams for myoglobin and hemoglobin

a. Myoglobin




Myoglobin consists of a single polypeptide chain with a heme unit shown in a ball-and-stick model.

b. Hemoglobin



Hemoglobin consists of two α and two β chains shown in red and blue, respectively, and four heme units shown in ball-and-stick models.

Fe^{2+} ion complexed with a nitrogen heterocycle called a **porphyrin.** The Fe^{2+} ion of hemoglobin and myoglobin binds oxygen in the blood. Hemoglobin, which is present in red blood cells, transports oxygen to wherever it is needed in the body, whereas myoglobin stores oxygen in tissues. Ribbon diagrams for myoglobin and hemoglobin are shown in Figure 29.19.





When red blood cells take on a "sickled" shape in persons with sickle cell disease, they occlude capillaries (causing organ injury) and they break easily (leading to profound anemia). This devastating illness results from the change of a single amino acid in hemoglobin. Note the single sickled cell surrounded by three red cells with normal morphology.

Myoglobin, the chapter-opening molecule, has 153 amino acid residues in a single polypeptide chain. It has eight separate α-helical sections that fold back on one another, with the prosthetic heme group held in a cavity inside the polypeptide. Most of the polar residues are found on the outside of the protein so that they can interact with the water solvent. Spaces in the interior of the protein are filled with nonpolar amino acids. Myoglobin gives cardiac muscle its characteristic red color.

Hemoglobin consists of four polypeptide chains (two α subunits and two β subunits), each of which carries a heme unit. Hemoglobin has more nonpolar amino acids than myoglobin. When each subunit is folded, some of these remain on the surface. The van der Waals attraction between these hydrophobic groups is what stabilizes the quaternary structure of the four subunits.

Carbon monoxide is poisonous because it binds to the Fe^{2+} of hemoglobin more strongly than does oxygen. Hemoglobin complexed with CO cannot carry O_2 from the lungs to the tissues. Without O_2 in the tissues for metabolism, cells cannot function, so they die.

The properties of all proteins depend on their three-dimensional shape, and their shape depends on their primary structure—that is, their amino acid sequence. This is particularly well exemplified by comparing normal hemoglobin with **sickle cell hemoglobin,** a mutant variation in which a single amino acid of both β subunits is changed from glutamic acid to valine. The replacement of one acidic amino acid (Glu) with one nonpolar amino acid (Val) changes the shape of hemoglobin, which has profound effects on its function. Deoxygenated red blood cells with sickle cell hemoglobin become elongated and crescent shaped, and they are unusually fragile. As a result, they do not

flow easily through capillaries, causing pain and inflammation, and they break open easily, leading to severe anemia and organ damage. The end result is often a painful and premature death.


This disease, called **sickle cell anemia,** is found almost exclusively among people originating from central and western Africa, where malaria is an enormous health problem. Sickle cell hemoglobin results from a genetic mutation in the DNA sequence that is responsible for the synthesis of hemoglobin. Individuals who inherit this mutation from both parents develop sickle cell anemia, whereas those who inherit it from only one parent are said to have the sickle cell trait. They do not develop sickle cell anemia and they are more resistant to malaria than individuals without the mutation. This apparently accounts for this detrimental gene being passed on from generation to generation.

KEY CONCEPTS


Amino Acids and Proteins

Synthesis of Amino Acids (29.2)

[1] From α-halo carboxylic acids by S_N2 reaction




[2] By alkylation of diethyl acetamidomalonate



- Alkylation works best with unhindered alkyl halides—that is, CH_3X and RCH_2X.

[3] Strecker synthesis




Preparation of Optically Active Amino Acids


[1] Resolution of enantiomers by forming diastereomers (29.3A)

- Convert a racemic mixture of amino acids into a racemic mixture of *N*-acetyl amino acids [(*S*)- and (*R*)-$CH_3CONHCH(R)COOH$].
- Treat the enantiomers with a chiral amine to form a mixture of diastereomers.
- Separate the diastereomers.
- Regenerate the amino acids by protonation of the carboxylate salt and hydrolysis of the *N*-acetyl group.

[2] Kinetic resolution using enzymes (29.3B)



[3] By enantioselective hydrogenation (29.4)



S enantiomer *S* amino acid


Rh* = chiral Rh hydrogenation catalyst

Summary of Methods Used for Peptide Sequencing (29.6)


- Complete hydrolysis of all amide bonds in a peptide gives the identity and amount of the individual amino acids.
- Edman degradation identifies the N-terminal amino acid. Repeated Edman degradations can be used to sequence a peptide from the N-terminal end.
- Cleavage with carboxypeptidase identifies the C-terminal amino acid.
- Partial hydrolysis of a peptide forms smaller fragments that can be sequenced. Amino acid sequences common to smaller fragments can be used to determine the sequence of the complete peptide.
- Selective cleavage of a peptide occurs with trypsin and chymotrypsin to identify the location of specific amino acids (Table 29.2).

Adding and Removing Protecting Groups for Amino Acids (29.7)


[1] Protection of an amino group as a Boc derivative




[2] Deprotection of a Boc-protected amino acid




[3] Protection of an amino group as an Fmoc derivative




[4] Deprotection of an Fmoc-protected amino acid



[5] Protection of a carboxy group as an ester




[6] Deprotection of an ester group



Synthesis of Dipeptides (29.7)

[1] Amide formation with DCC



[2] Four steps are needed to synthesize a dipeptide:

a. **Protect** the amino group of one amino acid with a Boc or Fmoc group.
b. **Protect** the carboxy group of the second amino acid as an ester.
c. Form the amide bond with **DCC.**
d. **Remove both protecting groups** in one or two reactions.


Summary of the Merrifield Method of Peptide Synthesis (29.8)

[1] Attach an Fmoc-protected amino acid to a polymer derived from polystyrene.
[2] Remove the Fmoc protecting group.
[3] Form the amide bond with a second Fmoc-protected amino acid by using DCC.
[4] Repeat steps [2] and [3].
[5] Remove the protecting group and detach the peptide from the polymer.


PROBLEMS

Problems Using Three-Dimensional Models


29.29 Draw the product formed when the following amino acid is treated with each reagent: (a) CH_3OH, H^+; (b) CH_3COCl, pyridine; (c) HCl (1 equiv); (d) NaOH (1 equiv); (e) $C_6H_5N{=}C{=}S$.



29.30 With reference to the following peptide: (a) Identify the N-terminal and C-terminal amino acids. (b) Name the peptide using one-letter abbreviations. (c) Label all the amide bonds in the peptide backbone.




29.31 Devise a synthesis of the following dipeptide from amino acid starting materials.



Amino Acids

29.32



penicillamine

a. (*S*)-Penicillamine, an amino acid that does not occur in proteins, is used as a copper chelating agent to treat Wilson's disease, an inherited defect in copper metabolism. (*R*)-Penicillamine is toxic, sometimes causing blindness. Draw the structures of (*R*)- and (*S*)-penicillamine.
b. What disulfide is formed from oxidation of (*S*)-penicillamine?

29.33 Histidine is classified as a basic amino acid because one of the N atoms in its five-membered ring is readily protonated by acid. Which N atom in histidine is protonated and why?

29.34 Tryptophan is not classified as a basic amino acid even though it has a heterocycle containing a nitrogen atom. Why is the N atom in the five-membered ring of tryptophan not readily protonated by acid?

29.35 What is the structure of each amino acid at its isoelectric point: (a) alanine; (b) methionine; (c) aspartic acid; (d) lysine?

29.36 What is the predominant form of each of the following amino acids at pH = 1? What is the overall charge on the amino acid at this pH? (a) threonine; (b) methionine; (c) aspartic acid; (d) arginine


29.37 What is the predominant form of each of the following amino acids at pH = 11? What is the overall charge on the amino acid? (a) valine; (b) proline; (c) glutamic acid; (d) lysine

29.38 a. Draw the structure of the tripeptide A–A–A, and label the two ionizable functional groups.
b. What is the predominant form of A–A–A at pH = 1?
c. The pK_a values for the two ionizable functional groups (3.39 and 8.03) differ considerably from the pK_a values of alanine (2.35 and 9.87; see Table 29.1). Account for the observed pK_a differences.


Synthesis and Reactions of Amino Acids

29.39 Draw the organic products formed in each reaction.


a.




b.



c.



d.




29.40 What alkyl halide is needed to synthesize each amino acid from diethyl acetamidomalonate: (a) Asn; (b) His; (c) Trp?


29.41 Devise a synthesis of threonine from diethyl acetamidomalonate.

29.42 Devise a synthesis of each amino acid from acetaldehyde (CH_3CHO): (a) glycine; (b) alanine.


29.43 Identify the lettered intermediates in the following reaction scheme. This is an alternative method to synthesize amino acids, based on the Gabriel synthesis of 1° amines (Section 25.7A).



29.44 Glutamic acid is synthesized by the following reaction sequence. Draw a stepwise mechanism for Steps [1]–[3].




29.45 Identify **A–E** in the following reaction sequence.




Resolution; The Synthesis of Chiral Amino Acids

29.46 Write out a scheme for the resolution of the two enantiomers of the antiplatelet drug clopidogrel with 10-camphorsulfonic acid.




clopidogrel




10-camphorsulfonic acid

29.47 Another strategy used to resolve amino acids involves converting the carboxy group to an ester and then using a *chiral carboxylic acid* to carry out an acid–base reaction at the free amino group. Using a racemic mixture of alanine enantiomers and (*R*)-mandelic acid as resolving agent, write out the steps showing how a resolution process would occur.




alanine
(two enantiomers)




(*R*)-mandelic acid

29.48 Brucine is a poisonous alkaloid obtained from *Strychnos nux vomica,* a tree that grows in India, Sri Lanka, and northern Australia. Write out a resolution scheme similar to the one given in Section 29.3A, which shows how a racemic mixture of phenylalanine can be resolved using brucine.




brucine

29.49 Draw the organic products formed in each reaction.



racemic mixture

29.50 What steps are needed to convert **A** to L-dopa, an uncommon amino acid that is effective in treating Parkinson's disease? These steps are the key reactions in the first commercial asymmetric synthesis using a chiral transition metal catalyst. This process was developed at Monsanto in 1974.



A L-dopa


Peptide Structure and Sequencing

29.51 Draw the structure for each peptide: (a) Phe–Ala; (b) Gly–Gln; (c) Lys–Gly; (d) R – H.


29.52 For the tetrapeptide Asp–Arg–Val–Tyr:

a. Name the peptide using one-letter abbreviations.
b. Draw the structure.
c. Label all amide bonds.
d. Label the N-terminal and C-terminal amino acids.

29.53 Name each peptide using both the three-letter and one-letter abbreviations of the component amino acids.




29.54 Gramicidin S, a topical antibiotic produced by the bacterium *Bacillus brevis*, is a cyclic decapeptide formed from five amino acids. Draw the structures of the amino acids that form gramicidin S, and explain why this compound possesses two unusual structural features.



gramicidin S

29.55 Draw the amino acids and peptide fragments formed when the decapeptide A–P–F–L–K–W–S–G–R–G is treated with each reagent or enzyme: (a) chymotrypsin; (b) trypsin; (c) carboxypeptidase; (d) $C_6H_5N{=}C{=}S$.

29.56 Consider the decapeptide angiotensin I.



angiotensin I

a. What products are formed when angiotensin I is treated with trypsin?
b. What products are formed when angiotensin I is treated with chymotrypsin?
c. Treatment of angiotensin I with ACE (the angiotensin-converting enzyme) cleaves only the amide bond with the carbonyl group derived from phenylalanine to afford two products. The larger polypeptide is angiotensin II, a hormone that narrows blood vessels and increases blood pressure. Give the amino acid sequence of angiotensin II using three-letter abbreviations. ACE inhibitors are drugs that lower blood pressure by inhibiting the ACE enzyme (Problem 5.15).


29.57 Give the amino acid sequence of each peptide using the fragments obtained by partial hydrolysis of the peptide with acid.
a. a tetrapeptide that contains Ala, Gly, His, and Tyr, which is hydrolyzed to the dipeptides His–Tyr, Gly–Ala, and Ala–His
b. a pentapeptide that contains Glu, Gly, His, Lys, and Phe, which is hydrolyzed to His–Gly–Glu, Gly–Glu–Phe, and Lys–His

29.58 Use the given experimental data to deduce the sequence of an octapeptide that contains the following amino acids: Ala, Gly (2 equiv), His (2 equiv), Ile, Leu, and Phe. Edman degradation cleaves Gly from the octapeptide, and carboxypeptidase forms Leu and a heptapeptide. Partial hydrolysis forms the following fragments: Ile–His–Leu, Gly, Gly–Ala–Phe–His, and Phe–His–Ile.

29.59 An octapeptide contains the following amino acids: Arg, Glu, His, Ile, Leu, Phe, Tyr, and Val. Carboxypeptidase treatment of the octapeptide forms Phe and a heptapeptide. Treatment of the octapeptide with chymotrypsin forms two tetrapeptides, **A** and **B.** Treatment of **A** with trypsin yields two dipeptides, **C** and **D.** Edman degradation cleaves the following amino acids from each peptide: Glu (octapeptide), Glu (**A**), Ile (**B**), Glu (**C**), and Val (**D**). Partial hydrolysis of tetrapeptide **B** forms Ile–Leu in addition to other products. Deduce the structure of the octapeptide and fragments **A–D.**

Peptide Synthesis

29.60 Draw the organic products formed in each reaction.



29.61 Draw all the steps in the synthesis of each peptide from individual amino acids: (a) Gly–Ala; (b) Ile–Ala–Phe.


29.62 Write out the steps for the synthesis of each peptide using the Merrifield method: (a) Ala–Leu–Phe–Phe; (b) Phe–Gly–Ala–Ile.

29.63 Besides the Boc and Fmoc protecting groups used in peptide synthesis, amines can also be protected by reaction with benzyl chloroformate ($C_6H_5CH_2OCOCl$). Draw the structure of the product formed by reaction of alanine with benzyl chloroformate.

29.64 Another method to form a peptide bond involves a two-step process:

[1] Conversion of a Boc-protected amino acid to a *p*-nitrophenyl ester.


[2] Reaction of the *p*-nitrophenyl ester with an amino acid ester.



p-nitrophenyl ester


a. Why does a *p*-nitrophenyl ester "activate" the carboxy group of the first amino acid to amide formation?

b. Would a *p*-methoxyphenyl ester perform the same function? Why or why not?



p-methoxyphenyl ester


29.65 Draw the mechanism for the reaction that removes an Fmoc group from an amino acid under the following conditions:



Proteins

29.66 Which of the following amino acids are typically found in the interior of a globular protein, and which are typically found on the surface: (a) phenylalanine; (b) aspartic acid; (c) lysine; (d) isoleucine; (e) arginine; (f) glutamic acid?

29.67 After the peptide chain of collagen has been formed, many of the proline residues are hydroxylated on one of the ring carbon atoms. Why is this process important for the triple helix of collagen?




Challenge Problems


29.68 Devise a stepwise synthesis of the tripeptide Val–Leu–Val from 3-methylbutanal [$(CH_3)_2CHCH_2CHO$] as the only organic starting material. You may also use any required inorganic or organic reagents.

29.69 Besides asymmetric hydrogenation (Section 29.4), several other methods are now available for the synthesis of optically active amino acids. How might a reaction like the Strecker synthesis be adapted to the preparation of chiral amino acids?

29.70 The anti-obesity drug orlistat works by irreversibly inhibiting pancreatic lipase, an enzyme responsible for the hydrolysis of triacylglycerols in the intestines, so they are excreted without metabolism. Inhibition occurs by reaction of orlistat with a serine residue of the enzyme, forming a covalently bound, inactive enzyme product. Draw the structure of the product formed during inhibition.



29.71 As shown in Mechanism 29.2, the final steps in the Edman degradation result in rearrangement of a thiazolinone to an *N*-phenylthiohydantoin. Draw a stepwise mechanism for this acid-catalyzed reaction.



thiazolinone *N*-phenylthiohydantoin

30 Synthetic Polymers

30.1 Introduction
30.2 Chain-growth polymers—Addition polymers
30.3 Anionic polymerization of epoxides
30.4 Ziegler–Natta catalysts and polymer stereochemistry
30.5 Natural and synthetic rubbers
30.6 Step-growth polymers—Condensation polymers
30.7 Polymer structure and properties
30.8 Green polymer synthesis
30.9 Polymer recycling and disposal



Polyethylene terephthalate (PET) is a synthetic polymer formed by the reaction of ethylene glycol ($HOCH_2CH_2OH$) and terephthalic acid. Because PET is lightweight and impervious to air and moisture, it is commonly used for transparent soft drink containers. PET is also used to produce synthetic fibers, sold under the trade name of Dacron. Of the six most common synthetic polymers, PET is the most easily recycled, in part because beverage bottles that bear the recycling code "1" are composed almost entirely of PET. Recycled polyethylene terephthalate is used for fleece clothing and carpeting. In Chapter 30, we learn about the preparation and properties of synthetic polymers like polyethylene terephthalate.

Chapter 30 discusses polymers, large organic molecules composed of repeating units—called **monomers**—that are covalently bonded together. Polymers occur naturally, as in the polysaccharides and proteins of Chapters 28 and 29, respectively, or they are synthesized in the laboratory.

This chapter concentrates on **synthetic polymers,** and expands on the material already presented in Chapters 15 and 22. Thousands of synthetic polymers have now been prepared. While some exhibit properties that mimic naturally occurring compounds, many others have unique properties. Although all polymers are large molecules, the size and branching of the polymer chain and the identity of the functional groups all contribute to determining an individual polymer's properties, thus making it suited for a particular product.

30.1 Introduction


A **polymer** is a large organic molecule composed of repeating units—called **monomers**—that are covalently bonded together. The word *polymer* is derived from the Greek words *poly* + *meros* meaning "many parts."

Polymerization is the joining together of monomers to make polymers.

Synthetic polymers are perhaps more vital to the fabric of modern society than any other group of compounds prepared in the laboratory. Nylon backpacks and polyester clothing, car bumpers and CD cases, milk jugs and grocery bags, artificial heart valves and condoms—all these products and innumerable others are made of synthetic polymers. Since 1976, the U.S. production of synthetic polymers has exceeded its steel production. Figure 30.1 illustrates several consumer products and the polymers from which they are made.


Synthetic polymers can be classified as chain-growth or step-growth polymers.

- **Chain-growth polymers, also called addition polymers, are prepared by chain reactions.**




These compounds are formed by adding monomers to the growing end of a polymer chain. The conversion of vinyl chloride to poly(vinyl chloride) is an example of chain-growth polymerization. These reactions were introduced in Section 15.14.

Figure 30.1 Polymers in some common consumer products



- We are surrounded by synthetic polymers in our daily lives. This cyclist rides on synthetic rubber tires, drinks from a polyethylene water bottle, wears a protective Lexan helmet and goggles, and uses a lightweight nylon backpack.

- **Step-growth polymers, also called condensation polymers, are formed when monomers containing two functional groups come together and lose a small molecule such as H_2O or HCl.**



monomers **polymer**


In this method, any two reactive molecules can combine, so the monomer is not necessarily added to the end of a growing chain. Step-growth polymerization is used to prepare polyamides and polyesters, as discussed in Section 22.16.

In contrast to many of the organic molecules encountered in Chapters 1–27, which have molecular weights much less than 1000 grams per mole (g/mol), polymers generally have high molecular weights, ranging from 10,000 to 1,000,000 grams per mole (g/mol). Synthetic polymers are really mixtures of individual polymer chains of varying lengths, so the reported molecular weight is an average value based on the average size of the polymer chain.

By convention, we often simplify the structure of a polymer by placing brackets around the repeating unit that forms the chain, as shown in Figure 30.2.


Problem 30.1 Give the shorthand structures of poly(vinyl chloride) and nylon 6,6 in Section 30.1.

Figure 30.2
Drawing a polymer in a shorthand representation



30.2 Chain-Growth Polymers—Addition Polymers

Chain-growth polymerization is a chain reaction that converts an organic starting material, usually an alkene, to a polymer via a reactive intermediate—a radical, cation, or anion.




new bonds shown in red


- **The alkene can be ethylene ($CH_2=CH_2$) or a derivative of ethylene ($CH_2=CHZ$ or $CH_2=CZ_2$).**
- **The substituent Z (in part) determines whether radicals, cations, or anions are formed as intermediates.**
- **An initiator—a radical, cation, or anion—is needed to begin polymerization.**
- **Since chain-growth polymerization is a chain reaction, the mechanism involves initiation, propagation, and termination (Section 15.4).**


In most chain-growth polymerizations, an initiator adds to the carbon–carbon double bond of one monomer to form a reactive intermediate, which then reacts with another molecule of monomer to build the chain. Polymerization of $CH_2{=}CHZ$ results in a carbon chain having the Z substituents on every other carbon atom.

Problem 30.2 What polymer is formed by chain-growth polymerization of each monomer?

a. 

b. 

c. 


d. 

30.2A Radical Polymerization

Radical polymerization of alkenes was first discussed in Section 15.14, and is included here to emphasize its relationship to other methods of chain-growth polymerization. The initiator is often a peroxy radical (RO·), formed by cleavage of the weak O−O bond in an organic peroxide, ROOR. Mechanism 30.1 is written with styrene ($CH_2{=}CHPh$) as the starting material.


Mechanism 30.1 Radical Polymerization of $CH_2{=}CHPh$

Part [1] Initiation



1–2 Initiation with ROOR occurs in two steps—homolysis of the weak O–O bond and addition of RO· to the alkene to form a carbon radical.


Part [2] Propagation



new C–C bond in red


3 Chain propagation consists of a single step. The carbon radical adds to another alkene to form a new C–C bond and another carbon radical. Addition forms the radical with the unpaired electron on the atom with the Z substituent. Step [3] occurs repeatedly to grow the polymer chain.

Part [3] Termination




4 Termination of the chain occurs when any two radicals combine to form a bond.

Radical polymerization of $CH_2=CHZ$ is favored by Z substituents that stabilize a radical by electron delocalization. Each addition step occurs to put the intermediate radical on the carbon bearing the Z substituent. With styrene as the starting material, the intermediate radical is benzylic and highly resonance stabilized. Figure 30.3 shows several monomers used in radical polymerization reactions.




five resonance structures for the benzylic radical

Figure 30.3
Monomers used in radical polymerization reactions




ethylene | vinyl chloride | styrene | vinyl acetate

Problem 30.3 What polymer is formed by the radical polymerization of each monomer?



Problem 30.4 Draw the mechanism for the radical polymerization of vinyl acetate ($CH_2=CHOCOCH_3$) using $(CH_3)_3CO-OC(CH_3)_3$ as the initiator.

Chain termination can occur by radical coupling, as shown in Mechanism 30.1. Chain termination can also occur by **disproportionation,** a process in which a hydrogen atom is transferred from one polymer radical to another, forming a new C−H bond on one polymer chain, and a double bond on the other.




new C–H bond and π bond in red

30.2B Chain Branching


HDPE is used in milk containers and water jugs, whereas LDPE is used in plastic bags and insulation.

The choice of reaction parameters greatly affects the properties of a synthetic polymer. In Section 15.14, we learned that there are two common types of polyethylene—**high-density polyethylene (HDPE) and low-density polyethylene (LDPE).** High-density polyethylene, which consists of long chains of CH_2 groups joined together in a linear fashion, is strong and hard because the linear chains pack well, resulting in strong van der Waals interactions. Low-density polyethylene, on the other hand, consists of long carbon chains with many branches along the

chain. Branching prohibits the chains from packing well, so LDPE has weaker intermolecular interactions, making it a much softer, pliable material.




linear polyethylene branched polyethylene



Linear polyethylene molecules pack well. Branched polyethylene molecules do *not* pack well.


Branching occurs when a radical on one growing polyethylene chain abstracts a hydrogen atom from a CH_2 group in another polymer chain, as shown in Mechanism 30.2. The new 2° radical then continues chain propagation by adding to another molecule of ethylene, thus forming a branch point.

Mechanism 30.2 Forming Branched Polyethylene During Radical Polymerization



1. Abstraction of a H atom from an existing polymer chain forms a 2° radical in the middle of the polymer chain.
2. Addition of the radical to another molecule of ethylene forms a new radical and a branch point along the polymer chain. Step [2] occurs repeatedly, and a long branch grows off the original polymer chain.

Problem 30.5 Explain why radical polymerization of styrene forms branched chains with 4° carbons as in **A,** but none with 3° carbons as in **B.**



A **B**


30.2C Ionic Polymerization

Chain-growth polymerization can also occur by way of cationic or anionic intermediates. **Cationic polymerization is an example of electrophilic addition to an alkene involving carbocations.** Cationic polymerization occurs with alkene monomers that have substituents capable of stabilizing intermediate carbocations, such as alkyl groups or other electron-donor groups. The initiator is an electrophile such as a proton source or Lewis acid.

Mechanism 30.3 illustrates cationic polymerization of the general monomer $CH_2{=}CHZ$ using $BF_3{\cdot}H_2O$, the Lewis acid–base complex formed from BF_3 and H_2O, as the initiator.


Mechanism 30.3 Cationic Polymerization of $CH_2{=}CHZ$

Part [1] Initiation



1 – 2 Electrophilic addition of H^+ from $BF_3{\cdot}H_2O$ forms a carbocation.


Part [2] Propagation



new C–C bond in red

3 The carbocation adds to another alkene to form a new C–C bond. Addition forms a carbocation stabilized by an electron-donor Z group. Step [3] occurs repeatedly to grow the polymer chain.

Part [3] Termination




4 Loss of a proton forms a new π bond and terminates the chain.


Since cationic polymerization involves carbocations, addition follows Markovnikov's rule to form the more stable, more substituted carbocation. Chain termination can occur by a variety of pathways, such as loss of a proton to form an alkene. Examples of alkene monomers that undergo cationic polymerization are shown in Figure 30.4.

Figure 30.4
Common polymers formed by ionic chain-growth polymerization


Monomer	⟶	Polymer	Consumer product


- Polymers formed by cationic polymerization





- Polymers formed by anionic polymerization





- A chain-growth polymer is named by adding the prefix *poly* to the name of the monomer from which it is made. When the name of the monomer contains two words, this name is enclosed in parentheses and preceded by the prefix *poly.*


Problem 30.6 Explain why cationic polymerization is an effective method of polymerizing $CH_2=C(CH_3)_2$ but not $CH_2=CH_2$.

Although alkenes readily react with electron-deficient radicals and electrophiles, alkenes do not generally react with anions and other nucleophiles. Consequently, **anionic polymerization takes place only with alkene monomers that contain electron-withdrawing groups** such as COR, COOR, or CN, which can stabilize an intermediate negative charge. The initiator is a strong

nucleophile, such as an organolithium reagent, RLi. Mechanism 30.4 illustrates anionic polymerization of the general monomer $CH_2{=}CHZ$.


Mechanism 30.4 Anionic Polymerization of $CH_2{=}CHZ$

Part [1] Initiation



1 Nucleophilic addition of RLi forms a carbanion stabilized by an electron-withdrawing group Z.


Part [2] Propagation



new C–C bond in red

2 The carbanion adds to another alkene to form a new C–C bond. Addition forms a new carbanion with the negative charge adjacent to the Z substituent. Step [2] occurs repeatedly to grow the polymer chain.


Part [3] Termination



3 An acid–base reaction with H_2O or another electrophile terminates the chain.

In contrast to other types of chain-growth polymerization, there are no efficient methods of terminating the chain mechanism in anionic polymerization. The reaction continues until all the initiator and monomer have been consumed, so that the end of each polymer chain contains a carbanion (Step [2] in Mechanism 30.4). Anionic polymerization is often called **living polymerization** because polymerization will begin again if more monomer is added at this stage. To terminate anionic polymerization an electrophile such as H_2O or CO_2 must be added. Examples of alkene monomers that undergo anionic polymerization are shown in Figure 30.4.

Problem 30.7 Which method of ionic polymerization—cationic or anionic—is preferred for each monomer? Explain your choices.




Problem 30.8 Draw a stepwise mechanism for the conversion of acrylonitrile ($CH_2{=}CHC{\equiv}N$) to polyacrylonitrile, $-[CH_2CHC{\equiv}N]_n-$, using butyllithium (BuLi) as the initiator and CO_2 as the electrophile to terminate the chain.

Problem 30.9 Explain why styrene ($CH_2{=}CHPh$) can be polymerized to polystyrene by all three methods of chain-growth polymerization.

30.2D Copolymers


All polymers discussed thus far are **homopolymers,** because they have been prepared by the polymerization of a single monomer. **Copolymers, on the other hand, are polymers prepared by joining two or more monomers (X and Y) together.**



- An *alternating copolymer* is formed when X and Y alternate regularly along the chain.
- A *random copolymer* is formed when X and Y are randomly distributed along the chain.


The structure of the copolymer depends on the relative amount and reactivity of **X** and **Y,** as well as the conditions used for polymerization.

Several copolymers are commercially important and used in a wide range of consumer products. The copolymer of vinyl chloride and vinylidene chloride forms **Saran,** the film used in the well-known plastic food wrap. Copolymerization of buta-1,3-diene and styrene forms **styrene–butadiene rubber (SBR),** the polymer used almost exclusively in automobile tires.



Problem 30.10 Draw the alternating copolymer formed from each set of monomers.

a.  and 

b.  and 

Problem 30.11 ABS, a widely produced copolymer used in crash helmets, small appliances, and toys, is formed from three monomers—acrylonitrile (CH_2=CHCN), buta-1,3-diene (CH_2=CH–CH=CH_2), and styrene (CH_2=CHPh). Draw a possible structure for ABS.




Lego bricks are made from the copolymer ABS (Problem 30.11).

30.3 Anionic Polymerization of Epoxides


Alkene monomers are the most common starting materials in chain-growth polymerizations, but epoxides can also serve as starting materials, forming **polyethers.** The strained three-membered ring of an epoxide is readily opened with a nucleophile (such as $^-$OH or $^-$OR) to form an alkoxide, which can then ring open another epoxide monomer to build the polymer chain. Unlike the other methods of chain-growth polymerization that join monomers together with C−C bonds, this process forms **new C−O bonds** in the polymer backbone.

For example, the ring opening of ethylene oxide with a $^-$OH initiator affords an alkoxide nucleophile, which propagates the chain by reacting with more ethylene oxide. This process yields **poly(ethylene glycol), PEG,** a polymer used in lotions and creams. The many C−O bonds in these polymers make them highly water soluble.




The ring opening of epoxides with nucleophiles was first discussed in Section 9.15.

Under anionic conditions, the ring opening follows an S_N2 mechanism. Thus, the ring opening of an unsymmetrical epoxide occurs at the more accessible, less substituted carbon, labeled in blue.




Problem 30.12 What polymer is formed by anionic polymerization of each monomer?



30.4 Ziegler–Natta Catalysts and Polymer Stereochemistry

Polymers prepared from monosubstituted alkene monomers (CH_2=CHZ) can exist in three different configurations, called **isotactic, syndiotactic,** and **atactic:**



- An *isotactic* polymer has all Z groups on the same side of the carbon backbone.
- A *syndiotactic* polymer has the Z groups alternating from one side of the carbon chain to the other.
- An *atactic* polymer has the Z groups oriented randomly along the polymer chain.

The more regular arrangement of the Z substituents in isotactic and syndiotactic polymers allows them to pack together better, making the polymer stronger and more rigid. In contrast, the chains of an atactic polymer tend to pack less closely together, resulting in a lower melting, softer polymer. Radical polymerization often affords an atactic polymer, but the particular reaction conditions can greatly affect the stereochemistry of the polymer formed.


In 1953, Karl Ziegler and Giulio Natta developed a new method of polymerizing alkene monomers using a metal catalyst to promote chain-growth polymerization. These catalysts, now called **Ziegler–Natta catalysts,** offer two advantages over other methods of chain-growth polymerization.

- The stereochemistry of the polymer is easily controlled. Polymerization affords isotactic, syndiotactic, or atactic polymers depending on the catalyst.
- Long, linear chains of polymer are prepared without significant branching. Radicals are not formed as reactive intermediates, so intermolecular hydrogen abstraction, which leads to chain branching, does not occur.

Ziegler and Natta received the 1963 Nobel Prize in Chemistry for their pioneering work on polymerization catalysts.

Many different Ziegler–Natta catalysts are used for polymerization, but most consist of an organoaluminum compound such as $(CH_3CH_2)_2AlCl$ and $TiCl_4$, a Lewis acid. The active catalyst is thought to be an alkyl titanium compound, formed by transfer of an ethyl group from $(CH_3CH_2)_2AlCl$ to $TiCl_4$, although many mechanistic details are not known with certainty. It is generally agreed that the alkene monomer coordinates to an alkyl titanium complex, and then inserts into the Ti−C bond to form a new carbon–carbon bond, as shown in Mechanism 30.5.

Mechanism 30.5 Ziegler–Natta Polymerization of $CH_2=CH_2$



1. Reaction of the organoaluminum compound with $TiCl_4$ forms the Ziegler–Natta catalyst with a Ti–C bond.
2. An alkene monomer coordinates with the Ti complex.
3. Insertion of $CH_2=CH_2$ into the Ti–C bond forms a new C–C bond. Repeating Steps [2] and [3] over and over yields the long polymer chain.



Dyneema, the strongest fabric known, is made of ultra high-density polyethylene, and is used for ropes, nets, bulletproof vests, and crash helmets.

The Ziegler–Natta polymerization of ethylene forms **high-density polyethylene, HDPE,** composed of long linear carbon chains that pack closely together, forming a rigid polymer. By using specialized manufacturing techniques that force the polymer chains to pack closely in the solid phase as a set of linear extended chains, this material is converted to ultra high-density polyethylene, a synthetic organic material stronger than steel.


Recently developed Ziegler–Natta polymerizations utilize zirconium complexes that are soluble in the reaction solvents typically used, and so they are **homogeneous catalysts.** Reactions that use these soluble catalysts are called **coordination polymerizations.**

30.5 Natural and Synthetic Rubbers


Natural rubber is composed of repeating five-carbon units, in which all the double bonds have the *Z* configuration. Because natural rubber is a hydrocarbon, it is water insoluble, and thus useful for waterproofing. The *Z* double bonds cause bends and kinks in the polymer chain, making it a soft material.



Natural rubber is obtained from latex that oozes from cuts made to the bark of the rubber tree. Waterproof latex is the rubber tree's natural protection, exuded in response to an injury. Although rubber was produced exclusively in Brazil until the late 1800s, today most of the world's rubber comes from plantations in Southeast Asia, Sri Lanka, and Indonesia.




The polymerization of isoprene under radical conditions forms a stereoisomer of natural rubber called **gutta-percha,** in which all the double bonds have the *E* configuration. Gutta-percha is also a naturally occurring polymer, although considerably less common than its *Z* stereoisomer. Polymerization of isoprene with a Ziegler–Natta catalyst forms natural rubber with all the double bonds having the desired *Z* configuration.



Natural rubber is too soft to be a useful material for most applications. Moreover, when natural rubber is stretched, the chains become elongated and slide past each other until the material pulls apart. In 1839, Charles Goodyear discovered that mixing hot rubber with sulfur produced a stronger and more elastic material. This process, called **vulcanization,** results in cross-linking of the hydrocarbon chains by disulfide bonds, as shown in Figure 30.5. When the polymer is stretched, the chains no longer can slide past each other and tearing does not occur. Vulcanized rubber is an ***elastomer,*** **a polymer that stretches when stressed but then returns to its original shape when the stress is alleviated.**

Figure 30.5
Vulcanized rubber




- Vulcanized rubber contains many disulfide bonds that cross-link the hydrocarbon chains together.

Gutta-percha, a much harder material than natural rubber obtained from latex, is used in golf ball casings.


The degree of cross-linking affects the rubber's properties. Harder rubber used for automobile tires has more cross-linking than the softer rubber used for rubber bands.

Other synthetic rubbers can be prepared by the polymerization of different 1,3-dienes using Ziegler–Natta catalysts. For example, the polymerization of buta-1,3-diene affords (*Z*)-poly(buta-1,3-diene), and the polymerization of 2-chlorobuta-1,3-diene yields neoprene, a polymer used in wet suits and tires.



Problem 30.13 Assign the *E* or *Z* configuration to the double bonds in neoprene. Draw a stereoisomer of neoprene in which all the double bonds have the opposite configuration.

Problem 30.14 The polymerization of $CH_2=CHCH=CH_2$ under radical conditions affords products **A** and **B.** Draw a mechanism that accounts for their formation.



30.6 Step-Growth Polymers—Condensation Polymers


Step-growth polymers, the second major class of polymers, are formed when monomers containing two functional groups come together and lose a small molecule such as H_2O or HCl. Commercially important step-growth polymers include:

- **Polyamides**
- **Polyesters**
- **Polyurethanes**
- **Polycarbonates**
- **Epoxy resins**

30.6A Polyamides


Nylon 6,6 is used in many products including parachutes and clothing.

Nylons are polyamides formed by step-growth polymerization. In Section 22.16A, we learned that **nylon 6,6** can be prepared by the reaction of a diacid chloride and a diamine. Nylon 6,6 can also be prepared by heating adipic acid and 1,6-diaminohexane. A Brønsted–Lowry acid–base reaction forms a diammonium salt, which loses H_2O at high temperature. In both methods, each starting material has two identical functional groups.



Nylon 6, trade name **Perlon,** is used to make rope and tire cord.


Nylon 6 is another polyamide, which is made by heating an aqueous solution of ε-caprolactam. The seven-membered ring of the lactam (a cyclic amide) is opened to form 6-aminohexanoic acid, the monomer that reacts with more lactam to form the polyamide chain. This step-growth polymerization thus begins with a single difunctional monomer that has two *different* functional groups, NH_2 and COOH.






Armadillo bicycle tires reinforced with Kevlar are hard to pierce with sharp objects, so a cyclist rarely gets a flat tire.

Kevlar is a polyamide formed from terephthalic acid and 1,4-diaminobenzene. The aromatic rings of the polymer backbone make the chains less flexible, resulting in a very strong material. Kevlar is light in weight compared to other materials that are similar in strength, so it is used in many products, such as bulletproof vests, army helmets, and the protective clothing used by firefighters.




Problem 30.15 What polyamide is formed from each monomer or pair of monomers?




30.6B Polyesters


Polyesters are formed by step-growth polymerization using nucleophilic acyl substitution reactions, as we learned in Section 22.16B. For example, the reaction of terephthalic acid and ethylene glycol forms **polyethylene terephthalate (PET),** the chapter-opening molecule.




Although PET is a very stable material, some polyesters are more readily hydrolyzed to carboxylic acids and alcohols in aqueous medium, making them suited for applications in which slow degradation is useful. For example, copolymerization of glycolic acid and lactic acid forms a copolymer used by surgeons in dissolving sutures. Within weeks, the copolymer is hydrolyzed to the monomers from which it was prepared, which are metabolized readily by the body. These sutures are used internally to hold tissues together while healing and scar formation occur.



Problem 30.16 Draw the structure of PEF, polyethylene furanoate, a condensation polymer formed from furandicarboxylic acid and ethylene glycol. PEF, which can be synthesized from precursors that are obtained from renewable resources, has many of the same properties as polyethylene terephthalate (PET).




furandicarboxylic acid



ethylene glycol


Problem 30.17 Polyethylene terephthalate is also prepared by the transesterification of dimethyl terephthalate with ethylene glycol. Draw the mechanism for this nucleophilic acyl substitution.




dimethyl terephthalate + ethylene glycol → polyethylene terephthalate (PET)

30.6C Polyurethanes

A **urethane** (also called a **carbamate**) is a compound that contains a carbonyl group bonded to both an OR group and an NHR (or NR_2) group (Section 29.7). Urethanes are prepared by the nucleophilic addition of an alcohol to the carbonyl group of an **isocyanate, RN=C=O.**




Polyurethanes are polymers formed by the reaction of a diisocyanate and a diol.



toluene 2,6-diisocyanate + ethylene glycol → a polyurethane

Spandex is a generic term for a strong and flexible polyurethane polymer that illustrates how the macroscopic properties of a polymer depend on its structure at the molecular level. Spandex was first used in women's corsets, girdles, and support hose, but is now routinely used in both men's and women's active wear. Spandex is strong and lends "support" to the wearer, but it also stretches. Spandex is lighter in weight than many other elastic polymers, and it does not break down when exposed to perspiration and detergents. On the molecular level, it has rigid regions that are joined together by soft, flexible segments. The flexible regions allow the polymer to expand and then recover its original shape. The rigid regions strengthen the polymer.




spandex
Trade name **Lycra**


30.6D Polycarbonates

Although it is not acutely toxic, bisphenol A (BPA) mimics the body's own hormones and disrupts normal endocrine functions. Concern over low-dose exposure by infants has led to a voluntary phase-out of BPA-based polymers in infant formula packaging.


A **carbonate** is a compound that contains a carbonyl group bonded to two OR groups. Carbonates can be prepared by the reaction of phosgene ($Cl_2C{=}O$) with two equivalents of an alcohol (ROH).



Polycarbonates are formed from phosgene and a diol. The most widely used polycarbonate is **Lexan,** a lightweight, transparent material that is formed from phosgene and bisphenol A, and used in bike helmets, goggles, catcher's masks, and bulletproof glass.




Problem 30.18 Lexan can also be prepared by the acid-catalyzed reaction of diphenyl carbonate with bisphenol A. Draw a stepwise mechanism for this process.




30.6E Epoxy Resins

Epoxy resins represent a class of step-growth polymer familiar to anyone who has used "epoxy" to glue together a broken object. An epoxy resin consists of two components: a fluid **prepolymer** composed of short polymer chains with reactive epoxides on each end, and a **hardener,** usually a diamine or triamine that ring opens the epoxides and cross-links the chains together. The prepolymer is formed by reacting two difunctional monomers, bisphenol A and epichlorohydrin.



Bisphenol A has two nucleophilic OH groups, while epichlorohydrin has polar C−O and C−Cl bonds that can react with two different nucleophiles. The general reaction of epichlorohydrin with nucleophiles is given in the accompanying equation. Nucleophilic attack on the strained epoxide ring affords an alkoxide that displaces chloride by an intramolecular S_N2 reaction, forming a new epoxide. Ring opening with a second nucleophile gives a 2° alcohol.







Figure 30.6 Formation of an epoxy resin from a prepolymer and a hardening agent

The polymer chains are cross-linked together.

When bisphenol A is treated with excess epichlorohydrin, this stepwise process continues until all the phenolic OH groups have been used in ring-opening reactions, leaving epoxy groups on both ends of the polymer chains. This constitutes the fluid **prepolymer,** as shown in Figure 30.6.

When the prepolymer is mixed with a diamine or triamine (the **hardener**), the reactive epoxide rings can be opened by the nucleophilic amino groups to cross-link polymer chains together, causing the polymer to harden. A wide range of epoxy resins is commercially prepared by this process, making them useful for adhesives and coatings. The longer and more extensively cross-linked the polymer chains, the harder the resin.


Problem 30.19 (a) Draw the structure of the prepolymer **A** formed from 1,4-dihydroxybenzene and excess epichlorohydrin. (b) Draw the structure of the cross-linked polymer **B** formed when **A** is treated with $H_2NCH_2CH_2CH_2NH_2$ as the hardening agent.



30.7 Polymer Structure and Properties

While the chemistry of polymer synthesis can be explained by the usual themes of organic reactions, the large size of polymer molecules gives them some unique physical properties compared to small organic molecules.

Linear and branched polymers do not form crystalline solids because their long chains prevent efficient packing in a crystal lattice. Most polymer chains have **crystalline regions** and **amorphous regions:**



- **Ordered crystalline regions,** called **crystallites,** are places where sections of the polymer chain lie in close proximity and are held together by intermolecular interactions. Ordered regions of polyethylene, $-[CH_2CH_2]_n-$, are held together by van der Waals interactions, whereas ordered regions of nylon chains are held together by intermolecular hydrogen bonding.
- **Amorphous regions** are places where the polymer chains are randomly arranged, resulting in weak intermolecular interactions.

Crystalline regions impart toughness to a polymer, while amorphous regions impart flexibility. The greater the crystallinity of a polymer—that is, the larger the percentage of ordered regions—the harder the polymer. Branched polymers are generally more amorphous and, since branching prevents chains from packing closely, they are softer, too.


Two temperatures, T_g and T_m, often characterize a polymer's behavior on heating:

- **T_g, the glass transition temperature, is the temperature at which a hard amorphous polymer becomes soft.**
- **T_m, the melt transition temperature, is the temperature at which the crystalline regions of the polymer melt to become amorphous. More ordered polymers have higher T_m values.**

Thermoplastics are polymers that can be melted and then molded into shapes that are retained when the polymer is cooled. Although they have high T_g values and are hard at room temperature, heating causes individual polymer chains to slip past each other, causing the material to soften. Polyethylene terephthalate and polystyrene are thermoplastic polymers.

Thermosetting polymers are complex networks of cross-linked polymers. Thermosetting polymers are formed by chemical reactions that occur when monomers are heated together to form a network of covalent bonds. Thermosetting polymers cannot be re-melted to form a liquid phase, because covalent bonds hold the network together. **Bakelite,** a thermosetting polymer prepared from phenol (PhOH) and formaldehyde ($H_2C{=}O$) in the presence of a Lewis acid, is formed by electrophilic aromatic substitution reactions. Since formaldehyde is a reactive electrophile and

Figure 30.7
The synthesis of Bakelite from phenol and formaldehyde






- Bakelite, the first totally synthetic polymeric material, was patented by Leo Baekeland in 1910. Bowling balls are made of Bakelite.

phenol contains a strongly electron-donating OH group, substitution occurs at all ortho and para positions to the OH group, resulting in a highly cross-linked polymer, shown in Figure 30.7.

Problem 30.20 Draw a stepwise mechanism for Step [2] in Figure 30.7 using $AlCl_3$ as the Lewis acid catalyst.

Sometimes a polymer is too stiff and brittle to be useful in many applications. In this case, a low molecular weight compound called a **plasticizer** is added to soften the polymer and give it flexibility. The plasticizer interacts with the polymer chains, replacing some of the intermolecular interactions between the polymer chains. This lowers the crystallinity of the polymer, making it more amorphous and softer.

Dibutyl phthalate is a plasticizer added to the poly(vinyl chloride) used in vinyl upholstery and garden hoses. Since plasticizers are more volatile than the high molecular weight polymers, they slowly evaporate with time, making the polymer brittle and easily cracked. Plasticizers like dibutyl phthalate that contain hydrolyzable functional groups are also slowly degraded by chemical reactions.



dibutyl phthalate

30.8 Green Polymer Synthesis

One hundred fifty years ago there were no chemical manufacturing plants and no synthetic polymers, and petroleum had little value. Synthetic polymers have transformed the daily lives of many in the modern world, but not without a hefty price. Polymer synthesis and disposal have a tremendous impact on the environment, creating two central issues:

- **Where do polymers come from?** What raw materials are used for polymer synthesis and what environmental consequences result from their manufacture?
- **What happens to polymers once they are used?** How does polymer disposal affect the environment, and what can be done to minimize its negative impact?

30.8A Environmentally Friendly Polymer Synthesis—The Feedstock

In Chapter 12, you were introduced to **green chemistry,** the use of environmentally benign methods to synthesize compounds. Given the billions of pounds of polymers manufactured worldwide each year, there is an obvious need for methods that minimize the environmental impact.

Recall from Section 4.7 that 3% of a barrel of crude oil is used as the feedstock for chemical synthesis.


To date, green polymer synthesis has been approached in a variety of ways:

- **Using starting materials that are derived from renewable sources, rather than petroleum. The starting materials for an industrial process are often called the chemical *feedstock*.**
- **Using safer, less toxic reagents that form fewer by-products.**
- **Carrying out reactions in the absence of solvent or in aqueous solution (instead of an organic solvent).**

Until recently, **the feedstock for all polymer synthesis has been petroleum;** that is, the monomers for virtually all polymer syntheses are made from crude oil, a nonrenewable raw material. As an example, nylon 6,6 is prepared industrially from adipic acid [$HOOC(CH_2)_4COOH$] and 1,6-diaminohexane [$H_2N(CH_2)_6NH_2$], both of which originate from benzene, a product of petroleum refining (Figure 30.8).

Figure 30.8

Synthesis of adipic acid and 1,6-diaminohexane for nylon 6,6 synthesis




- The synthesis of both monomers needed for nylon 6,6 synthesis begins with benzene, a petroleum product.

Besides beginning with a nonrenewable chemical feedstock, adipic acid synthesis has other problems. The use of benzene, a carcinogen and liver toxin, is undesirable, especially in a large-scale reaction. Moreover, oxidation with HNO_3 in Step [3] produces N_2O as a by-product. N_2O depletes ozone in the stratosphere in much the same way as the CFCs discussed in Chapter 15. In addition, N_2O also absorbs thermal energy from the earth's surface like CO_2, and may therefore contribute to global climate change, as discussed in Section 4.14.

As a result, several research groups are working to develop new methods of monomer synthesis that begin with renewable, more environmentally friendly raw materials and produce fewer hazardous by-products. As an example, chemists at Michigan State University have devised a two-step synthesis of adipic acid from D-glucose, a monosaccharide available from plant sources. The synthesis uses a genetically altered *E. coli* strain (called a **biocatalyst**) to convert D-glucose to (2*Z*,4*Z*)-hexa-2,4-dienoic acid, which is then hydrogenated to adipic acid. Methods such as this,

which avoid starting materials derived from petroleum, are receiving a great deal of attention in the chemical community.



Sorona, DuPont's trade name for **poly(trimethylene terephthalate),** is a large-volume polymer that can now be made at least in part from glucose derived from a renewable plant source such as corn. A biocatalyst converts D-glucose to propane-1,3-diol, which forms poly(trimethylene terephthalate) (PTT) on reaction with terephthalic acid, as shown in Figure 30.9.

In related chemistry, poly(lactic acid) (PLA) is a polymer used in bottles and packaging, and it can also be made into a synthetic fiber (trade name Ingeo) used in clothing and carpets. Poly(lactic acid) is prepared on a large scale by the fermentation of carbohydrates obtained from corn. Fermentation initially yields a cyclic lactone called lactide, derived from two molecules of lactic acid [$CH_3CH(OH)CO_2H$]. Heating lactide with acid forms poly(lactic acid). PLA is an especially attractive polymer choice, because it readily degrades in a landfill.






Figure 30.9 A swimsuit made (in part) from corn—The synthesis of poly(trimethylene terephthalate) from propane-1,3-diol derived from corn




- Poly(trimethylene terephthalate), sold as Sorona by the DuPont Corporation, is made into fibers used in clothing and other materials. Although propane-1,3-diol, one of the monomers needed for its synthesis, has been prepared from petroleum feedstocks in the past, it is now available from a renewable plant source such as corn.

30.8B Polymer Synthesis with Less Hazardous Reagents

Other approaches to green polymer synthesis have concentrated on using less hazardous reagents and avoiding solvents. For example, Lexan can now be prepared by the reaction of bisphenol A with diphenyl carbonate [$(PhO)_2C{=}O$] in the absence of solvent. This process avoids the use of phosgene ($Cl_2C{=}O$, Section 30.6D), an acutely toxic reagent that must be handled with extreme care, as well as the large volume of CH_2Cl_2 typically used as the solvent for the polymerization process.



Problem 30.21 Thermosetting resins similar to Bakelite (Section 30.7) have also been prepared from renewable feedstocks. One method uses cardinol, the major constituent of the liquid obtained from roasted cashew nutshells. What polymer is obtained when cardinol is treated with formaldehyde ($H_2C{=}O$) in the presence of a proton source?



30.9 Polymer Recycling and Disposal

The same desirable characteristics that make polymers popular materials for consumer products—durability, strength, and lack of reactivity—also contribute to environmental problems. Polymers do not degrade readily, and as a result, billions of pounds of polymers end up in landfills every year.







Two solutions to address the waste problem created by polymers are recycling existing polymer types to make new materials, and using biodegradable polymers that will decompose in a finite and limited time span.

30.9A Polymer Recycling

Although thousands of different synthetic polymers have now been prepared, six compounds, called the **"Big Six,"** account for 76% of the synthetic polymers produced in the United States each year. Each polymer is assigned a recycling code (1–6) that indicates its ease of recycling; **the lower the number, the easier to recycle.** Table 30.1 lists these six most common polymers, as well as the type of products made from each recycled polymer.


Recycling begins with sorting plastics by type, shredding the plastics into small chips, and washing the chips to remove adhesives and labels. After the chips are dried and any metal caps or rings are removed, the polymer chips are melted and molded for reuse.

Table 30.1 Recyclable Polymers


Recycling code	Polymer name	Structure	Recycled product
1	PET Polyethylene terephthalate		fleece jackets carpeting plastic bottles
2	HDPE High-density polyethylene		Tyvek insulation sports clothing
3	PVC Poly(vinyl chloride)		floor mats
4	LDPE Low-density polyethylene		trash bags
5	PP Polypropylene		furniture
6	PS Polystyrene		molded trays trash cans

Of the Big Six, only the polyethylene terephthalate (PET) in soft drink bottles and the high-density polyethylene (HDPE) in milk jugs and juice bottles are recycled to any great extent. Since recycled polymers are often still contaminated with small amounts of adhesives and other materials, these recycled polymers are generally not used for storing food or drink products. Recycled HDPE is converted to Tyvek, an insulating wrap used in new housing construction, and recycled PET is used to make fibers for fleece clothing and carpeting. Currently about 23% of all plastics are recycled in the United States.

An alternative recycling process is to re-convert polymers back to the monomers from which they were made, a process that has been successful with acyl compounds that contain C−O or C−N bonds in the polymer backbone. For example, heating polyethylene terephthalate with CH_3OH cleaves the esters of the polymer chain to give ethylene glycol ($HOCH_2CH_2OH$) and dimethyl terephthalate. These monomers then serve as starting materials for more PET. This chemical recycling process is a transesterification reaction that occurs by nucleophilic acyl substitution, as discussed in Chapter 22.



Similarly, treatment of discarded nylon 6 polymer with NH_3 cleaves the polyamide backbone, forming ε-caprolactam, which can be purified and re-converted to nylon 6.




Problem 30.22 Why can't chemical recycling—that is, the conversion of polymer to monomers and re-conversion of monomers to polymer—be done easily with HDPE and LDPE?

30.9B Biodegradable Polymers


Another solution to the accumulation of waste polymers in landfills is to design and use polymers that are biodegradable.

- **Biodegradable polymers are polymers that can be degraded by microorganisms—bacteria, fungi, or algae—naturally present in the environment.**

Several biodegradable polyesters have now been developed. For example, the **polyhydroxyalkanoates (PHAs)** are polymers of 3-hydroxy carboxylic acids, such as 3-hydroxybutyric acid or 3-hydroxyvaleric acid.




polyhydroxyalkanoate
PHA




3-hydroxy carboxylic acid
R = CH_3, 3-hydroxybutyric acid
R = CH_2CH_3, 3-hydroxyvaleric acid
monomer

The two most common PHAs are **polyhydroxybutyrate (PHB)** and a copolymer of **polyhydroxybutyrate** and **polyhydroxyvalerate (PHBV).** PHAs can be used as films, fibers, and coatings for hot beverage cups made of paper.



PHB




PHBV


Bacteria in the soil readily degrade PHAs, and in the presence of oxygen, the final degradation products are CO_2 and H_2O. The rate of degradation depends on moisture, temperature, and pH. Degradation is slower in enclosed landfills that are lined and covered.

An additional advantage of the polyhydroxyalkanoates is that the polymers can be produced by fermentation. Certain types of bacteria produce PHAs for energy storage when they are grown in glucose solution in the absence of specific nutrients. The polymer forms as discrete granules within the bacterial cell, and it is then removed by extraction to give a white powder that can be melted and modified into a variety of different products.

Biodegradable polyamides have also been prepared from amino acids. For example, aspartic acid can be converted to polyaspartate, abbreviated as **TPA** (thermal polyaspartate). TPA is commonly used as an alternative to poly(acrylic acid), which is used to line the pumps and boilers of wastewater treatment facilities.



Problem 30.23 What polymers are formed from each monomer?




KEY CONCEPTS

Synthetic Polymers

Chain-Growth Polymers—Addition Polymers

[1] Chain-growth polymers with alkene starting materials (30.2)


- General reaction:



- Mechanism—three possibilities, depending in part on the identity of Z:

Type	Identity of Z	Initiator	Comments
[1] radical polymerization	Z stabilizes a radical. Z = R, Ph, Cl, etc.	A source of radicals (ROOR)	Termination occurs by radical coupling or disproportionation. Chain branching occurs.
[2] cationic polymerization	Z stabilizes a carbocation. Z = R, Ph, OR, etc.	H – A or a Lewis acid ($BF_3 + H_2O$)	Termination occurs by loss of a proton.
[3] anionic polymerization	Z stabilizes a carbanion. Z = Ph, CO_2R, COR, CN, etc.	An organolithium reagent (R – Li)	Termination occurs only when an acid or other electrophile is added.


[2] Chain-growth polymers with epoxide starting materials (30.3)



- The mechanism is S_N2.
- Ring opening occurs at the less substituted carbon of the epoxide.


Examples of Step-Growth Polymers—Condensation Polymers (30.6)

Polyamides




nylon 6


Polyesters



polyethylene terephthalate




Kevlar




copolymer of
glycolic and lactic acids

Polyurethanes



a polyurethane

Polycarbonates



Lexan

[Key functional groups are indicated in red.]


Structure and Properties


- Polymers prepared from monomers having the general structure $CH_2=CHZ$ can be isotactic, syndiotactic, or atactic depending on the identity of Z and the method of preparation (30.4).
- Ziegler–Natta catalysts form polymers without significant branching. Polymers can be isotactic, syndiotactic, or atactic depending on the catalyst. Polymers prepared from 1,3-dienes have the *E* or *Z* configuration depending on the catalyst (30.4, 30.5).
- Most polymers contain ordered crystalline regions and less ordered amorphous regions (30.7). The greater the crystallinity, the harder the polymer.
- Elastomers are polymers that stretch and can return to their original shape (30.5).
- Thermoplastics are polymers that can be molded, shaped, and cooled such that the new form is preserved (30.7).
- Thermosetting polymers are composed of complex networks of covalent bonds, so they cannot be re-melted to form a liquid phase (30.7).

PROBLEMS


Problems Using Three-Dimensional Models


30.24 Draw the structure of the polymer formed by chain-growth polymerization of each monomer.

a. 


b. 



30.25 What monomer(s) are used to prepare each polymer or copolymer?

a. 

b. 

30.26 Draw the structure of the polymer formed by step-growth polymerization of each monomer or pair of monomers.


a. 


b.  + 


Polymer Structure and Properties

30.27 Draw the structure of the polymer formed by chain-growth polymerization of each monomer.

a. 


b. 


c. 

d. 

30.28 Draw the structure of the alternating copolymer formed from each pair of monomers.


a.  and 


b.  and 


c.  and 


d.  and 

30.29 What monomer(s) are used to prepare each polymer or copolymer?

a. 

c. 

b. 



d. 



30.30 Draw each polymer in Problem 30.29 using the shorthand representation shown in Figure 30.2.



30.31 Draw a short segment of each polymer: (a) isotactic poly(vinyl chloride); (b) syndiotactic polyacrylonitrile; (c) atactic polystyrene.


30.32 Draw the structure of the polymer that results from anionic polymerization of *p*-trichloromethylstyrene ($CCl_3C_6H_4CH{=}CH_2$) using ethylene oxide as the electrophile to terminate the chain.

30.33 Draw the structure of the polymer formed by step-growth polymerization of each monomer or pair of monomers.


a.  and 


c.  and 

b.  and 


d. 

30.34 Draw the structures of **Quiana** and **Nomex,** two commercially available step-growth polymers formed from the given monomers. Nomex is a strong polymer used in aircraft tires and microwave transformers. Quiana has been used to make wrinkle-resistant fabrics.

a. 

b. 


30.35 Draw the structure of the three monomers used to prepare polybutyrate adipate terephthalate (PBAT), a biodegradable copolymer sold under the trade name of Ecoflex. Because PBAT has properties similar to low-density polyethylene, it can be used in biodegradable food packaging and plastic bags.



PBAT

30.36 Kevlar (Section 30.6A) is a very stiff polymer because its backbone contains many aromatic rings and its polymer chains are extensively hydrogen bonded to each other. Draw a short segment of two Kevlar chains, and indicate how the chains are hydrogen bonded to each other.

30.37 Explain the differences observed in the T_g and T_m values for each pair of polymers: (a) polyester **A** and PET; (b) polyester **A** and nylon 6,6. (c) How would you expect the T_m value for Kevlar (Section 30.6A) to compare with the T_m value for nylon 6,6? Explain your prediction.




polyester **A**
T_g < 0 °C
T_m = 50 °C

PET
T_g = 70 °C
T_m = 265 °C

nylon 6,6
T_g = 53 °C
T_m = 265 °C

30.38 Explain why diester **A** is now often used as a plasticizer in place of dibutyl phthalate.




A

dibutyl phthalate

Mechanism


30.39 Draw a stepwise mechanism for the polymerization of isoprene to gutta-percha using $(CH_3)_3CO-OC(CH_3)_3$ as the initiator.



isoprene

gutta-percha


30.40 Cationic polymerization of 3-phenylpropene ($CH_2=CHCH_2Ph$) affords **A** as the major product rather than **B.** Draw a stepwise mechanism to account for this observation.




30.41 Explain why acrylonitrile ($CH_2=CHCN$) undergoes cationic polymerization more slowly than but-3-enenitrile ($CH_2=CHCH_2CN$).

30.42 Draw a stepwise mechanism for the anionic polymerization of styrene ($CH_2=CHPh$) to form polystyrene $-[CH_2CHPh]_n-$ using BuLi as the initiator. Use CO_2 as the electrophile that terminates the chain mechanism.


30.43 Although styrene undergoes both cationic and anionic polymerization equally well, one method is often preferred with substituted styrenes. Which method is preferred with each compound? Explain.




30.44 Rank the following compounds in order of increasing ability to undergo anionic chain-growth polymerization.




30.45 In the presence of H_3O^+, 2-methylpropene oxide undergoes chain-growth polymerization such that nucleophilic attack occurs at the more substituted end of the epoxide. Draw a stepwise mechanism for this process, and explain this regioselectivity.




30.46 Draw a stepwise mechanism for the conversion of dihalide **A** and cyclohexane-1,4-diol to polyether **B** in the presence of $AlCl_3$.



30.47 Draw a stepwise mechanism for the following reaction, which is used to prepare bisphenol A (BPA), a widely used monomer in polymer synthesis.




30.48 Draw a stepwise mechanism for the reaction of an alcohol with an isocyanate to form a urethane.



a urethane


Reactions and Synthesis

30.49 Draw the products of each reaction.




30.50 Explain why aqueous NaOH solution can be stored indefinitely in polyethylene bottles, but spilling aqueous base on a polyester shirt or nylon stockings quickly makes a hole.

30.51 What epoxy resin is formed by the following reaction sequence?




30.52 (a) Explain why poly(vinyl alcohol) cannot be prepared by the radical polymerization of vinyl alcohol ($CH_2=CHOH$). (b) Devise a stepwise synthesis of poly(vinyl alcohol) from vinyl acetate ($CH_2=CHOCOCH_3$). (c) How can poly(vinyl alcohol) be converted to poly(vinyl butyral), a polymer used in windshield safety glass?



poly(vinyl alcohol) poly(vinyl butyral)

30.53 Devise a synthesis of terephthalic acid and ethylene glycol, the two monomers needed for polyethylene terephthalate synthesis, from the given starting materials.



terephthalic acid ethylene glycol


30.54 The reaction of *p*-cresol with $CH_2=O$ resembles the reaction of phenol (PhOH) with $CH_2=O$, except that the resulting polymer is thermoplastic but not thermosetting. Draw the structure of the polymer formed, and explain why the properties of these two polymers are so different.



p-cresol


Biological Applications

30.55 In addition to glycolic and lactic acids (Section 30.6B), dissolving sutures can also be prepared from each of the following lactone monomers. Draw the structure of the polymer formed from each monomer.




ε-caprolactone *p*-dioxanone

30.56 Compound **A** is a novel poly(ester amide) copolymer that can be used as a bioabsorbable coating for the controlled release of drugs. **A** is a copolymer of four monomers, two of which are amino acids or amino acid derivatives. The body's enzymes recognize the naturally occurring amino acids in the polymer backbone, allowing for controlled enzymatic breakdown of the polymer and steady release of an encapsulated drug. Identify the four monomers used to synthesize **A;** then use Figure 29.2 to name the two amino acids.




poly(ester amide) **A**

30.57 Researchers at Rutgers University have developed biocompatible polymers that degrade into nonsteroidal anti-inflammatory drugs. For example, the reaction of two equivalents of benzyl salicylate and one equivalent of sebacoyl chloride forms a poly(anhydride ester) called PolyAspirin, which hydrolyzes to salicylic acid (an anti-inflammatory agent) and sebacic acid, which is excreted. This technology can perhaps be used for localized drug delivery at specific sites of injury. What is the structure of PolyAspirin?



Challenge Problems


30.58 Melmac, a thermosetting polymer formed from melamine and formaldehyde ($CH_2=O$), is used to make dishes and countertops. Draw a stepwise mechanism for the condensation of one mole of formaldehyde with two moles of melamine, which begins the synthesis of Melmac.



melamine

Melmac

30.59 Although chain branching in radical polymerizations can occur by intermolecular H abstraction as shown in Mechanism 30.2, chain branching can also occur by intramolecular H abstraction to form branched polyethylene that contains butyl groups as branches.



a. Draw a stepwise mechanism that illustrates which H must be intramolecularly abstracted to form butyl substituents.

b. Suggest a reason why the abstraction of this H is more facile than the abstraction of other H's.

30.60 The reaction of urea [$(NH_2)_2C=O$] and formaldehyde ($CH_2=O$) forms a highly cross-linked polymer used in foams. Suggest a structure for this polymer. [Hint: Examine the structures of Bakelite (Figure 30.7) and Melmac (Problem 30.58).]

Lipids

31



31.1 Introduction
31.2 Waxes
31.3 Triacylglycerols
31.4 Phospholipids
31.5 Fat-soluble vitamins
31.6 Eicosanoids
31.7 Terpenes
31.8 Steroids

Cholesterol is the most prominent member of the steroid family, a group of organic lipids that contains a tetracyclic structure. Cholesterol is synthesized in the liver and is found in almost all body tissues. It is a vital component for healthy cell membranes and serves as the starting material for the synthesis of all other steroids. But, as the general public now knows well, elevated cholesterol levels can lead to coronary artery disease. For this reason, consumer products are now labeled with their cholesterol content. In Chapter 31, we learn about the properties of cholesterol and other lipids. (Photo: © Jill Braaten)

We conclude the discussion of the organic molecules in biological systems by turning our attention to **lipids,** biomolecules that are soluble in organic solvents. Unlike the carbohydrates in Chapter 28 and the amino acids and proteins in Chapter 29, lipids contain many carbon–carbon and carbon–hydrogen bonds and few functional groups.

Since lipids are the biomolecules that most closely resemble the hydrocarbons we studied in Chapters 4 and 10, we have already learned many facts that directly explain their properties. Since there is no one functional group that is present in all lipids, however, the chemistry of lipids draws upon knowledge learned in many prior chapters.

31.1 Introduction

The word *lipid* comes from the Greek word ***lipos*** for "fat."

- ***Lipids*** **are biomolecules that are soluble in organic solvents.**


Lipids are unique among organic molecules because their identity is defined on the basis of a *physical property* and not by the presence of a particular functional group. Because of this, lipids come in a wide variety of structures and they have many different functions in the cell. Three examples are given in Figure 31.1.

The large number of **carbon–carbon and carbon–hydrogen σ bonds in lipids makes them very soluble in organic solvents and insoluble in water.** Monosaccharides (from which carbohydrates are formed) and amino acids (from which proteins are formed), on the other hand, are very polar, so they tend to be water soluble. Because lipids share many properties with hydrocarbons, several features of lipid structure and properties have already been discussed. Table 31.1 summarizes sections of the text where aspects of lipid chemistry were covered previously.


Table 31.1 Summary of Lipid Chemistry Discussed Prior to Chapter 31

Topic	Section	Topic	Section
• Vitamin A	3.5	• Lipid oxidation	15.11
• Soap	3.6	• Vitamin E	15.12
• Phospholipids, the cell membrane	3.7	• Steroid synthesis	16.14
• Lipids Part 1	4.15	• Prostaglandins	19.6
• Leukotrienes	9.17	• Lipid hydrolysis	22.12A
• Fats and oils	10.6	• Soap	22.12B
• Oral contraceptives	11.4	• Cholesteryl esters	22.17
• Hydrogenation of oils	12.4	• Steroid synthesis	24.8


Figure 31.1 Three examples of lipids



$PGF_{2\alpha}$
a prostaglandin



a triacylglycerol




progesterone
a steroid


- All lipids have many C – C and C – H bonds, but there is no one functional group common to all lipids.

Lipids can be categorized as hydrolyzable or nonhydrolyzable.

[1] ***Hydrolyzable lipids*** **can be cleaved into smaller molecules by hydrolysis with water.** Most hydrolyzable lipids contain an ester unit. We will examine three subgroups: waxes, triacylglycerols, and phospholipids.



[2] ***Nonhydrolyzable lipids*** **cannot be cleaved into smaller units by aqueous hydrolysis.** Nonhydrolyzable lipids tend to be more varied in structure. We will examine four different types: fat-soluble vitamins, eicosanoids, terpenes, and steroids.



31.2 Waxes




Water beads up on the surface of a leaf because of the leaf's waxy coating. (Photo: © Daniel C. Smith)



The seeds of the jojoba plant grown in the southwestern United States are rich in waxes used in cosmetics and personal care products. (Photo: USDA, ARS, National Genetic Resources Program)

Waxes are the simplest hydrolyzable lipids. **Waxes are esters (RCOOR') formed from a high molecular weight alcohol (R'OH) and a fatty acid (RCOOH).**

Because of their long hydrocarbon chains, **waxes are very hydrophobic.** They form a protective coating on the feathers of birds to make them water repellent, and on leaves to prevent water evaporation. **Lanolin,** a wax composed of a complex mixture of high molecular weight esters, coats the wool fibers of sheep. **Spermaceti wax,** isolated from the heads of sperm whales, is largely $CH_3(CH_2)_{14}COO(CH_2)_{15}CH_3$. The three-dimensional structure of this compound shows how small the ester group is compared to the long hydrocarbon chains.



spermaceti wax (from sperm whales)


Problem 31.1 One component of jojoba oil is a wax formed from eicosenoic acid [$CH_3(CH_2)_7CH=CH(CH_2)_9CO_2H$] and $CH_3(CH_2)_7CH=CH(CH_2)_8OH$. Draw the structure of the wax, including the cis geometry of both carbon–carbon double bonds.

31.3 Triacylglycerols

Triacylglycerols, or triglycerides, are the most abundant lipids, and for this reason we have already discussed many of their properties in earlier sections of this text.

- **Triacylglycerols are triesters that produce glycerol and three molecules of fatty acid upon hydrolysis.**

Line structures of stearic, oleic, linoleic, and linolenic acids can be found in Table 10.2. Ball-and-stick models of these fatty acids are shown in Figure 10.6.



triacylglycerol
the most common type of lipid

Simple triacylglycerols are composed of three identical fatty acid side chains, whereas **mixed triacylglycerols** have two or three different fatty acids. Table 31.2 lists the most common fatty acids used to form triacylglycerols.

Table 31.2 The Most Common Fatty Acids in Triacylglycerols

Number of C atoms	Number of C=C bonds	Structure	Name	Mp (°C)
		Saturated fatty acids		
12	0	$CH_3(CH_2)_{10}CO_2H$	lauric acid	44
14	0	$CH_3(CH_2)_{12}CO_2H$	myristic acid	58
16	0	$CH_3(CH_2)_{14}CO_2H$	palmitic acid	63
18	0	$CH_3(CH_2)_{16}CO_2H$	stearic acid	69
20	0	$CH_3(CH_2)_{18}CO_2H$	arachidic acid	77
		Unsaturated fatty acids		
16	1	$CH_3(CH_2)_5CH=CH(CH_2)_7CO_2H$	palmitoleic acid	1
18	1	$CH_3(CH_2)_7CH=CH(CH_2)_7CO_2H$	oleic acid	4
18	2	$CH_3(CH_2)_4(CH=CHCH_2)_2(CH_2)_6CO_2H$	linoleic acid	–5
18	3	$CH_3CH_2(CH=CHCH_2)_3(CH_2)_6CO_2H$	linolenic acid	–11
20	4	$CH_3(CH_2)_4(CH=CHCH_2)_4(CH_2)_2CO_2H$	arachidonic acid	–49

What are the characteristics of these fatty acids?

The most common saturated fatty acids are palmitic and stearic acids. The most common unsaturated fatty acid is oleic acid.

- **All fatty acid chains are unbranched, but they may be saturated or unsaturated.**
- **Naturally occurring fatty acids have an even number of carbon atoms.**
- **Double bonds in naturally occurring fatty acids generally have the *Z* configuration.**
- **The melting point of a fatty acid depends on the degree of unsaturation.**


Linoleic and linolenic acids are called **essential fatty acids** because we cannot synthesize them and must acquire them in our diets.

Fats and oils are triacylglycerols; that is, they are triesters of glycerol and these fatty acids.

- **Fats have higher melting points, making them solids at room temperature.**
- **Oils have lower melting points, making them liquids at room temperature.**


Figure 31.2 Three-dimensional structures of a saturated and unsaturated triacylglycerol

a. A saturated triacylglycerol



- Three saturated side chains lie parallel to each other, making a compact lipid.

b. An unsaturated triacylglycerol



- One *Z* double bond in a fatty acid side chain produces a twist so the lipid is no longer so compact.



Unlike other vegetable oils, oils from palm and coconut trees are very high in saturated fats. Considerable evidence currently suggests that diets high in saturated fats lead to a greater risk of heart disease. For this reason, the demand for coconut and palm oils has decreased considerably in recent years, and many coconut plantations previously farmed in the South Pacific are no longer in commercial operation. (Photos: top, © Corbis RF; bottom, © McGraw-Hill Education/ Elite Images)

This melting point difference correlates with the number of degrees of unsaturation present in the fatty acid side chains. **As the number of double bonds *increases,* the melting point *decreases,* as it does for the constituent fatty acids as well.**

Three-dimensional structures of a saturated and unsaturated triacylglycerol are shown in Figure 31.2. With no double bonds, the three side chains of the saturated lipid lie parallel to each other, making it possible for this compound to pack relatively efficiently in a crystalline lattice, thus leading to a high melting point. In the unsaturated lipid, however, a single *Z* double bond places a kink in the side chain, making it more difficult to pack efficiently in the solid state, thus leading to a lower melting point.

Solid fats have a relatively high percentage of saturated fatty acids and are generally of animal origin. Liquid oils have a higher percentage of unsaturated fatty acids and are generally of vegetable origin. Table 31.3 lists the fatty acid composition of some common fats and oils.

Table 31.3 Fatty Acid Composition of Some Fats and Oils

Source	% Saturated fatty acids	% Oleic acid	% Linoleic acid
beef	49–62	37–43	2–3
milk	37	33	3
coconut	86	7	—
corn	11–16	19–49	34–62
olive	11	84	4
palm	43	40	8
safflower	9	13	78
soybean	15	20	52

Data from *Merck Index,* 10th ed. Rahway, NJ: Merck and Co.; and Wilson, et al., 1967, *Principles of Nutrition,* 2nd ed. New York: Wiley.




Fish oils, such as cod liver and herring oils, are very rich in polyunsaturated triacylglycerols. These triacylglycerols pack so poorly that they have very low melting points; thus, they remain liquids even in the cold water inhabited by these fish. (Photo: © Rick Price/Corbis)




The hydrolysis, hydrogenation, and oxidation of triacylglycerols—reactions originally discussed in Chapters 12, 15, and 22—are summarized here for your reference.

[1] Hydrolysis of triacylglycerols (Section 22.12A)




Hydrolysis of a triacylglycerol with water in the presence of either acid, base, or an enzyme yields glycerol and three fatty acids. This cleavage reaction follows the same mechanism as any other ester hydrolysis (Section 22.11). This reaction is the first step in triacylglycerol metabolism.

[2] Hydrogenation of unsaturated fatty acids (Section 12.4)



The double bonds of an unsaturated fatty acid can be hydrogenated by using H_2 in the presence of a transition metal catalyst. Hydrogenation converts a liquid oil to a solid fat. This process, sometimes called **hardening,** is used to prepare margarine from vegetable oils.


[3] Oxidation of unsaturated fatty acids (Section 15.11)



Allylic C−H bonds are weaker than other C−H bonds and are thus susceptible to oxidation with molecular oxygen by a radical process. The hydroperoxide formed by this process is unstable, and it undergoes further oxidation to products that often have a disagreeable odor. This oxidation process turns an oil rancid.

In the cell, the principal function of triacylglycerols is energy storage. Complete metabolism of a triacylglycerol yields CO_2 and H_2O, and a great deal of energy. This overall reaction is reminiscent of the combustion of alkanes in fossil fuels, a process that also yields CO_2 and H_2O and provides energy to heat homes and power automobiles (Section 4.14B). Fundamentally both processes convert C–C and C–H bonds to C–O bonds, a highly exothermic reaction.

The average body fat content of men and women is ~20% and ~25%, respectively. (For elite athletes, however, the averages are more like <10% for men and <15% for women.) This stored fat can fill the body's energy needs for two or three months.






(Photo: © AP/Wide World Photos)

Carbohydrates provide an energy boost, but only for the short term, such as during strenuous exercise. Our long-term energy needs are met by triacylglycerols, because they store ~38 kJ/g, whereas carbohydrates and proteins store only ~16 kJ/g.

Because triacylglycerols release heat on combustion, they can in principle be used as fuels for vehicles. In fact, coconut oil was used as a fuel during both World War I and World War II, when gasoline and diesel supplies ran short. Since coconut oil is more viscous than petroleum products and freezes at 24 °C, engines must be modified to use it and it can't be used in cold climates. Nonetheless, a limited number of trucks and boats can now use vegetable oils, sometimes blended with diesel, as a fuel source. When the price of crude oil is high, the use of these **biofuels** becomes economically attractive.

Problem 31.2 How would you expect the melting point of eicosapentaenoic acid [$CH_3CH_2(CH{=}CHCH_2)_5(CH_2)_2COOH$] to compare with the melting points of the fatty acids listed in Table 31.2?

Problem 31.3 Draw the products formed when triacylglycerol **A** is treated with each reagent. Rank compounds **A, B,** and **C** in order of increasing melting point.



a. H_2O, H^+
b. H_2 (excess), Pd-C → **B**
c. H_2 (1 equiv), Pd-C → **C**



The cocoa butter in chocolate is rich in triacylglycerols derived from at least one molecule of oleic acid. (Photo: © PhotoDisc/Getty RF)


Problem 31.4 The main fatty acid component of the triacylglycerols in coconut oil is lauric acid, $CH_3(CH_2)_{10}COOH$. Explain why coconut oil is a liquid at room temperature even though it contains a large fraction of this saturated fatty acid.

Problem 31.5 Unlike many fats and oils, the cocoa butter used to make chocolate is remarkably uniform in composition. All triacylglycerols contain oleic acid esterified to the 2° OH group of glycerol, and either palmitic acid or stearic acid esterified to the 1° OH groups. Draw the structures of two possible triacylglycerols that compose cocoa butter.

31.4 Phospholipids

Phospholipids are hydrolyzable lipids that contain a phosphorus atom. There are two common types of phospholipids: **phosphoacylglycerols** and **sphingomyelins.** Both classes are found almost exclusively in the cell membranes of plants and animals, as discussed in Section 3.7.


Phospholipids are organic derivatives of phosphoric acid, formed by replacing two of the H atoms by R groups. This type of functional group is called a **phosphodiester,** or a **phosphoric acid diester.** These compounds are phosphorus analogues of carboxylic esters. In cells, the remaining OH group on phosphorus loses its proton, giving the phosphodiester a net negative charge.



phosphoric acid H_3PO_4 | phosphodiester | This form exists in cells.

31.4A Phosphoacylglycerols


Phosphoacylglycerols (or phosphoglycerides) are the second most abundant type of lipid. They form the principal lipid component of most cell membranes. Their structure resembles the triacylglycerols of Section 31.3 with one important difference. In phosphoacylglycerols, only two of the hydroxy groups of glycerol are esterified with fatty acids. The third OH group is part of a phosphodiester, which is also bonded to another low molecular weight alcohol.



phosphoacylglycerol

There are two prominent types of phosphoacylglycerols. They differ in the identity of the R" group in the phosphodiester.

- When R'' = $CH_2CH_2NH_3^+$, the phosphoacylglycerol is called a phosphatidylethanolamine or cephalin.
- When R'' = $CH_2CH_2N(CH_3)_3^+$, the phosphoacylglycerol is called a phosphatidylcholine or lecithin.




phosphatidylethanolamine or cephalin | phosphatidylcholine or lecithin

The middle carbon of the glycerol backbone of all of these compounds is a stereogenic center (labeled in blue), usually with the *R* configuration.

The phosphorus side chain of a phosphoacylglycerol makes it different from a triacylglycerol. The two fatty acid side chains form two nonpolar "tails" that lie parallel to each other, while the phosphodiester end of the molecule is a charged or polar "head." A three-dimensional structure of a phosphoacylglycerol is shown in Figure 31.3.

Figure 31.3
Three-dimensional structure of a phosphoacylglycerol



- A phosphoacylglycerol has two distinct regions: two nonpolar tails due to the long-chain fatty acids, and a very polar head from the charged phosphodiester.

As discussed in Section 3.7, when these phospholipids are mixed with water, they assemble in an arrangement called a **lipid bilayer.** The ionic heads of the phospholipid are oriented on the outside and the nonpolar tails on the inside. The identity of the fatty acids in the phospholipid determines the rigidity of this bilayer. When the fatty acids are saturated, they pack well in the interior of the lipid bilayer, and the membrane is quite rigid. When there are many unsaturated fatty acids, the nonpolar tails cannot pack as well and the bilayer is more fluid. Thus, important characteristics of this lipid bilayer are determined by the three-dimensional structure of the molecules that comprise it.

Cell membranes are composed of these lipid bilayers (see Figure 3.7). Proteins and cholesterol are embedded in the membranes as well, but the phospholipid bilayer forms the main fabric of the insoluble barrier that protects the cell.


Problem 31.6 Draw the structure of a lecithin containing oleic acid and palmitic acid as the fatty acid side chains.

Problem 31.7 Phosphoacylglycerols should remind you of soaps (Section 3.6). In what ways are these compounds similar?

31.4B Sphingomyelins

Sphingomyelins, the second major class of phospholipids, are derivatives of the amino alcohol **sphingosine,** in much the same way that triacylglycerols and phosphoacylglycerols are derivatives of glycerol. Other notable features of a sphingomyelin include:

- **A phosphodiester at C1.**
- **An amide formed with a fatty acid at C2.**




Like phosphoacylglycerols, **sphingomyelins are also a component of the lipid bilayer of cell membranes.** The coating that surrounds and insulates nerve cells, the **myelin sheath,** is particularly rich in sphingomyelins, and is vital for proper nerve function. Deterioration of the myelin sheath as seen in multiple sclerosis leads to disabling neurological problems.


Figure 31.4 compares the structural features of the most common hydrolyzable lipids: a triacylglycerol, a phosphoacylglycerol, and a sphingomyelin.

Problem 31.8 Why are phospholipids, but not triacylglycerols, found in cell membranes?

Figure 31.4
A comparison of a triacylglycerol, a phosphoacylglycerol, and a sphingomyelin




- A triacylglycerol has three nonpolar side chains.
- The three OH groups of glycerol are esterified with three fatty acids.



R" = H, CH_3

- A phosphoacylglycerol has two nonpolar side chain tails and one ionic head.
- Two OH groups of glycerol are esterified with fatty acids.
- A phosphodiester is located on a terminal carbon.



R = $CH_3(CH_2)_{12}$

R" = H, CH_3


- A sphingomyelin has two nonpolar side chain tails and one ionic head.
- A sphingomyelin is formed from sphingosine, not glycerol. One of the nonpolar tails is an amide.
- A phosphodiester is located on a terminal carbon.

31.5 Fat-Soluble Vitamins

Vitamins are organic compounds required in small quantities for normal metabolism (Section 3.5). Because our cells cannot synthesize these compounds, they must be obtained in the diet. Vitamins can be categorized as fat soluble or water soluble. **The fat-soluble vitamins are lipids.**


The four fat-soluble vitamins—**A, D, E,** and **K**—are found in fruits and vegetables, fish, liver, and dairy products. Although fat-soluble vitamins must be obtained from the diet, they do not have to be ingested every day. Excess vitamins are stored in fat cells, and then used when needed. Figure 31.5 shows the structure of these vitamins and summarizes their functions.

Figure 31.5 The fat-soluble vitamins




vitamin A

- **Vitamin A** (retinol, Section 3.5) is obtained from fish liver oils and dairy products, and is synthesized from β-carotene, the orange pigment in carrots.
- In the body, vitamin A is converted to 11-*cis*-retinal, the light-sensitive compound responsible for vision in all vertebrates (Section 21.11B). It is also needed for healthy mucous membranes.
- A deficiency of vitamin A causes night blindness, as well as dry eyes and skin.




vitamin D_3

- **Vitamin D_3** is the most abundant of the D vitamins. Strictly speaking, it is not a vitamin because it can be synthesized in the body from cholesterol. Nevertheless, it is classified as such, and many foods (particularly milk) are fortified with vitamin D_3 so that we get enough of this vital nutrient.
- Vitamin D helps regulate both calcium and phosphorus metabolism.
- A deficiency of vitamin D causes rickets, a bone disease characterized by knock-knees, spinal curvature, and other deformities.



vitamin E
(α-tocopherol)


- The term **vitamin E** refers to a group of structurally similar compounds, the most potent being α-tocopherol (Section 15.12).
- Vitamin E is an antioxidant, so it protects unsaturated side chains in fatty acids from oxidation.
- A deficiency of vitamin E causes numerous neurologic problems.



vitamin K

- **Vitamin K** (phylloquinone) regulates the synthesis of prothrombin and other proteins needed for blood to clot.
- A deficiency of vitamin K leads to excessive and sometimes fatal bleeding because of inadequate blood clotting.

Figure 31.6
Electrostatic potential plots of vitamins A and E



vitamin A vitamin E

- The electron density is distributed fairly evenly among the carbon atoms of these vitamins due to their many nonpolar C–C and C–H bonds.


Electrostatic potential plots of vitamins A and E (Figure 31.6) show that the electron density is virtually uniform in these compounds. The large regions of nonpolar C–C and C–H bonds tend to obscure small dipoles that occur in the one or two polar bonds, making these vitamins nonpolar and hydrophobic.

Problem 31.9 Explain why regularly ingesting a large excess of a fat-soluble vitamin can lead to severe health problems, whereas ingesting a large excess of a water-soluble vitamin often causes no major health problems.


31.6 Eicosanoids

The word ***eicosanoid*** is derived from the Greek word *eikosi,* meaning **20.**


The **eicosanoids** are a group of biologically active compounds containing 20 carbon atoms derived from arachidonic acid. The **prostaglandins** (Section 19.6) and the **leukotrienes** (Section 9.17) are two types of eicosanoids. Two others are the **thromboxanes** and **prostacyclins.**




$PGF_{2\alpha}$
a prostaglandin



LTC_4
a leukotriene




TXA_2
a thromboxane



PGI_2
a prostacyclin

All eicosanoids are very potent compounds present in low concentration in cells. They are **local mediators,** meaning that they perform their function in the environment in which they are synthesized. This distinguishes them from **hormones,** which are first synthesized and then transported in the bloodstream to their site of action. Eicosanoids are not stored in cells; rather, they are synthesized from arachidonic acid in response to an external stimulus.

Figure 31.7
The conversion of arachidonic acid to prostaglandins, thromboxanes, prostacyclins, and leukotrienes



Other details of the biosynthesis of leukotrienes and prostaglandins were given in Sections 9.17 and 19.6, respectively.


The synthesis of prostaglandins, thromboxanes, and prostacyclins begins with the oxidation of arachidonic acid with O_2 by a **cyclooxygenase** enzyme, which forms an unstable cyclic intermediate, PGG_2. PGG_2 is then converted via different pathways to these three classes of compounds. Leukotrienes are formed by a different pathway, using an enzyme called a **lipoxygenase.** These four paths for arachidonic acid are summarized in Figure 31.7.

Each eicosanoid is associated with specific types of biological activity (Table 31.4). In some cases, the effects oppose one another. For example, thromboxanes are vasoconstrictors that trigger blood platelet aggregation, whereas prostacyclins are vasodilators that inhibit platelet aggregation. The levels of these two eicosanoids must be in the right balance for cells to function properly.

Table 31.4 Biological Activity of the Eicosanoids


Eicosanoid	Effect	Eicosanoid	Effect
Prostaglandins	• Lower blood pressure • Inhibit blood platelet aggregation • Control inflammation • Lower gastric secretions • Stimulate uterine contractions • Relax smooth muscles of the uterus	**Thromboxanes**	• Constrict blood vessels • Trigger blood platelet aggregation
		Prostacyclins	• Dilate blood vessels • Inhibit blood platelet aggregation
		Leukotrienes	• Constrict smooth muscle, especially in the lungs

Because of their wide range of biological functions, prostaglandins and their analogues have found several clinical uses. For example, **dinoprostone,** the generic name for **PGE_2,** is administered to relax the smooth muscles of the uterus when labor is induced, and to terminate pregnancies in the early stages.




PGE_2
(dinoprostone)

Because prostaglandins themselves are unstable in the body, often having half-lives of only minutes, more stable analogues have been developed that retain their important biological activity longer. Misoprostol is a prostaglandin analogue administered to prevent gastric ulcers in patients who are at high risk of developing them, and unoprostone isopropyl is used to decrease eye pressure in glaucoma patients.



(+ epimer at C16)

misoprostol




unoprostone isopropyl

Studying the biosynthesis of eicosanoids has led to other discoveries as well. For example, aspirin and other nonsteroidal anti-inflammatory drugs (**NSAIDs**) inactivate the cyclooxygenase enzyme needed for prostaglandin synthesis. In this way, NSAIDs block the synthesis of the prostaglandins that cause inflammation (Section 19.6).


More recently, it has been discovered that two *different* cyclooxygenase enzymes, called **COX-1** and **COX-2,** are responsible for prostaglandin synthesis. COX-1 is involved with the usual production of prostaglandins, but COX-2 is responsible for the synthesis of additional prostaglandins in inflammatory diseases like arthritis. **NSAIDs like aspirin and ibuprofen inactivate both the COX-1 and COX-2 enzymes.** This activity also results in an increase in gastric secretions, making an individual more susceptible to ulcer formation.

A group of anti-inflammatory drugs that block only the COX-2 enzyme was developed in the 1990s. These drugs—**rofecoxib, valdecoxib,** and **celecoxib**—do not cause an increase in gastric


secretions, and thus were touted as especially effective NSAIDs for patients with arthritis, who need daily doses of these medications. Unfortunately, both rofecoxib and valdecoxib have now been removed from the market, because their use has been associated with an increased risk of heart attack and stroke.



Generic name rofecoxib
Trade name Vioxx



Generic name valdecoxib
Trade name Bextra




Generic name celecoxib
Trade name Celebrex

The discovery of drugs that block prostaglandin synthesis illustrates how basic research in organic chemistry can lead to important practical applications. Elucidating the structure and biosynthesis of prostaglandins began as a project in basic research. It has now resulted in a number of applications that benefit many individuals with various illnesses.

31.7 Terpenes


***Terpenes* are lipids composed of repeating five-carbon units called isoprene units.** An **isoprene unit** has five carbons: four in a row, with a one-carbon branch on a middle carbon.




Terpenes are hydrocarbons that may be acyclic or have one or more rings. The term ***terpenoid*** is used for compounds that contain isoprene units as well as an oxygen heteroatom. Many **essential oils,** a group of compounds isolated from plant sources by distillation, are terpenes and terpenoids. Examples include myrcene from bayberry and menthol from peppermint.



bayberry plant
(Photo: © Henriette Kress)



myrcene



menthol




peppermint plant
(Photo: © Mark Turner/
Getty Images)

31.7A Locating Isoprene Units

How do we identify the isoprene units in these molecules? Start at one end of the molecule near a branch point. Then **look for a four-carbon chain with a one-carbon branch.** This forms one isoprene unit. Continue along the chain or around the ring until all the carbons are part of an isoprene unit. Keep in mind the following:

- An isoprene unit may be composed of C – C σ bonds only, or there may be π bonds at any position.
- Isoprene units are always connected by one or more carbon–carbon bonds.
- Each carbon atom is part of one isoprene unit only.
- Every isoprene unit has five carbon atoms. Heteroatoms may be present but their presence is ignored in locating isoprene units.

Myrcene and menthol, for example, each have 10 carbon atoms, so they are composed of two isoprene units.




Terpenes and terpenoids are classified by the number of isoprene units they contain. A ***monoterpene* (or *monoterpenoid*) contains 10 carbons** and has two isoprene units, a ***sesquiterpene* (or *sesquiterpenoid*) contains 15 carbons** and has three isoprene units, and so forth. The different terpene classes are summarized in Table 31.5.

Several examples, with the isoprene units labeled in red, are given in Figure 31.8.

Table 31.5 Classes of Terpenes and Terpenoids


Name	Number of C atoms	Number of isoprene units
Monoterpene (Monoterpenoid)	10	2
Sesquiterpene (Sesquiterpenoid)	15	3
Diterpene (Diterpenoid)	20	4
Sesterterpene (Sesterterpenoid)	25	5
Triterpene (Triterpenoid)	30	6
Tetraterpene (Tetraterpenoid)	40	8

Figure 31.8
Examples of some common terpenes and terpenoids



- Isoprene units are labeled in red, with C–C bonds (in black) joining two units.
- The source of each terpene or terpenoid is given in parentheses.


Problem 31.10 Locate the isoprene units in each compound.






Amber, fossilized resin that oozed from trees long ago, contains biformene, as well as many other terpenoids called labdanoids. (Photo: © Thomas J. Abercrombie/Getty Images)

Problem 31.11 Locate the isoprene units in biformene, a component of amber, and classify biformene as a monoterpene, sesquiterpene, etc.




biformene

31.7B The Biosynthesis of Terpenes and Terpenoids



organic diphosphate





diphosphate leaving group


PP_i

Terpene and terpenoid biosynthesis is an excellent example of how syntheses in nature occur with high efficiency. There are two ways this is accomplished.


[1] The same reaction is used over and over again to prepare progressively more complex compounds.

[2] Key intermediates along the way serve as the starting materials for a wide variety of other compounds.

All terpenes and terpenoids are synthesized from dimethylallyl diphosphate and isopentenyl diphosphate. Both of these five-carbon compounds are organic diphosphates (Section 16.2B) with a good leaving group (diphosphate, $P_2O_7^{4-}$, PP_i).




dimethylallyl diphosphate



isopentenyl diphosphate

The overall strategy of biosynthesis from dimethylallyl diphosphate and isopentenyl diphosphate is summarized in Figure 31.9.

Figure 31.9
An outline of terpene and terpenoid biosynthesis



There are three basic parts:

[1] The two C_5 diphosphates are converted to **geranyl diphosphate, a C_{10} monoterpenoid.** Geranyl diphosphate is the starting material for all other monoterpenes and monoterpenoids.


[2] Geranyl diphosphate is converted to **farnesyl diphosphate, a C_{15} sesquiterpenoid,** by addition of a five-carbon unit. Farnesyl diphosphate is the starting material for all sesquiterpenes, diterpenes, and related terpenoids.

[3] Two molecules of farnesyl diphosphate are converted to **squalene, a C_{30} triterpene.** Squalene is the starting material for all triterpenes and steroids.

The biological formation of geranyl diphosphate from the two five-carbon diphosphates was shown in Mechanism 16.1. The biological conversion of geranyl diphosphate to farnesyl diphosphate involves a similar pathway, as shown in Mechanism 31.1.

Mechanism 31.1 Biological Formation of Farnesyl Diphosphate


Part [1] Formation of a new carbon–carbon σ bond



1 Loss of the diphosphate leaving group forms a resonance-stabilized carbocation.


2 Nucleophilic attack of isopentenyl diphosphate on the allylic carbocation forms the new C–C σ bond.

Part [2] Formation of a π bond by loss of a proton




3 Loss of a proton (shown with the general base B:) forms a new π bond and farnesyl diphosphate.

Two molecules of farnesyl diphosphate react to form squalene, from which all other triterpenes and steroids are synthesized.




Aqueous hydrolysis of geranyl and farnesyl diphosphates forms the monoterpenoid geraniol and the sesquiterpenoid farnesol, respectively.



All other terpenes and terpenoids are biologically derived from geranyl and farnesyl diphosphates by a series of reactions. Cyclic compounds are formed by intramolecular reactions involving nucleophilic attack of π bonds on intermediate carbocations. To form some cyclic compounds, the *E* double bond in geranyl diphosphate must first isomerize to an isomeric diphosphate with a *Z* double bond, neryl diphosphate, by the process illustrated in Mechanism 31.2. Isomerization forms a substrate with a leaving group and nucleophilic double bond in close proximity, so that an intramolecular reaction can occur.

Mechanism 31.2 Isomerization of Geranyl Diphosphate to Neryl Diphosphate




1 – 2 Loss of diphosphate forms a resonance-stabilized carbocation, which reacts with the diphosphate anion to form linalyl diphosphate.


3 Bond rotation of the single bond shown in red and loss of diphosphate forms a resonance-stabilized carbocation.

4 Nucleophilic attack with diphosphate forms neryl diphosphate, which has the leaving group and the double bond at the other end of the chain in close proximity for intramolecular cyclization.


In the synthesis of α-terpineol or limonene, for example, geranyl diphosphate isomerizes to form neryl diphosphate (Step [1] in the following reaction sequence). Neryl diphosphate then cyclizes to a 3° carbocation by intramolecular attack (Steps [2]–[3]). Nucleophilic attack of water on this carbocation yields the monoterpenoid α-terpineol (Step [4]) or loss of a proton yields the monoterpene limonene (Step [5]).



Problem 31.12 Write a stepwise mechanism for the following reaction.



Problem 31.13 Draw a stepwise mechanism for the conversion of geranyl diphosphate to α-terpinene.




31.8 Steroids


The steroids are a group of tetracyclic lipids, many of which are biologically active.

31.8A Steroid Structure

Steroids are composed of three six-membered rings and one five-membered ring, joined together as drawn. Many steroids also contain two methyl groups, called **angular methyl groups,** at the two ring junctions indicated. The steroid rings are lettered **A, B, C,** and **D,** and the 17 ring carbons are numbered as shown. The two angular methyl groups are numbered C18 and C19.



Whenever two rings are fused together, the substituents at the ring fusion can be arranged cis or trans. To see more easily why this is true, consider **decalin,** which consists of two six-membered rings fused together. ***trans*-Decalin** has the two hydrogen atoms at the ring fusion on opposite sides, whereas ***cis*-decalin** has them on the same side.




decalin
Two six-membered rings share a C–C bond.

***trans*-decalin**
2 H's on opposite sides

***cis*-decalin**
2 H's on the same side


Three-dimensional structures of these molecules show how different these two possible arrangements actually are. The two rings of *trans*-decalin lie roughly in the same plane, whereas the two

rings of *cis*-decalin are almost perpendicular to each other. **The trans arrangement is lower in energy and therefore more stable.**



In steroids, each ring fusion could theoretically have the cis or trans configuration, but, by far the most common arrangement is all trans. Because of this, **all four rings of the steroid skeleton lie in the same plane,** and the ring system is fairly rigid. The two angular methyl groups are oriented perpendicular to the plane of the molecule. These methyl groups make one side of the steroid skeleton significantly more hindered than the other, as shown in Figure 31.10.

Figure 31.10
The three-dimensional structure of the steroid nucleus




Although steroids have the same fused-ring arrangement of carbon atoms, they differ in the identity and location of the substituents attached to that skeleton.

Problem 31.14 (a) Draw a skeletal structure of the anabolic steroid 4-androstene-3,17-dione, also called "andro," from the following description. Andro contains the tetracyclic steroid skeleton with carbonyl groups at C3 and C17, a double bond between C4 and C5, and methyl groups bonded to C10 and C13. (b) Add wedges and dashed wedges for all stereogenic centers with the following information: the configuration at C10 is *R,* the configuration at C13 is *S,* and all substituents at ring fusions are trans to each other.

31.8B Cholesterol

Cholesterol has also been discussed in Sections 3.4C and 4.15. The role of cholesterol in plaque formation and atherosclerosis was discussed in Section 22.17.

Cholesterol, the chapter-opening molecule, has the tetracyclic carbon skeleton characteristic of steroids. It also has eight stereogenic carbons (seven on rings and one on a side chain), so there are $2^8 = 256$ possible stereoisomers. In nature, however, only the following stereoisomer exists:




cholesterol

Cholesterol is essential to life because it forms an important component of cell membranes and is the starting material for the synthesis of all other steroids. Humans do not have to ingest cholesterol, because it is synthesized in the liver and then transported to other tissues through the bloodstream. Because cholesterol has only one polar OH group and many nonpolar C−C and C−H bonds, it is insoluble in water (and, thus, in the aqueous medium of the blood).

Konrad Bloch and Feodor Lynen shared the 1964 Nobel Prize in Physiology or Medicine for unraveling the complex transformation of squalene to cholesterol.

Cholesterol is synthesized in the body from squalene, a C_{30} triterpene that is itself prepared from smaller terpenes, as discussed in Section 31.7B. Because the biosynthesis of all terpenes begins with acetyl CoA, every one of the 27 carbon atoms of cholesterol comes from the same two-carbon precursor. The major steps in the conversion of squalene to cholesterol are given in Figure 31.11.

Figure 31.11
The biosynthesis of cholesterol



The conversion of squalene to cholesterol consists of five different parts:

[1] **Epoxidation** of squalene with an enzyme, squalene epoxidase, gives squalene oxide, which contains a single epoxide on one of the six double bonds.

[2] **Cyclization** of squalene oxide yields a carbocation, called the protosterol cation. This reaction results in the formation of four new C−C bonds and the tetracyclic ring system.


[3] **The protosterol carbocation rearranges** by a series of 1,2-shifts of either a hydrogen or methyl group to form another 3° carbocation.

[4] **Loss of a proton** gives an alkene called **lanosterol.** Although lanosterol has seven stereogenic centers, a single stereoisomer is formed.

[5] Lanosterol is then converted to cholesterol by a multistep process that results in removal of three methyl groups.

Several drugs called statins are now available to reduce the level of cholesterol in the bloodstream. These compounds act by blocking the biosynthesis of cholesterol at its very early stages. Two examples include atorvastatin (Lipitor) and simvastatin (Zocor), whose structures appear in Figure 31.12.

Figure 31.12
Two cholesterol-lowering drugs




Generic name atorvastatin
Trade name Lipitor

Generic name simvastatin
Trade name Zocor

Problem 31.15 Draw the enantiomer and any two diastereomers of cholesterol. Does the OH group of cholesterol occupy an axial or equatorial position?

Problem 31.16 Treatment of cholesterol with mCPBA results in formation of a single epoxide **A,** with the stereochemistry drawn. Why isn't the isomeric epoxide **B** formed to any extent?








31.8C Other Steroids


Many other important steroids are hormones secreted by the endocrine glands. Two classes are the **sex hormones** and the **adrenal cortical steroids.**


There are two types of female sex hormones, **estrogens** and **progestins.** The male sex hormones are called **androgens.** The most important members of each hormone type are given in Table 31.6.

Table 31.6 The Female and Male Sex Hormones

Structure	Properties
 estradiol  estrone	• **Estradiol** and **estrone** are estrogens synthesized in the ovaries. They control the development of secondary sex characteristics in females and regulate the menstrual cycle.
 progesterone	• **Progesterone** is often called the "pregnancy hormone." It is responsible for the preparation of the uterus for implantation of a fertilized egg.
 testosterone  androsterone	• **Testosterone** and **androsterone** are androgens synthesized in the testes. They control the development of secondary sex characteristics in males.

Synthetic analogues of these steroids have found important uses, such as ethynylestradiol and norethindrone in oral contraceptives, first mentioned in Section 11.4.


ethynylestradiol



norethindrone





Some body builders use anabolic steroids to increase muscle mass. Long-term or excessive use can cause many health problems, including high blood pressure, liver damage, and cardiovascular disease. (Photo: © Comstock/ JupiterImages RF)

Synthetic androgen analogues, called **anabolic steroids,** promote muscle growth. They were first developed to help individuals whose muscles had atrophied from lack of use following surgery. They have since come to be used by athletes and body builders, although their use is not permitted in competitive sports. Many physical and psychological problems result from their prolonged use.


Anabolic steroids, such as stanozolol, nandrolone, and tetrahydrogestrinone have the same effect on the body as testosterone, but they are more stable, so they are not metabolized as quickly. Tetrahydrogestrinone (also called THG or The Clear), the performance-enhancing drug used by track star Marion Jones during the 2000 Sydney Olympics, was considered a "designer steroid" because it was initially undetected in urine tests for doping. After its chemical structure and properties were determined, it was added to the list of banned anabolic steroids in 2004.



stanozolol



nandrolone


tetrahydrogestrinone

A second group of steroid hormones includes the **adrenal cortical steroids.** Three examples of these hormones are **cortisone, cortisol,** and **aldosterone.** All of these compounds are synthesized in the outer layer of the adrenal gland. Cortisone and cortisol serve as anti-inflammatory agents and they also regulate carbohydrate metabolism. Aldosterone regulates blood pressure and volume by controlling the concentration of Na^+ and K^+ in body fluids.


cortisone


cortisol



aldosterone

KEY CONCEPTS


Lipids

Hydrolyzable Lipids

[1] **Waxes** (31.2)—Esters formed from a long-chain alcohol and a long-chain carboxylic acid.



R, R' = long chains of C's

[2] **Triacylglycerols** (31.3)—Triesters of glycerol with three fatty acids.



R, R', R'' = alkyl groups with 11–19 C's

[3] **Phospholipids** (31.4)


a. Phosphatidylethanolamine (cephalin)


R, R' = long carbon chain

b. Phosphatidylcholine (lecithin)


R, R' = long carbon chain

c. Sphingomyelin


R = $CH_3(CH_2)_{12}$
R' = long carbon chain
R'' = H or CH_3

Nonhydrolyzable Lipids


[1] **Fat-soluble vitamins** (31.5)—Vitamins A, D, E, and K.

[2] **Eicosanoids** (31.6)—Compounds containing 20 C's derived from arachidonic acid. There are four types: prostaglandins, thromboxanes, prostacyclins, and leukotrienes.

[3] **Terpenes** (31.7)—Lipids composed of repeating 5 C units called isoprene units.

Isoprene unit	Types of terpenes			
	[1] monoterpene	10 C's	[4] sesterterpene	25 C's
	[2] sesquiterpene	15 C's	[5] triterpene	30 C's
	[3] diterpene	20 C's	[6] tetraterpene	40 C's


[4] **Steroids** (31.8)—Tetracyclic lipids composed of three six-membered and one five-membered ring.




PROBLEMS


Problems Using Three-Dimensional Models

31.17 Locate the isoprene units in each compound.

a. 
α-pinene


b. 
humulene

31.18 Convert each ball-and-stick model to a skeletal structure that clearly shows the stereochemistry at the ring fusion of these decalin derivatives.

a. 

b. 

31.19 Convert the ball-and-stick model of androsterone to (a) a skeletal structure using wedges and dashed wedges around all stereogenic centers; and (b) a three-dimensional representation using chair cyclohexane rings.



androsterone


Waxes, Triacylglycerols, and Phospholipids

31.20 One component of lanolin, the wax that coats sheep's wool, is derived from cholesterol and stearic acid. Draw its structure, including the correct stereochemistry at all stereogenic centers.


31.21 Draw all possible constitutional isomers of a triacylglycerol formed from one mole each of palmitic, oleic, and linoleic acids. Locate the tetrahedral stereogenic centers in each constitutional isomer.

31.22 What is the structure of an optically inactive triacylglycerol that yields two moles of oleic acid and one mole of palmitic acid when hydrolyzed in aqueous acid?


31.23 Triacylglycerol **L** yields compound **M** when treated with excess H_2, Pd-C. Ozonolysis of **L** ([1] O_3; [2] $(CH_3)_2S$) affords compounds **N – P.** What is the structure of **L?**




M



N



O



P

31.24 Draw the structure of the following phospholipids:

a. a cephalin formed from two molecules of stearic acid

b. a sphingomyelin formed from palmitic acid


Prostaglandins

31.25 A difficult problem in the synthesis of $PGF_{2\alpha}$ is the introduction of the OH group at C15 in the desired configuration.


a. Label this stereogenic center as *R* or *S*.

b. A well known synthesis of $PGF_{2\alpha}$ involves reaction of **A** with $Zn(BH_4)_2$, a metal hydride reagent similar in reactivity to $NaBH_4$, to form two isomeric products, **B** and **C.** Draw their structures and indicate their stereochemical relationship.

c. Suggest a reagent to convert **A** to the single stereoisomer **X.**




$PGF_{2\alpha}$



A

[1] $Zn(BH_4)_2$
[2] H_2O


B and **C**




X


Terpenes and Terpenoids

31.26 Locate the isoprene units in each compound.


a. 

e. 


patchouli alcohol

g. 


dextropimaric acid

b. 


carvone


f. 

periplanone B

h. 


β-amyrin

c. 
lycopene

d. 
β-carotene


31.27 Classify each terpene and terpenoid in Problem 31.26 (e.g., as a monoterpene, sesquiterpene, etc.).

31.28 Crocin, which occurs naturally in crocus and gardenia flowers, is primarily responsible for the color of saffron. (a) What lipid and monosaccharides are formed by the hydrolysis of crocin? (b) Classify the lipid as a monoterpenoid, diperpenoid, etc., and locate the isoprene units.




crocin

31.29 An isoprene unit can be thought of as having a head and a tail. The "head" of the isoprene unit is located at the end of the chain nearest the branch point, and the "tail" is located at the end of the carbon chain farthest from the branch point. Most isoprene units are connected together in a "head-to-tail" fashion, as illustrated. For both lycopene (Problem 31.26), and squalene (Figure 31.9), decide which isoprene units are connected in a head-to-tail fashion and which are not.




These two isoprene units are connected in a head-to-tail fashion.

31.30 Draw a stepwise mechanism for the conversion of neryl diphosphate to α-pinene. α-Pinene is a component of pine oil and rosemary oil.



neryl diphosphate α-pinene

31.31 Flexibilene is a terpene isolated from *Sinularia flexibilis,* a soft coral found in the Indian Ocean. Draw a stepwise mechanism for the formation of flexibilene from farnesyl diphosphate and isopentenyl diphosphate. What is unusual about the cyclization that forms the 15-membered ring of flexibilene?




flexibilene

31.32 The biosynthesis of lanosterol from squalene has intrigued chemists since its discovery. It is now possible, for example, to synthesize polycyclic compounds from acyclic or monocyclic precursors by reactions that form several C – C bonds in a single reaction mixture.

a. Draw a stepwise mechanism for the following reaction.


b. Show how **X** can be converted to 16,17-dehydroprogesterone. (Hint: See Figure 24.5 for a related conversion.)




X 16,17-dehydroprogesterone

Steroids

31.33 Draw three-dimensional structures for each alcohol. Label the OH groups as occupying axial or equatorial positions.




31.34 Axial alcohols are oxidized faster than equatorial alcohols by PCC and other Cr^{6+} oxidants. Which OH group in each compound is oxidized faster?



31.35 (a) Draw a skeletal structure of the anabolic steroid methenolone from the following description. Methenolone contains the tetracyclic steroid skeleton with a carbonyl group at C3, a hydroxyl at C17, a double bond between C1 and C2, and methyl groups bonded to C1, C10, and C13. (b) Add wedges and dashed wedges for all stereogenic centers with the following information: the configuration at C10 is *R,* the configuration at C13 is *S,* the configuration at C17 is *S*, and all substituents at ring fusions are trans to each other. (c) Draw the structure of Primobolan, the product formed when methenolone is treated with $CH_3(CH_2)_5COCl$ and pyridine. Primobolan is an anabolic steroid that can be taken orally or by injection and has been used illegally by well-known Major League Baseball players.

31.36 Betamethasone is a synthetic anti-inflammatory steroid used as a topical cream for itching. Betamethasone is derived from cortisol, with the following structural additions: a C=C between C1 and C2, a fluorine at C9, and a methyl group at C16. The configuration at C9 is *R*, and the configuration at C16 is *S*. Draw the structure of betamethasone.

31.37 a. Draw a three-dimensional structure for the following steroid.
b. What is the structure of the single stereoisomer formed by reduction of this ketone with H_2, Pd-C? Explain why only one stereoisomer is formed.




31.38 Draw the products formed when cholesterol is treated with each reagent. Indicate the stereochemistry around any stereogenic centers in the product.

a. CH_3COCl
b. H_2, Pd-C
c. PCC
d. oleic acid, H^+
e. [1] BH_3•THF; [2] H_2O_2, ^-OH


Challenge Problems

31.39 Draw a stepwise mechanism for the following conversion, which forms camphene. Camphene is a component of camphor and citronella oils.




camphene

31.40 Draw a stepwise mechanism for the following reaction.



31.41 Farnesyl diphosphate is cyclized to sesquiterpene **A,** which is then converted to the bicyclic product epi-aristolochene. Write a stepwise mechanism for both reactions.

















farnesyl diphosphate **A** epi-aristolochene









APPENDIX




A

pK_a Values for Selected Compounds

Compound	pK_a
HI	–10
HBr	–9
H_2SO_4	–9
	–7.3
	–7
HCl	–7
$[(CH_3)_2OH]^+$	–3.8
$(CH_3OH_2)^+$	–2.5
H_3O^+	–1.7
CH_3SO_3H	–1.2
	0.0
CF_3CO_2H	0.2
CCl_3CO_2H	0.6
	1.0
Cl_2CHCO_2H	1.3
H_3PO_4	2.1
FCH_2CO_2H	2.7
$ClCH_2CO_2H$	2.8
$BrCH_2CO_2H$	2.9
ICH_2CO_2H	3.2
HF	3.2
	3.4
HCO_2H	3.8

Compound	pK_a
	3.9
	4.0
	4.2
	4.3
	4.5
	4.6
CH_3CO_2H	4.8
$(CH_3)_3CCO_2H$	5.0
	5.1
	5.3
	5.3
H_2CO_3	6.4
H_2S	7.0
	7.1
	7.8

Compound	pK_a
	8.9
$HC\equiv N$	9.1
	9.4
NH_4^+	9.4
$H_3\overset{+}{N}CH_2CO_2^-$	9.8
	10.0
	10.2
HCO_3^-	10.2
CH_3NO_2	10.2
	10.3
CH_3CH_2SH	10.5
$[(CH_3)_3NH]^+$	10.6
	10.7
$(CH_3NH_3)^+$	10.7
	10.7
$[(CH_3)_2NH_2]^+$	10.7
CF_3CH_2OH	12.4
	13.3

Compound	pK_a
	15
CH_3OH	15.5
H_2O	15.7
CH_3CH_2OH	16
CH_3CONH_2	16
CH_3CHO	17
$(CH_3)_3COH$	18
$(CH_3)_2C=O$	19.2
$CH_3CO_2CH_2CH_3$	24.5
$HC\equiv CH$	25
$CH_3C\equiv N$	25
$CHCl_3$	25
$CH_3CON(CH_3)_2$	30
H_2	35
NH_3	38
CH_3NH_2	40
	41
	43
$CH_2=CHCH_3$	43
$CH_2=CH_2$	44
	46
CH_4	50
CH_3CH_3	50

APPENDIX

B

Nomenclature

Although the basic principles of nomenclature are presented in the body of this text, additional information is often needed to name many complex organic compounds. Appendix B concentrates on three topics:


- **Naming alkyl substituents that contain branching**
- **Naming polyfunctional compounds**
- **Naming bicyclic compounds**

Naming Alkyl Substituents That Contain Branching


Alkyl groups that contain any number of carbons and no branches are named as described in Section 4.4A: change the *-ane* ending of the parent alkane to the suffix *-yl*. Thus the seven-carbon alkyl group $CH_3CH_2CH_2CH_2CH_2CH_2CH_2-$ is called *heptyl*.

When an alkyl substituent also contains branching, follow a stepwise procedure:

[1] Identify the longest carbon chain of the alkyl group that begins at the point of attachment to the parent. Begin numbering at the point of attachment and use the suffix *-yl* to indicate an alkyl group.




4 C's in the chain ⟶ butyl group



5 C's in the chain ⟶ pentyl group

[2] Name all branches off the main alkyl chain and use the numbers from Step [1] to designate their location.

methyl at C3




3-methylbutyl

methyls at C1 and C3




1,3-dimethylpentyl

[3] Set the entire name of the substituent in parentheses, and alphabetize this substituent name by the first letter of the complete name.



(3-methylbutyl)cyclohexane



1-(1,3-dimethylpentyl)-2-methylcyclohexane

- Alphabetize the **d** of **d**imethylpentyl before the **m** of **m**ethyl.
- Number the ring to give the lower number to the first substituent alphabetically: place the dimethylpentyl group at C1.

Naming Polyfunctional Compounds

Many organic compounds contain more than one functional group. When one of those functional groups is halo (X−) or alkoxy (RO−), these groups are named as substituents as described in Sections 7.2 and 9.3B. To name other polyfunctional compounds, we must learn which functional group is assigned a higher priority in the rules of nomenclature. Two steps are usually needed:

[1] **Name a compound using the suffix of the highest priority group,** and name other functional groups as *substituents.* Table B.1 lists the common functional groups in order of decreasing priority, as well as the prefixes needed when a functional group must be named as a substituent.


[2] Number the carbon chain to give the lower number to the highest priority functional group that can be named as a suffix, and then follow all other rules of nomenclature. Examples are shown in Figure B.1.

Table B.1 Summary of Functional Group Nomenclature

Functional group	Suffix	Substituent name (prefix)
Carboxylic acid	-oic acid	carboxy
Ester	-oate	alkoxycarbonyl
Amide	-amide	amido
Nitrile	-nitrile	cyano
Aldehyde	-al	oxo (=O) or formyl (– CHO)
Ketone	-one	oxo
Alcohol	-ol	hydroxy
Amine	-amine	amino
Alkene	-ene	alkenyl
Alkyne	-yne	alkynyl
Alkane	-ane	alkyl
Ether	—	alkoxy
Halide	—	halo


Increasing priority

Figure B.1
Examples of nomenclature of polyfunctional compounds




3-amino-2-hydroxybutanal

Name as a derivative of an aldehyde, because CHO is the highest priority functional group.




o-cyanobenzoic acid

Name as a derivative of benzoic acid, because COOH is the higher priority functional group.



methyl 4-oxohexanoate

Name as a derivative of an ester, because COOR is the higher priority functional group.






4-formyl-3-methoxycyclohexanecarboxamide

Name as a derivative of an amide, because $CONH_2$ is the highest priority functional group.


Polyfunctional compounds that contain C−C double and triple bonds have characteristic suffixes to identify them, as shown in Table B.2. The higher priority functional group is assigned the lower number.

Table B.2 Naming Polyfunctional Compounds with C–C Double and Triple Bonds

Functional groups	Suffix	Example
C = C and OH	enol	 5-methylhex-4-en-1-ol
C = C + C = O (ketone)	enone	 (*E*)-hept-4-en-3-one
C = C + C ≡ C	enyne	 hex-1-en-5-yne

Naming Bicyclic Compounds


Bicyclic ring systems—compounds that contain two rings that share one or two carbon atoms—can be bridged, fused, or spiro.




bridged ring fused ring spiro ring

- A bridged ring system contains two rings that share two non-adjacent carbons.
- A fused ring system contains two rings that share a common carbon–carbon bond.
- A spiro ring system contains two rings that share one carbon atom.


Fused and bridged ring systems are named as bicyclo[*x.y.z*]alkanes, where the parent alkane corresponds to the total number of carbons in both rings. The numbers *x, y,* and *z* refer to the number of carbons that join the shared carbons together, written in order of *decreasing* size. For a fused ring system, *z* always equals zero, because the two shared carbons are directly joined together. The shared carbons in a bridged ring system are called the **bridgehead carbons.**



Rings are numbered beginning at a *shared* carbon, and continuing around the *longest* bridge first, then the next longest, and so forth.



3,3-dimethylbicyclo[3.2.1]octane




7,7-dimethylbicyclo[2.2.1]heptane

Spiro ring systems are named as spiro[*x*.*y*]alkanes where the parent alkane corresponds to the total number of carbons in both rings, and *x* and *y* refer to the number of carbons that join the shared carbon (the spiro carbon), written in order of *increasing* size. When substituents are present, the rings are numbered beginning with a carbon *adjacent* to the spiro carbon in the *smaller* ring.



10 C's in the ring system

Name: spiro[4.5]decane



8 C's in the ring system

Name: 2-methylspiro[3.4]octane

APPENDIX

C

Bond Dissociation Energies for Some Common Bonds [A–B → A• + •B]

Bond	ΔH° kJ/mol	(kcal/mol)
H – Z bonds		
H – F	569	(136)
H – Cl	431	(103)
H – Br	368	(88)
H – I	297	(71)
H – OH	498	(119)
Z – Z bonds		
H – H	435	(104)
F – F	159	(38)
Cl – Cl	242	(58)
Br – Br	192	(46)
I – I	151	(36)
HO – OH	213	(51)
R – H bonds		
CH_3 – H	435	(104)
CH_3CH_2 – H	410	(98)
$CH_3CH_2CH_2$ – H	410	(98)
$(CH_3)_2CH$ – H	397	(95)
$(CH_3)_3C$ – H	381	(91)
CH_2=CH – H	435	(104)
HC≡C – H	523	(125)
CH_2=$CHCH_2$ – H	364	(87)
C_6H_5 – H	460	(110)
$C_6H_5CH_2$ – H	356	(85)
R – R bonds		
CH_3 – CH_3	368	(88)
CH_3 – CH_2CH_3	356	(85)
CH_3 – CH=CH_2	385	(92)
CH_3 – C≡CH	489	(117)

Bond	ΔH° kJ/mol	(kcal/mol)
R – X bonds		
CH_3 – F	456	(109)
CH_3 – Cl	351	(84)
CH_3 – Br	293	(70)
CH_3 – I	234	(56)
CH_3CH_2 – F	448	(107)
CH_3CH_2 – Cl	339	(81)
CH_3CH_2 – Br	285	(68)
CH_3CH_2 – I	222	(53)
$(CH_3)_2CH$ – F	444	(106)
$(CH_3)_2CH$ – Cl	335	(80)
$(CH_3)_2CH$ – Br	285	(68)
$(CH_3)_2CH$ – I	222	(53)
$(CH_3)_3C$ – F	444	(106)
$(CH_3)_3C$ – Cl	331	(79)
$(CH_3)_3C$ – Br	272	(65)
$(CH_3)_3C$ – I	209	(50)
R – OH bonds		
CH_3 – OH	389	(93)
CH_3CH_2 – OH	393	(94)
$CH_3CH_2CH_2$ – OH	385	(92)
$(CH_3)_2CH$ – OH	401	(96)
$(CH_3)_3C$ – OH	401	(96)
Other bonds		
CH_2=CH_2	635	(152)
HC=CH	837	(200)
O=C=O	535	(128)
O_2	497	(119)

APPENDIX

D Reactions That Form Carbon–Carbon Bonds

Section	Reaction
11.11A	S_N2 reaction of an alkyl halide with an acetylide anion, $^-C{\equiv}CR$
11.11B	Opening of an epoxide ring with an acetylide anion, $^-C{\equiv}CR$
15.14	Radical polymerization of an alkene
16.12	Diels–Alder reaction
18.5	Friedel–Crafts alkylation
18.5	Friedel–Crafts acylation
20.10	Reaction of an aldehyde or ketone with a Grignard or organolithium reagent
20.13A	Reaction of an acid chloride with a Grignard or organolithium reagent
20.13A	Reaction of an ester with a Grignard or organolithium reagent
20.13B	Reaction of an acid chloride with an organocuprate reagent
20.14A	Reaction of a Grignard reagent with CO_2
20.14B	Reaction of an epoxide with an organometallic reagent
20.15	Reaction of an α,β-unsaturated carbonyl compound with an organocuprate reagent
21.9	Cyanohydrin formation
21.10	Wittig reaction to form an alkene
22.18	S_N2 reaction of an alkyl halide with NaCN
22.18C	Reaction of a nitrile with a Grignard or organolithium reagent
23.8	Direct enolate alkylation using LDA and an alkyl halide
23.9	Malonic ester synthesis to form a carboxylic acid
23.10	Acetoacetic ester synthesis to form a ketone
24.1	Aldol reaction to form a β-hydroxy carbonyl compound or an α,β-unsaturated carbonyl compound
24.2	Crossed aldol reaction
24.3	Directed aldol reaction
24.5	Claisen reaction to form a β-keto ester
24.6	Crossed Claisen reaction to form a β-dicarbonyl compound
24.7	Dieckmann reaction to form a five- or six-membered ring
24.8	Michael reaction to form a 1,5-dicarbonyl compound
24.9	Robinson annulation to form a cyclohex-2-enone
25.14	Reaction of a diazonium salt with CuCN
26.1	Coupling of an organocuprate reagent (R_2CuLi) with an organic halide (R'X)
26.2	The palladium-catalyzed Suzuki reaction of an organic halide with an organoborane
26.3	The palladium-catalyzed Heck reaction of a vinyl or aryl halide with an alkene
26.4	Addition of a dihalocarbene to an alkene to form a cyclopropane
26.5	Simmons–Smith reaction of an alkene with CH_2I_2 and Zn(Cu) to form a cyclopropane
26.6	Olefin metathesis
27.3	Electrocyclic reactions
27.4	Cycloaddition reactions
27.5	Sigmatropic rearrangements
28.10B	Kiliani–Fischer synthesis of an aldose
29.2B	Alkylation of diethyl acetamidomalonate to form an amino acid
29.2C	Strecker synthesis of an amino acid
30.2	Chain-growth polymerization
30.4	Polymerization using Ziegler–Natta catalysts

APPENDIX

E



Characteristic IR Absorption Frequencies






Bond	Functional group	Wavenumber (cm^{-1})	Comment
O – H			
	• ROH	3600–3200	broad, strong
	• RCO_2H	3500–2500	very broad, strong
N – H			
	• RNH_2	3500–3300	two peaks
	• R_2NH	3500–3300	one peak
	• $RCONH_2$, RCONHR	3400–3200	one or two peaks; N – H bending also observed at 1640 cm^{-1}
C – H			
	• C_{sp} – H	3300	sharp, often strong
	• C_{sp^2} – H	3150–3000	medium
	• C_{sp^3} – H	3000–2850	strong
	• C_{sp^2} – H of RCHO	2830–2700	one or two peaks
C ≡ C		2250	medium
C ≡ N		2250	medium
C = O			strong
	• RCOCl	1800	
	• $(RCO)_2O$	1800, 1760	two peaks
	• RCO_2R	1745–1735	increasing $\tilde{\nu}$ with decreasing ring size
	• RCHO	1730	
	• R_2CO	1715	increasing $\tilde{\nu}$ with decreasing ring size
	• R_2CO, conjugated	1680	
	• RCO_2H	1710	
	• $RCONH_2$, RCONHR, $RCONR_2$	1680–1630	increasing $\tilde{\nu}$ with decreasing ring size
C = C			
	• Alkene	1650	medium
	• Arene	1600, 1500	medium
C = N		1650	medium

APPENDIX




F Characteristic NMR Absorptions

^{1}H NMR Absorptions

Compound type	Chemical shift (ppm)
Alcohol	
	1–5
	3.4–4.0
Aldehyde	
	9–10
Alkane	0.9–2.0
RCH_3	~0.9
R_2CH_2	~1.3
R_3CH	~1.7
Alkene	
 sp^2 C–H	4.5–6.0
 allylic sp^3 C–H	1.5–2.5
Alkyl halide	
	4.0–4.5
	3.0–4.0
	2.7–4.0
	2.2–4.0
Alkyne	
	~2.5

Compound type	Chemical shift (ppm)
Amide 	7.5–8.5
Amine 	0.5–5.0
	2.3–3.0
Aromatic compound  sp^2 C–H	6.5–8
 benzylic sp^3 C–H	1.5–2.5
Carbonyl compound  sp^3 C–H on the α carbon	2.0–2.5
Carboxylic acid 	10–12
Ether 	3.4–4.0

^{13}C NMR Absorptions

Carbon type	Structure	Chemical shift (ppm)
Alkyl, sp^3 hybridized C		5–45
Alkyl, sp^3 hybridized C bonded to N, O, or X	 Z = N, O, X	30–80
Alkynyl, *sp* hybridized C		65–100
Alkenyl, sp^2 hybridized C		100–140
Aryl, sp^2 hybridized C		120–150
Carbonyl C		160–210


APPENDIX

G General Types of Organic Reactions

Substitution Reactions

[1] Nucleophilic substitution at an sp^3 hybridized carbon atom


a. Alkyl halides (Chapter 7)




b. Alcohols (Section 9.11)



c. Ethers (Section 9.14)




d. Epoxides (Section 9.16)




[2] Nucleophilic acyl substitution at an sp^2 hybridized carbon atom

Carboxylic acids and their derivatives (Chapter 22)




[3] Radical substitution at an sp^3 hybridized C – H bond

Alkanes (Section 15.3)




[4] Electrophilic aromatic substitution

Aromatic compounds (Chapter 18)



[5] Nucleophilic aromatic substitution


Aromatic compounds (Chapter 18)




Elimination Reactions

β Elimination at an sp^3 hybridized carbon atom

a. Alkyl halides (Chapter 8)




b. Alcohols (Section 9.8)




Addition Reactions

[1] Electrophilic addition to carbon–carbon multiple bonds

a. Alkenes (Chapter 10)




b. Alkynes (Section 11.6)



[2] Nucleophilic addition to carbon–oxygen multiple bonds

Aldehydes and ketones (Chapter 21)



R' = H or alkyl

APPENDIX

H How to Synthesize Particular Functional Groups

Acetals

- Reaction of an aldehyde or ketone with two equivalents of an alcohol (21.14)

Acid chlorides

- Reaction of a carboxylic acid with thionyl chloride (22.10)

Alcohols

- Nucleophilic substitution of an alkyl halide with ^{-}OH or H_2O (9.6)
- Hydration of an alkene (10.12)
- Hydroboration–oxidation of an alkene (10.16)
- Reduction of an epoxide with $LiAlH_4$ (12.6)
- Reduction of an aldehyde or ketone (20.4)
- Hydrogenation of an α,β-unsaturated carbonyl compound with H_2 + Pd-C (20.4C)
- Enantioselective reduction of an aldehyde or ketone with the chiral CBS reagent (20.6)
- Reduction of an acid chloride with $LiAlH_4$ (20.7)
- Reduction of an ester with $LiAlH_4$ (20.7)
- Reduction of a carboxylic acid with $LiAlH_4$ (20.7)
- Reaction of an aldehyde or ketone with a Grignard or organolithium reagent (20.10)
- Reaction of an acid chloride with a Grignard or organolithium reagent (20.13)
- Reaction of an ester with a Grignard or organolithium reagent (20.13)
- Reaction of an organometallic reagent with an epoxide (20.14B)

Aldehydes

- Hydroboration–oxidation of a terminal alkyne (11.10)
- Oxidative cleavage of an alkene with O_3 followed by Zn or $(CH_3)_2S$ (12.10)
- Oxidation of a 1° alcohol with PCC (12.12)
- Oxidation of a 1° alcohol with $HCrO_4^{-}$, Amberlyst A-26 resin (12.13)
- Reduction of an acid chloride with $LiAlH[OC(CH_3)_3]_3$ (20.7)
- Reduction of an ester with DIBAL-H (20.7)
- Hydrolysis of an acetal (21.14B)
- Hydrolysis of an imine or enamine (21.12B)
- Reduction of a nitrile (22.18B)

Alkanes

- Catalytic hydrogenation of an alkene with H_2 + Pd-C (12.3)
- Catalytic hydrogenation of an alkyne with two equivalents of H_2 + Pd-C (12.5A)
- Reduction of an alkyl halide with $LiAlH_4$ (12.6)

- Reduction of a ketone to a methylene group (CH_2)—the Wolff–Kishner or Clemmensen reaction (18.15B)
- Protonation of an organometallic reagent with H_2O, ROH, or acid (20.9)
- Coupling of an organocuprate reagent (R_2CuLi) with an alkyl halide, R'X (26.1)
- Simmons–Smith reaction of an alkene with CH_2I_2 and Zn(Cu) to form a cyclopropane (26.5)

Alkenes

- Dehydrohalogenation of an alkyl halide with base (8.3)
- Dehydration of an alcohol with acid (9.8)
- Dehydration of an alcohol using $POCl_3$ and pyridine (9.10)
- β Elimination of an alkyl tosylate with base (9.13)
- Catalytic hydrogenation of an alkyne with H_2 + Lindlar catalyst to form a cis alkene (12.5B)
- Dissolving metal reduction of an alkyne with Na, NH_3 to form a trans alkene (12.5C)
- Wittig reaction (21.10)
- β Elimination of an α-halo carbonyl compound with Li_2CO_3, LiBr, and DMF (23.7C)
- Hofmann elimination of an amine (25.12)
- Coupling of an organocuprate reagent (R_2CuLi) with an organic halide, R'X (26.1)
- The palladium-catalyzed Suzuki reaction of a vinyl or aryl halide with a vinyl- or arylborane (26.2)
- The palladium-catalyzed Heck reaction of a vinyl or aryl halide with an alkene (26.3)
- Olefin metathesis (26.6)

Alkyl halides

- Reaction of an alcohol with HX (9.11)
- Reaction of an alcohol with $SOCl_2$ or PBr_3 (9.12)
- Cleavage of an ether with HBr or HI (9.14)
- Hydrohalogenation of an alkene with HX (10.9)
- Halogenation of an alkene with X_2 (10.13)
- Hydrohalogenation of an alkyne with two equivalents of HX (11.7)
- Halogenation of an alkyne with two equivalents of X_2 (11.8)
- Radical halogenation of an alkane (15.3)
- Radical halogenation at an allylic carbon (15.10)
- Radical addition of HBr to an alkene (15.13)
- Electrophilic addition of HX to a 1,3-diene (16.10)
- Radical halogenation of an alkyl benzene (18.14)
- Halogenation α to a carbonyl group (23.7)
- Addition of a dihalocarbene to an alkene to form a dihalocyclopropane (26.4)

Alkynes

- Dehydrohalogenation of an alkyl dihalide with base (11.5)
- S_N2 reaction of an alkyl halide with an acetylide anion, $^-C{\equiv}CR$ (11.11)

Amides

- Reaction of an acid chloride with NH_3 or an amine (22.8)
- Reaction of an anhydride with NH_3 or an amine (22.9)
- Reaction of a carboxylic acid with NH_3 or an amine and DCC (22.10)
- Reaction of an ester with NH_3 or an amine (22.11)

Amines

- Nucleophilic aromatic substitution (18.13)
- Reduction of a nitro group (18.15C)
- Reduction of an amide with $LiAlH_4$ (20.7B)
- Reduction of a nitrile (22.18B)
- S_N2 reaction using NH_3 or an amine (25.7A)
- Gabriel synthesis (25.7A)
- Reductive amination of an aldehyde or ketone (25.7C)

Amino acids

- S_N2 reaction of an α-halo carboxylic acid with excess NH_3 (29.2A)
- Alkylation of diethyl acetamidomalonate (29.2B)
- Strecker synthesis (29.2C)
- Enantioselective hydrogenation using a chiral catalyst (29.4)

Anhydrides

- Reaction of an acid chloride with a carboxylate anion (22.8)
- Dehydration of a dicarboxylic acid (22.10)

Aryl halides

- Halogenation of benzene with $X_2 + FeX_3$ (18.3)
- Reaction of a diazonium salt with CuCl, CuBr, HBF_4, NaI, or KI (25.14A)

Carboxylic acids

- Oxidative cleavage of an alkyne with ozone (12.11)
- Oxidation of a 1° alcohol with CrO_3 (or a similar Cr^{6+} reagent), H_2O, H_2SO_4 (12.12B)
- Oxidation of an alkyl benzene with $KMnO_4$ (18.15A)
- Oxidation of an aldehyde (20.8)
- Reaction of a Grignard reagent with CO_2 (20.14A)
- Hydrolysis of a cyanohydrin (21.9)
- Hydrolysis of an acid chloride (22.8)
- Hydrolysis of an anhydride (22.9)
- Hydrolysis of an ester (22.11)
- Hydrolysis of an amide (22.13)
- Hydrolysis of a nitrile (22.18A)
- Malonic ester synthesis (23.9)

Cyanohydrins

- Addition of HCN to an aldehyde or ketone (21.9)

1,2-Diols

- Anti dihydroxylation of an alkene with a peroxyacid, followed by ring opening with ^-OH or H_2O (12.9A)
- Syn dihydroxylation of an alkene with $KMnO_4$ or OsO_4 (12.9B)

Enamines

- Reaction of an aldehyde or ketone with a 2° amine (21.12)

Epoxides

- Intramolecular S_N2 reaction of a halohydrin using base (9.6)
- Epoxidation of an alkene with mCPBA (12.8)
- Enantioselective epoxidation of an allylic alcohol with the Sharpless reagent (12.15)

Esters

- S_N2 reaction of an alkyl halide with a carboxylate anion, RCO_2^- (7.18)
- Reaction of an acid chloride with an alcohol (22.8)
- Reaction of an anhydride with an alcohol (22.9)
- Fischer esterification of a carboxylic acid with an alcohol (22.10)

Ethers

- Williamson ether synthesis—S_N2 reaction of an alkyl halide with an alkoxide, ^-OR (9.6)
- Reaction of an alkyl tosylate with an alkoxide, ^-OR (9.13)
- Addition of an alcohol to an alkene in the presence of acid (10.12)
- Anionic polymerization of epoxides to form polyethers (30.3)

Halohydrins

- Reaction of an epoxide with HX (9.16)
- Addition of X and OH to an alkene (10.15)

Imine

- Reaction of an aldehyde or ketone with a 1° amine (21.11)

Ketones

- Hydration of an alkyne with H_2O, H_2SO_4, and $HgSO_4$ (11.9)
- Oxidative cleavage of an alkene with O_3 followed by Zn or $(CH_3)_2S$ (12.10)
- Oxidation of a 2° alcohol with any Cr^{6+} reagent (12.12, 12.13)
- Friedel–Crafts acylation (18.5)
- Reaction of an acid chloride with an organocuprate reagent (20.13)
- Hydrolysis of an imine or enamine (21.12B)
- Hydrolysis of an acetal (21.14B)
- Reaction of a nitrile with a Grignard or organolithium reagent (22.18C)
- Acetoacetic ester synthesis (23.10)

Nitriles

- S_N2 reaction of an alkyl halide with NaCN (7.18, 22.18)
- Reaction of an aryl diazonium salt with CuCN (25.14A)

Phenols

- Reaction of an aryl diazonium salt with H_2O (25.14A)
- Nucleophilic aromatic substitution (18.13)

Sulfides

- Reaction of an alkyl halide with ^-SR (9.15)

Thiols


- Reaction of an alkyl halide with ^-SH (9.15)

This page intentionally left blank

MISSING

MISSING

Axial bonds (Section 4.12A): Bonds located above or below and perpendicular to the plane of the chair conformation of cyclohexane. Three axial bonds point upwards (on the up carbons) and three axial bonds point downwards (on the down carbons).



Azo compound (Section 25.15): A compound having the general structure RN=NR'.

B

Backside attack (Section 7.11C): Approach of a nucleophile from the side opposite the leaving group.

Barrier to rotation (Section 4.10): The energy difference between the lowest and highest energy conformations of a molecule.

Base peak (Section 13.1): The peak in the mass spectrum having the greatest abundance value.

Basicity (Section 7.8): A measure of how readily an atom donates its electron pair to a proton.

Benedict's reagent (Section 28.9B): A reagent for oxidizing aldehydes to carboxylic acids using a Cu^{2+} salt, forming brick-red Cu_2O as a side product.

Benzoyl group (Section 21.2E): A substituent having the structure $-COC_6H_5$.

Benzyl group (Section 17.3D): A substituent having the structure $C_6H_5CH_2-$.

Benzylic halide (Sections 7.1, 18.14): A compound such as $C_6H_5CH_2X$, containing a halogen atom X bonded to a carbon that is bonded to a benzene ring.

Benzyne (Section 18.13B): A reactive intermediate formed by elimination of HX from an aryl halide.

Beta (β) carbon (Sections 8.1, 19.2B): In an elimination reaction, the carbon adjacent to the carbon with the leaving group. In a carbonyl compound, the carbon located two carbons from the carbonyl carbon.

Bimolecular reaction (Sections 6.9B, 7.10, 7.11A): A reaction in which the concentration of both reactants affects the reaction rate and both terms appear in the rate equation. In a bimolecular reaction, two reactants are involved in the only step or the rate-determining step.

Biodegradable polymer (Section 30.9B): A polymer that can be degraded by microorganisms naturally present in the environment.

Biomolecule (Section 3.9): An organic compound found in a biological system.

Boat conformation of cyclohexane (Section 4.12B): An unstable conformation adopted by cyclohexane that resembles a boat. The instability of the boat conformation results from torsional strain and steric strain. The boat conformation of cyclohexane is 30 kJ/mol less stable than the chair conformation.



Boiling point (Section 3.4A): The temperature at which molecules in the liquid phase are converted to the gas phase. Molecules with stronger intermolecular forces have higher boiling points. Boiling point is abbreviated as bp.

Bond dissociation energy (Section 6.4): The amount of energy needed to homolytically cleave a covalent bond.

Bonding (Section 1.2): The joining of two atoms in a stable arrangement. Bonding is a favorable process that leads to lowered energy and increased stability.

Bonding molecular orbital (Section 17.9A): A low-energy molecular orbital formed when two atomic orbitals of similar phase overlap.

Bond length (Section 1.7A): The average distance between the centers of two bonded nuclei. Bond lengths are reported in picometers (pm).

Branched-chain alkane (Section 4.1A): An acyclic alkane that has alkyl substituents bonded to the parent carbon chain.

Bridged ring system (Section 16.13D): A bicyclic ring system in which the two rings share non-adjacent carbon atoms.

Bromination (Sections 10.13, 15.6, 18.3): The reaction of a compound with bromine.

Bromohydrin (Section 10.15): A compound having a bromine and a hydroxy group on adjacent carbon atoms.

Brønsted–Lowry acid (Section 2.1): A proton donor, symbolized by HA. A Brønsted–Lowry acid must contain a hydrogen atom.

Brønsted–Lowry base (Section 2.1): A proton acceptor, symbolized by :B. A Brønsted–Lowry base must be able to form a bond to a proton by donating an available electron pair.

C

^{13}C NMR spectroscopy (Section 14.1): A form of nuclear magnetic resonance spectroscopy used to determine the type of carbon atoms in a molecule.

Cahn–Ingold–Prelog system of nomenclature (Section 5.6): The system of designating a stereogenic center as either *R* or *S* according to the arrangement of the four groups attached to the center.

Carbamate (Sections 29.7, 30.6): A functional group containing a carbonyl group bonded to both an oxygen and a nitrogen atom. A carbamate is also called a urethane.

Carbanion (Section 2.5D): An ion with a negative charge on a carbon atom.

Carbene (Section 26.4): A neutral reactive intermediate having the general structure $:CR_2$. A carbene contains a divalent carbon surrounded by six electrons, making it a highly reactive electrophile that adds to C−C double bonds.

Carbinolamine (Section 21.7B): An unstable intermediate having a hydroxy group and an amine group on the same carbon. A carbinolamine is formed during the addition of an amine to a carbonyl group.

Carbocation (Section 7.13C): A positively charged carbon atom. A carbocation is sp^2 hybridized and trigonal planar, and contains a vacant *p* orbital.

Carbohydrate (Sections 21.17, 28.1): A polyhydroxy aldehyde or ketone or a compound that can be hydrolyzed to a polyhydroxy aldehyde or ketone.

Carbonate (Section 30.6D): A compound having the general structure $(RO)_2C=O$.

Carbon backbone (Section 3.1): The C−C and C−H σ bond framework that makes up the skeleton of an organic molecule.

Carbon NMR spectroscopy (Section 14.1): A form of nuclear magnetic resonance spectroscopy used to determine the type of carbon atoms in a molecule.

Carbonyl group (Sections 3.2C, 11.9, 20.1): A functional group that contains a carbon–oxygen double bond (C=O). The polar carbon–oxygen bond makes the carbonyl carbon electrophilic.

Carboxy group (Section 19.1): A functional group having the structure COOH.

Carboxylate anion (Section 19.2C): An anion having the general structure RCO_2^-, formed by deprotonating a carboxylic acid with a Brønsted–Lowry base.

Carboxylation (Section 20.14): The reaction of an organometallic reagent with CO_2 to form a carboxylic acid after protonation.

Carboxylic acid (Section 19.1): A compound having the general structure RCO_2H.

Carboxylic acid derivatives (Section 20.1): Compounds having the general structure RCOZ, which can be synthesized from carboxylic acids. Common carboxylic acid derivatives include acid chlorides, anhydrides, esters, and amides.

Catalyst (Section 6.10): A substance that speeds up the rate of a reaction, but is recovered unchanged at the end of the reaction and does not appear in the product.

Catalytic hydrogenation (Section 12.3): A reduction reaction involving the addition of H_2 to a π bond in the presence of a metal catalyst.

Cation (Section 1.2): A positively charged ion that results from a neutral atom losing one or more electrons.

Cationic polymerization (Section 30.2C): Chain-growth polymerization of alkene monomers involving carbocation intermediates.

CBS reagent (Section 20.6A): A chiral reducing agent formed by reacting an oxazaborolidine with BH_3. CBS reagents predictably give one enantiomer as the major product of ketone reduction.

Cephalin (Section 31.4A): A phosphoacylglycerol in which the phosphodiester alkyl group is $-CH_2CH_2NH_3^+$. Cephalins are also called phosphatidylethanolamines.

Chain-growth polymer (Section 30.1): A polymer prepared by a chain reaction that adds a monomer to the growing end of a polymer chain. Chain-growth polymers are also called addition polymers.

Chain mechanism (Section 15.4A): A reaction mechanism that involves repeating steps.

Chair conformation of cyclohexane (Section 4.12A): A stable conformation adopted by cyclohexane that resembles a chair. The stability of the chair conformation results from the elimination of angle strain (all C−C−C bond angles are 109.5°) and torsional strain (all groups on adjacent carbon atoms are staggered).



Chemical shift (Section 14.1B): The position of an absorption signal on the *x* axis in an NMR spectrum relative to the reference signal of tetramethylsilane.

Chirality center (Section 5.3): A carbon atom bonded to four different groups. A chirality center is also called a chiral center, a stereogenic center, and an asymmetric center.

Chiral molecule (Section 5.3): A molecule that is not superimposable upon its mirror image.

Chlorination (Sections 10.14, 15.5, 18.3): The reaction of a compound with chlorine.

Chlorofluorocarbons (Sections 7.4, 15.9): Synthetic alkyl halides having the general molecular formula CF_xCl_{4-x}. Chlorofluorocarbons, abbreviated as CFCs, were used as refrigerants and aerosol propellants and contribute to the destruction of the ozone layer.

Chlorohydrin (Section 10.15): A compound having a chlorine and a hydroxy group on adjacent carbon atoms.

Chromate ester (Section 12.12A): An intermediate in the chromium-mediated oxidation of an alcohol having the general structure $R-O-CrO_3H$.

***s*-Cis** (Sections 16.6, 29.5B): The conformation of a 1,3-diene that has the two double bonds on the same side of the single bond that joins them.

Cis isomer (Sections 4.13B, 8.2B): An isomer of a ring or double bond that has two groups on the same side of the ring or double bond.

Claisen reaction (Section 24.5): A reaction between two molecules of an ester in the presence of base to form a β-keto ester.

Claisen rearrangement (Section 27.5): A [3,3] sigmatropic rearrangement of an unsaturated ether to a γ,δ-unsaturated carbonyl compound.

α Cleavage (Section 13.3B): A fragmentation in mass spectrometry that results in cleavage of a carbon–carbon bond. With aldehydes and ketones, α cleavage results in breaking the bond between the carbonyl carbon and the carbon adjacent to it. With alcohols, α cleavage occurs by breaking a bond between an alkyl group and the carbon that bears the OH group.

Clemmensen reduction (Section 18.15B): A method to reduce aryl ketones to alkyl benzenes using Zn(Hg) in the presence of a strong acid.

Coenzyme (Section 12.13): A compound that acts with an enzyme to carry out a biochemical process.

Combustion (Section 4.14B): An oxidation–reduction reaction, in which an alkane or other organic compound reacts with oxygen to form CO_2 and H_2O, releasing energy.

Common name (Section 4.6): The name of a molecule that was adopted prior to and therefore does not follow the IUPAC system of nomenclature.

Compound (Section 1.2): The structure that results when two or more elements are joined together in a stable arrangement.

Concerted reaction (Sections 6.3, 7.11B): A reaction in which all bond forming and bond breaking occurs in one step.

Condensation polymer (Sections 22.16A, 30.1): A polymer formed when monomers containing two functional groups come together with loss of a small molecule such as water or HCl. Condensation polymers are also called step-growth polymers.

Condensation reaction (Section 24.1B): A reaction in which a small molecule, often water, is eliminated during the reaction process.

Condensed structure (Section 1.8A): A shorthand representation of the structure of a compound in which all atoms are drawn in but bonds and lone pairs are usually omitted. Parentheses are used to denote similar groups bonded to the same atom.

Configuration (Section 5.2): A particular three-dimensional arrangement of atoms.

Conformations (Section 4.9): The different arrangements of atoms that are interconverted by rotation about single bonds.

Conjugate acid (Section 2.2): The compound that results when a base gains a proton in a proton transfer reaction.

Conjugate addition (Sections 16.10, 20.15): An addition reaction that adds groups to the atoms in the 1 and 4 positions of a conjugated system. Conjugate addition is also called 1,4-addition.

Conjugate base (Section 2.2): The compound that results when an acid loses a proton in a proton transfer reaction.

Conjugated diene (Section 16.1A): A compound that contains two carbon–carbon double bonds joined by a single σ bond. Pi (π) electrons are delocalized over both double bonds. Conjugated dienes are also called 1,3-dienes.

Conjugated protein (Section 29.10C): A structure composed of a protein unit and a non-protein molecule.

Conjugation (Section 16.1): The overlap of *p* orbitals on three or more adjacent atoms.

Conrotatory rotation (Section 27.3): Rotation of *p* orbitals in the same direction during electrocyclic ring closure or ring opening.

Constitutional isomers (Sections 1.4, 4.1A, 5.2): Two compounds that have the same molecular formula, but differ in the way the atoms are connected to each other. Constitutional isomers are also called structural isomers.

Coordination polymerization (Section 30.4): A polymerization reaction that uses a homogeneous catalyst that is soluble in the reaction solvents typically used.

Cope rearrangement (Section 27.5): A [3,3] sigmatropic rearrangement of a 1,5-diene to an isomeric 1,5-diene.

Copolymer (Section 30.2D): A polymer prepared by joining two or more different monomers together.

Counterion (Section 2.1): An ion that does not take part in a reaction and is opposite in charge to the ion that does take part in the reaction. A counterion is also called a spectator ion.

Coupling constant (Section 14.6A): The frequency difference, measured in Hz, between the peaks in a split NMR signal.

Coupling reaction (Section 25.15): A reaction that forms a bond between two discrete molecules.

Covalent bond (Section 1.2): A bond that results from the sharing of electrons between two nuclei. A covalent bond is a two-electron bond.

Crossed aldol reaction (Section 24.2): An aldol reaction in which the two reacting carbonyl compounds are different. A crossed aldol reaction is also called a mixed aldol reaction.

Crossed Claisen reaction (Section 24.6): A Claisen reaction in which the two reacting esters are different.

Crown ether (Section 3.7B): A cyclic ether containing multiple oxygen atoms. Crown ethers bind specific cations depending on the size of their central cavity.

Curved arrow notation (Section 1.6A): A convention that shows the movement of an electron pair. The tail of the arrow begins at the electron pair and the head points to where the electron pair moves.

Cyanide anion (Section 21.9A): An anion having the structure $^{-}C{\equiv}N$.

Cyano group (Section 22.1): A functional group consisting of a carbon–nitrogen triple bond (C≡N).

Cyanohydrin (Section 21.9): A compound having the general structure RCH(OH)C≡N. A cyanohydrin results from the addition of HCN across the carbonyl of an aldehyde or a ketone.

Cycloaddition (Section 27.1): A pericyclic reaction between two compounds with π bonds to form a cyclic product with two new σ bonds.


Cycloalkane (Sections 4.1, 4.2): A compound that contains carbons joined in one or more rings. Cycloalkanes with one ring have the general formula C_nH_{2n}.

Cyclopropanation (Section 26.4): An addition reaction to a carbon–carbon double bond that forms a cyclopropane.

D

D-Sugar (Section 28.2C): A sugar with the hydroxy group on the stereogenic center farthest from the carbonyl on the right side in the Fischer projection formula.

Decalin (Section 31.8A): Two fused six-membered rings. *cis*-Decalin has the hydrogen atoms at the ring fusion on the same side of the rings, whereas *trans*-decalin has the hydrogen atoms at the ring fusion on opposite sides of the rings.



Decarboxylation (Section 23.9A): Loss of CO_2 through cleavage of a carbon–carbon bond.

Degenerate orbitals (Section 17.9B): Orbitals (either atomic or molecular) having the same energy.

Degree of unsaturation (Section 10.2): A ring or a π bond in a molecule. The number of degrees of unsaturation compares the number of hydrogens in a compound to that of a saturated hydrocarbon containing the same number of carbons.

Dehydration (Sections 9.8, 22.10B): A reaction that results in the loss of the elements of water from the reaction components.

Dehydrohalogenation (Section 8.1): An elimination reaction in which the elements of hydrogen and halogen are lost from a starting material.

Delta (δ) scale (Section 14.1B): A common scale of chemical shifts used in NMR spectroscopy in which the absorption due to tetramethylsilane (TMS) occurs at zero parts per million.

Deoxy (Section 28.13B): A prefix that means without oxygen.

Deoxyribonucleoside (Section 28.13B): An *N*-glycoside formed by the reaction of D-2-deoxyribose with certain amine heterocycles.

Deoxyribonucleotide (Section 28.13B): A DNA building block having a deoxyribose and either a purine or pyrimidine base joined together by an *N*-glycosidic linkage, and a phosphate bonded to a hydroxy group of the sugar nucleus.

Deprotection (Section 20.12): A reaction that removes a protecting group, regenerating a functional group.

Deshielding effects (Section 14.3A): An effect in NMR caused by a decrease in electron density, thus increasing the strength of the magnetic field felt by the nucleus. Deshielding shifts an absorption downfield.

Dextrorotatory (Section 5.12A): Rotating plane-polarized light in the clockwise direction. The rotation is labeled *d* or (+).

1,3-Diacid (Section 23.9A): A compound containing two carboxylic acids separated by a single carbon atom. 1,3-Diacids are also called β-diacids.

Dialkylamide (Section 23.3B): An amide base having the general structure R_2N^{-}.

Diastereomers (Section 5.7): Stereoisomers that are not mirror images of each other. Diastereomers have the same *R,S* designation for at least one stereogenic center and the opposite *R,S* designation for at least one of the other stereogenic centers.

Diastereotopic protons (Section 14.2C): Two hydrogen atoms on the same carbon such that substitution of either hydrogen with a group

Z forms diastereomers. The two hydrogen atoms are not equivalent and give two NMR signals.

1,3-Diaxial interaction (Section 4.13A): A steric interaction between two axial substituents of the chair form of cyclohexane. Larger axial substituents create unfavorable 1,3-diaxial interactions, destabilizing a cyclohexane conformation.

Diazonium salt (Section 25.13A): An ionic salt having the general structure $(R-N\equiv N)^+Cl^-$.

Diazotization reaction (Section 25.13A): A reaction that converts 1° alkylamines and arylamines to diazonium salts.

1,3-Dicarbonyl compound (Section 23.2): A compound containing two carbonyl groups separated by a single carbon atom.

1,4-Dicarbonyl compound (Section 24.4): A dicarbonyl compound in which the carbonyl groups are separated by three single bonds. 1,4-Dicarbonyl compounds can undergo intramolecular reactions to form five-membered rings.

1,5-Dicarbonyl compound (Section 24.4): A dicarbonyl compound in which the carbonyl groups are separated by four single bonds. 1,5-Dicarbonyl compounds can undergo intramolecular reactions to form six-membered rings.

Dieckmann reaction (Section 24.7): An intramolecular Claisen reaction of a diester to form a ring, typically a five- or six-membered ring.

Diels–Alder reaction (Section 16.12): An addition reaction between a 1,3-diene and a dienophile to form a cyclohexene ring.

1,3-Diene (Section 16.1A): A compound containing two carbon–carbon double bonds joined by a single σ bond. Pi (π) electrons are delocalized over both double bonds. 1,3-Dienes are also called conjugated dienes.

Dienophile (Section 16.12): The alkene component in a Diels–Alder reaction that reacts with a 1,3-diene.

Dihedral angle (Section 4.9): The angle that separates a bond on one atom from a bond on an adjacent atom.

Dihydroxylation (Section 12.9): Addition of two hydroxy groups to a double bond to form a 1,2-diol.

Diol (Section 9.3A): A compound possessing two hydroxy groups. Diols are also called glycols.

Dipeptide (Section 29.5): Two amino acids joined together by one amide bond.

Diphosphate (Section 7.16): A good leaving group that is often used in biological systems. Diphosphate ($P_2O_7^{4-}$) is abbreviated as PP_i.

Dipole (Section 1.12): A partial separation of electronic charge.

Dipole–dipole interaction (Section 3.3B): An attractive intermolecular interaction between the permanent dipoles of polar molecules. The dipoles of adjacent molecules align so that the partial positive and partial negative charges are in close proximity.

Directed aldol reaction (Section 24.3): A crossed aldol reaction in which the enolate of one carbonyl compound is formed, followed by addition of the second carbonyl compound.

Disaccharide (Section 28.11): A carbohydrate containing two monosaccharide units joined together by a glycosidic linkage.

Disproportionation (Section 30.2): A method of chain termination in radical polymerization involving the transfer of a hydrogen atom from one polymer radical to another, forming a new C−H bond on one polymer chain and a new double bond on the other.

Disrotatory rotation (Section 27.3): Rotation of *p* orbitals in opposite directions during electrocyclic ring closure or ring opening.

Dissolving metal reduction (Section 12.2): A reduction reaction using alkali metals as a source of electrons and liquid ammonia as a source of protons.

Disubstituted alkene (Section 8.2A): An alkene that has two alkyl groups and two hydrogens bonded to the carbons of the double bond ($R_2C=CH_2$ or $RCH=CHR$).

Disulfide (Sections 9.15A, 29.5C): A compound having the general structure RSSR', often formed between the side chain of two cysteine residues.

Diterpene (Section 31.7A): A terpene that contains 20 carbons and four isoprene units. A diterpenoid contains at least one oxygen atom as well.

Doublet (Section 14.6): An NMR signal that is split into two peaks of equal area, caused by one nearby nonequivalent proton.

Doublet of doublets (Section 14.8): A splitting pattern of four peaks observed when a signal is split by two different nonequivalent protons.

Downfield shift (Section 14.1B): In an NMR spectrum, a term used to describe the relative location of an absorption signal. A downfield shift means the signal is shifted to the left in the spectrum to higher chemical shift on the δ scale.

E

E1 mechanism (Sections 8.3, 8.6): An elimination mechanism that goes by a two-step process involving a carbocation intermediate. E1 is an abbreviation for "Elimination Unimolecular."

E1cB mechanism (Section 24.1B): A two-step elimination mechanism that goes by a carbanion intermediate. E1cB stands for "Elimination Unimolecular, Conjugate Base."

E2 mechanism (Sections 8.3, 8.4): An elimination mechanism that goes by a one-step concerted process, in which both reactants are involved in the transition state. E2 is an abbreviation for "Elimination Bimolecular."

Eclipsed conformation (Section 4.9): A conformation of a molecule where the bonds on one carbon are directly aligned with the bonds on the adjacent carbon.



Edman degradation (Section 29.6B): A procedure used in peptide sequencing in which amino acids are cleaved one at a time from the N-terminal end, the identity of the amino acid determined, and the process repeated until the entire sequence is known.

Eicosanoids (Section 31.6): A group of biologically active compounds containing 20 carbon atoms derived from arachidonic acid.

Elastomer (Section 30.5): A polymer that stretches when stressed but then returns to its original shape.

Electrocyclic ring closure (Section 27.1): An intramolecular pericyclic reaction that forms a cyclic product containing one more σ bond and one fewer π bond than the reactant.

Electrocyclic ring-opening reaction (Section 27.1): A pericyclic reaction in which a σ bond of a cyclic reactant is cleaved to form a conjugated product with one more π bond.

Electromagnetic radiation (Section 13.5): Radiant energy having dual properties of both waves and particles. The electromagnetic spectrum contains the complete range of electromagnetic radiation, arbitrarily divided into different regions.

Electron-donating inductive effect (Section 7.13A): An inductive effect in which an electropositive atom or polarizable group donates electron density through σ bonds to another atom.

Electronegativity (Section 1.12): A measure of an atom's attraction for electrons in a bond. Electronegativity indicates how much a particular atom "wants" electrons.

Electron-withdrawing inductive effect (Sections 2.5, 7.13A): An inductive effect in which a nearby electronegative atom pulls electron density towards itself through σ bonds.

Electrophile (Section 2.8): An electron-deficient compound, often symbolized by E^+, which can accept a pair of electrons from an electron-rich compound, forming a covalent bond. Lewis acids are electrophiles.

Electrophilic addition reaction (Section 10.9): An addition reaction in which the first step of the mechanism involves addition of the electrophilic end of a reagent to a π bond.

Electrophilic aromatic substitution (Section 18.1): A characteristic reaction of benzene in which a hydrogen atom on the ring is replaced by an electrophile.

Electrospray ionization (Section 13.4C): A method for ionizing large biomolecules in a mass spectrometer. Electrospray ionization is abbreviated as ESI.

Electrostatic potential map (Section 1.12): A color-coded map that illustrates the distribution of electron density in a molecule. Electron-rich regions are indicated in red and electron-deficient regions are indicated in blue. Regions of intermediate electron density are shown in orange, yellow, and green.

α Elimination (Section 26.4): An elimination reaction involving the loss of two elements from the same atom.

β Elimination (Section 8.1): An elimination reaction involving the loss of elements from two adjacent atoms.

Elimination reaction (Sections 6.2B, 8.1): A chemical reaction in which elements of the starting material are "lost" and a π bond is formed.

Enamine (Section 21.12): A compound having an amine nitrogen atom bonded to a carbon–carbon double bond [$R_2C{=}CH(NR'_2)$].

Enantiomeric excess (Section 5.12D): A measurement of how much one enantiomer is present in excess of the racemic mixture. Enantiomeric excess (*ee*) is also called optical purity; *ee* = % of one enantiomer – % of the other enantiomer.

Enantiomers (Section 5.3): Stereoisomers that are mirror images but are not superimposable upon each other. Enantiomers have the exact opposite *R,S* designation at every stereogenic center.

Enantioselective reaction (Sections 12.15, 20.6A, 29.4): A reaction that affords predominantly or exclusively one enantiomer. Enantioselective reactions are also called asymmetric reactions.

Enantiotopic protons (Section 14.2C): Two hydrogen atoms on the same carbon such that substitution of either hydrogen with a group Z forms enantiomers. The two hydrogen atoms are equivalent and give a single NMR signal.

Endo position (Section 16.13D): A position of a substituent on a bridged bicyclic compound in which the substituent is closer to the longer bridge that joins the two carbons common to both rings.

Endothermic reaction (Section 6.4): A reaction in which the energy of the products is higher than the energy of the reactants. In an endothermic reaction, energy is absorbed and the $\Delta H°$ is a positive value.

Energy diagram (Section 6.7): A schematic representation of the energy changes that take place as reactants are converted to products. An energy diagram indicates how readily a reaction proceeds, how many steps are involved, and how the energies of the reactants, products, and intermediates compare.

Energy of activation (Section 6.7): The energy difference between the transition state and the starting material. The energy of activation, symbolized by E_a, is the minimum amount of energy needed to break bonds in the reactants.

Enolate (Sections 20.15, 23.3): A resonance-stabilized anion formed when a base removes an α hydrogen from the α carbon to a carbonyl group.

Enol tautomer (Sections 9.1, 11.9, 20.15): A compound having a hydroxy group bonded to a carbon–carbon double bond. An enol tautomer [such as $CH_2{=}C(OH)CH_3$] is in equilibrium with its keto tautomer [$(CH_3)_2C{=}O$].

Enthalpy change (Section 6.4): The energy absorbed or released in a reaction. Enthalpy change is symbolized by $\Delta H°$ and is also called the heat of reaction.

Entropy (Section 6.6): A measure of the randomness in a system. The more freedom of motion or the more disorder present, the higher the entropy. Entropy is denoted by the symbol $S°$.

Entropy change (Section 6.6): The change in the amount of disorder between reactants and products in a reaction. The entropy change is denoted by the symbol $\Delta S°$. $\Delta S° = S°_{products} - S°_{reactants}$.

Enzyme (Section 6.11): A biochemical catalyst composed of at least one chain of amino acids held together in a very specific three-dimensional shape.

Enzyme–substrate complex (Section 6.11): A structure having a substrate bonded to the active site of an enzyme.

Epoxidation (Section 12.8): Addition of a single oxygen atom to an alkene to form an epoxide.

Epoxide (Section 9.1): A cyclic ether having the oxygen atom as part of a three-membered ring. Epoxides are also called oxiranes.

Epoxy resin (Section 30.6E): A step-growth polymer formed from a fluid prepolymer and a hardener that cross-links polymer chains together.

Equatorial bonds (Section 4.12A): Bonds located in the plane of the chair conformation of cyclohexane (around the equator). Three equatorial bonds point slightly upward (on the down carbons) and three equatorial bonds point slightly downward (on the up carbons).



Equilibrium constant (Section 6.5A): A mathematical expression, denoted by the symbol K_{eq}, which relates the amount of starting material and product at equilibrium. K_{eq} = [products]/[starting materials].

Essential oil (Section 31.7): A class of terpenes isolated from plant sources by distillation.

Ester (Sections 20.1, 22.1): A compound having the general structure RCOOR'.

Esterification (Section 22.10C): A reaction that converts a carboxylic acid or a derivative of a carboxylic acid to an ester.

Ether (Section 9.1): A functional group having the general structure ROR'.

Ethynyl group (Section 11.2): An alkynyl substituent having the structure $-C{\equiv}C-H$.

Excited state (Sections 1.9B, 16.15A): A high-energy electronic state in which one or more electrons have been promoted to a higher energy orbital by absorption of energy.

Exo position (Section 16.13D): A position of a substituent on a bridged bicyclic compound in which the substituent is closer to the shorter bridge that joins the two carbons common to both rings.

Exothermic reaction (Section 6.4): A reaction in which the energy of the products is lower than the energy of the reactants. In an exothermic reaction, energy is released and the $\Delta H°$ is a negative value.

Extraction (Section 19.12): A laboratory method to separate and purify a mixture of compounds using solubility differences and acid–base principles.

***E,Z* System of nomenclature** (Section 10.3B): A system for unambiguously naming alkene stereoisomers by assigning priorities to the two groups on each carbon of the double bond. The *E* isomer has the two higher priority groups on opposite sides of the double bond, and the *Z* isomer has them on the same side.

F

Fat (Sections 10.6B, 31.3): A triacylglycerol that is solid at room temperature and composed of fatty acid side chains with a high degree of saturation.

Fatty acid (Sections 10.6A, 19.6): A long-chain carboxylic acid having between 12 and 20 carbon atoms.

Fehling's reagent (Section 28.9B): A reagent for oxidizing aldehydes to carboxylic acids using a Cu^{2+} salt as an oxidizing agent, forming brick-red Cu_2O as a by-product.


Fibrous proteins (Section 29.10): Long linear polypeptide chains that are bundled together to form rods or sheets.

Fingerprint region (Section 13.6B): The region in an IR spectrum at $< 1500\ cm^{-1}$. The region often contains a complex set of peaks and is unique for every compound.

First-order rate equation (Sections 6.9B, 7.10): A rate equation in which the reaction rate depends on the concentration of only one reactant.

Fischer esterification (Section 22.10C): An acid-catalyzed esterification reaction between a carboxylic acid and an alcohol to form an ester.

Fischer projection formula (Section 28.2A): A method for representing stereogenic centers with the stereogenic carbon at the intersection of vertical and horizontal lines. Fischer projections are also called cross formulas.



Fishhook (Section 6.3B): A half-headed curved arrow used in a reaction mechanism to denote the movement of a single electron.

Flagpole hydrogens (Section 4.12B): Hydrogens in the boat conformation of cyclohexane that are on either end of the "boat" and are forced into close proximity to each other.

Formal charge (Section 1.3C): The electronic charge assigned to individual atoms in a Lewis structure. The formal charge is calculated by subtracting an atom's unshared electrons and half of its shared electrons from the number of valence electrons that a neutral atom would possess.

Formyl group (Section 21.2E): A substituent having the structure $-CHO$.

Four-centered transition state (Section 10.16): A transition state that involves four atoms.

Fragment (Section 13.1): Radicals and cations formed by the decomposition of the molecular ion in a mass spectrometer.

Freons (Sections 7.4, 15.9): Chlorofluorocarbons consisting of simple halogen-containing organic compounds that were once commonly used as refrigerants.

Frequency (Section 13.5): The number of waves passing a point per unit time. Frequency is reported in cycles per second (s^{-1}), which is also called hertz (Hz). Frequency is abbreviated with the Greek letter nu (ν).

Friedel–Crafts acylation (Section 18.5A): An electrophilic aromatic substitution reaction in which benzene reacts with an acid chloride in the presence of a Lewis acid to give a ketone.

Friedel–Crafts alkylation (Section 18.5A): An electrophilic aromatic substitution reaction in which benzene reacts with an alkyl halide in the presence of a Lewis acid to give an alkyl benzene.

Frontside attack (Section 7.11C): Approach of a nucleophile from the same side as the leaving group.

Full-headed curved arrow (Section 6.3B): An arrow used in a reaction mechanism to denote the movement of a pair of electrons.

Functional group (Section 3.1): An atom or group of atoms with characteristic chemical and physical properties. The functional group is the reactive part of the molecule.

Functional group interconversion (Section 11.12): A reaction that converts one functional group into another.

Functional group region (Section 13.6B): The region in an IR spectrum at $\geq 1500\ cm^{-1}$. Common functional groups show one or two peaks in this region, at a characteristic frequency.

Furanose (Section 28.6): A cyclic five-membered ring of a monosaccharide containing an oxygen atom.


Fused ring system (Section 16.13D): A bicyclic ring system in which the two rings share one bond and two adjacent atoms.

G

Gabriel synthesis (Section 25.7A): A two-step method that converts an alkyl halide into a primary amine using a nucleophile derived from phthalimide.

Gas chromatography (Section 13.4B): An analytical technique that separates the components of a mixture based on their boiling points and the rate at which their vapors travel through a column.

Gauche conformation (Section 4.10): A staggered conformation in which the two larger groups on adjacent carbon atoms have a dihedral angle of 60°.



GC–MS (Section 13.4B): An analytical instrument that combines a gas chromatograph (GC) and a mass spectrometer (MS) in sequence.

***gem*-Diol** (Section 21.13): A compound having the general structure $R_2C(OH)_2$. *gem*-Diols are also called hydrates.

Geminal dihalide (Section 8.10): A compound that has two halogen atoms on the same carbon atom.

Gibbs free energy (Section 6.5A): The free energy of a molecule. Gibbs free energy is denoted by the symbol $G°$.

Gibbs free energy change (Section 6.5A): The overall energy difference between reactants and products. The Gibbs free energy change is denoted by the symbol $\Delta G°$. $\Delta G° = G°_{products} - G°_{reactants}$.

Globular proteins (Section 29.10): Polypeptide chains that are coiled into compact shapes with hydrophilic outer surfaces that make them water soluble.

Glycol (Section 9.3A): A compound possessing two hydroxy groups. Glycols are also called diols.

Glycosidase (Section 28.12B): An enzyme that hydrolyzes glycosidic linkages. An α-glycosidase hydrolyzes only α-glycosidic linkages.

Glycoside (Section 28.7A): A monosaccharide with an alkoxy group bonded to the anomeric carbon.

***N*-Glycoside** (Section 28.13B): A monosaccharide containing a nitrogen bonded to the anomeric carbon.

Glycosidic linkage (Section 28.11): An acetal linkage formed between an OH group on one monosaccharide and the anomeric carbon on a second monosaccharide.

Green chemistry (Sections 12.13, 30.8): The use of environmentally benign methods to synthesize compounds.

Grignard reagent (Section 20.9): An organometallic reagent having the general structure RMgX.

Ground state (Sections 1.9B, 16.15A): The lowest energy arrangement of electrons for an atom.

Group number (Section 1.1): The number above a particular column in the periodic table. Group numbers are represented by either an Arabic (1 to 8) or Roman (I to VIII) numeral followed by the letter A or B. The group number of a second-row element is equal to the number of valence electrons in that element.

Grubbs catalyst (Section 26.6): A widely used ruthenium catalyst for olefin metathesis that has the structure $Cl_2(Cy_3P)_2Ru{=}CHPh$.

Guest molecule (Section 9.5B): A small molecule that can bind to a larger host molecule.

H

^{1}H NMR spectroscopy (Section 14.1): A form of nuclear magnetic resonance spectroscopy used to determine the number and type of hydrogen atoms in a molecule. ^{1}H NMR is also called proton NMR spectroscopy.

Half-headed curved arrow (Section 6.3B): An arrow used in a reaction mechanism to denote the movement of a single electron. A half-headed curved arrow is also called a fishhook.

α-Halo aldehyde or ketone (Section 23.7): An aldehyde or ketone with a halogen atom bonded to the α carbon.

Haloform reaction (Section 23.7B): A halogenation reaction of a methyl ketone ($RCOCH_3$) with excess halogen, which results in formation of RCO_2^- and CHX_3 (haloform).

Halogenation (Sections 10.13, 15.3, 18.3): The reaction of a compound with a halogen.

Halohydrin (Sections 9.6, 10.15): A compound that has a hydroxy group and a halogen atom on adjacent carbon atoms.

Halonium ion (Section 10.13): A positively charged halogen atom. A bridged halonium ion contains a three-membered ring and is formed in the addition of a halogen (X_2) to an alkene.

Hammond postulate (Section 7.14): A postulate that states that the transition state of a reaction resembles the structure of the species (reactant or product) to which it is closer in energy.

Haworth projection (Section 28.6A): A representation of the cyclic form of a monosaccharide in which the ring is drawn flat.

Head-to-tail polymerization (Section 15.14B): A mechanism of radical polymerization in which the more substituted radical of the growing polymer chain always adds to the less substituted end of the new monomer.

Heat of hydrogenation (Section 12.3A): The $\Delta H°$ of a catalytic hydrogenation reaction equal to the amount of energy released by hydrogenating a π bond.

Heat of reaction (Section 6.4): The energy absorbed or released in a reaction. Heat of reaction is symbolized by $\Delta H°$ and is also called the change in enthalpy.

Heck reaction (Section 26.3): The palladium-catalyzed coupling of a vinyl or aryl halide with an alkene to form a more highly substituted alkene with a new carbon–carbon bond.

α-Helix (Section 29.9B): A secondary structure of a protein formed when a peptide chain twists into a right-handed or clockwise spiral.

Heme (Section 29.10C): A complex organic compound containing an Fe^{2+} ion coordinated with a porphyrin.

Hemiacetal (Section 21.14A): A compound that contains an alkoxy group and a hydroxy group bonded to the same carbon atom.

Hertz (Section 13.5): A unit of frequency measuring the number of waves passing a point per second.

Heteroatom (Sections 1.6, 3.1): An atom other than carbon or hydrogen. Common heteroatoms in organic chemistry are nitrogen, oxygen, sulfur, phosphorus, and the halogens.

Heterocycle (Section 9.3B): A cyclic compound containing a heteroatom as part of the ring.

Heterolysis (Section 6.3A): The breaking of a covalent bond by unequally dividing the electrons between the two atoms in the bond. Heterolysis generates charged intermediates. Heterolysis is also called heterolytic cleavage.

Hexose (Section 28.2): A monosaccharide containing six carbons.

Highest occupied molecular orbital (Section 17.4B): The molecular orbital with the highest energy that also contains electrons. The highest occupied molecular orbital is abbreviated as HOMO.

High-resolution mass spectrometer (Section 13.4A): A mass spectrometer that can measure mass-to-charge ratios to four or more decimal places. High-resolution mass spectra are used to determine the molecular formula of a compound.

Hofmann elimination (Section 25.12): An E2 elimination reaction that converts an amine into a quaternary ammonium salt as the leaving group. The Hofmann elimination gives the less substituted alkene as the major product.

Homologous series (Section 4.1B): A group of compounds that differ by only a CH_2 group in the chain.

Homolysis (Section 6.3A): The breaking of a covalent bond by equally dividing the electrons between the two atoms in the bond. Homolysis generates uncharged radical intermediates. Homolysis is also called homolytic cleavage.

Homopolymer (Section 30.2D): A polymer prepared from a single monomer.

Homotopic protons (Section 14.2C): Two equivalent hydrogen atoms such that substitution of either hydrogen with a group Z forms the same product. The two hydrogen atoms give a single NMR signal.

Hooke's law (Section 13.7): A physical law that can be used to calculate the frequency of a bond vibration from the strength of the bond and the masses of the atoms attached to it.

Host–guest complex (Section 9.5B): The complex that is formed when a small guest molecule binds to a larger host molecule.

Host molecule (Section 9.5B): A large molecule that can bind a smaller guest molecule.

Hückel's rule (Section 17.7): A principle that states for a compound to be aromatic, it must be cyclic, planar, completely conjugated, and have $4n + 2$ π electrons.

Hybridization (Section 1.9B): The mathematical combination of two or more atomic orbitals (having different shapes) to form the same number of hybrid orbitals (all having the same shape).

Hybrid orbital (Section 1.9B): A new orbital that results from the mathematical combination of two or more atomic orbitals. The hybrid orbital is intermediate in energy compared to the atomic orbitals that were combined to form it.

Hydrate (Sections 12.12B, 21.13): A compound having the general structure $R_2C(OH)_2$. Hydrates are also called *gem*-diols.

Hydration (Sections 10.12, 21.9A): Addition of the elements of water to a molecule.

Hydride (Section 12.2): A negatively charged hydrogen ion ($H:^-$).

1,2-Hydride shift (Section 9.9): Rearrangement of a less stable carbocation to a more stable carbocation by the shift of a hydrogen atom from one carbon atom to an adjacent carbon atom.

Hydroboration (Section 10.16): The addition of the elements of borane (BH_3) to an alkene or alkyne.

Hydrocarbon (Sections 3.2A, 4.1): A compound made up of only the elements of carbon and hydrogen.

Hydrogen bonding (Section 3.3B): An attractive intermolecular interaction that occurs when a hydrogen atom bonded to an O, N, or F atom is electrostatically attracted to a lone pair of electrons on an O, N, or F atom in another molecule.

Hydrogenolysis (Section 29.7): A reaction that cleaves a σ bond using H_2 in the presence of a metal catalyst.

α Hydrogens (Section 23.1): The hydrogen atoms on the carbon bonded to the carbonyl carbon atom (the α carbon).

Hydrohalogenation (Section 10.9): An electrophilic addition of hydrogen halide (HX) to an alkene or alkyne.

Hydrolysis (Section 21.9A): A cleavage reaction with water.

Hydroperoxide (Section 15.11): An organic compound having the general structure ROOH.

Hydrophilic (Section 3.4C): Attracted to water. The polar portion of a molecule that interacts with polar water molecules is hydrophilic.

Hydrophobic (Section 3.4C): Not attracted to water. The nonpolar portion of a molecule that is not attracted to polar water molecules is hydrophobic.

β-Hydroxy carbonyl compound (Section 24.1A): An organic compound having a hydroxy group on the carbon β to the carbonyl group.

Hydroxy group (Section 9.1): The OH functional group.

Hyperconjugation (Section 7.13B): The overlap of an empty *p* orbital with an adjacent σ bond.

I

Imide (Section 25.7A): A compound having a nitrogen atom between two carbonyl groups.

Imine (Sections 21.7B, 21.11A): A compound with the general structure $R_2C{=}NR'$. Imines are also called Schiff bases.

Iminium ion (Section 21.11A): A resonance-stabilized cation having the general structure $(R_2C{=}NR'_2)^+$, where R' = H or alkyl.

Inductive effect (Sections 2.5B, 7.13A): The pull of electron density through σ bonds caused by electronegativity differences of atoms.

Infrared (IR) spectroscopy (Section 13.6): An analytical technique used to identify the functional groups in a molecule based on their absorption of electromagnetic radiation in the infrared region.

Initiation (Section 15.4A): The initial step in a chain mechanism that forms a reactive intermediate by cleavage of a bond.

Inscribed polygon method (Section 17.10): A method to predict the relative energies of cyclic, completely conjugated compounds to determine which molecular orbitals are filled or empty. The inscribed polygon is also called a Frost circle.

Integration (Section 14.5): The area under an NMR signal that is proportional to the number of absorbing nuclei that give rise to the signal.

Intermolecular forces (Section 3.3): The types of interactions that exist between molecules. Functional groups determine the type and strength of these forces. Intermolecular forces are also called noncovalent interactions or nonbonded interactions.

Internal alkene (Section 10.1): An alkene that has at least one carbon atom bonded to each end of the double bond.

Internal alkyne (Section 11.1): An alkyne that has one carbon atom bonded to each end of the triple bond.

Inversion of configuration (Section 7.11C): The opposite relative stereochemistry of a stereogenic center in the starting material and product of a chemical reaction. In a nucleophilic substitution reaction, inversion results when the nucleophile and leaving group are in the opposite position relative to the three other groups on carbon.

Iodoform test (Section 23.7B): A test for the presence of methyl ketones, indicated by the formation of the yellow precipitate, CHI_3, via the haloform reaction.

Ionic bond (Section 1.2): A bond that results from the transfer of electrons from one element to another. Ionic bonds result from strong electrostatic interactions between ions with opposite charges. The transfer of electrons forms stable salts composed of cations and anions.

Ionophore (Section 3.7B): An organic molecule that can form a complex with cations so they may be transported across a cell membrane. Ionophores have a hydrophobic exterior and a hydrophilic central cavity that complexes the cation.

Isocyanate (Section 30.6C): A compound having the general structure $RN{=}C{=}O$.

Isoelectric point (Sections 19.14C, 29.1A): The pH at which an amino acid exists primarily in its neutral zwitterionic form. Isoelectric point is abbreviated as p*I*.

Isolated diene (Section 16.1A): A compound containing two carbon–carbon double bonds joined by more than one σ bond.

Isomers (Sections 1.4A, 4.1A, 5.1): Two different compounds that have the same molecular formula.

Isoprene unit (Section 31.7): A five-carbon unit with four carbons in a row and a one-carbon branch on one of the middle carbons.

Isotactic polymer (Section 30.4): A polymer having all the substituents on the same side of the carbon backbone of an elongated polymer chain.

Isotope (Section 1.1): Two or more atoms of the same element having the same number of protons in the nucleus but a different number of neutrons. Isotopes have the same atomic number but different mass numbers.

IUPAC system of nomenclature (Section 4.3): A systematic method for naming compounds developed by the International Union of Pure and Applied Chemistry.

K

***K*$_a$** (Section 2.3): The symbol that represents the acidity constant of an acid HA. The larger the K_a, the stronger the acid.

$$K_a = \frac{[H_3O^+][A:^-]}{[H{-}A]}$$

***K*$_{eq}$** (Section 2.3): The equilibrium constant. K_{eq} = [products]/[starting materials].

Kekulé structures (Section 17.1): Two equilibrating structures for benzene. Each structure contains a six-membered ring and three π bonds alternating with σ bonds around the ring.

Ketal (Section 21.14): A compound having the general structure $R_2C(OR')_2$, where R = alkyl or aryl. Ketals are derived from ketones and constitute a subclass of acetals.

β-Keto ester (Section 23.10): A compound containing a ketone carbonyl on the carbon β to the ester carbonyl group.

Ketone (Section 11.9): A compound with two alkyl groups bonded to the $C{=}O$ carbon atom, having the general structures $R_2C{=}O$ or RCOR'.

Ketose (Section 28.2): A monosaccharide comprised of a polyhydroxy ketone.

Keto tautomer (Section 11.9): A tautomer of a ketone that has a C=O and a hydrogen bonded to the α carbon. The keto tautomer is in equilibrium with the enol tautomer.

Kiliani–Fischer synthesis (Section 28.10B): A reaction that lengthens the carbon chain of an aldose by adding one carbon to the carbonyl end.

Kinetic enolate (Section 23.4): The enolate that is formed the fastest—generally the less substituted enolate.

Kinetic product (Section 16.11): In a reaction that can give more than one product, the product that is formed the fastest.

Kinetic resolution (Section 29.3B): The separation of two enantiomers by a chemical reaction that selectively occurs for only one of the enantiomers.

Kinetics (Section 6.5): The study of chemical reaction rates.

L

L-Sugar (Section 28.2C): A sugar with the hydroxy group on the stereogenic center farthest from the carbonyl on the left side in the Fischer projection formula.

Lactam (Section 22.1): A cyclic amide in which the carbonyl carbon–nitrogen σ bond is part of a ring. A β-lactam contains the carbon–nitrogen σ bond in a four-membered ring.

Lactol (Section 21.16): A cyclic hemiacetal.

Lactone (Section 22.1): A cyclic ester in which the carbonyl carbon–oxygen σ bond is part of a ring.

Leaving group (Section 7.6): An atom or group of atoms (Z) that is able to accept the electron density of the C−Z bond during a substitution or elimination reaction.

Leaving group ability (Section 7.7): A measure of how readily a leaving group (Z) can accept the electron density of the C−Z bond during a substitution or elimination reaction.

Le Châtelier's principle (Section 9.8D): The principle that a system at equilibrium will react to counteract any disturbance to the equilibrium.

Lecithin (Section 31.4A): A phosphoacylglycerol in which the phosphodiester alkyl group is $-CH_2CH_2N(CH_3)_3^+$. Lecithins are also called phosphatidylcholines.

Leukotriene (Section 9.16): An unstable and potent biomolecule synthesized in cells by the oxidation of arachidonic acid. Leukotrienes are responsible for biological conditions such as asthma.

Levorotatory (Section 5.12A): Rotating plane-polarized light in the counterclockwise direction. The rotation is labeled *l* or (–).

Lewis acid (Section 2.8): An electron pair acceptor.

Lewis acid–base reaction (Section 2.8): A reaction that results when a Lewis base donates an electron pair to a Lewis acid.

Lewis base (Section 2.8): An electron pair donor.

Lewis structure (Section 1.3): A representation of a molecule that shows the position of covalent bonds and nonbonding electrons. In Lewis structures, unshared electrons are represented by dots and a two-electron covalent bond is represented by a solid line. Lewis structures are also called electron dot structures.

Ligand (Section 26.2A): A group coordinated to a metal, which donates electron density to or sometimes withdraws electron density from the metal.

"Like dissolves like" (Section 3.4C): The principle that compounds dissolve in solvents having similar kinds of intermolecular forces; that is, polar compounds dissolve in polar solvents and nonpolar compounds dissolve in nonpolar solvents.

Lindlar catalyst (Section 12.5B): A catalyst for the hydrogenation of an alkyne to a cis alkene. The Lindlar catalyst is Pd adsorbed onto $CaCO_3$ with lead(II) acetate and quinoline.

Lipid (Sections 4.15, 31.1): A biomolecule with a large number of C−C and C−H σ bonds that is soluble in organic solvents and insoluble in water.

Lone pair of electrons (Section 1.2): A pair of valence electrons that is not shared with another atom in a covalent bond. Lone pairs are also called unshared or nonbonded pairs of electrons.

Lowest unoccupied molecular orbital (Section 17.9B): The molecular orbital with the lowest energy that does not contain electrons. The lowest unoccupied molecular orbital is abbreviated as the LUMO.

M

M peak (Section 13.1): The peak in the mass spectrum that corresponds to the mass of the molecular ion. The M peak is also called the molecular ion peak or the parent peak.

M + 1 peak (Section 13.1): The peak in the mass spectrum that corresponds to the mass of the molecular ion plus one. The M + 1 peak is caused by the presence of isotopes that increase the mass of the molecular ion.

M + 2 peak (Section 13.2): The peak in the mass spectrum that corresponds to the mass of the molecular ion plus two. The M + 2 peak is caused by the presence of isotopes, typically of a chlorine or a bromine atom.

Macrocyclic lactone (Section 22.6A): A cyclic ester contained in a large ring. Macrocyclic lactones are also called macrolides.

Macrolide (Section 22.6A): A cyclic ester contained in a large ring. Macrolides are also called macrocyclic lactones.

Magnetic resonance imaging (MRI) (Section 14.12): A form of NMR spectroscopy used in medicine.

Malonic ester synthesis (Section 23.9A): A stepwise method that converts diethyl malonate into a carboxylic acid having one or two carbons bonded to the α carbon.

Markovnikov's rule (Section 10.10): The rule that states in the addition of HX to an unsymmetrical alkene, the H atom bonds to the less substituted carbon atom.

Mass number (Section 1.1): The total number of protons and neutrons in the nucleus of a particular atom.

Mass spectrometry (Section 13.1): An analytical technique used for measuring the molecular weight and determining the molecular formula of an organic molecule.

Mass-to-charge ratio (Section 13.1): A ratio of the mass to the charge of a molecular ion or fragment. Mass-to-charge ratio is abbreviated as *m*/*z*.

Megahertz (Section 14.1A): A unit used for the frequency of the RF radiation in NMR spectroscopy. Megahertz is abbreviated as MHz; 1 MHz = 10^6 Hz.

Melting point (Section 3.4B): The temperature at which molecules in the solid phase are converted to the liquid phase. Molecules with stronger intermolecular forces and higher symmetry have higher melting points. Melting point is abbreviated as mp.

Merrifield method (Section 29.8): A method for synthesizing polypeptides using insoluble polymer supports.

Meso compound (Section 5.8): An achiral compound that contains two or more tetrahedral stereogenic centers.

Meta director (Section 18.7): A substituent on a benzene ring that directs a new group to the meta position during electrophilic aromatic substitution.

Meta isomer (Section 17.3B): A 1,3-disubstituted benzene ring. Meta substitution is abbreviated as *m*-.

Metal hydride reagent (Section 12.2): A reagent containing a polar metal–hydrogen bond that places a partial negative charge on the hydrogen and acts as a source of hydride ions ($H:^-$).

Metathesis (Section 26.6): A reaction between two alkene molecules that results in the interchange of the carbons of their double bonds.

Methylation (Section 7.16): A reaction in which a CH_3 group is transferred from one compound to another.

Methylene group (Sections 4.1B, 10.3C): A CH_2 group bonded to a carbon chain ($-CH_2-$) or part of a double bond ($CH_2=$).

1,2-Methyl shift (Section 9.9): Rearrangement of a less stable carbocation to a more stable carbocation by the shift of a methyl group from one carbon atom to an adjacent carbon atom.

Micelles (Section 3.6): Spherical droplets formed by soap molecules having the ionic heads on the surface and the nonpolar tails packed together in the interior. Grease and oil dissolve in the interior nonpolar region.

Michael acceptor (Section 24.8): The α,β-unsaturated carbonyl compound in a Michael reaction.

Michael reaction (Section 24.8): A reaction in which a resonance-stabilized carbanion (usually an enolate) adds to the β carbon of an α,β-unsaturated carbonyl compound.

Mixed aldol reaction (Section 24.2): An aldol reaction between two different carbonyl compounds. A mixed aldol reaction is also called a crossed aldol reaction.

Mixed anhydride (Section 22.1): An anhydride with two different alkyl groups bonded to the carbonyl carbon atoms.

Molecular ion (Section 13.1): The radical cation having the general structure $M^{+\bullet}$, formed by the removal of an electron from an organic molecule. The molecular ion is also called the parent ion.

Molecular orbital theory (Section 17.9A): A theory that describes bonds as the mathematical combination of atomic orbitals to form a new set of orbitals called molecular orbitals. Molecular orbital theory is also called MO theory.

Molecular recognition (Section 9.5B): The ability of a host molecule to recognize and bind specific guest molecules.

Molecule (Section 1.2): A compound containing two or more atoms bonded together with covalent bonds.

Monomers (Sections 5.1, 15.14): Small organic compounds that can be covalently bonded to each other (polymerized) in a repeating pattern.

Monosaccharide (Section 28.2): A simple sugar having three to seven carbon atoms.

Monosubstituted alkene (Section 8.2A): An alkene that has one alkyl group and three hydrogens bonded to the carbons of the double bond ($RCH=CH_2$).

Monoterpene (Section 31.7A): A terpene that contains 10 carbons and two isoprene units. A monoterpenoid also contains at least one oxygen atom.

Multiplet (Section 14.6C): An NMR signal that is split into more than seven peaks.


Mutarotation (Section 28.6A): The process by which a pure anomer of a monosaccharide equilibrates to a mixture of both anomers when placed in solution.

N

***n* + 1 rule** (Section 14.6C): The rule that an NMR signal for a proton with *n* nearby nonequivalent protons will be split into *n* + 1 peaks.

Natural product (Section 7.18): A compound isolated from a natural source.

Newman projection (Section 4.9): An end-on representation of the conformation of a molecule. The Newman projection shows the three groups bonded to each carbon atom in a particular C−C bond, as well as the dihedral angle that separates the groups on each carbon.



Nitration (Section 18.4): An electrophilic aromatic substitution reaction in which benzene reacts with $^+NO_2$ to give nitrobenzene, $C_6H_5NO_2$.

Nitrile (Sections 22.1, 22.18): A compound having the general structure $RC\equiv N$.

Nitronium ion (Section 18.4): An electrophile having the structure $^+NO_2$.

***N*-Nitrosamine** (Sections 7.16, 25.13B): A compound having the general structure $R_2N-N=O$. Nitrosamines are formed by the reaction of a secondary amine with ^+NO.

Nitrosonium ion (Section 25.13): An electrophile having the structure ^+NO.

NMR peak (Section 14.6A): The individual absorptions in a split NMR signal due to nonequivalent nearby protons.

NMR signal (Section 14.6A): The entire absorption due to a particular kind of proton in an NMR spectrum.

NMR spectrometer (Section 14.1A): An analytical instrument that measures the absorption of RF radiation by certain atomic nuclei when placed in a strong magnetic field.

Nonbonded pair of electrons (Section 1.2): A pair of valence electrons that is not shared with another atom in a covalent bond. Nonbonded electrons are also called unshared or lone pairs of electrons.

Nonbonding molecular orbital (Section 17.10): A molecular orbital having the same energy as the atomic orbitals that formed it.

Nonnucleophilic base (Section 7.8B): A base that is a poor nucleophile due to steric hindrance resulting from the presence of bulky groups.

Nonpolar bond (Section 1.12): A covalent bond in which the electrons are equally shared between the two atoms.

Nonpolar molecule (Section 1.13): A molecule that has no net dipole. A nonpolar molecule has either no polar bonds or multiple polar bonds whose dipoles cancel.

Nonreducing sugar (Section 28.9B): A carbohydrate that cannot be oxidized by Tollens, Benedict's, or Fehling's reagent.

Normal alkane (Section 4.1A): An acyclic alkane that has all of its carbons in a row. A normal alkane is an "*n*-alkane" or a straight-chain alkane.

Nuclear magnetic resonance spectroscopy (Section 14.1): A powerful analytical tool that can help identify the carbon and hydrogen framework of an organic molecule.

Nucleophile (Sections 2.8, 7.6): An electron-rich compound, symbolized by $:Nu^-$, which donates a pair of electrons to an electron-deficient compound, forming a covalent bond. Lewis bases are nucleophiles.

Nucleophilic acyl substitution (Sections 20.2B, 22.1): Substitution of a leaving group by a nucleophile at a carbonyl carbon.

Nucleophilic addition (Section 20.2A): Addition of a nucleophile to the electrophilic carbon of a carbonyl group followed by protonation of the oxygen.

Nucleophilic aromatic substitution (Section 18.13): A substitution reaction of an aryl halide with a strong nucleophile.

Nucleophilicity (Section 7.8A): A measure of how readily an atom donates an electron pair to other atoms.

Nucleophilic substitution (Section 7.6): A reaction in which a nucleophile replaces the leaving group in a molecule.

Nucleoside (Section 28.13B): A biomolecule having a sugar and either a purine or pyrimidine base joined together by an *N*-glycosidic linkage.

Nucleotide (Section 28.13B): A biomolecule having a sugar and either a purine or pyrimidine base joined together by an *N*-glycosidic linkage, and a phosphate bonded to a hydroxy group of the sugar nucleus.

O

Observed rotation (Section 5.12A): The angle that a sample of an optically active compound rotates plane-polarized light. The angle is denoted by the symbol α and is measured in degrees (°).

Octet rule (Section 1.2): The general rule governing the bonding process for second-row elements. Through bonding, second-row elements attain a complete outer shell of eight valence electrons.

Oil (Sections 10.6B, 31.3): A triacylglycerol that is liquid at room temperature and composed of fatty acid side chains with a high degree of unsaturation.

Olefin (Section 10.1): An alkene; a compound possessing a carbon–carbon double bond.

Optically active (Section 5.12A): Able to rotate the plane of plane-polarized light as it passes through a solution of a compound.

Optically inactive (Section 5.12A): Not able to rotate the plane of plane-polarized light as it passes through a solution of a compound.

Optical purity (Section 5.12D): A measurement of how much one enantiomer is present in excess of the racemic mixture. Optical purity is also called enantiomeric excess (*ee*); *ee* = % of one enantiomer – % of the other enantiomer.

Orbital (Section 1.1): A region of space around the nucleus of an atom that is high in electron density. There are four different kinds of orbitals, called *s, p, d,* and *f*.

Order of a rate equation (Section 6.9B): The sum of the exponents of the concentration terms in the rate equation of a reaction.

Organoborane (Section 10.16): A compound that contains a carbon–boron bond. Organoboranes have the general structure RBH_2, R_2BH, or R_3B.

Organocopper reagent (Section 20.9): An organometallic reagent having the general structure R_2CuLi. Organocopper reagents are also called organocuprates.

Organolithium reagent (Section 20.9): An organometallic reagent having the general structure RLi.

Organomagnesium reagent (Section 20.9): An organometallic reagent having the general structure RMgX. Organomagnesium reagents are also called Grignard reagents.

Organometallic reagent (Section 20.9): A reagent that contains a carbon atom bonded to a metal.

Organopalladium compound (Section 26.2): An organometallic compound that contains a carbon–palladium bond.

Organophosphorus reagent (Section 21.10A): A reagent that contains a carbon–phosphorus bond.

Ortho isomer (Section 17.3B): A 1,2-disubstituted benzene ring. Ortho substitution is abbreviated as *o*-.

Ortho, para director (Section 18.7): A substituent on a benzene ring that directs a new group to the ortho and para positions during electrophilic aromatic substitution.

Oxaphosphetane (Section 21.10B): An intermediate in the Wittig reaction consisting of a four-membered ring containing a phosphorus–oxygen bond.

Oxazaborolidine (Section 20.6A): A heterocycle possessing a boron, a nitrogen, and an oxygen. An oxazaborolidine can be used to form a chiral reducing agent.

Oxidation (Sections 4.14A, 12.1): A process that results in a loss of electrons. For organic compounds, oxidation results in an increase in the number of C−Z bonds or a decrease in the number of C−H bonds; Z = an element more electronegative than carbon.

Oxidative addition (Section 26.2A): The addition of a reagent to a metal, often increasing the number of groups around the metal by two.

Oxidative cleavage (Section 12.10): An oxidation reaction that breaks both the σ and π bonds of a multiple bond to form two oxidized products.

Oxime (Section 28.10A): A compound having the general structure $R_2C{=}NOH$.

Oxirane (Section 9.1): A cyclic ether having the oxygen atom as part of a three-membered ring. Oxiranes are also called epoxides.

Oxy-Cope rearrangement (Section 27.5): A [3,3] sigmatropic rearrangement of a 1,5-dien-3-ol to a δ,ε-unsaturated carbonyl compound.

Ozonolysis (Section 12.10): An oxidative cleavage reaction in which a multiple bond reacts with ozone (O_3) as the oxidant.

P

Para isomer (Section 17.3B): A 1,4-disubstituted benzene ring. Para substitution is abbreviated as *p*-.

Parent ion (Section 13.1): The radical cation having the general structure $M^{+\bullet}$, formed by the removal of an electron from an organic molecule. The parent ion is also called the molecular ion.

Parent name (Section 4.4): The portion of the IUPAC name of an organic compound that indicates the number of carbons in the longest continuous chain in the molecule.

Pentose (Section 28.2): A monosaccharide containing five carbons.

Peptide bond (Section 29.5): The amide bond in peptides and proteins.

Peptides (Sections 22.6B, 29.5): Low molecular weight polymers of less than 40 amino acids joined together by amide linkages.

Percent *s*-character (Section 1.11B): The fraction of a hybrid orbital due to the *s* orbital used to form it. As the percent *s*-character increases, a bond becomes shorter and stronger.

Percent transmittance (Section 13.6B): A measure of how much electromagnetic radiation passes through a sample of a compound and how much is absorbed.

Pericyclic reaction (Section 27.1): A concerted reaction that proceeds through a cyclic transition state.

Peroxide (Section 15.2): A reactive organic compound with the general structure ROOR. Peroxides are used as radical initiators by homolysis of the weak O−O bond.

Peroxyacid (Section 12.7): An oxidizing agent having the general structure RCO_3H.

Peroxy radical (Section 15.11): A radical having the general structure ROO·.

Petroleum (Section 4.7): A fossil fuel containing a complex mixture of compounds, primarily hydrocarbons with 1 to 40 carbon atoms.

Phenol (Sections 9.1, 15.12): A compound such as C_6H_5OH, which contains a hydroxy group bonded to a benzene ring.

Phenyl group (Section 17.3D): A group formed by removal of one hydrogen from benzene, abbreviated as C_6H_5- or Ph−.

Pheromone (Section 4.1): A chemical substance used for communication in an animal or insect species.

Phosphate (Section 7.16): A PO_4^{3-} anion.

Phosphatidylcholine (Section 31.4A): A phosphoacylglycerol in which the phosphodiester alkyl group is $-CH_2CH_2N(CH_3)_3^+$. Phosphatidylcholines are also called lecithins.

Phosphatidylethanolamine (Section 31.4A): A phosphoacylglycerol in which the phosphodiester alkyl group is $-CH_2CH_2NH_3^+$. Phosphatidylethanolamines are also called cephalins.

Phosphoacylglycerols (Section 31.4A): A lipid having a glycerol backbone with two of the hydroxy groups esterified with fatty acids and the third hydroxy group as part of a phosphodiester.

Phosphodiester (Section 31.4): A functional group having the general formula $ROPO_2OR'$ formed by replacing two of the H atoms in phosphoric acid (H_3PO_4) with alkyl groups.

Phospholipid (Sections 3.7A, 31.4): A hydrolyzable lipid that contains a phosphorus atom.

Phosphonium salt (Section 21.10A): An organophosphorus reagent with a positively charged phosphorus and a suitable counterion; for example, $R_4P^+X^-$. Phosphonium salts are converted to ylides upon treatment with a strong base.

Phosphorane (Section 21.10A): A phosphorus ylide; for example, $Ph_3P{=}CR_2$.

Photon (Section 13.5): A particle of electromagnetic radiation.

Pi (π) bond (Section 1.10B): A bond formed by side-by-side overlap of two *p* orbitals where electron density is not concentrated on the axis joining the two nuclei. Pi (π) bonds are generally weaker than σ bonds.

pK_a (Section 2.3): A logarithmic scale of acid strength. p$K_a = -\log K_a$. The smaller the pK_a, the stronger the acid.

Plane of symmetry (Section 5.3): A mirror plane that cuts a molecule in half, so that one half of the molecule is the mirror reflection of the other half.

Plane-polarized light (Section 5.12A): Light that has an electric vector that oscillates in a single plane. Plane-polarized light, also called polarized light, arises from passing ordinary light through a polarizer.

Plasticizer (Section 30.7): A low molecular weight compound added to a polymer to give it flexibility.

β-Pleated sheet (Section 29.9B): A secondary structure of a protein formed when two or more peptide chains line up side by side.

Poisoned catalyst (Section 12.5B): A hydrogenation catalyst with reduced activity that allows selective reactions to occur. The Lindlar catalyst is a poisoned Pd catalyst that converts alkynes to cis alkenes.

Polar aprotic solvent (Section 7.8C): A polar solvent that is incapable of intermolecular hydrogen bonding because it does not contain an O−H or N−H bond.

Polar bond (Section 1.12): A covalent bond in which the electrons are unequally shared between the two atoms. Unequal sharing of electrons results from bonding between atoms of different electronegativity values, usually with a difference of ≥ 0.5 units.

Polarimeter (Section 5.12A): An instrument that measures the degree that a compound rotates plane-polarized light.

Polarity (Section 1.12): A characteristic that results from a dipole. The polarity of a bond is indicated by an arrow with the head of the arrow pointing toward the negative end of the dipole and the tail with a perpendicular line through it at the positive end of the dipole. The polarity of a bond can also be indicated by the symbols δ+ and δ–.

Polarizability (Section 3.3B): A measure of how the electron cloud around an atom responds to changes in its electronic environment.

Polar molecule (Section 1.13): A molecule that has a net dipole. A polar molecule has either one polar bond or multiple polar bonds whose dipoles reinforce.

Polar protic solvent (Section 7.8C): A polar solvent that is capable of intermolecular hydrogen bonding because it contains an O−H or N−H bond.

Polyamide (Sections 22.16A, 30.6A): A step-growth polymer that contains many amide bonds. Nylon 6,6 and nylon 6 are polyamides.

Polycarbonate (Section 30.6C): A step-growth polymer that contains many $-OC(=O)O-$ bonds in its backbone, often formed by reaction of $Cl_2C{=}O$ with a diol.

Polycyclic aromatic hydrocarbon (Sections 9.17, 17.5): An aromatic hydrocarbon containing two or more benzene rings that share carbon–carbon bonds. Polycyclic aromatic hydrocarbons are abbreviated as PAHs.

Polyene (Section 16.7): A compound that contains three or more double bonds.

Polyester (Sections 22.16B, 30.6B): A step-growth polymer consisting of many ester bonds between diols and dicarboxylic acids.

Polyether (Sections 9.5B, 30.3): A compound that contains two or more ether linkages.

Polymer (Sections 5.1, 15.14): A large molecule composed of smaller monomer units covalently bonded to each other in a repeating pattern.

Polymerization (Section 15.14A): The chemical process that joins together monomers to make polymers.

Polysaccharide (Section 28.12): A carbohydrate containing three or more monosaccharide units joined together by glycosidic linkages.

Polyurethane (Section 30.6C): A step-growth polymer that contains many $-NHC(=O)O-$ bonds in its backbone, formed by reaction of a diisocyanate and a diol.

Porphyrin (Section 29.10C): A nitrogen-containing heterocycle that can complex metal ions.

Primary (1°) alcohol (Section 3.2): An alcohol having the general structure RCH_2OH.

Primary (1°) alkyl halide (Section 3.2): An alkyl halide having the general structure RCH_2X.

Primary (1°) amide (Section 3.2): An amide having the general structure $RCONH_2$.

Primary (1°) amine (Section 3.2): An amine having the general structure RNH_2.

Primary (1°) carbocation (Section 7.13): A carbocation having the general structure RCH_2^+.

Primary (1°) carbon (Section 3.2): A carbon atom that is bonded to one other carbon atom.

Primary (1°) hydrogen (Section 3.2): A hydrogen that is bonded to a 1° carbon.

Primary protein structure (Section 29.9A): The particular sequence of amino acids joined together by peptide bonds.

Primary (1°) radical (Section 15.1): A radical having the general structure $RCH_2\cdot$.

Propagation (Section 15.4A): The middle part of a chain mechanism in which one reactive particle is consumed and another is generated. Propagation repeats until a termination step occurs.

Prostaglandin (Section 4.15): A class of lipids containing 20 carbons, a five-membered ring, and a CO_2H group. Prostaglandins possess a wide range of biological activities.

Prosthetic group (Section 29.10C): The non-protein unit of a conjugated protein.

Protecting group (Section 20.12): A blocking group that renders a reactive functional group unreactive, so that it does not interfere with another reaction.

Protection (Section 20.12): The reaction that blocks a reactive functional group with a protecting group.


Proteins (Sections 22.6B, 29.5): High molecular weight polymers of 40 or more amino acids joined together by amide linkages.

Proton (Section 2.1): A positively charged hydrogen ion (H^+).

Proton NMR spectroscopy (Section 14.1): A form of nuclear magnetic resonance spectroscopy used to determine the number and type of hydrogen atoms in a molecule.

Proton transfer reaction (Section 2.2): A Brønsted–Lowry acid–base reaction; a reaction that results in the transfer of a proton from an acid to a base.

Purine (Sections 25.3, 28.13B): A bicyclic aromatic heterocycle having two nitrogens in each of the rings.



Pyranose (Section 28.6): A cyclic six-membered ring of a monosaccharide containing an oxygen atom.

Pyrimidine (Sections 25.3, 28.13B): A six-membered aromatic heterocycle having two nitrogens in the ring.



Q

Quantum (Section 13.5): The discrete amount of energy associated with a particle of electromagnetic radiation (i.e., a photon).

Quartet (Section 14.6C): An NMR signal that is split into four peaks having a relative area of 1:3:3:1, caused by three nearby nonequivalent protons.

Quaternary (4°) carbon (Section 3.2): A carbon atom that is bonded to four other carbon atoms.

Quaternary protein structure (Section 29.9C): The shape adopted when two or more folded polypeptide chains aggregate into one protein complex.

Quintet (Section 14.6C): An NMR signal that is split into five peaks caused by four nearby nonequivalent protons.

R

Racemic mixture (Section 5.12B): An equal mixture of two enantiomers. A racemic mixture, also called a racemate, is optically inactive.

Racemization (Section 7.12C): The formation of equal amounts of two enantiomers from an enantiomerically pure starting material.

Radical (Sections 6.3B, 15.1): A reactive intermediate with a single unpaired electron, formed by homolysis of a covalent bond.

Radical anion (Section 12.5C): A reactive intermediate containing both a negative charge and an unpaired electron.

Radical cation (Section 13.1): A species with an unpaired electron and a positive charge, formed in a mass spectrometer by the bombardment of a molecule with an electron beam.

Radical inhibitor (Section 15.2): A compound that prevents radical reactions from occurring. Radical inhibitors are also called radical scavengers.

Radical initiator (Section 15.2): A compound that contains an especially weak bond that serves as a source of radicals.

Radical polymerization (Section 15.14B): A radical chain reaction involving the polymerization of alkene monomers by adding a radical to a π bond.

Radical scavenger (Section 15.2): A compound that prevents radical reactions from occurring. Radical scavengers are also called radical inhibitors.

Rate constant (Section 6.9B): A constant that is a fundamental characteristic of a reaction. The rate constant, symbolized by *k,* is a complex mathematical term that takes into account the dependence of a reaction rate on temperature and the energy of activation.

Rate-determining step (Section 6.8): In a multistep reaction mechanism, the step with the highest energy transition state.

Rate equation (Section 6.9B): An equation that shows the relationship between the rate of a reaction and the concentration of the reactants. The rate equation depends on the mechanism of the reaction and is also called the rate law.

Reaction coordinate (Section 6.7): The *x* axis in an energy diagram that represents the progress of a reaction as it proceeds from reactant to product.

Reaction mechanism (Section 6.3): A detailed description of how bonds are broken and formed as a starting material is converted to a product.

Reactive intermediate (Sections 6.3, 10.18): A high-energy unstable intermediate formed during the conversion of a stable starting material to a stable product.

Reciprocal centimeter (Section 13.6A): The unit for wavenumber, which is used to report frequency in IR spectroscopy.

Reducing sugar (Section 28.9B): A carbohydrate that can be oxidized by Tollens, Benedict's, or Fehling's reagent.

Reduction (Sections 4.14A, 12.1): A process that results in the gain of electrons. For organic compounds, reduction results in a decrease in the number of C−Z bonds or an increase in the number of C−H bonds; Z = an element more electronegative than carbon.

Reductive amination (Section 25.7C): A two-step method that converts aldehydes and ketones into amines.

Reductive elimination (Section 26.2A): The elimination of two groups that surround a metal, often forming new carbon–hydrogen or carbon–carbon bonds.

Regioselective reaction (Section 8.5): A reaction that yields predominantly or exclusively one constitutional isomer when more than one constitutional isomer is possible.

Resolution (Section 29.3): The separation of a racemic mixture into its component enantiomers.

Resonance (Section 14.1A): In NMR spectroscopy, when an atomic nucleus absorbs RF radiation and spin flips to a higher energy state.

Resonance hybrid (Sections 1.6C, 16.4): A structure that is a weighted composite of all possible resonance structures. The resonance hybrid shows the delocalization of electron density due to the different locations of electrons in individual resonance structures.

Resonance structures (Sections 1.6, 16.2): Two or more structures of a molecule that differ in the placement of π bonds and nonbonded electrons. The placement of atoms and σ bonds stays the same.

Retention of configuration (Section 7.11C): The same relative stereochemistry of a stereogenic center in the reactant and the product of a chemical reaction.

Retention time (Section 13.4B): The length of time required for a component of a mixture to travel through a chromatography column.

Retro Diels–Alder reaction (Section 16.14B): The reverse of a Diels–Alder reaction in which a cyclohexene is cleaved to give a 1,3-diene and an alkene.

Retrosynthetic analysis (Section 10.18): Working backwards from a product to determine the starting material from which it is made.

RF radiation (Section 14.1A): Radiation in the radiofrequency region of the electromagnetic spectrum, characterized by long wavelength and low frequency and energy.

Ribonucleoside (Section 28.13B): An *N*-glycoside formed by the reaction of D-ribose with certain amine heterocycles.

Ribonucleotide (Section 28.13B): An RNA building block having a ribose and either a purine or pyrimidine base joined together by an *N*-glycosidic linkage, and a phosphate bonded to a hydroxy group of the sugar nucleus.

Ring-closing metathesis (Section 26.6): An intramolecular olefin metathesis reaction using a diene starting material, which results in ring closure.

Ring current (Section 14.4): A circulation of π electrons in an aromatic ring caused by the presence of an external magnetic field.

Ring-flipping (Section 4.12B): A stepwise process in which one chair conformation of cyclohexane interconverts with a second chair conformation.

Ring-opening metathesis polymerization (Problem 26.33): An olefin metathesis reaction that forms a high molecular weight polymer from certain cyclic alkenes.

Robinson annulation (Section 24.9): A ring-forming reaction that combines a Michael reaction with an intramolecular aldol reaction to form a cyclohex-2-enone.

***R,S* System of nomenclature** (Section 5.6): A system of nomenclature that distinguishes the stereochemistry at a tetrahedral stereogenic center by assigning a priority to each group connected to the stereogenic center. *R* indicates a clockwise orientation of the three highest priority groups and *S* indicates a counterclockwise orientation of the three highest groups. The system is also called the Cahn–Ingold–Prelog system.

Rule of endo addition (Section 16.13D): The rule that the endo product is preferred in a Diels–Alder reaction.

S

Sandmeyer reaction (Section 25.14A): A reaction between an aryl diazonium salt and a copper(I) halide to form an aryl halide (C_6H_5Cl or C_6H_5Br).

Saponification (Section 22.11B): Basic hydrolysis of an ester to form an alcohol and a carboxylate anion.

Saturated fatty acid (Section 10.6A): A fatty acid having no carbon–carbon double bonds in its long hydrocarbon chain.

Saturated hydrocarbon (Section 4.1): A compound that contains only C−C and C−H σ bonds and no rings, thus having the maximum number of hydrogen atoms per carbon.

Schiff base (Section 21.11A): A compound having the general structure $R_2C{=}NR'$. A Schiff base is also called an imine.

Secondary (2°) alcohol (Section 3.2): An alcohol having the general structure R_2CHOH.

Secondary (2°) alkyl halide (Section 3.2): An alkyl halide having the general structure R_2CHX.

Secondary (2°) amide (Section 3.2): An amide having the general structure RCONHR'.

Secondary (2°) amine (Section 3.2): An amine having the general structure R_2NH.

Secondary (2°) carbocation (Section 7.13): A carbocation having the general structure R_2CH^+.

Secondary (2°) carbon (Section 3.2): A carbon atom that is bonded to two other carbon atoms.

Secondary (2°) hydrogen (Section 3.2): A hydrogen that is attached to a 2° carbon.

Secondary protein structure (Section 29.9B): The three-dimensional conformations of localized regions of a protein.

Secondary (2°) radical (Section 15.1): A radical having the general structure $R_2CH{\cdot}$.

Second-order rate equation (Sections 6.9B, 7.10): A rate equation in which the reaction rate depends on the concentration of two reactants.

Separatory funnel (Section 19.12): An item of laboratory glassware used for extractions.

Septet (Section 14.6C): An NMR signal that is split into seven peaks caused by six nearby nonequivalent protons.

Sesquiterpene (Section 31.7A): A terpene that contains 15 carbons and three isoprene units. A sesquiterpenoid also contains at least one oxygen atom.

Sesterterpene (Section 31.7A): A terpene that contains 25 carbons and five isoprene units. A sesterterpenoid also contains at least one oxygen atom.

Sextet (Section 14.6C): An NMR signal that is split into six peaks caused by five nearby nonequivalent protons.

Sharpless asymmetric epoxidation (Section 12.15): An enantioselective oxidation reaction that converts the double bond of an allylic alcohol to a predictable enantiomerically enriched epoxide.

Sharpless reagent (Section 12.15): The reagent used in the Sharpless asymmetric epoxidation. The Sharpless reagent consists of *tert*-butyl hydroperoxide, a titanium catalyst, and one enantiomer of diethyl tartrate.

Shielding effects (Section 14.3A): An effect in NMR caused by small induced magnetic fields of electrons in the opposite direction to the applied magnetic field. Shielding decreases the strength of the magnetic field felt by the nucleus and shifts an absorption upfield.

1,2-Shift (Section 9.9): Rearrangement of a less stable carbocation to a more stable carbocation by the shift of a hydrogen atom or an alkyl group from one carbon atom to an adjacent carbon atom.

Sigma (σ) bond (Section 1.9A): A cylindrically symmetrical bond that concentrates the electron density on the axis that joins two nuclei. All single bonds are σ bonds.

Sigmatropic rearrangement (Section 27.1): A pericyclic reaction in which a σ bond is broken in the reactant, the π bonds rearrange, and a σ bond is formed in the product.

Silyl ether (Section 20.12): A common protecting group for an alcohol in which the O−H bond is replaced by an O−Si bond.

Simmons–Smith reaction (Section 26.5): Reaction of an alkene with CH_2I_2 and Zn(Cu) to form a cyclopropane.

Singlet (Section 14.6A): An NMR signal that occurs as a single peak.

Skeletal structure (Section 1.8B): A shorthand representation of the structure of an organic compound in which carbon atoms and the hydrogen atoms bonded to them are omitted. All heteroatoms and the hydrogens bonded to them are drawn in. Carbon atoms are assumed to be at the junction of any two lines or at the end of a line.

S_N1 mechanism (Sections 7.10, 7.12): A nucleophilic substitution mechanism that goes by a two-step process involving a carbocation intermediate. S_N1 is an abbreviation for "Substitution Nucleophilic Unimolecular."

S_N2 mechanism (Sections 7.10, 7.11): A nucleophilic substitution mechanism that goes by a one-step concerted process, where both reactants are involved in the transition state. S_N2 is an abbreviation for "Substitution Nucleophilic Bimolecular."

Soap (Sections 3.6, 22.12B): The carboxylate salts of long-chain fatty acids prepared by the basic hydrolysis or saponification of a triacylglycerol.

Solubility (Section 3.4C): A measure of the extent to which a compound dissolves in a liquid.

Solute (Section 3.4C): The compound that is dissolved in a liquid solvent.

Solvent (Section 3.4C): The liquid component into which the solute is dissolved.

Specific rotation (Section 5.12C): A standardized physical constant for the amount that a chiral compound rotates plane-polarized light. Specific rotation is denoted by the symbol [α] and defined using a specific sample tube length (l in dm), concentration (c in g/mL), temperature (25 °C), and wavelength (589 nm). $[\alpha] = \alpha/(l \times c)$

Spectator ion (Section 2.1): An ion that does not take part in a reaction and is opposite in charge to the ion that does take part in a reaction. A spectator ion is also called a counterion.

Spectroscopy (Section 13.1): An analytical method using the interaction of electromagnetic radiation with molecules to determine molecular structure.

Sphingomyelin (Section 31.4B): A hydrolyzable phospholipid derived from sphingosine.

Spin flip (Section 14.1A): In NMR spectroscopy, when an atomic nucleus absorbs RF radiation and its magnetic field flips relative to the external magnetic field.

Spin–spin splitting (Section 14.6): Splitting of an NMR signal into peaks caused by nonequivalent protons on the same carbon or adjacent carbons.

Spiro ring system (Problem 23.61, Appendix B): A compound having two rings that share a single carbon atom.

Staggered conformation (Section 4.9): A conformation of a molecule in which the bonds on one carbon bisect the R−C−R bond angle on the adjacent carbon.



Step-growth polymer (Sections 22.16A, 30.1): A polymer formed when monomers containing two functional groups come together with loss of a small molecule such as water or HCl. Step-growth polymers are also called condensation polymers.

Stereochemistry (Sections 4.9, 5.1): The three-dimensional structure of molecules.

Stereogenic center (Section 5.3): A site in a molecule at which the interchange of two groups forms a stereoisomer. A carbon bonded to four different groups is a tetrahedral stereogenic center. A tetrahedral stereogenic center is also called a chirality center, a chiral center, or an asymmetric center.

Stereoisomers (Sections 4.13B, 5.1): Two isomers that differ only in the way the atoms are oriented in space.


Stereoselective reaction (Section 8.5): A reaction that yields predominantly or exclusively one stereoisomer when two or more stereoisomers are possible.

Stereospecific reaction (Section 10.14): A reaction in which each of two stereoisomers of a starting material yields a particular stereoisomer of a product.

Steric hindrance (Section 7.8B): A decrease in reactivity resulting from the presence of bulky groups at the site of a reaction.

Steric strain (Section 4.10): An increase in energy resulting when atoms in a molecule are forced too close to one another.

Steroid (Sections 16.14C, 31.8): A tetracyclic lipid composed of three six-membered rings and one five-membered ring.



Straight-chain alkane (Section 4.1A): An acyclic alkane that has all of its carbons in a row. Straight-chain alkanes are also called normal alkanes.

Strecker amino acid synthesis (Section 29.2C): A reaction that converts an aldehyde into an α-amino acid by way of an α-amino nitrile.

Structural isomers (Sections 4.1A, 5.2): Two compounds that have the same molecular formula but differ in the way the atoms are connected to each other. Structural isomers are also called constitutional isomers.

Substituent (Section 4.4): A group or branch attached to the longest continuous chain of carbons in an organic molecule.

Substitution reaction (Section 6.2A): A reaction in which an atom or a group of atoms is replaced by another atom or group of atoms. Substitution reactions involve σ bonds: one σ bond breaks and another is formed at the same atom.

Substrate (Section 6.11): An organic molecule that is transformed by the action of an enzyme.

Sulfide (Section 9.15): A compound having the general structure RSR'.

Sulfonate anion (Section 19.13): An anion having the general structure RSO_3^-, formed by deprotonating a sulfonic acid with a Brønsted–Lowry base.

Sulfonation (Section 18.4): An electrophilic aromatic substitution reaction in which benzene reacts with $^+SO_3H$ to give a benzenesulfonic acid, $C_6H_5SO_3H$.

Sulfonic acid (Section 19.13): A compound having the general structure RSO_3H.

Suprafacial reaction (Section 27.4): A pericyclic reaction that occurs on the same side of the two ends of the π electron system.

Suzuki reaction (Section 26.2): The palladium-catalyzed coupling of an organic halide (R'X) with an organoborane (RBY_2) to form a product R−R'.

Symmetrical anhydride (Section 22.1): An anhydride that has two identical alkyl groups bonded to the carbonyl carbon atoms.

Symmetrical ether (Section 9.1): An ether with two identical alkyl groups bonded to the oxygen.

Syn addition (Section 10.8): An addition reaction in which two parts of a reagent are added from the same side of a double bond.

Syn dihydroxylation (Section 12.9B): The addition of two hydroxy groups to the same face of a double bond.

Syndiotactic polymer (Section 30.4): A polymer having the substituents alternating from one side of the backbone of an elongated polymer chain to the other.

Syn periplanar (Section 8.8): In an elimination reaction, a geometry in which the β hydrogen and the leaving group are on the same side of the molecule.

Systematic name (Section 4.3): The name of a molecule indicating the compound's chemical structure. The systematic name is also called the IUPAC name.

T

Target compound (Section 11.12): The final product of a synthetic scheme.

Tautomerization (Sections 11.9, 23.2A): The process of converting one tautomer into another.

Tautomers (Section 11.9): Constitutional isomers that are in equilibrium and differ in the location of a double bond and a hydrogen atom.

Terminal alkene (Section 10.1): An alkene that has the double bond at the end of the carbon chain.

Terminal alkyne (Section 11.1): An alkyne that has the triple bond at the end of the carbon chain.

C-Terminal amino acid (Section 29.5A): The amino acid at the end of a peptide chain with a free carboxy group.

N-Terminal amino acid (Section 29.5A): The amino acid at the end of a peptide chain with a free amino group.

Termination (Section 15.4A): The final step of a chain reaction. In a radical chain mechanism, two radicals combine to form a stable bond.

Terpene (Section 31.7): A hydrocarbon composed of repeating five-carbon isoprene units.

Terpenoid (Section 31.7): A lipid that contains isoprene units as well as at least one oxygen heteroatom.

Tertiary (3°) alcohol (Section 3.2): An alcohol having the general structure R_3COH.

Tertiary (3°) alkyl halide (Section 3.2): An alkyl halide having the general structure R_3CX.

Tertiary (3°) amide (Section 3.2): An amide having the general structure $RCONR'_2$.

Tertiary (3°) amine (Section 3.2): An amine having the general structure R_3N.

Tertiary (3°) carbocation (Section 7.13): A carbocation having the general structure R_3C^+.

Tertiary (3°) carbon (Section 3.2): A carbon atom that is bonded to three other carbon atoms.

Tertiary (3°) hydrogen (Section 3.2): A hydrogen that is attached to a 3° carbon.

Tertiary protein structure (Section 29.9C): The three-dimensional shape adopted by an entire peptide chain.

Tertiary (3°) radical (Section 15.1): A radical having the general structure $R_3C\cdot$.

Tesla (Section 14.1A): A unit used to measure the strength of a magnetic field. Tesla is denoted with the symbol "T."

Tetramethylsilane (Section 14.1B): An internal standard used as a reference in NMR spectroscopy. The tetramethylsilane (TMS) reference peak occurs at 0 ppm on the δ scale.

Tetrasubstituted alkene (Section 8.2A): An alkene that has four alkyl groups and no hydrogens bonded to the carbons of the double bond ($R_2C{=}CR_2$).

Tetraterpene (Section 31.7A): A terpene that contains 40 carbons and eight isoprene units. A tetraterpenoid contains at least one oxygen atom as well.

Tetrose (Section 28.2): A monosaccharide containing four carbons.

Thermodynamic enolate (Section 23.4): The enolate that is lower in energy—generally the more substituted enolate.

Thermodynamic product (Section 16.11): In a reaction that can give more than one product, the product that predominates at equilibrium.

Thermodynamics (Section 6.5): A study of the energy and equilibrium of a chemical reaction.

Thermoplastics (Section 30.7): Polymers that can be melted and then molded into shapes that are retained when the polymer is cooled.

Thermosetting polymer (Section 30.7): A complex network of cross-linked polymer chains that cannot be re-melted to form a liquid phase.

Thioester (Section 22.17): A compound with the general structure RCOSR'.

Thiol (Section 9.15): A compound having the general structure RSH.

Tollens reagent (Sections 20.8, 28.9B): A reagent that oxidizes aldehydes, and consists of silver(I) oxide in aqueous ammonium hydroxide. A Tollens test is used to detect the presence of an aldehyde.

***p*-Toluenesulfonate** (Section 9.13): A very good leaving group having the general structure $CH_3C_6H_4SO_3^-$ and abbreviated as TsO^-. Compounds containing a *p*-toluenesulfonate leaving group are called alkyl tosylates and are abbreviated ROTs.

Torsional energy (Section 4.9): The energy difference between the staggered and eclipsed conformations of a molecule.

Torsional strain (Section 4.9): An increase in the energy of a molecule caused by eclipsing interactions between groups attached to adjacent carbon atoms.

Tosylate (Section 9.13): A very good leaving group having the general structure $CH_3C_6H_4SO_3^-$, and abbreviated as TsO^-.

***s*-Trans** (Sections 16.6, 29.5B): The conformation of a 1,3-diene that has the two double bonds on opposite sides of the single bond that joins them.

Trans diaxial (Section 8.8B): In an elimination reaction of a cyclohexane, a geometry in which the β hydrogen and the leaving group are trans with both in the axial position.

Trans isomer (Sections 4.13B, 8.3B): An isomer of a ring or double bond that has two groups on opposite sides of the ring or double bond.

Transition state (Section 6.7): An unstable energy maximum as a chemical reaction proceeds from reactants to products. The transition state is at the top of an energy "hill" and can never be isolated.

Triacylglycerol (Sections 10.6, 22.12A, 31.3): A lipid consisting of the triester of glycerol with three long-chain fatty acids. Triacylglycerols are the lipids that comprise animal fats and vegetable oils. Triacylglycerols are also called triglycerides.

Triose (Section 28.2): A monosaccharide containing three carbons.

Triphosphate (Section 7.16): A good leaving group used in biological systems. Triphosphate ($P_3O_{10}^{5-}$) is abbreviated as PPP_i.

Triplet (Section 14.6): An NMR signal that is split into three peaks having a relative area of 1:2:1, caused by two nearby nonequivalent protons.

Trisubstituted alkene (Section 8.2A): An alkene that has three alkyl groups and one hydrogen bonded to the carbons of the double bond ($R_2C{=}CHR$).

Triterpene (Section 31.7A): A terpene that contains 30 carbons and six isoprene units. A triterpenoid contains at least one oxygen atom as well.

U

Ultraviolet (UV) light (Section 16.15): Electromagnetic radiation with a wavelength from 200–400 nm.

Ultraviolet (UV) spectrum (Section 16.15): A plot of the absorbance of ultraviolet light versus wavelength, often recorded for conjugated systems.

Unimolecular reaction (Sections 6.9B, 7.10, 7.12A): A reaction that has only one reactant involved in the rate-determining step, so the concentration of only one reactant appears in the rate equation.

α,β-Unsaturated carbonyl compound (Section 20.15): A conjugated compound containing a carbonyl group and a carbon–carbon double bond separated by a single σ bond.

Unsaturated fatty acid (Section 10.6A): A fatty acid having one or more carbon–carbon double bonds in its hydrocarbon chain. In natural fatty acids, the double bonds generally have the *Z* configuration.

Unsaturated hydrocarbon (Section 10.2): A hydrocarbon that has fewer than the maximum number of hydrogen atoms per carbon atom. Hydrocarbons with π bonds or rings are unsaturated.

Unsymmetrical ether (Section 9.1): An ether in which the two alkyl groups bonded to the oxygen are different.

Upfield shift (Section 14.1B): In an NMR spectrum, a term used to describe the relative location of an absorption signal. An upfield shift means a signal is shifted to the right in the spectrum to lower chemical shift.

Urethane (Section 30.6C): A compound that contains a carbonyl group bonded to both an OR group and an NHR (or NR_2) group. A urethane is also called a carbamate.

V

Valence bond theory (Section 17.9A): A theory that describes covalent bonding as the overlap of two atomic orbitals with the electron pair in the resulting bond being shared by both atoms.

Valence electrons (Section 1.1): The electrons in the outermost shell of orbitals. Valence electrons determine the properties of a given element. Valence electrons are loosely held and participate in chemical reactions.

van der Waals forces (Section 3.3B): Very weak intermolecular interactions caused by momentary changes in electron density in molecules. The changes in electron density cause temporary dipoles, which are attracted to temporary dipoles in adjacent molecules. van der Waals forces are also called London forces.

Vicinal dihalide (Section 8.10): A compound that has two halogen atoms on adjacent carbon atoms.

Vinyl group (Section 10.3C): An alkene substituent having the structure $-CH=CH_2$.

Vinyl halide (Section 7.1): A molecule containing a halogen atom bonded to the sp^2 hybridized carbon of a carbon–carbon double bond.

Vitamins (Sections 3.5, 31.5): Organic compounds needed in small amounts by biological systems for normal cell function.

VSEPR theory (Section 1.7B): Valence shell electron pair repulsion theory. A theory that determines the three-dimensional shape of a molecule by the number of groups surrounding a central atom. The most stable arrangement keeps the groups as far away from each other as possible.

W

Walden inversion (Section 7.11C): The inversion of a stereogenic center involved in an S_N2 reaction.

Wavelength (Section 13.5): The distance from one point of a wave to the same point on the adjacent wave. Wavelength is abbreviated with the Greek letter lambda (λ).

Wavenumber (Section 13.6A): A unit for the frequency of electromagnetic radiation that is inversely proportional to wavelength. Wavenumber, reported in reciprocal centimeters (cm^{-1}), is used for frequency in IR spectroscopy.

Wax (Sections 4.15, 31.2): A hydrolyzable lipid consisting of an ester formed from a high molecular weight alcohol and a fatty acid.

Williamson ether synthesis (Section 9.6): A method for preparing ethers by reacting an alkoxide (RO^-) with a methyl or primary alkyl halide.

Wittig reaction (Section 21.10): A reaction of a carbonyl group and an organophosphorus reagent that forms an alkene.

Wittig reagent (Section 21.10A): An organophosphorus reagent having the general structure $Ph_3P=CR_2$.

Wohl degradation (Section 28.10A): A reaction that shortens the carbon chain of an aldose by removing one carbon from the aldehyde end.

Wolff–Kishner reduction (Section 18.15B): A method to reduce aryl ketones to alkyl benzenes using hydrazine (NH_2NH_2) and strong base (KOH).

Woodward–Hoffmann rules (Section 27.3): A set of rules based on orbital symmetry used to explain the stereochemical course of pericyclic reactions.

Y

Ylide (Section 21.10A): A chemical species that contains two oppositely charged atoms bonded to each other, and both atoms have octets of electrons.

Z

Zaitsev rule (Section 8.5): In a β elimination reaction, a rule that states that the major product is the alkene with the most substituted double bond.

Ziegler–Natta catalysts (Section 30.4): Polymerization catalysts prepared from an organoaluminum compound and a Lewis acid such as $TiCl_4$, which afford polymer chains without significant branching and with controlled stereochemistry.

Zwitterion (Sections 19.14B, 29.1B): A neutral compound that contains both a positive and negative charge.

This page intentionally left blank

Credits

Front Matter

Page iii: © Daniel C. Smith. Table of Contents 1: © Walter B. McKenzie/ Getty Images RF; 2: © Jill Braaten; 3: © S. Aubert/Univ.Grenoble; 4: © Vince Bevan/Alamy; 5: © George Ostertag/Alamy RF; 6: © ninikas/Getty Images RF; 7: © McGraw-Hill Education/Mark Dierker, photographer; 8: © Comstock/ PunchStock RF; 9: © Daniel C. Smith; 10: © Inga Spence/Science Source; 11: © McGraw-Hill Education/Chris Kerrigan, photographer; 12: © Royalty Free/ Corbis; 13: Library of Congress; 14: © Daniel C. Smith; 15: © Jamie Grill/Getty Images RF; 16: © John Foxx/Getty Images RF; 17: © iodrakon/Getty Images RF; 18: © Jill Braaten; 19: © Sarka Babicka/Getty Images RF; 20: © Daniel C. Smith; 21: © Richo Cech, Horizon Herbs, LLC.; 22: © Jill Braaten; 23: © iStock/ Getty Images Plus RF; 24: © Jill Braaten; 25: © James Forte/Getty Images RF; 26: © De Agostini Picture Library/Getty Images; 27: © Nature Picture Library/ Alamy; 28: © Phil Degginger/Alamy; 29: © Royalty-Free/Corbis; 30: © Stuart Burford/Alamy; 31: © Jill Braaten.

Prologue

Figure 1a: © Comstock Images/PictureQuest RF; 1b: © Corbis Premium RF/ Alamy; 1c: © Steven May/Alamy; 1d: © Layne Kennedy/Corbis; 2(quinine): US Geological Survey-Photo by Forest&Kim Starr; (chloroquine): © Chris Howes/Wild Places Photography/Alamy; (artemisinin): © Gregory G. Dimijian, M.D./Science Source.

Chapter 1

Opener: © Walter B. McKenzie/Getty Images RF; p. 11: © Jill Braaten: p. 19(both): © Steve Gschmeissner/Science Source; p. 30: Courtesy John Somerville; p. 34(top): © vast natalia/Alamy RF; (bottom): © McGraw-Hill Education/Elite Images; p. 40: © Corbis RF; p. 41: © Arthur Tilley/Getty Images RF; p. 42: © Phillip Spears/Getty Images RF; p. 45: © McGraw-Hill Education/Mark Dierker, photographer; p. 50: © Cristina Pedrazzini/Science Source.

Chapter 2

Opener: © Jill Braaten; p. 62: © Bon Appetit/Alamy; p. 72&74: © McGraw-Hill Education/Jill Braaten, photographer; p. 79: © Wilmer Stratton; 2.7: © National Geographic/Getty Images.

Chapter 3

Opener: © S. Aubert/Univ.Grenoble; p. 95: © Michael Pettigrew/Getty Images RF; p. 96: Courtesy of Gebauer Company, Cleveland, Ohio; p. 97&99: © Jill Braaten; p. 100(top): © Andrew Syred/Science Source; (bottom): © Martin Harvey/Corbis; p. 110(top): © Comstock/PunchStock RF; (bottom): © McGraw-Hill Education/ Mary Reeg, photographer; p. 111: © Digital Vision/Getty RF.

Chapter 4

Opener: © Vince Bevan/Alamy; p. 129(top): © God of Insects; (bottom): © Douglas Peebles/Corbis; p. 132: © Daniel C. Smith; p. 133: © Jill Braaten; 4.5: © Glow Images RF; p. 142: © McGraw-Hill Education/Jill Braaten, photographer; p. 164: © Riorita/Getty Images RF; p. 165: © Eric Couture, courtesy Canadian Shark Research Laboratory.

Chapter 5

Opener: © George Ostertag/Alamy RF; 5.1(starch): © Jim Erickson/Corbis; (wheat): © Pixtal/age fotostock RF; (cotton plant): © Corbis RF; (cotton fabric): © Daniel C. Smith; p. 178: © Daniel C. Smith; p. 180: © McGraw-Hill Education/ Jill Braaten, photographer; p. 181: © Mark W. Skinner; p. 184: © Shallenberger Photography; p. 190: © Robert Brook/Science Source; p. 195: © McGraw-Hill Education/Jill Braaten, photographer; p. 203&p. 204(caraway): © McGraw-Hill Education/Elite Images; (spearmint): © DAJ/Getty Images RF; (celery): © Aaron Roeth Photography RF.

Chapter 6

Opener: © ninikas/Getty Images RF; p. 216: © Getty Images RF; p. 234(top): © Corbis. All Rights Reserved; (bottom): © McGraw-Hill Education/Jill Braaten, photographer.

Chapter 7

Opener: © McGraw-Hill Education/Mark Dierker, photographer; p. 252: © Michael Guiry; p. 255&284: © Jill Braaten; p. 287: © McGraw-Hill Education/Mark Dierker, photographer.

Chapter 8

Opener: © Comstock/PunchStock RF; p. 300: © McGraw-Hill Education/Jill Braaten, photographer; p. 301: © Steven P. Lynch RF; p. 307: © Forest&Kim Starr.

Chapter 9

Opener: © Daniel C. Smith; p. 333: © Peter J.S. Franks, Scripps Institution of Oceanography; 9.4(left): © Pixtal/age fotostock RF; (right): Photo by Mark Wilson/Getty Images; p. 352: © Henriette Kress; p. 365: © Purestock/SuperStock RF; p. 371: © McGraw-Hill Education/Jill Braaten, photographer; p. 373: © McGraw-Hill Education/John Thoeming, photographer.

Chapter 10

Opener: © Inga Spence/Science Source; p. 390: © Daniel C. Smith; 10.5(carrots): © Corbis RF ; (ginger): © McGraw-Hill Education/Jill Braaten, photographer; (apples): © Pixtal/AGE Fotostock RF; (oranges): USDA; p. 393&395: © Jill Braaten; p. 405: © Andrew Lambert Photography/Science Source; p. 415: © Henriette Kress.

Chapter 11

Opener: © McGraw-Hill Education/Chris Kerrigan, photographer; 11.3: © Michael SewelL/Getty Images; p. 443: © Michael G. Moye.

Chapter 12

Opener: © Corbis RF; p. 462(peanut butter&label): © McGraw-Hill Education/Elite Images; (vegetable oil&butter): © McGraw-Hill Education/Jill Braaten, photographer; p. 464: Charlotte BjÃ¶rnstrÃ¶m/EyeEm/Getty Images RF; p. 465; © 1998 Richard Megna, Fundamental Photographers, NYC; p. 470: USDA APHIS PPQ Archive, USDA APHIS PPQ, Bugwood.org; p. 474: © Balazs Kovacs/Getty Images RF; 12.9b: © McGraw-Hill Education/Elite Images; p. 483: © Image Source RF.

Chapter 13

Opener: Library of Congress; p. 500: © MizC/Getty Images RF; p. 505: © Ted Kinsman/Science Source; p. 517: © Andrew McClenaghan/Science Source.

Chapter 14

Opener: © Daniel C. Smith; p. 558: © Brand X Pictures/PunchStock RF; 14.15a: © Peter Noyce PLB/Alamy RF; 14.15b: With permission, Daniel C. Smith. Spectra courtesy of the Chemistry Department at Rutgers University.

Chapter 15

Opener: © Jamie Grill/Getty Images RF; p. 584(top): © McGraw-Hill Education/ Jill Braaten, photographer; (bottom): NASA Ozone Watch; p. 589: © McGraw-Hill Education/Elite Images; p. 590(top): © Stockbyte/Corbis RF; (bottom): © Pixtal/SuperStock RF; p. 593: © McGraw-Hill Education/Jill Braaten, photographer; p. 594(top): © Image Source/Getty RF; (middle): © Corbis RF; (bottom): © Jill Braaten.

Chapter 16

Opener: © John Foxx/Getty Images RF; p. 614: © Franz-Marc-Frei/Corbis; p. 622: © Stephen Frink/Getty Images RF; p. 623: © Alvis Upitis/Getty Images RF; p. 631: © C Squared Studios/Getty Images RF; p. 632: © McGraw-Hill Education/Jill Braaten, photographer.

Chapter 17

Opener: © iodrakon/Getty Images RF; 17.3(left): © C. Zachariasen/PhotoAlto RF; (right): Photo by Bob Nichols, USDA Natural Resources Conservation Service; p. 657: © Daniel C. Smith; p. 658(top): © Forest&Kim Starr; (bottom): © Bloomberg/Getty Images; p. 666(top): © Charles O'Rear/Corbis; (bottom): © Lester V. Bergman/Corbis; p. 667: © Photodisc/Getty Images RF.

Chapter 18

Opener: © Jill Braaten; p. 682: © Time Life Pictures/Getty Images; p. 689: © Rene Dulhoste/Science Source; p. 690: © Bill Aron/Photo Edit, Inc.; p. 714: © McGraw-Hill Education/Jill Braaten, photographer.

Chapter 19

Opener: © Sarka Babicka/Getty Images RF; p. 732: © Dr. Steven J. Wolf; p. 737(top): © Ryman Cabannes/Photocuisine/Corbis; p. 737(bottom): © Biopix.dx http://www.biopix.dk; p. 749: © Ken Samuelsen/Getty Images RF; p. 752: © FoodCollection RF

Chapter 20

Opener: © Daniel C. Smith; p. 766: © Gerald D. Carr, PhD; p. 773: © M.K. Ranjitsinh/Science Source; p. 776: © John Kaprielian/Science Source; p. 777: © C Squared Studios/Getty Images RF; p. 782: © McGraw-Hill Education/Charles D. Winters, photographer p. 789: © George Grall/National Geographic/Getty Images.

Chapter 21

Opener: © Richo Cech, Horizon Herbs, LLC; 21.6&p. 834(top): © McGraw-Hill Education/Jill Braaten, photographer; (bottom): © Daniel C. Smith; p. 851(both): © iStock/Getty Images RF.

Chapter 22

Opener&p. 880(top): © Jill Braaten; (bottom): © blickwinkel/Alamy RF; p. 885: © McGraw-Hill Education/Joe Franek, photographer; p. 886: Photo by Scott Bauer, USDA-ARS; p. 887: © Coby Schal/NC State University; p. 897: © McGraw-Hill Education/Jill Braaten, photographer p. 898: © Jill Braaten; p. 902: © Jeff Morgan 14/Alamy; p. 903: © Jill Braaten.

Chapter 23

Opener: © iStock/Getty Images Plus RF; p. 932&939: © McGraw-Hill Education/Joe Franek, photographer; p. 944: © McGraw-Hill Education/Mary Reeg, photographer; p. 951: © iStock/Getty Images Plus RF.

Chapter 24

Opener: © Jill Braaten; p. 972(cockroach): © James H. Robinson/Science Source; (tumeric)& p. 973(Donepezil): © Jill Braaten; p. 978: © McGraw-Hill Education/Jill Braaten, photographer.

Chapter 25

Opener: © James Forte/Getty Images RF; 25.6(nightshade): © Werner Arnold; (tobacco): © Creatas Images RF; (hemlock): © Steven P. Lynch RF; p. 1006: © Daniel C. Smith; p. 1016: © McGraw-Hill Education/Jill Braaten, photographer; p. 1022(top): James Gathany/CDC; (bottom): © Daniel C. Smith; p. 1036(indigo): © Kirsten Soderlind/Corbis; (sea snail): © Andrew J. Martinez/Science Source; (madder): © Bob Gibbons/Alamy; (purple shawl): © Science&Society Picture Library.

Chapter 26

Opener: © De Agostini Picture Library/Getty Images; p. 1055: © Pascal Goetgheluck/Science Source; p. 1059: © McGraw-Hill Education/John Thoeming, photographer; p. 1066: © Ross Warner/Alamy.

Chapter 27

Opener: © Nature Picture Library/Alamy; p. 1086: © McGraw-Hill Education/Mary Reeg, photographer; p. 1090: © blickwinkel/Alamy; p. 1097: © Marina Khaytarova TopTropicals.com.

Chapter 28

Opener: © Phil Degginger/Alamy; p. 1108(top): © McGraw-Hill Education/Jill Braaten, photographer; (bottom): © McGraw-Hill Education/Elite Images; p. 1123: © Brand X Pictures/PunchStock RF; p. 1126: © Nature Picture Library/Alamy; p. 1127: © inga spence/Alamy; p. 1135: © Vol. 3/PhotoDisc/Getty RF; p. 1136: © McGraw-Hill Education/Elite Images; 28.9(left): © McGraw-Hill Education/Elite Images; (right): © David Muench/Corbis; 28.10: © McGraw-Hill Education/Jill Braaten, photographer; p. 1140(top): © McGraw-Hill Education/Jill Braaten, photographer; (bottom): © Comstock Images/PictureQuest RF.

Chapter 29

Opener: © Corbis RF; p. 1155: © Jill Braaten; p. 1188: © Eye of Science/Science Source.

Chapter 30

Opener: © Stuart Burford/Alamy; 30.1: © Getty Images; 30.1: © Getty Images; 30.4(balls): © Dynamicgraphics/JupiterImages RF; (paints): © Beathan/Corbis RF; (fibers): © Fernando Bengoe/Corbis; (Super glue): © McGraw-Hill Education/John Thoeming, photographer; p. 1207: © Savushkin/Getty Images RF; p. 1209: © DSM Dyneema; p. 1210: © Bob Krist/Corbis; p. 1212: © McGraw-Hill Education/John Thoeming, photographer ; p. 1217: © Rim Light/PhotoLink/Getty RF; 30.9(left): © Morey Milradt/Brand X/Corbis RF; (right): © E.I. du Pont de Nemours and Company.

Index

Page numbers followed by f indicate figures; those followed by t indicate tables. An A– before page numbers indicates appendix pages.

A

ABS, 1207
Abyssomicin C, 995
Acebutolol, 125
Acetal(s)
  conversion to hemiacetals, 850–851
  cyclic, 847
    in carbohydrates, 855–856
    conversion of cyclic hemiacetal to, 854–855
  ether compared, 847
  glycosides, 1124–1127
  hemiacetals compared to, 1124
  hydrolysis of, 850–851
  as protecting groups, 851–852
  structure of, 1124
  synthesis of
    addition of alcohols to aldehydes and ketones, 847–851
    from hemiacetals, 848–849, 854–855
  terminology of, 847
Acetaldehyde
  aldol reaction of, 963, 966, 968
  ethanol oxidized to, 481–482
  hydrate formed from, 845
  structure of, 820
Acetamide, 23–24, 59, 89
  structure of, 869, 874, 877, 877f
Acetaminophen, 59, 888, 900, 919, 1046
Acetanilide, 760, 1024, 1024f
Acetate
  bond length in, 615
  formation from acetaldehyde oxidation, 481
  as nucleophile, 262–264, 269
  resonance structures of, 74–75, 609, 742, 744f
Acetic acid, 736
  acidity of, 74–75, 77, 79, 742, 744f, 745
  as Brønsted–Lowry acid, 62f
  deprotonation of, 741
  dimers, 734f
  electrostatic potential plot of, 75f, 730, 730f
  in ethyl acetate formation, 236–237
  glacial, 736
  halogenation of carbonyl compounds in, 938
  melting point of, 106
  NMR spectrum of, 556
  pK_a of, 741
  as polar protic solvent, 259f
  structure of, 732t, 869
Acetic anhydride, 635, 869, 874, 1128
Acetic benzoic anhydride, 874
Acetoacetic ester synthesis, 945, 949–952, 979
Acetone
  acidity of, 75
  aldol reactions of, 968
  butan-2-one synthesis, 951
  dipoles in adjacent molecules, 101
  hydrate formed from, 845
  odor of, 825
  as polar aprotic solvent, 260, 260f
  solubility of, 107
  structure and bonding of, 44, 598, 821, 825
  surface area and boiling point of, 103, 104f
  uses of, 825
Acetone enolate, 930f, 936f
Acetonitrile, 58
  acidity of, 75
  deprotonation of, 69
  as polar aprotic solvent, 260f
  structure of, 875f
p-Acetophenetidin, 919
Acetophenone, 775
  aldol reaction of, 965
  structure of, 821
N-Acetyl amino acids, 1160–1162
Acetylation, 888
Acetyl chloride, 244, 1128
  structure of, 869, 873
Acetylcholine, 904
Acetyl coenzyme A, 904, 906, 1252
1-Acetylcyclohexene, 955
2-Acetylcyclopentanone, 993
1-Acetylcyclopentene, 991
Acetylene
  acetylide anion formation from, 446–447
  in acid–base reactions, 68–69
  acidity of, 75–76, 76f
  bond angles, 427
  bond dissociation energies of, 427
  bonds in, 42–43, 43f, 44t
  combustion of, 430
  deprotonation of, 446
  disparlure synthesis from, 471, 471f
  electrostatic potential map of, 432, 433f
  functional group in, 93, 94t
  Lewis structure, 16
  molecular shape of, 27
  nomenclature, 429
N-Acetyl-D-glucosamine (NAG), 906, 1140
Acetyl group, 1128
  abbreviation for, 1053
  structure, 822
Acetylide anions
  conversion of terminal alkynes to, 441
  formation by deprotonation of terminal alkynes, 433
  as organometallic compounds, 784
  reactions with alkyl halides, 441–444, 443f
  reactions with epoxides, 444
  retrosynthetic analysis with, 446–447
  synthesis from acetylene, 446–447
  in synthesis of disparlure, 471, 471f
Acetylsalicylic acid, 61, 737, 888, 900. *See also* Aspirin
Achiral enolate, 936
Achiral epoxides, 472
Achiral intermediate, 406
Achiral meso compound, 193, 195, 407, 408f, 470, 624, 1060
Achiral molecules
  definition of, 178–180, 180f
  halogenation of, 582, 582t
  optical inactivity of, 198
Achiral product, 402
Achiral reactant, 368, 402
Achiral starting material, 470, 482, 582, 773, 788
Acid(s)
  amine reactions with, 1014–1016
  amino. *See* Amino acids
  Brønsted–Lowry
    carboxylic acids, 740–745
    description of, 62–63
    reactions of, 63–65
  carboxylic. *See* Carboxylic acid(s)
  as catalysts, 236–237
  common, 78–79
  conjugate, 63–65, 256–258, 256t
  dehydration of alcohols to alkenes, 345–350
  as electron pair acceptors, 62, 65, 81
  in epoxide ring opening reactions, 369–371, 370f
  halogenation in, 938, 940f
  Lewis, 81–83
  origin of word, 62
  as proton donors, 62, 69
  reaction of ethers with strong, 362–363
  strength, 66–78
  sulfonic, 751–752
Acid–base reactions. *See also* Nucleophilic substitution reactions
  amines, 1014–1016
  Brønsted–Lowry, 63–65, 254, 272
  of carboxylic acids, 889–894
  description, 63–65
  Lewis, 82–83, 254
  organometallic reagents and, 784–785
  predicting the outcome of, 68–69
Acid chlorides
  acidity/basicity of, 872, 872t
  boiling point of, 878t
  functional group in, 98t
  IR spectra of, 878, 879t
  leaving group, 768, 796
  melting point of, 878t
  nomenclature of, 873, 876t
  nucleophilic substitution reactions, 768
  odor of, 885
  physical properties of, 878t
  pK_a, 872t
  reactions of, 885–887, 901t
  with ammonia or amines to form amides, 1023
  conversion to amides, 886
  conversion to anhydrides, 885, 886
  conversion to carboxylic acids, 885, 887
  conversion to esters, 885
  Friedel–Crafts acylation reactions, 684
  with organometallic reagents, 785, 795–797
  reactivity, 883
  reduction of, 777–778
  solubility of, 878t
  stability of, 872
  structure and bonding of, 765, 869, 872
  synthesis from carboxylic acids, 889f, 890
Acidity
  basicity, inverse relationship with, 67, 67t
  of carbon–hydrogen bonds, 71
  of carboxylic acids, 740, 741–746, 744f
  determining the relative acidity of protons, 77–78
  electronegativity and, 70–73
  factors affecting, 70–78
    element effects, 70–72, 77f
    hybridization effects, 75–76, 76f, 77f
    inductive effects, 72–74, 73f, 77f
    resonance effects, 74–75, 75f, 77f
  periodic trends in, 70–72
Acidity constant (K_a), 66
Acid rain, 79
Acrolein, 624f
Acrylic acid, 594
Acrylonitrile, 563, 1205f
Activation energy. *See* Energy of activation (E_a)
Active methylene compounds, 970, 981
Active site, enzyme, 238, 238f
Acyclic alkanes
  branched-chain, 130
  conformations of, 144–151
    butane, 148–151, 148f, 150f
    ethane, 144–147, 146f, 147f
  constitutional isomers, 130–132, 131t
  having more than five carbon atoms, 131–132, 131t
  having one to five carbon atoms, 129–131, 131t
  homologous series, 131, 131t
  molecular formula, 129, 131, 131t
  naming, 135–138, 137f
  normal (*n*) (straight-chain), 130–131, 131t
  structure, 129–130
  torsional and steric strain energies in, 150t
Acyclovir, 955
Acylases, 1162

Acylation
  definition of, 684
  Friedel–Crafts. *See* Friedel–Crafts acylation
Acyl derivatives, 765
Acyl group, 684, 869, 873, 874
  naming, 822
Acylium ion, 503, 686
Acyl transfer reactions, 904–905, 905f
Addition polymers. *See* Chain-growth polymers
Addition reactions
  in alkenes, 384, 396–418, A–13
    anti addition, 397, 403, 403t, 405, 406, 409
    halogenation, 397f, 405–407, 408f
    halohydrin formation, 397f, 408–410, 410t
    hydration, 397f, 404–405, 415
    hydroboration–oxidation, 397f, 411–415, 412f, 413f, 414t, 415f
    hydrohalogenation, 397–404, 397f, 399f, 401f, 403t
    overview of, 396–397, 397f
    stereochemistry of, 397, 402–404, 403t
    syn addition, 397, 403, 403t, 411, 413
  allylic bromination, 587
  anti additions, 463, 465, 472, 473
  carbene addition, 1060
  conjugate, 981, 983
  in conjugated dienes
    Diels–Alder reaction, 621–630, 622f
    electrophilic, 616–618
    kinetic versus thermodynamic product, 618–621
  description of, 216
  dihydroxylation, 472–474
  epoxidation, 469–472, 482–484
  exothermic reactions, 398, 398f
  ionic addition, 592
  in nucleophilic aromatic substitution, 706–709
  oxidative, 1053, 1055, 1058
  radical reactions, 587, 590–593
  syn additions, 458, 460, 463, 469, 471, 471f, 472, 473–474, 1060
Adenine, 1142, 1143f
Adenosine triphosphate (ATP), 284, 366
S-Adenosylmethionine (SAM), 284–285, 366
-adiene (suffix), 387
Adipic acid, 1211, 1218, 1218f, 1219
Adipoyl chloride, 902
Adrenal cortical steroids, 1253, 1255
Adrenaline
  structure of, 1005
  synthesis of, 285–286, 285f
Advair. *See* Fluticasone
Advil. *See* Ibuprofen
Aerosol propellant, 252, 584
Agent Orange, 682f
Aglycon, 1127
-al (suffix), 819
Alanine, 118
  abbreviation for, 753t
  acid–base reactions of, 753–754, 754f
  in dipeptide synthesis, 1172–1173, 1176
  enantiomers of, 183f
    N-acetyl, 1160–1162
  isoelectric point of, 755, 1156t
  in peptide formation, 1165
  physical characteristics, 200
  pK_a values for, 1156t
  structure of, 1154f
  synthesis of, 1158
Albuterol, 181–182, 371, 371f, 1048
Alcohol dehydrogenase, 481–482
Alcoholic beverages, 338, 455
Alcohols, 331–382
  alkene preparation by acid-catalyzed dehydration, 395
  allylic, epoxidation of, 482–484
  boiling point of, 338t
  bonding in, 333
  classification of, 96, 332
  dehydration reactions of, 345–352
    alkene synthesis in, 345–352
    carbocation rearrangements in, 348–350
    dehydration to alkenes, 345–352, 348f
    E1 mechanism in, 346–347
    E2 mechanism in, 347, 351
    enthalpy change in, 347, 348f
    Le Châtelier's principle in, 347
    in phosphorus oxychloride and pyridine, 351, 352f
    in strong acid, 345–350
  electrostatic potential maps, 333f
  elimination reactions, 343–352, 361, 362f
  ether formation from, 406
  fragmentation pattern in mass spectrum, 503
  Friedel–Crafts alkylation of, 688
  functional group, 93, 96t
  β-hydroxy carbonyl compounds, 963–975
  infrared (IR) spectra of, 514
  interesting examples, 338–339, 339f
  leaving groups of, 344, 346–347, 349–354, 356–361
  melting point of, 338t
  NMR spectra of, 551–552
  nomenclature, 334–335
    common names, 335
    for cyclic alcohols, 334, 334f
    IUPAC system, 334, 335
  oxidation of, 468f, 476–482
  physical properties of, 337–338, 338t
  as polar protic solvents, 259, 259f
  primary (1°), 415
    classification as, 332
    conversion to alkyl halides with HX, 352–354, 357t
    conversion to alkyl halides with phosphorus tribromide, 357, 357t
    conversion to alkyl halides with thionyl chloride, 356, 357t
    dehydration by E2 mechanism, 347
    hydrogen bonding extent, 337
    from nucleophilic addition to aldehyde, 778
    oxidation to carboxylic acids, 477, 478–479, 739
    from reactions of aldehydes with organometallic reagents, 785–787
    from reduction of acid chlorides, 777–778
    from reduction of aldehydes, 461, 770, 771, 772
    from reduction of carboxylic acid derivatives, 770
    from reduction of carboxylic acids, 780
    from reduction of esters, 777–778
    from reduction of monosaccharides, 1129
  products of S_N2 substitution reactions, 286, 287t
  racemic mixture of, 414
reactions
  conversion to alkyl halides with HX, 352–355, 357t
    conversion to alkyl halides with thionyl chloride and phosphorus tribromide, 356–357, 357t
    conversion to alkyl tosylates, 359
    elimination reactions, 343–352
    general features, 343–344
    nucleophilic substitution reactions, 344, 352–361, 362f
    summary of, 361, 362f
  secondary (2°), 415
    classification as, 332
    conversion to alkyl halides with HX, 352–355, 357t
    conversion to alkyl halides with phosphorus tribromide, 357, 357t
    conversion to alkyl halides with thionyl chloride, 356, 357t
    dehydration by an E1 mechanism, 346–347
    hydrogen bonding extent, 337
    oxidation to ketones, 477–478
    from reactions of aldehydes with organometallic reagents, 785–787
    from reduction of ketones, 461, 770, 771, 772
  solubility, 338t
  structure of, 332, 333
  substitution reactions, 343–344, 352–361, 362f
  synthesis of, 341–342
    from alkyl halides, 341–342
    hydration of alkenes, 404–405
    hydroboration–oxidation of alkenes, 411–415
    by nucleophilic substitution reaction, 341–342
    reactions of organometallic reagents with aldehydes and ketones, 785–787, 832–833
    reactions of organometallic reagents with epoxides, 798–799
    reduction of epoxides, 466–467, 467f
  tertiary (3°)
    classification as, 332
    conversion to alkyl halides with HX, 352–355, 357t
    dehydration by an E1 mechanism, 346–347
    hydrogen bonding extent, 337
    from reaction of esters and acid chlorides with organometallic reagents, 794–796
    from reaction of ketones with organometallic reagents, 785–787
Aldaric acids, 1129, 1130, 1133–1134
  D-aldaric acids, 1133
-aldehyde (suffix), 820
Aldehyde dehydrogenase, 481–482
Aldehyde group, as electron-withdrawing group, 692–693
Aldehydes. *See also* Carbonyl compounds
  aldol reactions of, 963–975
  boiling point of, 823t
  carbohydrates, 855–856
  electrostatic potential map of, 766, 766f
  fragmentation pattern in mass spectrum, 503
  functional group in, 98t
  hydration of, 844–847
    kinetics of hydrate formation, 846–847
    thermodynamics of hydrate formation, 845
  hydroxy
    cyclization of, 853, 855–856
    β-hydroxy, synthesis in aldol reactions, 963–975
  interesting examples, 825–826, 826f
  IR spectra, 823–824, 823f, 824f
  melting point of, 823t
  NMR spectra, 825, 825f
  nomenclature, 819–820, 822
    common names, 820, 820f
    IUPAC system, 819, 820f
  nucleophilic addition reactions, 767, 772, 778, 817–867
    carbanion addition, 832
    cyanide addition, 833–834
    hydride addition, 831–832
    mechanism of, 829–830
    nucleophiles in, 830, 831f
    primary amine addition, 840–842, 843f
    secondary amine addition, 842–844, 843f
  odors of, 826, 826f
  oxidation reactions, 770, 782
    to carboxylic acid, 1129–1130
  physical properties of, 822, 823t
  pK_a, 931t
  polyhydroxy, 855–856
  protecting groups for, 851–852
  reactions of
    with amines, 1023
    at the α carbon, 828–829
    at the carbonyl carbon, 828, 829
    halogenation at the α carbon, 937–941, 940f
    with organometallic reagents, 785–787, 832
  reactivity of, 767
  reduction of, 831–832
    to alcohols, 461, 770, 771, 772
  reductive amination to amines, 1011–1013
  solubility of, 823t
  structure of, 765, 818
  synthesis of, 826–828
    alcohol oxidation, 468f, 477, 478–479
    carboxylic acid reduction, 770
    hydroboration–oxidation of alkynes, 440
    hydroboration–oxidation of alkynes, 827
    hydrolysis of acetals, 850–851
    hydrolysis of imines and enamines, 844
    oxidation of primary alcohols, 827

oxidative cleavage of alkenes, 474–475, 827
reduction of acid chlorides, 777
reduction of acid chlorides and esters, 827
reduction of esters, 777
reduction of nitriles, 909
unreactivity to nucleophilic substitution, 769
Wittig reaction, 835–839
Alder, Kurt, 621
Alditol, 1129
Aldohexoses, 1109
D-aldohexoses, 1114, 1115f
Haworth projections of, 1119–1121
pyranose ring of, 1117
stereogenic centers of, 1111, 1114
Aldol reactions
crossed, 967–971
synthetically useful reactions, 968–970, 970f
useful transformations of aldol products, 970, 971f
dehydration of aldol product, 965–967
directed, 971–973, 972f
features of, 963–965, 964f
intramolecular, 973–975, 975f
in Robinson annulation, 982–986
mechanism of, 964
retrosynthetic analysis using, 966–967
Aldonic acid, 1129, 1130
D-Aldopentoses, 1114, 1115f, 1133
Aldoses
D-aldose family, 1113–1115, 1115*f*
in Kiliani–Fischer synthesis, 1131, 1132–1134
oxidation of, 1129–1131
reduction of, 1129
structure of, 1108–1109
in Wohl degradation, 1131–1132
Aldosterone, 1255
Aldotetroses
D-aldotetroses, 1114, 1114f, 1115f, 1133, 1134
stereoisomers of, 1114, 1114f
Aldrin, 637
Alendronic acid (Fosamax), 19
Aleve. *See* Naproxen
Aliphatic carboxylic acids, 744–746
Aliphatic hydrocarbons, 93, 129. *See also* Alkanes; Alkenes; Alkynes
Aliskiren, 182, 492
Alizarin, 1036
Alizarine yellow R, 1037
Alkaloids, 996, 1004, 1004f
Alkanes, 128–173
acyclic
branched-chain, 130
conformations of, 144–151, 146f, 147f, 148f, 150f
constitutional isomers, 130–132, 131t
having more than five carbon atoms, 131–132, 131t
having one to five carbon atoms, 129–131, 131t
homologous series, 131, 131t
molecular formula, 129, 131, 131t
normal (*n*) (straight-chain), 130–131, 131t
structure of, 129–130
torsional and steric strain energies in, 150t
boiling point of, 143, 143t
bonds in, 93
combustion of, 162–163
cycloalkanes
molecular formula, 129, 132
naming, 138–140, 140f
structure, 129, 132, 152f
substituted, 156–161, 157f, 160f
definition of, 129
fossil fuels, 141–142
halo, 249
halogenation, 574–583
of achiral starting materials, 582, 582t
bromination, 578–580, 580f
of chiral starting materials, 582t, 583
chlorination, 574–583, 581f
energy changes during, 576–577, 577f
mechanism of, 575–576
in organic synthesis, 581
stereochemistry of, 582–583, 582t
infrared (IR) spectra of, 513
melting point of, 143, 143t
nomenclature, 133–141, 137f, 140f
common names, 141, 141f
IUPAC system, 135–140, 137f
substituents, 134
nonreactivity of, 94
oxidation of, 161–163
physical properties, 143–144, 143t
radical reactions, 573
solubility of, 143, 143t
synthesis of
reduction of alkenes, 456, 456f, 458–461
reduction of alkyl halides, 466, 785
reduction of alkynes, 456f, 463–464, 466f
Alkene oxides, 336, 337
Alkenes, 383–425
addition reactions, 384, 396–418
anti addition, 397, 403, 403t, 405, 406, 409
halogenation, 397f, 405–407, 408f
halohydrin formation, 397f, 408–410, 410t
hydration, 397f, 404–405, 415
hydroboration–oxidation, 397f, 411–415, 412f, 413f, 414t, 415f
hydrohalogenation, 397–404, 397f, 398f, 399f, 401f, 403t
overview of, 396–397, 397f
stereochemistry of, 397, 402–404, 403t
syn addition, 397, 403, 403t, 406, 411, 413
amino acid synthesis from, 1163–1164
boiling point of, 391
bonds in, 93
classification of, 300, 300f
constitutional isomers of, 345
conversion to alkynes, 431–432
conversion to cyclopropanes, 1058–1061
in coupling reactions with organic halides (Heck reaction), 1050, 1056–1058
cycloaddition between, 1088
cyclopropane synthesis in Simmons–Smith reaction, 1061–1062
degrees of unsaturation, calculating, 385–387, 460–461
dehydration of alcohols to, 345–352
diastereomers of, 301
dihydroxylation of, 472–474
disubstituted, 300–303, 300f, 306, 308–309, 312, 316–317
double bonds in, 298–302, 384, 385t
restricted rotation around, 300–301, 301f
electrophiles, reactions with, 117
electrostatic potential plots, 396, 396f
epoxidation of, 468f, 469–470
fatty acids, 383, 393–395, 394f, 394t
Friedel–Crafts alkylation of, 688
functional group in, 93, 94t
hybrid orbitals in, 299–300, 302
hydrogenation of, 458–463
alkene stability and, 458–459, 459f
cis and trans isomers compared, 458, 459f
degrees of unsaturation, determination of, 460–461
mechanism of catalytic hydrogenation, 459–460
infrared (IR) spectra of, 513
interesting examples, 391, 392f
internal, 384
lipids, 393–395
melting point of, 391, 393, 394t
metathesis, 1062–1067, 1063f, 1065f
molecular formula of, 385
monomers in polymerization, 593–595, 594t
monosubstituted, 300, 300f, 306
NMR spectra of, 533, 549–551, 549f, 550f
nomenclature, 387–390
common names, 390, 390f
IUPAC system, 387–388
for stereoisomers, 388–390
organic synthesis, 417–418
oxidative cleavage of, 468f, 474–476, 827
physical properties of, 391
polymerization of, 1200–1207, 1208–1209
radical reactions, 573, 590–593
reduction of, 456, 456f, 458–461
solubility of, 391
stability of, 302–303, 458–459, 459f
stereoisomers of, 300–302
alkyne reduction and, 463–466
synthesis of, 395–396
amine conversion, 1024–1027
comparison of methods, 839
dehydration of alcohols, 395
dehydrohalogenation of alkyl halides with strong base, 395
by elimination reactions, 298–299, 395–396
by Hofmann elimination reaction, 1024–1027
reduction of alkynes to cis alkenes, 463, 464, 466f
reduction of alkynes to trans alkenes, 463, 465–466, 466f
substituted alkenes in Heck reaction, 1056–1057
Wittig reaction, 835–839
terminal, 384
tetrasubstituted, 300
trisubstituted, 300, 300f, 308–309, 312, 316–317
Alkenols, 388
Alkoxides, 298
electron density plot, 930, 930f
protonation of, 774, 786, 787, 790, 795, 799
Alkoxide salt, in ether synthesis, 342
Alkoxy group, 96t, 336
Alkylamines
basicity of, 1017–1019, 1021t
reaction with nitrous acid, 1028–1029
Alkylation reactions
in acetoacetic ester synthesis, 945, 949–952
definition of, 684
of a diethyl malonate derivative, 1157–1158, 1159f
of enolates, 942–945, 951
general features of, 942–943
tamoxifen synthesis, 944
of unsymmetrical ketones, 943–944
Friedel–Crafts. *See* Friedel–Crafts alkylation
in malonic ester synthesis, 945–949
polyalkylation, 702
Alkyl benzenes
electron density of, 692–693
halogenation of, 709–711
oxidation of, 712, 739
sulfonation of, 683
synthesis of
Friedel–Crafts alkylation, 679f, 684–685, 713, 713f
by reduction of aryl ketones, 712–713
Alkylboranes
formation by hydroboration, 411–415
oxidation of, 413–414
Alkyl bromides
mass spectrum of, 500, 501f
synthesis of, 590–593
Alkyl chloride
in Friedel–Crafts alkylation reactions, 685, 687–688
mass spectrum of, 500, 501f
Alkyl diazonium salts, 1028
Alkyl group, 134–137, 249, 253, 267, 272–275, 390, 684, 873–875
in alkyl tosylates, 359
carbon radical stability and, 571–572, 572f
as electron-donating group, 691–693
in ethers, 332, 335–336
nomenclature
in alcohols, 335
for branching molecules, A–3
in ethers, 335–336
polarizable, 691
stabilization of carbonyl group by, 845
Alkyl halides
acetylide anion reactions with, 441–444, 443f
alkane formation from, 785
allylic halides, 248, 248f
benzylic halides, 248, 248f
classification of, 248, 320
conversion of alcohols to, 352–358
with HX, 352–355, 357t
importance of, 358

Alkyl halides (*Continued*)
with thionyl chloride and phosphorus tribromide, 356–357, 357t
conversion to carboxylic acids, 798
conversion to ethers of monosaccharides, 1127
conversion to primary amines, 1009, 1010
coupling reaction with organocuprates, 1050–1051
dehydrohalogenation with strong base, 395
in direct enolate alkylation, 942–944
elimination reactions, 253, 297–330
E1 mechanism, 312–313, 319–322
E2 mechanism, 303–307, 307t, 314, 316–317, 319–322, 1027, 1027f
examples, 248f, 251–252, 251f
functional group, 96, 96t
mass spectrometry of, 500–501, 501f
molecular formula of, 248
nomenclature, 249–250, 250f
common names, 249
IUPAC, 249, 250f
nucleophiles, reactions with, 117
nucleophilic substitution reactions, 254–288
alcohol synthesis, 341–342
alkyl halide structure and, 267, 273, 279, 279f
ether synthesis, 341–342
to form amines, 1007–1009
S_N1 mechanism, 319–323, 320t
S_N2 mechanism, 319, 320t, 321–322
physical properties, 250, 250t
polar carbon–halogen bond, 252–253, 253f
primary (1°)
acetylide anion reactions with, 441–442
classification as, 248
E2 elimination reactions, 306, 320–321
S_N2 substitution reactions, 267, 279, 279f, 320t, 321
synthesis of, 352
reaction mechanism, determining type of, 319–323, 320t
reduction of, 466–467, 467f
secondary (2°)
acetylide anion reactions with, 441–442
classification as, 248
E1 elimination reactions, 311, 320f, 321
E2 elimination reactions, 306, 320t, 321
S_N1 substitution reactions, 273, 279, 279f, 320t, 321
S_N2 substitution reactions, 267, 279, 279f, 320f, 321
synthesis of, 352
synthesis of
halogenation of alkanes, 574, 579–581
hydrohalogenation of alkenes, 397–404, 403t
tertiary (3°)
acetylide anion reaction with, 441–442
classification as, 248
E1 elimination reactions, 311, 313, 320, 320t
E2 elimination reactions, 306, 320, 320t
S_N1 substitution reactions, 271, 273, 279, 279f, 313, 320, 320t
synthesis of, 352, 355
unreactive, 248, 248f
vinyl halides, 248, 248f, 286
1,2-Alkyl shift, 348
Alkylthio group, 96t, 366
Alkyl titanium compound, 1209
Alkyl tosylates, 359–361, 362f, 395
conversion of alcohols to, 359, 361
reactions of, 360
Alkynes, 426–454
addition reactions
halogenation, 433f, 436–437
hydration, 433f, 437–439
hydroboration–oxidation, 433f, 439–441
hydrohalogenation, 433f, 434–436
overview of, 432
boiling point of, 429
bond dissociation energies of, 427
bonds in, 93
cycloalkynes, 428
degrees of unsaturation, 427
deprotonation of, 433–434, 433t
examples, 430–431, 430f, 431f
functional group in, 93, 94t
hydroboration of, 1054, 1056
hydroboration–oxidation of, 433f, 439–441
infrared (IR) spectra of, 514
internal
definition of, 427
hydration of, 437
hydroboration–oxidation of, 440
oxidative cleavage of, 476, 739
synthesis in acetylide anion reactions with alkyl halides, 441–443
melting point of, 429
molecular formula of, 427
nomenclature, 428–429
oxidative cleavage of, 468f, 476, 739
physical properties of, 429
reduction of, 456, 456f, 463–466, 466f
to alkanes, 456f, 463–464, 466f
to cis alkenes, 463, 464, 466f
to trans alkenes, 463, 465–466, 466f
solubility of, 429
structure of, 427–428
synthesis of, 431–432, 446–447
from alkenes, 431–432
by dehydrohalogenation, 318–319, 319f
from dihalides, 431–432
E2 elimination reactions, 298, 318–319, 319f
multistep, 446–447
terminal
definition of, 427
hydration of, 437
hydroboration–oxidation, 440
hydrohalogenation, 434
oxidative cleavage of, 476, 739
reaction as an acid, 433–434
synthesis in acetylide anion reactions with alkyl halides, 441–443
triple bond of, 427–428
D-Allaric acid, 1131
Allegra. *See* Fexofenadine
Allenes, 211, 639
Allicin, 132
D-Allose, 1115f, 1131
Allyl aryl ether, 1096
2-Allylcycloheptanol, 390f
Allyl group, 390
Allylic alcohol
aldol reactions of, 971f
epoxidation of, 482–484
synthesis from α, β-unsaturated carbonyl compounds, 799
Allylic carbocations, 618
in biological reactions, 608
as conjugated system, 605, 606
electrostatic potential plot, 607, 607f
resonance of, 607–609
resonance-stabilized, 618
stability, 607
Allylic carbon
oxidation of, 462, 462f
radical halogenation at, 585–588
bromination, 586–588
product mixtures in, 587–588
in triacylglycerols, 462, 462f
Allylic halides, 248, 248f
Allylic sigma bonds, 1092
Allylic substitution, 587
Allyl radical, 585–588, 589f
Allyl vinyl ether, 1096
Alpha (α) carbon, 298–299, 732, 735, 752
of aldehydes and ketones, 820, 825, 828–829
substitution reactions of carbonyl compounds at the α carbon, 924–961
Alpha (α) cleavage, 503–504
Alpha (α) elimination reaction, 1059
Alpha (α) helix, 1180, 1181f, 1182, 1183f, 1185f
Alpha (α) hydrogens, 925, 929, 938–939, 968–969, 971, 977
Altocid. *See* Methoprene
D-Altrose, 1115f
Aluminum chloride, in Friedel–Crafts alkylations, 684–690, 702
Alzheimer's disease, 99, 973, 1022, 1097
Amber, 1247
Amberlyst A-26 resin, 479–480
Ambident nucleophile, 933
-amide (suffix), 874–875, 876t
Amide bond, 1164–1166, 1180
Amides
for alkyne synthesis by dehydrohalogenation, 319
basicity of, 872t, 1019, 1021t
boiling point of, 878t
classification, 97
cyclic, 870
functional group in, 97–98, 98t
hydrolysis of, 899
infrared (IR) spectra of, 515
interesting examples, 880–881
IR spectra of, 878, 879t
leaving group, 768
melting point of, 878t
nitrogen bases, 79, 79f
NMR spectra of, 879
nomenclature, 874–875, 876t
nucleophilic substitution reactions, 768
physical properties of, 877, 877f, 878t
pK_a, 872t, 931t
primary (1°), 870, 874, 877–878, 878t, 879, 883, 886, 888, 894
tautomers, 907
as a protecting group for an amine, 1024, 1024f
proteins, 880–881
reactivity, 883
reduction to amines, 780, 1010
secondary (2°), 870, 874–875, 877–878, 878t, 879, 886, 888, 894
solubility of, 878t
stability of, 872
structure and bonding of, 765, 869, 870
synthesis of
from acid chlorides, 886, 1023
from anhydrides, 888, 1023
from carboxylic acids, 889f, 892–894
from esters, 894
tautomers, 907–908
term usage, 932
tertiary (3°), 870, 874–875, 886, 888, 894
enolate formation from, 930
Amide tautomer, 907–908
Amikacin, 1140
-amine (suffix), 999
Amine oxide, 473
Amines, 287t, 996–1048
aromatic
basicity of heterocyclic, 1020, 1021t
nomenclature, 1000
basicity of, 997, 1014–1022, 1021t
aromaticity effects on, 1020, 1021t
hybridization effects on, 1020–1021, 1021t
inductive effects on, 1021t
pK_a values, 1014, 1016–1017, 1021t
relative to other compounds, 1016–1022, 1021*t*
resonance effects of, 1021t
boiling point of, 1001t
classification of, 97, 997
electrostatic potential plots of, 998, 998*f*
functional group, 96t, 97
general molecular formula for, 1002
infrared (IR) spectra of, 515
interesting examples, 1004–1007
derivatives of 2-phenylethanamine, 1005–1006, 1006f
histamine and antihistamines, 1004–1005
simple amines and alkaloids, 1004, 1004f
IR spectra of, 1002, 1003*f*
mass spectra of, 1002, 1002*f*
melting point of, 1001*t*
NMR spectra of, 1003, 1003*f*
nomenclature of, 999–1001
as nucleophiles, 997, 1014, 1022–1024
odors of, 1004
physical properties of, 1001, 1001*t*
primary (1°)
addition to aldehydes and ketones, 840–842, 842f, 843f
amino acids, 1155
from direct nucleophilic substitution, 1007–1008
Gabriel synthesis, 1008–1009

IR spectra of, 1002, 1003f
nomenclature, 999
reactions with acid chlorides and anhydrides, 1023
reactions with aldehydes and ketones, 1023
reactions with nitrous acid, 1028–1029
from reduction of amides, 1010
from reduction of nitriles, 909, 1010
from reduction of nitro compounds, 1009
structure of, 997
protecting group for, 1024, 1024*f*
reactions of
with acid chlorides, 886, 1023
with acids, 1014–1016
with aldehydes and ketones, 1023
with anhydrides, 888, 1023
with carboxylic acids, 889f, 892–894
conversion to alkenes, 1024–1027
conversion to amides, 1023–1024
with esters, 894
general features, 1014
Hofmann elimination, 1024–1027
with monosaccharides, 1141
with nitrous acid, 1028–1029
secondary (2°)
addition to aldehydes and ketones, 842–844, 843f
from direct nucleophilic substitution, 1007–1008
IR spectra of, 1002, 1003f
nomenclature, 999–1000
proline, 1155
reactions with acid chlorides and anhydrides, 1023
reactions with aldehydes and ketones, 1023
reaction with nitrous acid, 1029
from reduction of amides, 1010
structure of, 997
separation from other organic compounds, 1015, 1015f
solubility of, 1001t
spectroscopic properties of, 1002–1003, 1002f, 1003f
structure and bonding of, 997–999
synthesis of, 1007–1013
direct nucleophilic substitution reactions, 1007–1008
Gabriel synthesis, 1008–1009
reduction of amides, 780, 1010
reduction of nitriles, 1010
reduction of nitro compounds, 1009
reductive amination of aldehydes and ketones, 1011–1013
tertiary (3°)
from direct nucleophilic substitution, 1007–1008
IR spectra of, 1002, 1003f
nomenclature, 999–1000
from reduction of amides, 1010
structure of, 997
Amino acids, 117–118, 752–755, 1152–1197. *See also* Peptides; Proteins
abbreviations for, 753t, 1154f
N-acetyl, 1160–1162
acid–base behavior of, 753–754, 754f, 1155–1156, 1155f, 1156t
α-amino acids, 752, 1153, 1153f, 1156t
D-amino acids, 752, 1153, 1153f
L-amino acids, 752, 1153, 1153f
analysis of peptide composition, 1169
enantiomers of, 1153, 1153f
separation of, 1159–1162
essential, 752, 1155
general features of, 1153, 1153f, 1155
isoelectric points, 755
naturally occurring, 1153, 1154f
peptide synthesis from, 1172–1179
pK_a values, 1155, 1156, 1156t
representative examples, 753t
resolution of
by conversion to diastereomers, 1160–1162, 1160f
kinetic, using enzymes, 1162–1163
separation of, 1159–1163
stereogenic center of, 1153, 1155
structure of, 752, 1153, 1153f, 1154f, 1155
synthesis of, 1156–1159
alkylation of a diethyl malonate derivative, 1157–1158, 1159f
enantioselective hydrogenation, 1163–1164
reaction of α-halo acids with ammonia, 1156–1157, 1159f
Strecker synthesis, 1158–1159, 1159f
zwitterion form, 753, 1155
Aminobenzene, 645. *See also* Aniline
p-Aminobenzoic acid (PABA), 632, 919, 1037–1038
Aminoglycoside antibiotics, 1140
Amino group, 96t, 1000, 1019, 1024
as activating group, 702
as electron-donating group, 691–692, 693f, 701
inductive effect of, 691
nitro reduction to, 714
as ortho, para director, 699
resonance effect of, 691
6-Aminohexanoic acid, 1212
α-Amino nitrile, 1158–1159
Amino sugars, 1140
Ammonia
basicity of, 1017
boiling point, 465
in dissolving metal reductions, 457, 465
hybrid orbitals in, 39, 39f
molecular shape, 28
pK_a of, 68
reactions with
acid chlorides, 886, 1023
alkyl halide to form amines, 1007–1008
anhydrides, 888, 1023
carboxylic acids, 892
esters, 894
α-halo acids, 1156–1157
Ammonium cation, of amino acids, 754
Ammonium hydroxide, 782
Ammonium salt, 892
in amine extraction process, 1015, 1015f
enantiomers, 998
in Hofmann elimination reaction, 1024–1026
synthesis of, 997, 1008
Amoxicillin, 3, 58, 207, 881
Amphetamine, 81, 85, 207, 1006, 1011
Amygdalin, 210, 834
Amylopectin, 1139
Amylose, 175f, 1139
β-Amyrin, 1258
Anabolic steroids, 1254
Analgesics, 604, 737, 888, 943, 952, 957, 958
Anastrozole, 906, 959
Androgens, 1254
Androsterone, 1254t, 1257
-ane (suffix), 33, 131, 134, 135, 139, 141, 387
Anesthetics, 95–96, 114, 251f, 340, 714, 726
Angiotensin, 190, 1195
Angle strain, 151–152
in epoxides, 333, 344
Angstrom (Å), 26
Angular methyl groups, 1250–1251
Anhydrides
acidity/basicity of, 872t
boiling point of, 878t
cyclic, 870, 889f, 890
derivation of term, 874
IR spectra of, 879t
melting point of, 878t
mixed, 870, 873
nomenclature, 873, 876t
physical properties of, 878t
pK_a, 872t
reactions of, 887–889, 901t
acetylation, 888
with ammonia or amines to form amides, 1023
conversion to amides, 888
conversion to carboxylic acids, 888
conversion to esters, 888
reactivity, 883
solubility of, 878t
stability of, 872
structure and bonding of, 869–870, 872
symmetrical, 870, 873
synthesis of
from acid chlorides, 885, 886
dehydration of dicarboxylic acids, 890
Aniline, 640, 1023
basicity of, 1017–1018, 1019f
electrophilic aromatic substitution reaction, 682, 699, 701
electrostatic potential plot of, 693f, 1019f
nomenclature and, 646
polyhalogenation of, 701
protection of, 1024, 1024f
resonance structures, 691
structure of, 645, 1000
synthesis of
electrophilic aromatic substitution reaction, 682
by nitrobenzene reduction, 714
Anionic oxy-Cope rearrangements, 1096
Anionic polymerization, 1205–1206, 1205f
of epoxides, 1207–1208
Anions
acetate, 74–75
carbanion, 76, 76f, 79, 218f, 219
conjugated, 612
cyclopentadienyl, 658–660, 664
definition of, 8
enolate, 22
naked, 260
radical, 465
salt formation, 12
solvation by hydrogen bonding, 259
Anisole, 363
Annulation
origin of word, 982
Robinson, 982–986
Annulenes, 654–655
18-annulene, 569
Anomeric carbon, 1117–1122, 1124, 1135–1137, 1142
Anomers, 1118–1123, 1122f
Antabuse, 482
Antarafacial bond formation, 1089, 1091t
Antarafacial rearrangement, 1093–1094, 1093t
Anthracene, 655
Anti addition, 397, 403, 403t, 405, 406, 409, 463, 465, 472, 473
Antiaromatic compounds, 652, 653, 659, 660
Antibiotics, 2f, 3
aminoglycoside, 1140
amoxicillin, 207
cephalosporins, 881
chloramphenicol, 181–182
ionophores, 115–116
β-lactam, 495, 517–518, 881, 900–901
penicillin(s), 495, 517–518, 881, 900–901
sulfa drugs, 1037–1038, 1038f
Antibonding molecular orbital, 661–662, 662f, 663f, 664, 666
Anticancer drugs
anastrozole, 959
aromatase inhibitors, 906
doxorubicin, 1107, 1107f
epothilone A, 1065f
illudin-S, 951
imatinib mesylate, 881
tamoxifen, 924, 944
Anti conformation, 148–150, 148f, 150f, 301f
Antidepressant, 268, 492, 682f, 690, 708, 1006
Antifreeze, 339f
Antihistamines, 644, 682f, 1005, 1016
Anti-inflammatory drugs
aspirin, 61, 80–81, 233, 287, 287f, 737–738, 1244
dexamethasone, 97
fluticasone, 247, 293
ibuprofen, 133, 164, 202, 713, 773f, 962, 979, 1244
nabumetone, 952
naproxen, 86, 203, 943
nonsteroidal anti-inflammatory drugs (NSAIDs), 1244–1245
Antimalarial drugs, 307, 307f, 414–415, 415f, 658, 941, 1022
Antimicrobials. *See* Antibiotics
Antioxidants, 246, 589–590
Anti periplanar geometry, 314–315, 314f
Antiseptics, 255
Antiviral agents, 99, 1065f
Applied magnetic field, 529, 531, 539, 543

Aprotic solvents, polar, 260, 260f, 281, 282t, 305, 307t, 934
D-Arabinose, 1114, 1115f, 1131–1132
Arachidic acid, 1234t
Arachidonic acid, 600, 603
  leukotriene synthesis by oxidation of, 372
  prostaglandin synthesis from, 738
  structure of, 109, 1234t
  synthesis of eicosanoids from, 1243, 1243f
Arene, 642, 679f
Arene oxide, 171
Arginine
  isoelectric point for, 1156t
  pK_a values for, 1156t
  structure of, 1154f
Aricept. *See* Donepezil
Arimidex. *See* Anastrozole
Aripiprazole, 1042
Aromatase inhibitors, 906
Aromatic compounds, 641–676
  background on, 642
  bonding and antibonding orbitals, 661–662
  buckminsterfullerene, 666–667
  charged, 658–660
  electrophilic substitution. *See* Electrophilic aromatic substitution reactions
  examples, 654–660
    charged compounds, 658–660
    heterocycles, 655–658, 657f
    multiple rings, 655
    single ring, 654–655
  Hückel's rule, 651–654
    basis of, 660–663
  interesting, 648–649, 648f, 649f
  molecular orbitals, 661–666, 662f, 663f
  nomenclature, 644–647
  predicting aromaticity using the inscribed polygon method, 663–666, 665f
  reactions of, 677–728
    electrophilic aromatic substitution, 678–706
    nucleophilic aromatic substitution, 706–709
  spectroscopic properties, 647, 647f, 647t
Aromatic hydrocarbons, 93–94, 94t
Aromaticity, effect on basicity of amines, 1020, 1021t
Arrows
  curved, 21–22, 60, 63, 64, 83, 218, 220–221, 220t, 254, 1081
  double-headed, 19, 63–64, 220t
  equilibrium, 63–64, 69
  half-headed, 465, 571
  open, 444
  use in describing organic reactions, 218, 220–221, 220t
Artemisinin, 211, 414–415, 415f
Artificial sweeteners, 1137, 1137f, 1168
Arylamines
  basicity of, 1017–1019, 1021t
  reaction with nitrous acid, 1028–1029
Arylboranes, 1054
Aryl bromides, synthesis from aryl diazonium salts, 1030
Aryl chlorides
  examples of biologically active, 682f
  reactivity of, 707
  synthesis from aryl diazonium salts, 1030
Aryl diazonium salt, 1028–1035
Aryl fluoride, 707
Aryl fluorides, synthesis from aryl diazonium salts, 1030
Aryl groups, 646
Aryl halides, 248, 248f, 286
  coupling reactions
    with alkene, 1056–1058
    with organoborane, 1054
    with organocuprates, 1050
  examples of biologically active, 682f
  nucleophilic aromatic substitution, 706–709
  reactivity of, 686
  synthesis of
    from aryl diazonium salts, 1030
    halogenation of benzene, 679f, 681–682
Aryl iodides, synthesis from aryl diazonium salts, 1030
Aryl ketones, reduction to alkyl benzenes, 712–713
Ascorbic acid, 90, 91, 110–111
  See also Vitamin C
Asparagine
  isoelectric point for, 1156t
  pK_a values for, 1156t
  structure of, 1154f
Aspartame, 119, 916–917, 1137, 1137f, 1168
Aspartic acid, 1155, 1223
  isoelectric point for, 1156t
  pK_a values for, 1156t
  structure of, 1154f
Aspirin, 58, 61, 80–81, 737–738, 888, 917
  crossing of cell membrane, 81
  derivation of word, 737
  mechanism of action, 233, 738
  mode of action, 1244
  proton transfer reaction, 80–81
  structure, 737
  synthesis of, 287, 287f
Asthma, 372
Asymmetric carbon. *See* Stereogenic centers
Asymmetric reaction, 482
Asymmetric reduction, 774
Atactic polymer, 1208–1209
-ate (suffix), 733, 873, 876, 876t
Atenolol, 88, 99
Atherosclerosis, 905, 905f
Atmosphere
  carbon dioxide concentration in, 163, 163f
  ozone layer destruction, 252
Atomic mass unit (amu), 8
Atomic number, 8
Atomic orbitals, number of, 1078
Atomic weight, 8
Atom mass, infrared (IR) absorption and, 510–511, 511f
Atoms
  components of, 8
  excited state, 37
  ground state, 36–37
Atorvastatin, 1253, 1253f
Atropine, 1004, 1004f
Automated peptide synthesis, 1177–1179
Avobenzone, 34, 124, 978
Avocado, 111
Axial bond, 176
Axial hydrogens, 153–154, 153f, 157f
Aziridines, 382
Azo compound
  applications of, 1036–1038
    dyes, 1036–1037
    sulfa drugs, 1037–1038, 1038f
  formation of, 1029, 1034–1035
Azo coupling, 1029, 1034–1035
AZT (azidodeoxythymidine), 3, 670
Azulene, 669

B

Backside attack, in nucleophilic substitution reactions, 265–266, 267t, 280
Baekeland, Leo, 1217f
Baeyer, Adolf von, 132
Baeyer strain theory, 151
Bakelite, 1216–1217, 1217f
Ball-and-stick model
  of acyclic alkanes, 129–130
  of molecular shape, 27–29
Barbituric acid, 132
Barrier to rotation, 150
Base(s)
  for alkyne deprotonation, 433, 433t
  for alkyne synthesis by dehydrohalogenation, 318–319
  amines as, 1014–1016
  Brønsted–Lowry
    description of, 62–63
    reactions of, 63–65
    substitution reactions with alkyl halides, 253
  common, 79–80, 79f
  conjugate, 63–65, 67, 67t
  dehydration of β-hydroxy carbonyl compounds with, 966
  as electron pair donors, 62, 65, 81
  in elimination reactions, 298, 298t, 317t
    in E1 elimination reactions, 312, 312t, 313, 317, 317t
    in E2 elimination reactions, 305, 305f, 307t, 314, 317, 317t
  enolate formation with, 931–933, 932t
  halogenation in, 938–939, 940f
  hydrides, 79, 79f
  Lewis, 81–83
    nucleophile as, 253, 257, 269
  nitrogen, 79, 79f
  nonnucleophilic, 258
  organometallic reagents, 783, 784–785
  oxygen, 79, 79f
  as proton acceptors, 62
  strong, 79
  used to deprotonate carboxylic acids, 741, 741t
  weak as good leaving group in nucleophilic substitution reactions, 255–257
Base pairs, 1144, 1144f
Base peak, in mass spectrum, 497
Bases, nucleoside, 1142, 1144
Basicity
  acidity, inverse relationship with, 67, 67t
  of amines, 997, 1014–1022, 1021t
  of carboxylic acid derivatives, 871–872, 872t
  of carboxylic acids, 740
  leaving group ability and, 255–257
  nucleophilicity compared to, 257–258
  periodic trends in, 70–72
Basketane, 141f
Beeswax, 164, 164f
Benadryl, 1016 *See also* Diphenhydramine
Bender, Myron, 922–923
Benedict's reagent, 1129–1130
Bent molecular shape
  alcohols, 333
  description of, 29
  ethers, 333
  polarity and, 49
Benzaldehyde
  aldol reactions of, 968, 969, 970, 970f
  electron density of, 693, 693f
  resonance structures of, 692
  secondary alcohol formed from, 787
  structure of, 818, 820
Benzamide, structure of, 874
Benzene, 58, 93, 94t, 123, 506
  as annulene, 654
  background on, 642
  in BTX mixture, 648
  p-chlorostyrene synthesis from, 716
  conjugation of, 652
  degrees of unsaturation, 642, 643
  electrostatic potential plot of, 644, 644f, 693f
  Friedel–Crafts acylation of, 679f, 684–690, 713, 713f, 718
  Friedel–Crafts alkylation of, 679f, 684–690, 713, 713f, 715–716
  halogenation of, 679f, 681–682, 705
  heat of hydrogenation, 650–651, 650f
  hybridization and orbitals, 644
  lack of addition reactivity, 642, 651, 678
  metabolism of, 171
  molecular orbitals of, 662–664, 663f
  nitration of, 679f, 682–683, 705, 715
  p-nitrobenzoic acid synthesis from, 715–716
  NMR spectrum of, 53, 539, 540f, 540t, 654
  pi (π) electrons of, 643, 644, 653, 660, 678
  in polymer synthesis, 1218, 1218f
  resonance, 643
  stability, 649–651, 653
  structure of, 643–644
  substitution reactions, 642, 651, 678–690. *See also* Electrophilic aromatic substitution reactions
  sulfonation of, 679f, 682–683
  synthesis from aryl diazonium salts, 1031
  synthesis of bromonitrobenzene from, 705
  synthesis of *m*-bromoaniline from, 714–715
  synthesis of *o*-nitrotoluene from, 705–706
  synthesis of 1,3,5-tribromobenzene from, 1031, 1031f
  trisubstituted benzene synthesis from, 717–718
Benzenecarbaldehyde. *See* Benzaldehyde
Benzene derivatives
  disubstituted, 703–704

electrophilic aromatic substitution reactions of, 703–704
halogenation of, 703–704
nitration of, 706
synthesis of, 705–706
drugs containing a benzene ring, 649f
interesting examples, 648–649, 648f, 649f
NMR spectra, 552–553, 552f
nomenclature, 644–647
aromatic rings as substituents, 646
disubstituted, 645
monosubstituted, 645
polysubstituted, 646
spectroscopic properties, 647–648, 647f, 647t
substituted
activating and deactivating substituents, 694–697, 697f
directing effects of substituents, 694–701, 701f
electrophilic aromatic substitution, 694–697, 697f, 701–703
electrostatic potential plot of, 693, 693f
Friedel–Crafts acylation of, 703
Friedel–Crafts alkylation of, 702
halogenation of, 701–702
inductive effects on, 691–693
multistep synthesis of, 715–718
orientation effects on, 698–701
oxidation of, 712
reduction of, 714–715
resonance effects on, 691–693
synthesis of, 706–707, 715–718
with diazonium salts, 1031–1033
trisubstituted, synthesis of, 717–718
Benzene ring
of arylamines, 1017
NMR spectra of, 539, 540t, 552–553, 552f
Benzenesulfonic acid, 679f, 682
Benzo[*a*]pyrene, 373, 648, 648f
Benzocaine, 714
Benzoic acid, 890
acidity of, 746–748, 747f
deprotonation of, 741
separation from cyclohexanol by extraction procedure, 749–751, 751f
structure of, 732t
substituted, 746–748, 747f
synthesis of
by arene oxidation, 712
by oxidation of alkyl benzenes, 739
Benzoic anhydride, 876t
Benzonitriles, 876t, 1031
Benzophenone, structure of, 821
Benzoquinone, 624f
Benzoyl chloride, 876t, 890
Benzoyl group, 822
Benzphetamine, 1043
2-Benzylbutanoic acid, 947
Benzyl ester, 1175
Benzyl group, 646, 822
Benzylic bromination, 710–711, 716
Benzylic C–H bond, 710, 712, 739
Benzylic halides, 248, 248f, 710–711
Benzylic radical, 710, 1202
Benzyl salicylate, 1229
Benzyne, 708–709
Bergstrom, Sune, 738
Beryllium, octet rule exceptions in, 19
Beryllium hydride
hybrid orbitals in, 39
molecular shape of, 27
Beta (β) blocker, 99
Beta (β) carbon, 298–299, 308, 315, 732
Beta (β) elimination reactions. *See also* Elimination reactions
of alcohols, 343–352
dehydrohalogenation as, 298
Zaitsev rule, 308–309
Beta (β) lactam(s), 330, 870, 881, 900–901
Beta (β) lactam family of antibiotics, 495, 517–518
Bextra. *See* Valdecoxib
BHA (butylated hydroxy anisole), 596
BHT (butylated hydroxy toluene), 589–590
Biaryls, 1073
Bicyclic compounds, naming, A–5–6
Bicyclic product, 625
Bicyclic ring system, bridged, 625–626, 626f
Biformene, 1247
"Big Six" synthetic polymers, 1220–1221, 1221t
Bile acid, 172
Bile salt, 172
Bilobalide, 95
Bimatoprost, 738
Bimolecular elimination. *See* E2 elimination reactions
Bimolecular reactions
elimination reactions, 303
in nucleophilic substitution reactions, 263
organic reactions, 235
Biocatalyst, 1218, 1219, 1219f
Biodegradable polymers, 1222–1223
Biofuels, 1237
Biomolecules, 117–119, 118f, 1107. *See also* Carbohydrates; DNA; Lipids; Proteins
definition of, 117
families of, 117
mass spectra of, 506
Birch reduction, 493
2,2'-Bis(diphenylphosphino)-1,1'-binaphthyl (BINAP), 1164
Bisphenol A (BPA), 1214, 1215f, 1220, 1227
Bisphenol F, 725
Blattellaquinone, 887
Bloch, Konrad, 1252
Blood alcohol content, measurement of, 479, 479f
Blood–brain barrier, 113–114
Boat form, of cyclohexane, 155, 155f
Boc (*tert*-butoxycarbonyl protecting group), 1174, 1176
Boiling point
of alcohols, 338t
of aldehydes and ketones, 823t
of alkanes, 143, 143t
of alkenes, 391
of alkyl halides, 250t
of alkynes, 429
of amines, 1001t
of carboxylic acids, 734t
definition of, 103
of enantiomers, 199, 199t
of epoxides, 338t
of ethers, 338t
intermolecular forces and, 103–104
polarizability and, 103, 104f
of racemic mixtures, 199, 199t
separation of liquids based on, 104, 105f
surface area and, 103, 104f
Bombykol, 1055, 1055f
Bond(s) and bonding, 11–13. *See also specific bond types*
bond angle, 26–29, 29t, 44t
bond length, 26, 26t, 45–46, 46t
breaking and making in organic reactions, 217–221, 218f
covalent, 11, 12–13
definition of, 11
double, 15–16, 21–25, 31, 31f, 42, 42f, 45–46, 46t
energy and entropy, 228
ionic, 11–12
Lewis structures, 13–19
in nucleophilic substitution reactions, 261–264, 269–270
octet rule, 11
π (pi), 42–43, 42f, 43f, 44t
polar, 48–49, 50f
rotation, 41, 42
σ (sigma), 36, 38, 41–43, 42f, 43f, 44t
triple, 16, 42–43, 45, 46t
Bond angles
in alkynes, 427
four groups around an atom, 26–29, 29t, 44t
molecular shape and, 26–29
three groups around an atom, 26, 27, 29t, 44t
two groups around an atom, 26, 27, 29t, 44t
Bond cleavage
heterolytic, 217–218
homolytic, 217–218
in nucleophilic substitution reactions, 261–264, 269–270
Bond dissociation energy, 221–225
in alkynes, 427
for benzylic C–H bond, 710
bond length and, 223
bond strength and, 223
carbon radical stability and, 571–572
of common bonds, 222t, A–7
definition of, 221
enthalpy change calculation from, 223–224
limitations of, 224
periodic table trend in, 223
Bonding (shared) electrons, 13
Bonding molecular orbital, 661–666, 662f, 663f, 665f
Bond length
average for common bonds, 26t
in benzene, 643
bond dissociation energy and, 223
bond strength and, 45–46, 46t
in conjugated dienes, 614–615
percent *s*-character and, 46, 615
periodic trends in, 26
Bond polarity, 48–50, 50f
Bond strength
bond dissociation energy and, 223–224
bond length and, 45–46, 46t
infrared (IR) absorption and, 510–512, 510f, 512t
percent *s*-character and, 46
9-Borabicyclo[3.3.1]nonane (9-BBN), 412
Borane, 411–414, 412f, 774
Boron, octet rule exceptions in, 19
Boron trifluoride, 27, 39
BPA (bisphenol A), 1214, 1215f, 1220, 1227
Bradykinin, 1167
Brevenal, 333
Brevicomin, 866
Bridged bicyclic ring system, 625–626, 626f
Bridged halonium ion, 406–410, 410t, 436
Bridgehead carbons, A–5
Bromination, 714
of alkanes, 578–580, 580f
of alkenes, 590–593
at allylic C–H bonds, 586–588
with NBS, 586–588
product mixtures in, 587–588
selective, 586–587
substitution versus addition, 587
of benzene, 681, 705
benzylic, 710–711, 716
chlorination compared, 578–581
as endothermic reaction, 579–580
as exothermic reaction, 591–592
polybromination, 701
of substituted benzenes, 701
as test for a pi (π) bond, 405
Bromine, isotopes of, 500
m-Bromoacetophenone, 822f
m-Bromoaniline, 714–715
o-Bromoaniline, 1000
p-Bromobenzaldehyde, synthesis of, 1032–1033
Bromobenzene, 363, 681
4-Bromobutanal, 862
1-Bromobutane, 306
2-Bromobutane, 299, 583
1-Bromobut-2-ene, 619–620
3-Bromobut-1-ene, 619
o-Bromochlorobenzene, 645
1-Bromo-1-chloroethylene, NMR spectrum of, 533
Bromocyclohexane, 402
cis-4-Bromocyclohexanol, 381
trans-4-Bromocyclohexanol, 381
2-Bromocyclohexanone, 941
1-Bromocyclohexene, 1052
3-Bromohexane, 181
enantiomers of, 183f
6-Bromohexan-2-one, 866
1-Bromohex-2-ene, 292
trans-1-Bromohex-1-ene, 1051
Bromohydrins, 409
Bromomethane, 342
in nucleophilic substitution reactions, 262–264, 267f
2-Bromo-2-methylbutane, 308
1-Bromo-3-methylbut-2-ene, 587
3-Bromo-3-methylbut-1-ene, 587
cis-1-Bromo-4-methylcyclohexane, 279
1-Bromo-1-methylcyclopentane, 312
2-Bromo-2-methylpropane, 299, 306, 442
Bromonitrobenzene, 705
Bromonium ion, 587
2-Bromopentane, in an E2 elimination reaction, 1027, 1027f
1-Bromo-1-phenylpropane, 296

1-Bromopropane
  NMR spectrum of, 548, 548f
  synthesis of, 591
2-Bromopropane, 342
  mass spectrum of, 500, 501f
  NMR spectrum of, 546, 546f
  synthesis of, 580, 591
2-Bromoprop-1-ene, 639
α-Bromopropiophenone, 939
(*Z*)-2-Bromostyrene, 1056, 1072
N-Bromosuccinimide (NBS), 409, 586–588
p-Bromotoluene, 645
Bronchodilator, 181, 341, 371, 371f
Brønsted–Lowry acid
  carboxylic acids, 740–745
  description of, 62–63
  reactions of, 63–65
Brønsted–Lowry base
  alkyl halide and elimination reactions, 253
  description of, 62–63
  in elimination reactions with alkyl halides, 298
  reactions of, 63–65
Brucine, 1194
BTX mixture, 93, 648
Buckminsterfullerene, 666–667
Bufotenin, 1006
Bufo toads, 1006
Bupivacaine, 88
Bupropion, 682f, 959
Buta-1,2-diene, 451
Buta-1,3-diene, 613, 1077, 1081, 1105
  conformations of, 1166
  conjugation in, 605–606, 606f, 613
  electrostatic potential plot for, 606, 606f
  HBr addition to, 617–620, 620f
  molecular orbitals of, 1079–1080, 1079f
  polymerization of, 1207, 1211
  resonance structures, 609, 615
  sigma (σ) bond length in, 614–615
  stability of, 653
  ultraviolet absorption, 630
Butanal
  boiling point of, 103
  intermolecular forces in, 102
  melting point of, 105
  naming, 334
Butan-2-ol
  (*S*)-butan-2-ol, conversion to its tosylate, 359
  enantiomers of, 186–188, 773–774
  infrared (IR) spectrum, 514
  stereogenic center of, 182
  synthesis using Grignard reaction, 790–791
tert-Butanol, as polar protic solvent, 259f
Butan-2-one, 449
  boiling point of, 822
  infrared (IR) spectrum, 514
  reduction of, 773–774
  structure of, 821
  synthesis of, 951
But-1-ene, 302, 326, 385t
  bond dissociation energy, 635
But-2-ene
  cis-but-2-ene, 385t, 391, 399, 407, 408f
    from alkyne reduction, 464
    carbene addition to, 1060
    epoxidation of, 470
    hydrogenation of, 458–459, 459f
  trans-but-2-ene, 385t, 391, 407, 408f, 409
    from but-2-yne reduction, 465
    carbene addition to, 1060
    epoxidation of, 470
    hydrogenation of, 458, 459f
  stereoisomers of, 300–302, 301f, 326
tert-Butoxide, nucleophilicity of, 258
tert-Butoxycarbonyl protecting group (Boc), 1174, 1176
tert-Butoxy group, 336
Butter, 395, 462
Butter yellow dye, 1034
Butylbenzene, 717–718
tert-Butylbenzene, 645
Butylcyclohexane, 140f
tert-Butylcyclohexane, 157f
tert-Butyl cyclohexanecarboxylate, 874
cis-4-*tert*-Butylcyclohexanol, 815
4-*tert*-Butylcyclohexanone, 815
tert-Butylcyclopentane, 139
1-*tert*-Butylcyclopentene, 381
tert-Butyldimethylsilyl ether (TBDMS), 793–794
sec-Butyl ethyl sulfide, 365
Butyl group, 134
  sec-butyl group, 134
  tert-butyl group, 134, 157
tert-Butyl hydroperoxide, 467, 483
tert-Butyl iodide, 249
Butyllithium, 79, 836, 933
Butylmagnesium chloride, 802
1-*sec*-Butyl-3-methylcyclohexane, 140f
sec-Butyl methyl ether, 335
tert-Butyl methyl ether (MTBE), 108
  formation of, 405
  as fuel additive, 405, 530
  NMR spectrum of, 530, 532, 535
5-*tert*-Butyl-3-methylnonane, 137
But-1-yne, 451, 454
  addition reactions of, 432, 433f
  hydrohalogenation of, 433f, 434, 435
But-2-yne, 451, 454, 465
  hydration of, 438
Butyric acid, structure of, 732t
γ-Butyrolactone, 917

C

Cadaverine, 1004
Caffeine, 58, 113–114, 124, 562
Cahn–Ingold–Prelog system, 185
Calcitriol, 614, 614f
Calorie (cal), 45
Camphene, 1260
Camphor, 1246
10-Camphorsulfonic acid, 1193
Cancer, 373. *See also* Anticancer drugs; Polycyclic aromatic hydrocarbons
Candlenuts, 383
Capnellene, 443, 443f
Caproic acid, 732, 732t
ε-Caprolactam, 1212, 1222
ε-Caprolactone, 1229
Capsaicin, 4, 56, 216, 517, 562, 641, 960
Caraway, odor of, 204
-carbaldehyde suffix, 819
Carbamate, 920, 1174, 1213
Carbanion(s), 218f, 219
  addition to aldehydes and ketones, 832
  defined, 76
  electrostatic potential plots, 76f
  formation in alkyne reduction, 465–466
  nucleophilic addition with, 767, 832
  nucleophilic substitution with, 768
  as strong bases, 79
  vinyl, 466
Carbenes
  intermediates in cyclopropane synthesis, 1058–1061
  metal–carbene intermediates in metathesis, 1064
  preparation of dihalocarbenes, 1059–1061
  structure of, 1058, 1059
Carbenoid, 1062
Carbidopa, 50, 51
Carbinolamines, 830, 831f, 840, 841, 843, 1023
Carbocation(s), 22, 218f, 219
  achiral, 403
  allylic, 618
    in biological reactions, 608
    as conjugated system, 605, 606–607
    electrostatic potential plot, 607, 607f
    resonance of, 607–609
    stability, 607
  classification, 273
  in electrophilic addition reactions, 399–404
  in electrophilic aromatic substitution reactions, 679–681, 696–697, 697f
  electrostatic potential maps, 274, 274f
  in the E1 mechanism, 310–313, 317t
  energy diagram for formation of, 278f
  formation in Lewis acid–base reactions, 83
  Hammond postulate, 400, 401f
  as Lewis acid, 399
  localized versus delocalized, 607f
  in nucleophilic substitution reactions, 262
  planar, 270–272, 280, 403, 1124–1125
  primary (1°), 273–275
  as reactive intermediate, 350
  resonance-stabilized, 435–436, 438, 503, 607, 607f, 617–618, 620, 679, 681, 696–697, 707, 927
  secondary (2°), 273–275, 685, 687, 688
  in S_N1 reactions, 313
  in S_N1 substitution reactions, 269–278, 278f, 280–281, 284
  tertiary (3°), 273–275, 685, 687, 688
  trigonal planar, 270–271
  vinyl, 286, 434–435, 438
Carbocation intermediates, 591
Carbocation rearrangements
  conversions of alcohols to alkyl halides, 355
  in dehydration of alcohols, 348–350
  Friedel–Crafts alkylation, 687–688
  in hydration reactions, 404
  1,2-shifts, 348–350
    1,2-hydride shift, 687–688
    1,2-methyl shift, 349
  in S_N1 reactions, 355
Carbocation stability, 273–275
  of allylic carbocations, 607
  conjugation and, 607
  in electrophilic aromatic substitution reactions, 687, 696–697, 697f
  Hammond postulate and, 275–278
  hyperconjugation and, 274–275
  inductive effects and, 273
  1,2-shifts and, 348
Carbohydrates, 855–856, 1106–1151
  amino sugars, 1140
  cellulose, 175–176, 175f, 176f, 902, 902f, 1107f, 1139
  disaccharides, 1134–1137, 1136*f*, 1137*f*
  energy in, 1107–1108
  examples of, 1107, 1107f
  glycosides, 1124–1127
    N-glycosides, 1141–1144
  molecular formula of, 1107
  monosaccharides, 1108–1134. *See also* Monosaccharides
  natural fibers, 902, 902f
  nomenclature, 1108–1109, 1114
  polysaccharides, 1138–1140
  starch, 175–176, 175f, 176f, 339, 339f, 1139
Carbon
  alpha (α), 298–299, 732, 735, 752
    of aldehydes and ketones, 820, 825, 828–829
    substitution reactions of carbonyl compounds at the α carbon, 924–961
  beta (β), 298–299, 308, 314, 315, 732
  bonding of, 12, 13f, 16, 17, 18t, 36–38
  charged atom, 35
  chiral. *See* Stereogenic centers
  classification of atoms, 94–95
  common nucleophiles, 261t
  delta (δ), 732
  electron-deficient, 73, 92, 95, 117, 219, 252–253, 253f
  electronegativity value, 783
  electrophilic, 253
  elemental forms of, 666
  excited state, 37
  formal charge of, 18t
  gamma (γ), 732
  ground state, 36–37

isotopes of, 8, 504t
mass of, 504t
periodic table entry, 9
radicals, 571–572, 572f
tetravalence of, 4, 28, 30, 32–34
valence electrons of, 36
Carbonate, 1214
Carbon backbone (skeleton), 92, 107
Carbon–carbon bonds
bond length and bond strength, 45–46, 46t
in condensed structures, 31, 31f
double. *See also* Double bonds
in alkenes, 299–301. *See also* Alkenes
properties of, 385t
radical reactions, 590–593
restricted rotation around, 300–301, 301f
double bond, 16
NMR spectra of, 539–540, 540f, 540t
nonpolar nature of, 48
reactions forming, 1049–1075, A–8
coupling reactions of organocuprates, 1050–1052
cyclopropane synthesis, 1058–1061
Heck reaction, 1050, 1056–1058
metathesis, 1062–1067, 1063f, 1065f
Simmons–Smith reaction, 1061–1062
Suzuki reaction, 1050, 1052–1056, 1055f
rotation, 41, 42
rotation around, 144–145
Carbon dioxide
atmospheric concentration of, 163, 163f
electrostatic potential plot for, 50f
global climate change, role in, 163
as nonpolar molecule, 49
reaction of Grignard reagents with, 798
release in combustion reactions, 162–163
Carbon–halogen bond, polar, 252–253, 253f
Carbon–hydrogen bonds
acidity of, 71
benzylic, 710, 712
bond length and bond strength, 46
chemical shifts, 538
halogenation of, 585–588
nonpolar nature of, 48
radical reaction with, 573
Carbon monoxide, 1188
Carbon NMR, 528, 556–559
of aldehydes, 825, 825f
of amines, 1003
basis of, 556
of benzene derivatives, 647, 647f, 647t
of carboxylic acid derivatives, 879
of carboxylic acids, 735, 736f
characteristic absorptions, A–11
chemical shifts in, 556, 558, 558t, 559f
examples of spectra, 559f
of ketones, 825
of nitriles, 879
number of signals in, 557–558
position of signals in, 558–559
Carbon–oxygen bonds
carbonyl group, 97, 98t
condensed structures containing a double, 31, 31f
in functional groups, 97–99, 98t
polar nature of, 48
Carbon skeleton (backbone), 92, 107, 108
Carbon tetrachloride, 251
as solvent, 106, 107
-carbonyl chloride (suffix), 873, 876t
Carbonyl compounds. *See also specific classes*
classes of, 765
condensation reactions, 962–995
conversion of ozonides to, 474
electron deficiency of, 766
electrostatic potential map of, 766, 766f
formation by oxidation of alcohols, 476–478
β-hydroxy
synthesis of, in aldol reactions, 963–975
useful transformations of, 970, 971f
leaving groups of, 765, 767, 768–769
pK_a, 929, 929t, 930t, 932
reactions at the α carbon, 924–961
acetoacetic ester synthesis, 949–952
enolate alkylation, 937, 942–945
halogenation, 937–941, 940f
malonic ester synthesis, 945–949
racemization, 936, 936f
reactions of
general, 766–769
nucleophilic addition, 767
nucleophilic substitution, 767, 768–769
with organometallic reagents, 785–789, 792–794, 793f
oxidation, 770, 782
reduction, 769–781
reactivity, 818
reduction of, 769–781
biological, 776–777
carboxylic acids and derivatives, 777–781
by catalytic hydrogenation, 772
enantioselective, 774–777, 775f
with metal hydride reagents, 771–773, 773f, 781t
stereochemistry of, 773–774
synthesis by Cope rearrangement, 1095
synthesis by hydroboration–oxidation of alkynes, 439–441
tautomers, 907–908
α,β-unsaturated compounds, 799–801
unsymmetrical, enolates of, 934–935
Carbonyl condensation reactions, 962–995
aldol reactions, 963–975. *See also* Aldol reactions
Claisen reaction, 975–979, 976f
Dieckmann reaction, 979–980
Michael reaction, 980–982, 982f
Robinson annulation, 982–986
Carbonyl groups, 437, 765
conjugated, 824, 824f
as electron-withdrawing group, 624
features of, 871
formation by oxidative cleavage of alkenes, 468f, 474–475
as functional group, 97, 98t
hydrogenation of, 461
IR absorption of, 735, 735f, 823–824, 823f, 824f
of monosaccharides, 1108
nucleophilic addition to, 828–830, 831f, 846
reactions of, 1128–1134
reactivity of, 97
stabilization by alkyl groups, 845
tautomers of, 926–927
Carboxy, derivation of term, 730
Carboxy group, 98t, 730
of benzene derivatives, 712, 715–716
in prostaglandins, 164
Carboxylate anion
of amino acids, 754
formation of, 741, 742
metal salt of, 733, 733f
synthesis of
hydrolysis of amides, 899
hydrolysis of esters, 894–896
hydrolysis of nitriles, 907–908
Carboxylation, 798
Carboxylic acid(s), 729–763
acidity of, 740, 741–746, 744f, 872t, 889
aliphatic, inductive effects in, 744–746
amino acids, 752–755
aspirin as, 737–738
basicity of, 740
benzene derivatives, 712
boiling point of, 734t
deprotonation of, 741–742, 741t
diacids, 733, 945–947
as dimers, 734, 734f
dipole–dipole interactions of, 734
electrostatic potential plot, 730, 730f
functional group in, 98t
hydrogen bonding, 734, 734f
interesting examples, 736–737
intermolecular forces, 734, 734f, 734t
IR spectra of, 735, 735f
leaving group, 768
melting point of, 734t
NMR spectra of, 735, 736f
nomenclature of, 731–734, 733f
common names, 732–733, 732t
IUPAC system, 731–734, 733f
nucleophilic acyl substitution reactions, 889–894, 889f, 901t
conversion to acid chlorides, 889f, 890
conversion to amides, 889f, 892–894
conversion to cyclic anhydrides, 889f, 890
conversion to esters, 889f, 891–892
nucleophilic substitution reactions, 768
oxidation of aldehydes to, 1129
physical properties, 734–735, 734t, 878t
pK_a of, 872t
protonation of, 740
reactions of
ester synthesis, 889f, 891–892
general features, 740–741
nucleophilic acyl substitution reactions, 889–894, 889f, 901t
reduction, 770, 778, 780
reactivity, 883
resonance stabilization of, 742–743, 743f, 744f
salts of, 737
separation from other organic compounds, 749–751, 751f
solubility of, 734t
spectroscopic properties, 735–736, 735f, 736f
stability of, 872
structure and bonding of, 730, 765, 869, 872
substituted benzoic acids, 746–748, 747f
synthesis of, 739–740
conversion of acid chlorides, 885, 887
conversion of anhydrides, 888
hydrolysis of amides, 899
hydrolysis of esters, 894–896
hydrolysis of nitriles, 907–908
malonic ester synthesis, 945–949
oxidation of aldehydes, 770, 782
oxidation of alkyl benzenes, 739
oxidation of alkynes, 739
oxidation of primary (1°) alcohols, 468f, 477, 478–479, 739
oxidative cleavage of alkynes, 468f, 476
reaction of organometallic reagents with carbon dioxide, 798
-carboxylic acid (suffix), 873
Carboxylic acid derivatives. *See also* Acid chlorides; Amides; Anhydrides; Esters
acidity/basicity of, 871–872, 872t
boiling point of, 878t
IR spectra of, 878–879, 879t
leaving groups of, 870, 872t, 882–883
melting point of, 878t
NMR spectra of, 879
nomenclature of, 873–877, 876t
nucleophilic acyl substitution reactions of, 768
mechanism of, 882–883
relative reactivity, 883–884, 901
summary of, 901, 901t
physical properties of, 877–878, 878t
reaction of organometallic reagents with, 785, 794–797
reactivity with nucleophiles, 768
reduction of, 770, 777–781
resonance structures of, 871
solubility of, 878t
stability of, 872
structure and bonding of, 765, 869–872
Carboxypeptidase, 1171, 1171t
Carcinogen, 373
Cardinol, 1220
Carotatoxin, 124
β-Carotene, 110, 392f, 837, 837f, 1258
Carroll rearrangement, 1104
Carson, Rachel, 252
Carvacrol, 675
Carvone, 204, 1258
(*R*)-Carvone, 675
Caryophyllene, 325
Cassava, 834

Catalysts, 236–237. *See also specific catalysts*
  acids as, 236–237
  biocatalyst, 1218, 1219, 1219f
  chiral, 1163–1164
  definition of, 236
  energy of activation lowered by, 237, 237f
  enzymes, 237–238, 238f
  homogeneous, 1209
  hydrogenation, 458
  Lewis acid, 354
  Lindlar, 464, 466f
  metal, 236–237, 458
  palladium, 236, 237, 772. *See also* Palladium-catalyzed reactions
  Ziegler–Natta, 1209, 1210
Catalytic amount, 237
Catalytic hydrogenation, 458–460
  of aldeyhdes and ketones, 772
Catecholborane, 1054, 1056
Cation(s)
  carbocation. *See* Carbocation(s)
  counterions, 62
  cyclopentadienyl, 659
  definition of, 8
  radical, 497, 501
  salt formation, 12
  solvation by ion–dipole interactions, 259–260, 280
  spectator ions, 62
  tropylium, 660, 664
Cationic polymerization, 1204–1205, 1205f
CBS reagent, 774–776, 775f
Cedrol, 1246f
Celecoxib (Celebrex), 862, 1244–1245
Celery ketone, 204
Cell membrane
  cholesterol in, 165, 165f, 1252
  functions, 113
  phospholipids in, 1238–1240
  structure of, 113–114, 114f
  transport across, 115–116, 116f
Cellobiose, 1136
Cells, 113
Cellulose, 902, 902f
  hydrolysis of, 175f, 176, 1138
  solubility of, 177
  structure of, 175–177, 175f, 176f, 1107f, 1138
Cellulose acetate, 1138
Cephalexin, 881
Cephalin, 1238
Cephalosporins, 881
Cetylpyridinium chloride (CPC), 255
CFCs (chlorofluorocarbons), 252, 584, 585f
Chain branching, in polymerization, 1202–1203
Chain-growth polymerization, 1200–1207
  anionic, 1205–1206, 1205f, 1207–1208
  cationic, 1204–1205, 1205f
  chain branching in, 1202–1203
  copolymers, 1207
  of epoxides, 1207–1208
  ionic, 1204–1206, 1205f
  radical, 1201–1202, 1202f
Chain-growth polymers, 1199, 1200–1207
Chain mechanism, 575, 584, 593
Chain termination, 1201–1202
Chair conformation
  of cyclohexanes, 152–161, 153f
  of glucose, 1122
Charge
  Brønsted–Lowry acids and bases, 62, 62f, 65
  on carbon atom, 35
  dipole, 48, 49, 50f
  formal, 16–18, 18t
  partial, 25, 48
  in resonance hybrid, 25
Charged ion, 8
Chauvin, Yves, 1062
Chemical shift
  in ^{13}C NMR, 556, 558, 558t, 559f
  in ^{1}H NMR
    equation for, 531
    predicting values of, 537–538
    protons on benzene rings, 539, 540t
    protons on carbon–carbon double bonds, 539, 540t
    protons on carbon–carbon triple bonds, 540, 540t
    scale, 530
    values for common proton types, 538t
Chiral catalysts, 1163–1164
Chiral drugs, 202–203
Chirality, 178–180
  basic principles of, 180f
  of naturally occurring objects, 180
  in substituted cycloalkanes, 196
  test for, 178
Chirality center, 179. *See also* Stereogenic centers
Chiral molecules, 178–180, 180f
  definition of, 178, 179
  halogenation of, 582t, 583
  optical activity of, 198
Chiral reagent, 1162, 1163
Chiral reducing agent, 774
Chitin, 1107, 1140
Chloral, 844, 845
Chloral hydrate, 844, 845
Chloramphenicol, 181–182
Chlorination
  of alkanes, 574–583, 577f, 581f
  of benzene, 682, 716
  bromination compared to, 578–581
  of ethane, 576–577, 577f
  as exothermic reaction, 577, 577f, 580–581
  mechanism of, 575–576
Chlorine
  isotopes of, 500
  mass spectra of alkyl chlorides, 500, 501f
2-Chloroacetic acid, 745
Chlorobenzene, 645, 681, 708
2-Chlorobuta-1,3-diene, 1211
1-Chlorobutane, synthesis of, 582
2-Chlorobutane
  NMR spectrum of, 534
  synthesis of, 582
Chlorobutanoic acid, 746
Chlorocyclohexane, chair conformation of, 315–316, 315f
Chlorocyclopentane, 109, 575–576
Chlorocyclopropane, NMR spectrum of, 533
Chloroethane, 57, 95–96
  NMR spectrum of, 557
Chloroethylene, NMR spectrum of, 533
4-Chloro-1-ethyl-2-propylbenzene, 646
Chlorofluorocarbons (CFCs), 3, 252, 584, 585f
Chloroform, 251
  dichlorocarbene preparation from, 1059
Chlorohydrin, 408, 410f
Chloromethane, 251f
1-Chloro-2-methylbutane, 574
1-Chloro-3-methylbutane, 574, 585
2-Chloro-2-methylbutane, 574, 585
2-Chloro-3-methylbutane, 574, 585
1-Chloro-2-methylcyclohexane
  cis-1-chloro-2-methylcyclohexane, 315
  trans-1-chloro-2-methylcyclohexane, 315
1-Chloro-1-methylcyclopropane, 327
2-Chloro-5-methylheptane, 249
1-Chloro-2-methylpropane, 250f
p-Chloronitrobenzene, 706
1-Chloropentane, 532
2-Chloropentane, 532
3-Chloropentane, 532
meta-Chloroperoxybenzoic acid, 468–471
m-Chlorophenol, 1032
2-Chloropropanal, 820f
1-Chloropropane, 400, 581, 713
2-Chloropropane, 400, 581
  mass spectrum of, 500, 501f
2-Chloropropene, 434
3-Chloropropenoic acid, NMR spectra of, 549, 549f
α-Chloropropionaldehyde, 820f
2-Chloropyridine, 724
Chloroquine, 1022
p-Chlorostyrene, 716
N-Chlorosuccinimide, 453
p-Chlorotoluene, 708–709
1-Chloro-1,3,3-trimethylcyclohexane, 402
Chlorpheniramine, 682f
Cholesterol, 629, 905, 905f, 1231, 1252–1253
  biosynthesis of, 1252–1253, 1252f
  in cell membranes, 165, 165f, 1252
  cholesterol-lowering drugs, 1253, 1253f
  solubility of, 107–108, 164
  specific rotation of, 201
  structure of, 164, 164f, 185, 1252, 1252f
  synthesis stimulation by saturated fats, 395
  trans fats and, 463
Cholesteryl esters, 904–905, 905f
Cholic acid, 172
Choline, 904
Chondrocole A, 252
Chromate ester, 477–478
Chromium
  anionic oxidant of, 480
  in blood alcohol screening, 479, 479f
  in oxidizing agents, 468, 477–480, 782
Chromium(VI) oxide, 468
Chrysanthemic acid, 558
Churchane, 141f
Churchill, Winston, 252
Chymotrypsin, 1171, 1171t
Ciguatoxin (CTX3C), 764, 779, 779f
Cimetidine, 1005
Cinnamaldehyde, 826f, 968
Cinnamoylcocaine, 916
Cis (prefix), 388–389
Cis-fused, 625
Cis geometry
  alkenes, 385, 385t, 388–389, 403
  cycloalkenes, 385
Cis isomers
  of but-2-ene, 301–302, 301f
  of cycloalkanes, 158–161
  definition of, 158, 301
  of disubstituted cycloalkanes, 195
  stability, 303
Cis protons, 549, 549f
Citalopram, 208
Citral, 1246f
Citric acid, 53
Citronellal, 826f
Citronellol, 526
Claisen reactions, 975–979, 976f
  crossed, 977–979
  intramolecular, 979–980
Claisen rearrangement, 1095, 1096–1097
Claritin. *See* Loratadine
Clavulanic acid, 420
Clemmensen reduction, 712
Clopidogrel, 190, 957, 1193
^{13}C NMR. *See* Carbon NMR
Cocaine, 87, 125, 880, 916, 1004, 1006
Cocoa butter, 1237
Coconut oil, 395, 1235, 1237
Codeine, 759, 1007
Coenzyme, 481
  NADH, 481, 776
Coffee, decaffeinated, 251f
Collagen, 1153
  structure of, 1187, 1187f
Column, in the periodic table, 9
Combustion
  of acetylene, 430
  of alkanes, 162–163
  definition of, 162
  of triacylglycerols, 1237
Common names
  of alcohols, 335
  of aldehydes, 820, 820f
  of alkanes, 141, 141f
  of alkenes, 390, 390f
  of alkyl halides, 249
  of amines, 999
  of benzene derivatives, 644–645
  of carbohydrates, 1108, 1114
  of carboxylic acid derivatives, 873
  of carboxylic acids, 732–733, 732t
  of epoxides, 337
  of ethers, 335
  of ketones, 821, 822f
  of nitriles, 876, 876f
Compounds
  examples of, 11
  formation of, 11
  ionic, 11–12
Concentration, reaction rate and, 234–236
Concerted reaction, 217, 304
Condensation polymer. *See also* Step-growth polymers
  nylon, 902
  polyesters, 903
Condensation reactions
  aldol. *See* Aldol reactions
  carbonyl, 962–995
  definition of, 965
Condensed structures, 30–32, 31f
Configuration
  inversion of
    in alkyl tosylate reactions, 360, 361
  retention of
    in chlorination, 583

in conversion of alcohols to alkyl tosylates, 359, 361
in oxidation of alkylboranes, 413
in S_N2 reactions, 265–266, 266f, 267t, 280
of stereoisomers, 177
Conformations
of acyclic alkanes, 144–151
butane, 148–151, 148f, 150f
ethane, 144–147, 146f
anti, 148–150, 148f, 150f
of cycloalkanes, 151–161
disubstituted, 158–161
monosubstituted, 156–158
of cyclohexane(s), 152–161
boat, 155, 155f
chair, 152–161, 153f
disubstituted cyclohexane, 158–161, 160f
monosubstituted cyclohexane, 156–158, 157f
ring-flipping, 155–156, 155f
definition of, 144
eclipsed, 144–151, 146f, 147f, 148f, 150f, 150t, 301, 314f
free energy change and, 227
gauche, 148–149, 148f, 150f
staggered, 144–151, 146f, 147f, 148f, 150f, 314, 314f
Congo red, 1037
Coniine, 1004, 1004f
Conivaptan, 867
Conjugate acid, 63–65, 256–257, 256t
Conjugate addition, 617, 981, 983
Conjugate base, 63–65, 67, 67t
Conjugated dienes, 613–632
addition reactions
Diels–Alder reaction, 621–630, 622f
electrophilic, 616–618
kinetic versus thermodynamic product, 618–621
carbon–carbon bond length in, 614–615
conformation of
s-cis, 613
s-trans, 613
electrostatic plot for, 606, 606f
features of, 614
hydrogenation of, 615–616
interesting examples of, 614
stability of, 615–616, 616f, 659
stereoisomers of, 613
synthesis in Suzuki reaction, 1056
ultraviolet (UV) light absorption, 630–632, 631f
Conjugated double bond, 966
Conjugated proteins, 1187
Conjugated systems
allylic carbocation, 605, 606–608
1,3-dienes, 605–606, 606f
Conjugation, 604–640
definition of, 605
β-dicarbonyl compounds, 926
effect on carbonyl group IR absorption, 824, 824f
Hückel's rule and, 652
Conrotatory, 1083–1087, 1087t
Constitutional isomers
of alkanes, 130–132, 131t
of alkenes, 345
definition of, 19
of dipeptides, 1165
in elimination reactions, 299
features of, 177, 202
ketone and aldehyde, 440
of monosaccharides, 1108
stereoisomers compared, 177f
tautomers, 437–439
Contraceptives, 2f. *See also* Oral contraceptives
Cope rearrangements, 1095–1096
Copolymers, 1207, 1212
Copper
electronegativity value of, 783
organocuprate reagents, 783, 784, 1050–1052
Copper(I) cyanide, 1031
Corey, E. J., 372, 918
Corn
ethanol production from, 339, 339f
use in green polymer synthesis, 1219, 1219f
Coronary artery disease, 164, 190, 395
Cortisol, 332, 1255
Cortisone, 826, 991, 1255
Diels–Alder reactions, 629
Cotton
cellulose in, 1138
structure of, 902, 902f
Coumarin, 992
Counterions, 62
Coupling constant, 543, 549
Coupling reactions
alkenes with organic halides (Heck reaction), 1050, 1056–1058
of aryl diazonium salts, 1029, 1034–1035
of organocuprate reagents, 1050–1052
Covalent bonding
description of, 11, 12–13
in Lewis structures, 13–16
Covalent molecules
boiling points of, 103
intermolecular forces, 99–102, 100f, 102t
melting points of, 105
solubility of, 108f
COX-1, 1243f, 1244
COX-2, 1243f, 1244
CPC (cetylpyridinium chloride), 255
Cracking, 391
Crafts, James, 679
Creatine, 54
p-Cresol, 727, 1229
Crocin, 1258
Crossed aldol reactions, 967–971
synthetically useful reactions, 968–970, 970f
with two different aldehydes, both having α H atoms, 968
useful transformations of aldol products, 970, 971f
Crossed Claisen reactions, 977–979
Cross formula, 1109
Crown ethers, 115, 340–341, 340f
Crude oil, 141–142, 142f, 144
Crutzen, Paul, 584
Crystallites, 1216
CTX3C (ciguatoxin), 764, 779, 779f
Cubane, 141f, 152
Cucumber aldehyde, 45, 822
Curcumin, 674
Curved arrow notation, 21–22, 60, 63, 64, 83, 218, 220–221, 220t, 254, 1081
Cyanide
addition to aldehydes and ketones, 833–834
addition to imines, 1158
α-Cyano carbonyl compounds, 970
2-Cyanocyclohexanecarboxylic acid, 875f
Cyano group, 834
of nitriles, 870, 875
Cyanohydrins
in Kiliani–Fischer synthesis, 1131, 1133
naturally occurring derivatives, 834
synthesis by cyanide addition to aldehydes and ketones, 833–834
in Wohl degradation, 1131, 1132
Cyclic anhydrides, 870, 889f, 890
Cyclic compounds
entropy decrease with formation of, 228
stereogenic centers in, 183–185
Cyclization, of hydroxy aldehydes, 853, 855–856
Cyclo- (prefix), 132, 138
Cycloaddition reactions, 1087–1091. *See also* Diels–Alder reaction
[2 + 2] cycloadditions, 1088, 1090–1091
[4 + 2] cycloadditions, 1088, 1089–1090
description of, 1077
orbital symmetry and, 1088–1089
Woodward–Hoffmann rules, 1091, 1091t
Cycloalkanes
conformations of, 151–161
disubstituted, 158–161
monosubstituted, 156–158
disubstituted, 195–196
molecular formula, 129, 132, 385
naming, 138–140, 140f
NMR spectra, 533–534
stereoisomers, 158–161
structure of, 129, 132, 152f
substituted, 156–161, 157f, 160f
Cycloalkenes, 384
cis and trans geometry of, 385
nomenclature, 388, 388f
Cycloalkynes, 428
Cyclobutadiene, 652–653, 660, 665–666
Cyclobutane
conformation, 151
structure of, 132, 152f
Cyclobutene, 1077, 1081, 1082, 1086, 1088
1-Cyclobutylhexane, 140f
Cyclodecane, structure of, 152f
α-Cyclogeraniol, 425
Cycloheptane, structure of, 152f
Cycloheptа-1,3,5-triene, 652, 659
Cycloheptatrienone, 1088
Cyclohexa-1,3-diene, 652, 1077, 1081
heat of hydrogenation, 650
stability of, 653
ultraviolet absorption, 630
Cyclohexadienone, 1096
Cyclohexanamine, 640, 999
ammonium salt of, 1015, 1015f
in Hofmann elimination reaction, 1025
separation from cyclohexanol, 1015, 1015f
Cyclohexane(s)
conformations of, 152–161
boat, 155, 155f
chair, 152–161, 153f, 315–316, 315f
disubstituted cyclohexane, 158–161, 160f
free energy change and, 227
monosubstituted cyclohexane, 156–158, 157f
ring-flipping, 155–156, 155f, 160
conversion to cyclohexene, 581
E2 elimination in, 315–317
formation from cyclohexene, 236–237
in mango, 129
NMR spectrum, 552
skeletal structures, 32
structure of, 129, 132, 139
Cyclohexanecarbaldehyde, 818
Cyclohexanecarbonyl chloride, structure of, 873
Cyclohexane-1,2-diols, 472
cis-cyclohexane-1,2-diol, 473
Cyclohexane-1,4-diol, 1227
Cyclohexane-1,3-dione, 760
Cyclohexanol, 523, 862
cyclohexane formation from, 418
dehydration of, 351, 417
1,2-dibromocyclohexane synthesis from, 417–418
separation from benzoic acid by extraction procedure, 749–751, 751f
separation of cyclohexanamine from, 1015, 1015f
synthesis from cyclohexene, 404
Cyclohexanone, 523, 526, 803
aldol reactions of, 970, 971f
alkylation of, 943
conversion to an alkene, 839
enolate formation from, 929
halogenation of, 941
nucleophilic addition reactions, 830, 831f
structure, 598
tertiary alcohol formed from, 787
Cyclohexene, 688
addition reactions of, 397f, 402, 404, 405
bromination of, 586
1,2-dibromocyclohexane synthesis from, 417
dihydroxylation of, 472
heat of hydrogenation, 650
as Lewis base, 83
reaction with hydrogen, 236–237
stereoisomers, 624
synthesis
cyclohexane conversion, 581
from cyclohexanol, 418
by dehydration of cyclohexanol, 351
as synthetic intermediate, 418
Cyclohex-2-en-1-ol, 381, 523
Cyclohex-3-en-1-ol, 381
Cyclohex-2-enone, 523, 526, 569, 865
formation in Robinson annulation, 982
reduction of, 772
synthesis of, 941
Cyclohexyl acetate, 915
Cyclohexylamine. *See* Cyclohexanamine
Cyclononatetraenyl anion, 672
Cyclononatetraenyl cation, 672
Cyclononatetraenyl radical, 672
Cycloocta-1,5-diene, 1095
Cyclooctane, 203, 203f
Cyclooctatetraene, 652, 672
NMR spectrum of, 654

Cyclooctene, cis and trans isomers of, 385
Cyclooctyne, 428
Cyclooxygenase, 738, 1243, 1243f, 1244
Cyclopentadiene, 625–626, 628, 659
Cyclopentadienyl anion, 658–660, 664
Cyclopentadienyl cation, 659
Cyclopentadienyl radical, 659
Cyclopentane, 575–576
  structure of, 132, 152f
Cyclopentanecarboxylic acid, 948
cis-Cyclopentane-1,2-diol, 473
trans-Cyclopentane-1,2-diol, 335
Cyclopentanol, 109
Cyclopentanone, 862
Cyclopentene, chlorination of, 406
Cyclopropane(s)
  conformation, 151
  dihalo to dialkyl conversion, 1061
  NMR spectrum of, 533
  structure of, 132
  synthesis of
    in Simmons–Smith reaction, 1061–1062
    using carbene intermediates, 1058–1061
Cyclopropenone, 864
Cyclopropenyl radical, 671
Cysteine
  abbreviation for, 753t
  isoelectric point for, 1156t
  in peptide formation, 1165
  pK_a values for, 1156t
  structure of, 1154f
Cytidine, 1142
Cytidine monophosphate, 1142
Cytosine, 1142, 1143f

D

2,4-D (2,4-dichlorophenoxyacetic acid herbicide), 682f
Darunavir, 867
Darvon, 121. *See also* Propoxyphene
DBN (1,5-diazabicyclo[4.3.0]non-5-ene), 305, 305f, 320
DBU (1,8-diazabicyclo[5.4.0]undec-7-ene), 89, 305, 320
DCC. *See* Dicyclohexylcarbodiimide
d (dextrorotatory) compounds, 199
DDE (dichlorodiphenyldichloroethylene), 297, 299
DDT (dichlorodiphenyltrichloroethane), 4, 233, 252
  biological effects of, 297
  DDE formed from, 297
  degradation of, 299
Dean–Stark trap, 848, 848f
(2*Z*,4*Z*)-2,4-Decadienoic acid, 1218–1219
Decalin, 172, 491, 1250–1251
Decamethrin, 1059
Decane, 128, 138
  constitutional isomers of, 131t
  molecular formula of, 131t
Decarboxylation, 1157
  in acetoacetic ester synthesis, 949–950
  of β-acids, 946
  of β-keto acids, 946
  in malonic ester synthesis, 946–948
Decongestants, 1016, 1046
DEET, 886
Dehydrating agent, 893
Dehydration reactions
  of alcohols to alkenes, 345–352, 395
    alkene synthesis in, 345–352
    carbocation rearrangements in, 348–350
    E1 mechanism in, 346–347
    E2 mechanism in, 347, 351
    enthalpy change in, 347, 348f
    Le Châtelier's principle in, 347
    in phosphorus oxychloride and pyridine, 351, 352f
    in strong acid, 345–350
  of aldol product, 965–967
  defined, 345
  dicarboxylic acid conversion to cyclic anhydrides, 890
  as β-elimination reaction, 345
  fragmentation pattern in mass spectrum, 503
7-Dehydrocholesterol, 1086
Dehydrohalogenation
  alkene synthesis by, 298
  of alkyl halides with strong base, 395
  alkyne synthesis by, 318–319, 319f
  bases commonly used in, 298, 298t
  drawing products of, 299
  E1 mechanism of, 310
  E2 mechanism of, 303, 309
16,17-Dehydroprogesterone, 1259
Delocalization, resonance, 75
Delta (δ) carbon, 732
Delta (δ) scale, 530, 531
Dementia, 99, 973
Demerol. *See* Meperidine
Dendrobates histrionicus (poison dart frog), 431, 431f
Deoxy (prefix), 1141
2-Deoxyadenosine, 1142
Deoxyadenosine monophosphate, 118, 1107, 1107f, 1142
2-Deoxy-D-ribose, 1141
Deoxyribonucleic acid (DNA), 118, 118f, 1141, 1142–1144, 1143f, 1144f
Deoxyribonucleosides, 1141–1142
Deoxyribonucleotides, 1142
Deprotection, 792
Deprotonation
  in acetoacetic ester synthesis, 950
  of acetylene, 446
  of alkyne, 433–434, 433t, 438–439
  of carboxylic acids, 741–742, 741t
  of diethyl acetamidomalonate, 1157
  in direct enolate alkylation, 942
  in glycoside formation, 1125
  in halogenation at the α carbon of carbonyl compounds, 938–940
  in malonic ester synthesis, 947
  racemization at the α carbon of carbonyl compounds, 936
  in tautomerization, 927
Deshielding effects, in NMR, 536–537, 536f, 537f, 540f, 540t
DET (diethyl tartrate), 483–484
Detergents, 113
Deuterium, 8
Deuterochloroform, 529f
Dexamethasone, 97
Dextropimaric acid, 1258
Dextrorotatory, 1113
DHA, 491
Di- (prefix), 136, 335, 875, 999
Diabetes, 825
Diacids, 733, 945–947
  formation from monosaccharide oxidation, 1129, 1130–1131
  in nylon synthesis, 902, 1211
Diacon. *See* Methoprene
Dialkylamides, 932
Dialkylborane, 412
Diamine, in nylon synthesis, 902, 1211
1,4-Diaminobenzene, 1212
1,6-Diaminohexane, 1211, 1218, 1218f
Diamond, 666
Diastereomers, 190–192, 192f, 832
  alkenes, 385t, 388–389
  of but-2-ene, 301
  definition of, 192
  disubstituted cycloalkanes, 195–196
  epimers, 1114
  of glucose, 1118, 1119f
  halogenation reactions and, 583
  of monosaccharides, 1114
  NMR spectrum of, 534–535
  nucleophilic substitution reactions and, 280
  physical properties of, 201–202, 201f
  separation of amino acid enantiomers into, 1160–1161, 1160f
Diastereotopic protons, 534–535
1,3-Diaxial interactions, 157
Diazepam, of structure, 649f
Diazomethane, 1061
Diazonium salts, 1027–1035
  alkyl, 1028
  aryl, 1028–1035
  coupling reactions of, 1029, 1034–1035
  explosiveness of, 1028
  formation from primary amines, 1028
  substitution reactions of, 1029–1033
    synthesis of aryl chlorides and bromides, 1030
    synthesis of aryl fluorides, 1030
    synthesis of aryl iodides, 1030
    synthesis of benzene, 1031
    synthesis of benzonitriles, 1031
    synthesis of phenols, 1030
  uses in synthesis, 1032–1033
Diazotization, 1028, 1031
DIBAL-H (diisobutylaluminum hydride), 777–779, 779f, 781t, 909–910
Dibromobenzene
  meta-, 645, 647, 647f
  ortho-, 645, 647, 647f
  para-, 645, 647, 647f
1,4-Dibromobutane, 948
2,2-Dibromobutane, 434–435, 451
2,3-Dibromobutane, 181, 193–194, 194f
Dibromocarbene, 1061
1,2-Dibromocyclohexane, synthesis from cyclohexanol, 417–418
1,3-Dibromocyclopentane, 195–196
2,3-Dibromopentane, stereoisomers of, 191–192, 192f, 194
Di-*tert*-butyl dicarbonate, 1174
Dibutyl phthalate, 1217, 1226
Dicarbonyl compounds, 475
  aldol reactions with, 973–975
  1,4-dicarbonyl compounds, 973
  1,5-dicarbonyl compounds, 974, 975f
    in Robinson annulation, 983
  β-dicarbonyl compounds, 969–970, 970f
    pK_a values, 930, 931t
    synthesis in crossed Claisen reactions, 978
    tautomers, 926
2,5-Dichloroaniline, 646
4,4'-Dichlorobiphenyl, 108
Dichlorocarbene, 1059
1,2-Dichlorocyclopentane, 406
Dichlorodifluoromethane, 584
Dichlorodiphenyldichloroethylene. *See* DDE
Dichlorodiphenyltrichloroethane. *See* DDT
1,1-Dichloroethylene, NMR spectrum of, 533
Dichloromethane, 251f, 477
  as solvent in extraction procedure, 749, 749f, 751f
2,4-Dichlorophenoxyacetic acid herbicide (2,4-D), 682f
trans-1,3-Dichloropropene, 551
Dictyopterene, 491
Dicyclohexylcarbodiimide (DCC), 889f, 893–894, 1173, 1176, 1177
Dicyclohexylurea, 893
Dicyclopentadiene, 628
Dieckmann reaction, 979–980
Dieldrin, 637
Diels, Otto, 621
Diels–Alder reaction, 244, 621–630
  cycloaddition, 1088, 1090
  diene reactivity in, 623
  dienophile reactivity in, 623–24
  drawing product of, 622–623
  features of, 621
  retro, 628
  retrosynthetic analysis of product, 627–628
  rule of endo addition, 625–627
  rules governing, 623–627
  stereospecificity of, 624–625
  in steroid synthesis, 629
  tetrodotoxin synthesis, 622, 622f
  as thermal reaction, 621
Dienes, 387
  conjugated, 613–632. *See also* Conjugated dienes
  Cope rearrangements, 1094, 1095
  cycloaddition, 1088
  isolated, 605–606, 606f, 614, 616, 616f, 617, 650
  in metathesis, 1065
  ozonolysis of, 475
  reactivity, 623
  stability, 650
1,3-Dienes, 605–606, 606f, 613. *See also* Conjugated dienes
  polymerization of, 1211
  synthesis in Suzuki reaction, 1056
Dienophiles, 1088
  cis, 622, 623–624, 625, 628
  cyclic, 623, 625
  Diels–Alder reaction, 621–628, 624f
  reactivity of, 623–624
  stereochemistry, 624–625
  trans, 623, 624
Diesel fuel, composition of hydrocarbons in, 142
Diesters
  in Dieckmann reaction, 979–980
  1,3-diesters, pK_a of, 931t
  1,6-diesters, in Dieckmann reaction, 979
  1,7-diesters, in Dieckmann reaction, 980

β-diesters
as active methylene compound, 970
synthesis in crossed Claisen reaction, 978
Diethyl acetamidomalonate, 1157
Diethylamine, 886
solubility of, 109
Diethylbenzene, 672
Diethyl carbonate, in crossed Claisen reaction, 978
cis-1,2-Diethylcyclopropane, 1062
Diethyl ether, 96, 1116
as anesthetic, 251f, 340
boiling point of, 822
in extraction procedures, 749
Grignard reagent preparation, 783
infrared (IR) spectrum, 515
naming, 335
solubility of, 109, 126
as solvent, 106, 362, 749
Diethyl maleate, 1074
Diethyl malonate, 969, 970, 970f, 979
alkylation of derivatives, 1157–1158, 1159f
conversion to a carboxylic acid, 945–949
N,*N*-Diethyl-*m*-toluamide, 886
Diethyl oxalate, 993
Diethyl sulfide, 365
Diethyl tartrate (DET), 483–484
Digoxin, 817, 855
Dihalides, 318–319, 319f
alkyne synthesis from, 431–432
Dihalocarbenes, 1059–1061
Dihedral angle, 145, 146f, 147f, 148, 150f, 314
Dihydroxyacetone, 1108, 1109, 1115, 1116f
Dihydroxyacetone phosphate, 928
1,4-Dihydroxybenzene, 1215
Dihydroxylation, 468f, 472–474
anti, 472–473
syn, 472, 473–474
1,4-Dihydroxymethylcyclohexane, 903
1,4-Dihydroxynapththoic acid, 690, 690f
6,7-Dihydroxynonan-2-one, 866
Diiodomethane, 1061
Diisobutylaluminum hydride (DIBAL-H), 777–779, 779f, 781t, 909–910
Diisocyanate, 1213
Diisopropylamine, 933, 999
β-Diketone, pK_a of, 931t
Dimers, carboxylic acids as, 734, 734f
2,2-Dimethoxycyclopentane-1,3-dicarbaldehyde, 828
1,2-Dimethoxyethane, 531
5,5-Dimethoxypentan-2-one, 863
Dimethylacetylene dicarboxylate, 640
Dimethylallyl diphosphate, 608, 1247
Dimethylamine, 1029
4-(*N*,*N*-Dimethylamino)pyridine, 1021
N,*N*-Dimethylbenzamide, 875
1,3-Dimethylbenzene, 704
2,3-Dimethylbutanal, 819
3,3-Dimethylbutan-2-ol, 348, 349
2,2-Dimethylbut-3-enal, 822
Dimethyl carbonate, 811
cis-3,4-Dimethylcyclobutene, 1082, 1084, 1086
trans-3,4-Dimethylcyclobutene, 1084
cis-5,6-Dimethylcyclohexa-1,3-diene, 1082–1083
trans-5,6-Dimethylcyclohexa-1,3-diene, 1082–1083
1,2-Dimethylcyclohexane, 140f
1,3-Dimethylcyclohexane, 139
1,4-Dimethylcyclohexane, 158–160, 160f
2,2-Dimethylcyclohexanone, 943, 944
2,6-Dimethylcyclohexanone, 943
1,6-Dimethylcyclohexene, 388f
1,2-Dimethylcyclopentane, 158
cis isomer, 177f
trans isomer, 177f
1,3-Dimethylcyclopentane, 202
(*S*,*S*)-1,2-Dimethylcyclopropane, 599
Dimethylcyclopropanes, 210, 533
Dimethyl ether
isomers of, 18–19
NMR spectrum of, 557
Dimethylformamide (DMF)
boiling point of, 104
as solvent, 260f, 941
2,5-Dimethylhept-3-yne, 429f
2,5-Dimethylhexa-2,3-diene, 454
4,5-Dimethylhexanoic acid, 731
2,4-Dimethylhexan-3-one, 804
2,4-Dimethylhexanoyl chloride, 876
2,5-Dimethylhex-3-yne, 454
6,6-Dimethyloct-3-yne, 429
2,2-Dimethyloxirane, 337, 370
2,3-Dimethyloxirane, 369
2,3-Dimethylpentane, 137f
mass spectrum of, 502–503
2,2-Dimethylpropane (neopentane), 100, 100f, 106, 130
2,2-Dimethylpropanoic acid, 745
Dimethyl succinate, 542
Dimethyl sulfide, conversion of ozonide to carbonyl compounds, 474–476
Dimethyl sulfoxide (DMSO), 19
aqueous, 409
as polar aprotic solvent, 260f, 282t
as solvent for alkyne synthesis, 319, 319f
Dimethyl terephthalate, 1213, 1221
1,3-Dinitriles
active methylene compounds, 970
pK_a, 931t
Dinoprostone, 1244
-dioic acid (suffix), 733
-diol (suffix), 335
Diol epoxide, 373
Diols, 335
1,2-diol formation by dihydroxylation, 468f, 472–474
gem-diol, 844–847
p-Dioxanone, 1229
Dioxybenzone, 124
Dipeptides
constitutional isomers of, 1165
synthesis of, 1164–1165, 1172–1176
Diphenhydramine, 293, 644, 1016
Diphenyl carbonate, 1214, 1220
Diphosphate, 283, 608
as leaving group, 608, 1247, 1249
Dipole, 48, 49, 50f
in alkenes, 391
temporary, 100–101
Dipole–dipole interactions, 101, 102t
of aldehydes and ketones, 822
of alkyl halides, 250
of carbonyl compounds, 877
of carboxylic acids, 734
of nitriles, 877
in polar aprotic solvents, 260, 281
in polar protic solvents, 259
Dipole moment, infrared (IR) absorption and, 512
Directed aldol reactions, 971–973, 972f
Disaccharides, 1134–1137
artificial sweeteners, 1137, 1137*f*
general features, 1134–1135
lactose, 1130, 1136
maltose, 1135
sucrose, 1136–1137, 1136f
Discodermolide, 212
Disorder, entropy as measure of, 227–228
Disparlure, synthesis of, 470–472, 471f
Disproportionation, 1202
Disrotatory, 1083–1087, 1087t
Dissolving metal reductions, 457, 465–466
Distillation, of crude petroleum, 142, 142f
Distillation apparatus, 104, 105f
trans-1,2-Disubstituted cycloalkanes, 368, 369
Disulfide bonds, 365
in α-keratins, 1186, 1187f
in peptides, 1168
in tertiary structure of proteins, 1184, 1184f, 1185f
in vulcanized rubber, 1210, 1210f
Diterpene, 1246t, 1248
cis-1,2-Divinylcyclobutane, 1095
Diynes, 428
DMAP, 1021
DMF. *See* Dimethylformamide
DMSO. *See* Dimethyl sulfoxide
DNA, 118, 118f, 1141, 1142–1144, 1142f, 1143f
Dobutamine, 999
Docetaxel, 920
Dodecahedrane, 141, 152, 640
Dodecane, 131t
Dolastatin, 98
Domagk, Gerhard, 1037
Donepezil, 99, 973
Dopamine, 7, 863, 1005–1006, 1006f
Double bonds. *See also* Alkenes
in alkenes, 93, 94t, 299–301
in alkyl halides, 248
bond length and bond strength, 45–46, 46t
components of carbon–carbon, 42, 42f
in condensed structures, 31, 31f
conjugated, 966
conjugated dienes, 613
degrees of unsaturation, calculating, 460–461
effect on melting point of fatty acids, 391
in fats and oils, 395
in functional group, 93, 94t
hydrogenation of, 461–462, 462f
Lewis structure, 15–16
metal–carbon, 1064
nitrogen–nitrogen, 1029, 1034
NMR spectra of, 539, 540t
pi (π) bonds, 93, 97
planar, 403, 406, 412, 414
properties of, 385t
radical reactions with, 573, 590–593
in resonance structure, 21–25
resonance structures
conjugated, 609
major contributor, 610–611, 615
minor contributors, 610–611, 615
with one atom more electronegative than the other, 610
rotation restriction, 42, 300–301, 301f, 385t
in trans fats, 462–463
in triacylglycerols, 1234–1236, 1234t
in unsaturated fatty acids, 383, 393–395
Z configuration, 393
Double dagger notation, 230
Double-headed arrow, 19, 63–64, 220t
Doublet, NMR spectrum, 542–543, 544t, 545t, 546, 547
Doublet of doublets, NMR spectrum, 550, 550f
Doxorubicin, 1107, 1107f
Dyes
azo compounds, 1034, 1036–1037
natural and synthetic, 1036–1037
sulfa drugs, 1037–1038, 1038f
Dyneema, 1209

E

E (*prefix*), 389–390
E1cB elimination reactions, 966, 983
Eclipsed conformation, 144–151, 146f, 147f, 148f, 150f, 150t, 191, 301f, 314f
E configuration, double bond, 1210
Edman degradation, 1169–1170
E1 elimination reactions, 303, 310–313
acid-catalyzed dehydration of alcohols, 395
base in, 312, 312t, 313, 317, 317t
characteristics of, 312t
dehydration of secondary and tertiary alcohols, 346–347
E2 mechanism compared, 310, 313, 317, 317t, 319–323, 320t
energy diagram for, 310, 311f
kinetics of, 310, 312t
leaving group in, 310, 312t
rate of reaction, 311–312, 312t
regioselectivity of, 312
S_N1 substitution reactions compared, 313, 319–323, 320t
S_N2 substitution reactions compared, 319–323, 320t
solvent for, 312t
two-step mechanism, 310–311, 312t
E2 elimination reactions
acetylide anion reactions with alkyl halides, 441–442
alkyl halide identity in, 306–307, 307t
alkyl tosylates reactions, 360
alkyne synthesis, 318–319, 431–432
anti periplanar geometry in, 314–315, 314f
base in, 305, 305f, 307t, 317, 317t
characteristics of, 307t
dehydration of alcohols, 347, 351
dehydrohalogenation of alkyl halides, 395
E1 mechanism compared, 310, 313, 317, 317t, 319–323, 320t
energy diagram of, 304, 304f
in Hofmann elimination, 1025–1027, 1027f
kinetics of, 303, 307t
leaving group in, 304–306, 307t
one-step mechanism, 304–305, 307t

E2 elimination reactions (*Continued*)
organic synthesis examples, 307f
rate of reaction, 305, 306
regioselectivity of, 308
S_N1 and S_N2 substitution reactions compared, 319–323, 320t
solvent for, 305, 307t
stereochemistry of, 314–317, 314f
stereoselectivity of, 309
syn periplanar geometry in, 314, 314f
trans diaxial geometry in, 315–316, 315f
Zaitsev rule, 308–309, 316
Effexor. *See* Venlafaxine
E-85 fuel, 339, 339f
Ehrlich, Paul, 1037
Eicosanoids, 1242–1245
biological activity of, 1242, 1244t
biosynthesis of, 1243, 1243f, 1244
derivation of word, 1242
E isomer, 389
Elaidic acid, 421
Elastomer, 1210
Electrocyclic reaction, 1081–1087
description of, 1077
photochemical, 1085–1086
stereochemistry and orbital symmetry, 1082–1083
thermal, 1083–1085
Woodward–Hoffmann rules for, 1086, 1087t
Electrocyclic ring closure, 1077, 1081–1086
Electrocyclic ring opening, 1077, 1081–1082, 1084
Electromagnetic radiation, 506–508, 507f
Electromagnetic spectrum, 507, 507f
Electron cloud, 8
Electron-deficient sites/species, 48–49, 50f, 65, 81–83, 92, 95, 116–117, 219, 766
Electron density, delocalization, 74
Electron-donating groups, 276, 691–693, 697, 702, 744–748, 748f
Electron dot representation. *See* Lewis structures
Electronegativity
acidity trends and, 70–73
bond polarity and, 48–49
definition of, 47
inductive effects on substituted benzenes, 691
oxidation and, 161
periodic trends in, 47, 47f
values for common elements, 47f
Electron pair acceptor, 62, 81
Electron pair donor, 62, 81
Electron pairs
in Brønsted–Lowry acid–base definition, 62
Lewis acids and bases, 81–83
movement of, 63, 218
Electron-poor sites/species, 82
Electron-rich sites/species, 48, 50f, 65, 82, 116–117, 396
Electrons
bonded (shared), 13
delocalized, 20, 24, 586, 607, 612, 616, 643, 655–657
density of, 9, 36, 43, 48–49, 49f, 100
excited state, 630–631
ground state, 630–631
in Lewis structure, 13–19
lone pairs. *See* Lone pairs
nonbonded (unshared), 13, 13f
number of, 8, 11
in resonance structures, 19–25
unpaired, 218
valence, 11–13, 18t
Electron-withdrawing groups, 276, 623–624, 691–693, 697, 702–703, 706–708, 744–748, 748f
Electrophiles, 97
alkene reaction with, 396
alkyne reactions with, 432
carbene, 1059
carbocations as, 219
Lewis acid, 82
meaning of term, 82
radicals as, 219
reactivity of, 116–117
Electrophilic addition reactions, A–13
of conjugated dienes
kinetic versus thermodynamic products, 618–621, 620f
1,2- versus 1,4-addition, 616–618
energy diagram for, 399, 399f
halogenation of alkynes, 436
hydration of alkenes, 404–405, 415
hydrohalogenation of alkenes, 397–404, 397f, 398f, 399f, 401f, 403t
hydrohalogenation of alkynes, 434–436
Markovnikov's rule on, 400–402, 401f, 403t
stereochemistry of, 402–404, 403t
Electrophilic aromatic substitution reactions, 678–706, A–12
azo coupling, 1034–1035
energy changes in, 680, 680f
examples of, 679f
Friedel–Crafts acylation, 679f, 684–690, 689f, 703, 712–713, 713f, 717–718
Friedel–Crafts alkylation, 679f, 684–690, 702, 712–713, 713f, 715–716
halogenation, 679f, 681–682, 701–704
mechanism of, 679–680
nitration, 679f, 682–683, 705–706
of substituted benzenes, 691–706
disubstituted benzenes, 703–704
limitations on reactions, 701–703
rate of reactions, 696, 697, 697f
sulfonation, 679f, 682–683
Electrophilic carbon, in alkyl halides, 253
Electrospray ionization (ESI), 506
Electrostatic interactions
in ionic compounds, 99
between ions, 11–12
Electrostatic potential plot
of acetic acid, 75f, 730, 730f
of acetylene, 432, 433f
of alcohol, 333f
of alkoxide, 930, 930f
of amines, 998, 998f
of benzene, 644, 644f
of carbanions, 76f
of carbocations, 274, 274f
localized and delocalized, 607, 607f
carbon dioxide, 50f
description of, 48
of dienes, 606, 606f
of enolate, 930, 930f
of epoxide, 333f
of ethanol, 73f, 75f
of ether, 333f
of ethylene, 396, 396f
of formaldehyde, 766, 766f
of halomethanes, 253f
of methyl chloride, 49, 49f
of pyridine and pyrrole, 657, 657f
of substituted anilines, 1018, 1019f
of 2,2,2-trifluroethanol, 73f
of vitamins A and E, 1242, 1242f
of water, 50f
Elements, periodic table, 8–10, 9f
Element symbol, 8
Elemicin, 121
Eleostearic acid, 421
Eletriptan, 1070
Elimination reactions, A–13
of alcohols, 343–352, 361, 362f
alkene synthesis, 298–299, 395–396
of alkyl halides, 253, 297–330
alkyne synthesis, 298, 318–319, 319f, 431–432
alpha (α), 1059
beta (β), 298
bimolecular, 303. *See also* E2 elimination reactions
description of, 215
elimination, unimolecular, conjugate base (E1cB) mechanism, 966, 983
features of, 298–299
Hofmann elimination, 1024–1027
mechanisms of, 303
E1. *See* E1 elimination reactions
E2. *See* E2 elimination reactions
E1 and E2 compared, 310, 317, 317t, 319–323, 320t
reductive, 1053, 1055, 1058
stereochemistry, 314–317
unimolecular, 303, 310. *See also* E1 elimination reactions
Zaitsev rule on, 308–309, 312
Enalapril, 1012
Enals, 822
Enamines, 450
hydrolysis of, 844
synthesis of
from aldehydes and ketones, 831f, 843–844, 843f, 1023
Enanthotoxin, 30
Enantiomeric excess, 200–201, 483, 775
Enantiomers
of amino acids, 752, 1153, 1153f
separation of, 1159–1163, 1160f
of an ammonium salt, 998
in carbonyl reductions, 773–774
chemical properties of, 202–204
chiral drugs, 202–203
of cyclic compounds, 184
cyclic hemiacetals, 853
definition of, 179, 190, 192
dienophile, 624
dihydroxylation, 472
disubstituted cycloalkanes, 196
drawing, 182–183, 183f
in electrophilic addition to alkenes, 402–403, 403t
enantioselective reactions, 482–483
enantiotopic protons, 534–535
in enolate, 936, 943
from epoxidation of alkenes, 469, 472
in epoxide ring opening reactions, 368–369
formation in halogenation of alkenes, 405–407
halogenation reactions and, 582–583
in halohydrin formation, 409
in hydroboration–oxidation reactions, 414
labeling stereogenic centers with *R* or *S*, 185–190, 187f, 189f, 194–195
of monosaccharides, 1109–1110, 1112–1114, 1114f
in nucleophilic substitution reactions, 265, 271–272
optical purity, 200–201
in organometallic reagent reactions, 788
physical properties, 197–202, 199t, 201f
racemic mixture of, 199, 199t, 402–403
sense of smell and, 203–204, 203f
three-dimensional representations for, 183f
Enantioselective reactions, 482–483
amino acid synthesis, 1163–1164
biological reduction, 776–777
carbonyl reductions, 774–777, 775f
Enantiotopic protons, NMR signals of, 534–535
Enclomiphene, 420
Endergonic reaction, 228
Endiandric acids, 1105
Endo addition, 625–627, 627f
Endo product, 625–626, 627f
Endothermic reaction
bond breakage and, 221–222
bromination, 579–580
dehydration of alcohols, 348f
energy diagram of, 230f
enthalpy change and, 228
reaction rate, 276
transition state in, 276, 276f
-ene (suffix), 387
Energy
nonrenewable, 142
of photons, 506, 507–508
release in bond formation, 219
stored in triacylglycerols, 1237
torsional, 147
units of measurement, 45
Energy barrier, 230–232, 234, 304, 310, 311f
Energy change
during the chlorination of ethane, 576–577, 577f
during radical addition of HBr, 592–593, 592f
Energy diagrams, 228–233, 233f
for carbocation formation, 278f
for chlorination of ethane, 577f
for E1 elimination reactions, 310, 311f
for E2 elimination reactions, 304, 304f
for electrophilic addition of HBr to buta-1,3-diene, 620, 620f
for electrophilic additions, 399, 399f
for electrophilic aromatic substitution, 680, 680f
for halogenations, 577f, 580f, 581f
rate of electrophilic aromatic substitution of substituted benzenes, 697, 697f

for S_N1 substitution reactions, 270, 270f, 277, 278f
for S_N2 substitution reactions, 264, 264f
for two-step reaction mechanisms, 231–233, 233f
Energy of activation (E_a)
catalysts and, 237, 237f
definition of, 229
energy barrier height determined by, 230
in energy diagrams, 229–233, 230f, 233f
in exothermic reactions, 277, 277f
rate constant, relationship to, 234
reaction rate and, 234
transition state energy effect on, 276
Enol(s), 925–928
hydroxy group of, 332
reaction mechanism, 928
resonance structures, 928
synthesis of, 800
hydration of alkynes, 437–439
hydroboration–oxidation of alkynes, 439–440
tautomers, 926–927
-enol (suffix), A–5
Enolate(s), 22, 828, 925, 928–935
achiral, 936
in aldol reactions, 963, 964, 964f, 965, 966, 968, 970, 970f, 971, 972, 974
alkylation, 942–945, 951
general features of, 942–943
tamoxifen synthesis, 944
of unsymmetrical ketones, 943–944
in Claisen reactions, 975–979
electrostatic potential plot of, 930, 930f
examples, 930
kinetic, 934, 935, 943, 971
in Michael reactions, 980–982, 982f
racemization, 936, 936f, 943
reactions of, 933, 937, 942–945
resonance-stabilized, 966, 976, 981
resonance structures, 928
in Robinson annulation, 982–986
synthesis of, 800
with bases, 931–933, 932t
from carbonyl compounds, 928–929
from esters, 930
from tertiary amides, 930
thermodynamic, 934, 935, 943, 944
of unsymmetrical carbonyl compounds, 934–935
Enol tautomers, 437–439, 908, 926–927
-enone (suffix), A–5
Enones, 822
Entacapone, 59
Enthalpy change ($\Delta H°$), 221–225
calculation from bond dissociation energies, 223–224
in dehydration of alcohols, 347, 348f
E1 elimination reactions, 310, 311f
in energy diagrams, 229–233, 230f, 233f
free energy change, relation to, 227–228
negative, 221, 228
positive, 221, 228
Entropy ($S°$), 227–228
Entropy change ($\Delta S°$), 227–228
-enyne (suffix), A–5
Enynes, 428
Enzymes, 237–238, 238f, 1153
active site of, 238, 238f
catalytic action of, 237–238, 238f
in kinetic resolution of amino acids, 1162
in oxidation reactions, 481–482
in peptide sequencing, 1171, 1171t
Enzyme–substrate complex, 238, 238f
Ephedrine, 181, 208
Epi-aristolochene, 1260
Epichlorohydrin, 380, 1214–1215, 1215f
Epilupinine, 1048
Epimers
definition of, 1114
of monosaccharides, 1114, 1132–1133
Epinephrine
structure of, 1005
synthesis of, 285–286, 285f
Eplerenone, 341
Epothilone A, 1065f
Epoxidation, 468f, 469–472
in anti dihydroxylation, 472
Sharpless, 482–484
of squalene, 1252f, 1253
stereochemistry of, 469–470
in synthesis of disparlure, 470–472, 471f
Epoxides, 296
achiral, 472
from alkene oxidation, 468f, 469–472
angle strain, 333, 344
anionic polymerization of, 1207–1208
applications, 371–373
boiling point of, 338t
bonding, 333
electrostatic potential plot of, 333f
interesting examples, 341
leaving group, 344
melting point of, 338t
nomenclature, 336–337
physical properties of, 337–338, 338t
reactions of, 367–371
with acetylide anions, 444
with acids HZ, 369–371, 370f
general features, 344
with organometallic reagents, 798–799
reduction, 466–467, 467f
with strong nucleophiles, 367–369
reduction of, 466–467, 467f
ring opening reactions, 1207–1208
with acids, 369–371
enantiomers in, 368–369
nucleophilic substitution reactions, 367–371
regioselectivity of, 369–370
solubility of, 338t
structure of, 332, 333
synthesis of, 343
from halohydrin, 343
Epoxy (prefix), 336
Epoxyalkane, 336
1,2-Epoxycyclohexane, 336, 368
1,2-Epoxy-2-methylpropane, 336
cis-2,3-Epoxypentane, 336
Epoxy resins, 1214–1215, 1215f
Equal. *See* Aspartame
Equations for organic reactions, writing, 214, 214f
Equatorial bond, 176
Equatorial hydrogens, 153–154, 153f, 157f
Equilibrium, 225–227, 226f, 226t
acid–base reactions, 68–69
direction of, 234
nucleophilic substitution and, 257
Equilibrium arrow, 63–64, 69, 220t
Equilibrium constant (K_{eq}), 225–227, 226f, 226t, 234
Ergot, 689f
Erlotinib, 448
Erythrose, 209
D-Erythrose, 1114, 1114f, 1115f
L-Erythrose, 1114, 1114f
D-Erythrulose, 1116f
ESI (electrospray ionization), 506
Esmolol, 89
Essential amino acids, 752
Essential fatty acids, 393
Essential oils, 1245
Esterification, 246
Fischer, 891–892, 903
Esters
acidity/basicity of, 872t
boiling point of, 878t
cholesteryl, 904–905, 905f
chromate, 477–478
cyclic, 870
enolates from, 930
functional group in, 98t
general structure of, 393
interesting examples, 880
IR spectra of, 879t
leaving group, 768
melting point of, 878t
nomenclature of, 873–874, 876t
nucleophilic substitution reactions, 768
odors of, 880
physical properties of, 878t
pK_a, 872t, 931t
polyesters, 903
reactions of, 894–898, 901t
with ammonia and amines to form amides, 894
carboxylic acid preparation by malonic ester synthesis, 945–949
Claisen reaction, 975–979, 976f
hydrolysis to carboxylic acids, 894–896
lipid hydrolysis, 896–898
methyl ketone preparation by acetoacetic ester synthesis, 945, 949–952
reaction with organometallic reagents, 785, 794–796
reactivity, 883
reduction of, 777–779
solubility of, 878t
stability of, 872
structure and bonding of, 765, 869, 872
synthesis of
from acid chlorides, 885
from anhydrides, 888
from carboxylic acids, 889f, 891–892
from monosaccharides, 1127–1128
thioesters, 904–905
in triacylglycerols, 393
waxes, 1233
Estradiol, 56
analogs of, 430
function in body, 1254t
structure of, 430, 1254t
synthesis of, 307, 307f
Estrogens, 1254, 1254t
Estrone, 728
function in the body, 1254t
structure of, 794, 1254t
synthesis of
from chlorohydrin, 410f
in Diels–Alder reactions, 629
using a Michael reaction, 982, 982f
Eth- (prefix), 133
Ethambutol, 56, 207, 268, 268f
Ethanal. *See* Acetaldehyde
Ethane
acidity of, 75–76, 76f
bond dissociation energy, 635
bonds in, 40–41, 44t
chlorination of, 576–577, 577f
conformations of, 144–147, 146f, 147f
energy, 147
molecular formula, 129, 131t
nonreactivity of, 92
structure of, 92, 93, 129
Ethanedioic acid, 733
Ethane-1,2-diol, 335
Ethanethiol, 364
Ethanol, 125
acidity of, 72–75, 742, 744f
in alcoholic beverages, 338, 455
blood alcohol concentration, 518
boiling point, 364
description, 3
electrostatic potential plot, 73f, 75f
in ethyl acetate formation, 236–237
isomers of, 18–19
NMR spectrum of, 532, 551, 551f
oxidation of, 481–482
pK_a of, 742, 744f
as polar protic solvent, 259f
preparation from ethylene, 392f
production by fermentation of carbohydrates, 338–339, 339f
reactivity of, 93
as renewable fuel source, 339, 339f
sodium ethoxide production from, 342
solubility of, 107–108
structure of, 93
Ethene, 390. *See also* Ethylene
Ethers
acetal compared, 847
boiling point of, 338t
bonding of, 333
Claisen rearrangement, 1096
crown, 115, 340–341, 340f
electrostatic potential maps, 333f
formation by alcohol addition to alkenes, 405
functional group, 96t
infrared (IR) spectra of, 515
interesting examples, 340–341, 340f
leaving group, 344
melting point of, 338t
nomenclature, 335–336
physical properties of, 337–338, 338t
preparation by nucleophilic substitution reaction, 286, 287t
reactions of
alkyl halide formation, 362–363
general features, 344
with strong acid, 362–363

Ethers (*Continued*)
simple and complex, 335–336
solubility of, 338t
structure of, 332, 333
symmetrical, 332, 341
synthesis of
from alkyl halides, 341–342
from monosaccharides, 1127–1128
by nucleophilic substitution reactions, 341–342
Williamson ether synthesis, 341, 342
unsymmetrical, 332, 341
Ethoxide, 74, 742, 744f
nucleophilicity of, 258
Ethoxy group, 336
4-Ethoxyoctane, 336
Ethyl acetate, 246
in Claisen reaction, 975–976
formation from acetic acid and ethanol, 236–237
as solvent, 891
structure of, 874
synthesis of, 891
Ethyl acetoacetate, 954
conversion to a ketone, 949–951
synthesis in Claisen reaction, 975
Ethyl *p*-aminobenzoate, 714
N-Ethylaniline, 1000
Ethylbenzene, 645
bromination of, 710
styrene synthesis from, 711
Ethyl benzoate, 876t, 923
in crossed Claisen reactions, 978
Ethyl butanoate, 880
Ethyl chloride, 95, 249
Ethyl chloroformate, in crossed Claisen reaction, 978
Ethyl chrysanthemate, 918
Ethyl α-cyanoacrylate, 1205f
Ethylcyclobutane, 140f
N-Ethylcyclohexanamine, 1013
2-Ethylcyclohexanecarbaldehyde, 819
2-Ethylcyclohexanone, 943
4-Ethyl-3,4-dimethyloctane, 137f
5-Ethyl-2,6-dimethyloctane, 136
Ethylene, 390
acidity of, 75–76, 76f
bond dissociation energies, 427
bonds of, 41–425, 42f, 44t, 384
bromine addition to, 642
electrostatic potential plot, 396, 396f
fruit ripening and, 300
functional group in, 93, 94t
in Heck reaction, 1057
as industrial starting material, 391, 392f
Lewis structure, 15–16
molecular orbitals of, 1078, 1079f
molecular shape of, 27
monomers in polymerization, 593
pK_a of, 68
polymerization of, 1203, 1209
preparation from petroleum, 391
structure, 299–300, 1202f
Ethylene diacetate, 542
Ethylene glycol, 850
acetal formation from, 847
in antifreeze, 339f
formation from PET recycling, 1221
naming, 335
in polyester synthesis, 903
in polyethylene terephthalate (PET), 1198, 1200f, 1212–1213
in polyurethane synthesis, 1213
preparation from ethylene, 392f
structure of, 339f
synthesis, 1228
Ethylene oxide, 337
polymerization of, 1208
ring opening of, 1208
1-Ethyl-2-fluorocyclopentane, 250f
N-Ethylformamide, 875
Ethyl formate, in crossed Claisen reactions, 978
Ethyl group, 134
2-Ethylhep-1-ene, 387
Ethyl hexa-2,4-dienoate, 992
Ethyl isopropyl ether, 342
Ethyl isopropyl ketone, 822f
1-Ethyl-3-methylcyclohexane, 139
Ethyl methyl ether, 545
Ethyl methyl ketone, 821
4-Ethyl-5-methyloctane, 137f
N-Ethyl-*N*-methylcyclopentanamine, 1000
Ethyl *p*-nitrobenzoate, 714
3-Ethylpentane, bromination of, 579
Ethyl propyl sulfide, 366
Ethyne, 429. *See also* Acetylene
Ethynylcyclohexane, 1069
Ethynylestradiol, 426, 430, 430f, 788, 789f, 794, 1254
Ethynyl group, 429
1-Ethynyl-2-isopropylcyclohexane, 429f
Etoposide, 860
Excited state
of atom, 37
of electron, 630–631
Exergonic reaction, 228
Exo product, 625–626, 627f
Exothermic reactions
addition reaction, 398, 398f
bond formation and, 221–222, 224
chlorination, 577, 577f, 580–581
energy diagram of, 229, 230f
enthalpy change and, 228
hydrogenation, 458
radical addition of HBr to alkene, 591–592
reaction rate, 277
transition state in, 276–277, 277f
Extraction, 749–751, 749f, 751f
for separation of amines, 1015, 1015f
Ezetimibe, 205, 776, 918, 994
E, *Z* system of nomenclature, 389–390

F

Faraday, Michael, 642
α-Farnesene, 392f
Farnesol, 1246f, 1248, 1249
Farnesyl diphosphate, 608, 1247f, 1248–1250, 1260
Fats, 395, 461–463, 896–898
body content of, 1237
degree of unsaturation, 461–462
fatty acid composition, 1235, 1235t
melting point of, 395, 1234–1235
oxidation, 589
saturated, 395, 589
trans, 462–463
Fat-soluble vitamins, 1241–1242, 1241f, 1242f
Fatty acids
characteristics of, 1234
composition in fats and oils, 1234–1235, 1235t
essential, 393, 1234
melting point of, 393, 394t
omega-3, 394, 395
oxidation of, 897
saturated, 393, 462–463, 1234–1235, 1234t, 1235t
synthesis in lipid hydrolysis, 897
thioesters of, 904
trans, 462–463
triacylglycerol formation and, 118
in triacylglycerols, 393, 395, 1234–1237, 1234t
unsaturated, 383, 393, 1234–1236, 1234t, 1235t
hydrogenation of, 1236
oxidation of, 1236
Feedstock, 1218
Fehling's reagent, 1129–1130
Femara. *See* Letrozole
Fenfluramine, 86
Fenn, Dr. John, 506
Fenofibrate, 896
Fentanyl, 127, 1044
Fermentation, alcohol production by, 338–339, 339f
Ferulic acid, 639
Fexofenadine, 56, 1005
Fibers
natural, 902, 902f
synthetic, 902–903
Fibrinogen, 1153
Fibrous proteins, 1186
Fingerprint, of IR spectrum, 509–511, 510f, 511f
First-order kinetics, 235
of E1 elimination reactions, 310, 312t
of nucleophilic substitution reactions, 263
of S_N1 substitution reactions, 269
First-order rate equation, 235, 262–263
Fischer esterification, 891–892, 903
Fischer projection formulas, 1109–1113, 1112f
Fishhook notation, 218, 220t
Five-membered rings, synthesis of
in Dieckmann reaction, 979
in intramolecular aldol reactions, 973–975
Flagpole hydrogens, 155, 155f
Flat helical ribbon, shorthand for α-helix, 1182
Flat polygons, 195
Flat wide arrow, shorthand for β-pleated sheet, 1182
Fleming, Sir Alexander, 495, 517, 881
Flexibilene, 1259
Flonase. *See* Fluticasone
Flosal, 969
9-Fluorenylmethoxycarbonyl protecting group (Fmoc), 1174, 1178–1179
Fluorine
polarizability, 100–101
valence electrons of, 12
Fluoroboric acid, 1030
Fluoromethane, polarizability and boiling point of, 103, 104f
m-Fluoronitrobenzene, 645
Fluoxetine, 3, 86, 202, 268, 268f, 708, 727, 1006
Fluticasone, 247, 293
Fmoc (9-fluorenylmethoxycarbonyl protecting group), 1174, 1178–1179
Folic acid, 1037–1038
Formal charge, 16–18, 18t
Formaldehyde, 16
aldol reactions of, 968, 970
Bakelite preparation from, 1216–1217, 1217f
electrostatic potential map of, 766, 766f
formation from methanol oxidation, 482
hydration of, 844–845
primary alcohol formed from, 787
structure of, 820, 825
uses of, 825
Formaldehyde hydrate, 844
Formalin, 825
Formic acid, 68, 132, 482, 732t, 736
Formyl group, 822
Fosamax (alendronic acid), 19
Fossil fuels, 79, 141–142, 163
Four-centered transition state, 412
Fragmentation, 497, 501–504, 502f
Free energy change (Δ$G°$), 225–228, 226f, 226t
Freon 11, 252, 584
Freon 12, 584
Frequency
definition of, 507
of electromagnetic radiation, 506–508, 507f
Hooke's law, 511, 511f
of infrared radiation, 508–509, 511
Friedel, Charles, 679
Friedel–Crafts acylation, 677, 679f, 684–690, 717–718
alkyl benzene synthesis, 713, 713f
of benzene, 679f, 713f
electrophilic aromatic substitution reaction, 679f
general features of, 684
intramolecular, 689–690, 689f
ketone synthesis, 679f, 684, 712
limitations of, 702–703
mechanism of, 686
Friedel–Crafts alkylation, 679f, 684–690, 715–716
of alcohols, 688
of alkenes, 688
in alkyl benzene synthesis, 684–685, 713, 713f
of benzene, 679f, 713f
carbocation rearrangements, 687–688
electrophilic aromatic substitution reaction, 679f
general features of, 684
intramolecular reactions, 690
limitations in, 702–703
mechanism of, 685
polyalkylation, 702
unreactive halides, 686–687
Friedel–Crafts reaction of amides, 1024, 1024f
Frontalin, 484, 484f
Frontside attack, in nucleophilic substitution reactions, 265
Frost circle, 663
Fructofuranose
α-D-fructofuranose, 1123
β-D-fructofuranose, 1123
Fructofuranose ring, 1137
Fructose
D-fructose, 1108, 1115, 1116f, 1123
stereogenic centers, 181
in sucrose, 1136–1137
Fucose, 1151
Fuels, renewable, 339, 339f
Fukui, Kenichi, 1078

Full-headed curved arrow, 218, 218f, 220–221, 220t
Fumaric acid, 127, 624
Functional groups, 92–99. *See also specific functional groups*
  compounds containing a C–O group, 97–99, 98t
  compounds containing a C–Z σ bond, 95–97, 96t
  definition of, 92
  fragmentation pattern in mass spectrum, 503–504
  grouping organic compounds by, 93
  hydrocarbons, 93–94, 94t
  importance of, 98
  infrared (IR) spectroscopy and, 508–512, 510f, 511f, 512t
  intermolecular forces and, 99
  nomenclature of, A–4–5
  nucleophilic substitutions and, 286, 287t
  reactivity and, 116–117
  solubility and, 107
  synthesis of, A–14–17
Funk, Casimir, 110
Funnel, separatory, 749, 749f
Furan, 670, 674
Furandicarboxylic acid, 1213
Furanose(s), 1122–1123
Furanose rings, 1117, 1122–1123
Fused bicyclic system, 626f
Fused ring system, 625–626, 626f

G

Gabapentin, 185
Gabriel synthesis of primary (1°) amines, 1008–1009
Galactose, 1136
  D-galactose, 856, 1114, 1115f, 1132
  in lactose, 238, 238f
  Wohl degradation, 1132
Gamma (γ) carbon, 732
Gamma (γ) hydroxybutyric acid (GHB), 737
Garlic, 132
Garsubellin A, 1097
Gas chromatography–mass spectrometry (GC–MS), 505–506, 505f, 506f
Gasohol, 339, 339f
Gasoline
  additives
    benzene, 93
    BTX mixture, 648
    ethanol, 339
    high-octane, 530, 648
    MTBE, 108
  combustion of, 162–163
  composition of hydrocarbons in, 142, 144
  octane and, 93, 108
  oxidation of isooctane, 224
  refining, 142f
Gauche conformation, 148–149, 148f, 150f
GC–MS (gas chromatography–mass spectrometry), 505–506, 505f, 506f
Geckos, 100
Geminal dichloride, 431
Geminal dihalide, 318
  alkyne synthesis from, 431–432
  synthesis in hydrohalogenation of alkynes, 434
Geminal protons, 549
Generic name, 133
Geranial, 826f, 1096
Geraniol, 284, 484, 1246, 1249
Geranyl diphosphate, 284, 608, 1247f, 1248–1249
GHB [gamma (γ) hydroxybutyric acid], 737
Gibberellic acid, 991
Gibbs free energy ($G°$), 225
Ginger, 623
Gingerol, 973
Ginkgo biloba, 95, 732
Ginkgolide B, 868, 896
Glacial acetic acid, 736
Gleevec, 881
Global climate change, 163
Globular proteins, 1186
D-Glucaric acid, 1130–1131
Glucitol, 1129
D-Gluconic acid, 1129
Glucopyranose
  α-D-glucopyranose, 1130f
  β-D-glucopyranose, 1118, 1141
Glucosamine, 906
  D-glucosamine, 1140
Glucose, 213, 855–856
  acyclic aldehyde, 1118, 1119f
  anomers of, 1118, 1119f, 1122, 1122f
  in complex carbohydrates, 118
  conversion to esters, 1127–1128
  conversion to ethers, 1127–1128
  diastereomers of, 1118, 1119f
  in disaccharides, 1135–1137
  drawing as cyclic hemiacetal, 1117–1119, 1119f
  energy in, 1108
  Fischer projection of, 1111–1112, 1112f
  D-glucose, 1108, 1111, 1112f, 1113, 1114, 1115f, 1117–1118, 1119f, 1122, 1122f, 1123, 1129–1132, 1150, 1218–1219
    α-D-glucose, 856, 1118, 1119f, 1122f, 1124–1126, 1128, 1135, 1138
    β-D-glucose, 855–856, 866, 1107f, 1118, 1119f, 1122, 1122f, 1124–1126, 1128, 1135, 1138
  L-glucose, 1113
  glycoside formation from, 1124–1125
  in honey, 1123
  from hydrolysis
    of cellulose, 1138
    of glycosides, 1125–1126
    of maltose, 1135
    of starch, 175f, 176, 1139
  from Kiliani–Fischer synthesis, 1132
  in lactose, 238, 238f
  oxidation of, 223, 224, 228, 1129–1130
  reduction of carbonyl group, 1129
  structure, 118, 175f
  in sucrose, 1136–1137
  three-dimensional representations for D-glucose, 1122, 1122f
  Wohl degradation, 1131–1132
Glucosidase
  α-glucosidase, 1139
  β-glucosidase, 1138
Glutamic acid, 763
  abbreviation for, 753t
  isoelectric point for, 1156t
  pK_a values for, 1156t
  structure of, 1154f
Glutamine
  isoelectric point for, 1156t
  pK_a values for, 1156t
  structure of, 1154f
Glutathione, 1168
Glyceraldehyde
  enantiomers of, 1109–1110, 1112–1113
  D-glyceraldehyde, 1108, 1113, 1115f
  L-glyceraldehyde, 1113
  (*R*)-glyceraldehyde, 1109, 1112–1113
  (*S*)-glyceraldehyde, 1109, 1112–1113
  optical activity, 199
Glyceraldehyde 3-phosphate, 928
Glycerol
  naming, 335
  synthesis in lipid hydrolysis, 897
  in triacylglycerols, 393, 1234
(*R*)-Glycerol phosphate, 489
Glycine, 752
  abbreviation for, 753t
  in dipeptide synthesis, 1172–1173, 1176
  isoelectric point for, 1156t
  pK_a values for, 1156t
  structure of, 1153f, 1154f
  zwitterion form, 90
Glycogen, 1139
Glycolic acid, 16, 74
  copolymerization with lactic acid, 1212
Glycols, 335
  formation by dihydroxylation, 472
Glycopeptide transpeptidase, 900
Glycoside(s), 1124–1127
  formation, 1124–1125
  α glycoside, 1124–1125
  β glycoside, 1124–1125
  hydrolysis of, 1125–1126
  naturally occurring, 1126–1127
N-Glycoside(s), 1141–1144, 1146f
  α-*N*-glycoside, 1141
  β-*N*-glycoside, 1141
Glycosidic bond/linkage
  in disaccharides, 1134–1137
  α-glycosidic linkage, 1137, 1139
    1→4-α-glycosidic bond, 1135, 1139
    1→6-α-glycosidic bond, 1139
  β-glycosidic linkage, 1134, 1136, 1140, 1142
    1→4-β-glycosidic bond, 1134, 1136, 1140
  in polysaccharides, 1138–1140
Goodyear, Charles, 1210
Gramicidin S, 1195
Grandisol, 1246
Grapefruit mercaptan, 365
Graphite, 666
Green chemistry, 479–480
  Amberlyst A-26 resin and, 479–480
  definition of, 479
  green polymer synthesis, 1217–1220, 1218f, 1219f
Grignard, Victor, 783
Grignard products, retrosynthetic analysis of, 790–792
Grignard reagents, 783
  addition to nitriles, 910–911
  organic synthesis with, 802–805
  preparation of, 783
  reaction as base, 785, 792
  reaction with aldehydes and ketones, 786–789, 832
  reaction with carbon dioxide, 798
  reaction with epoxides, 798–799
  reaction with esters and acid chlorides, 795–797
  reaction with α,β-unsaturated carbonyl compounds, 801
Ground state
  of atom, 36–37
  of electron, 630–631
Group, in periodic table, 9
Group number, 9, 10
Grubbs, Robert, 1062
Grubbs catalyst, 1062–1065, 1065f
Guanidine, 1048
Guanine, 1142, 1143f
D-Gulose, 1115f
Gutta-percha, 1210, 1211, 1226
Gypsy moth, 470, 471f

H

Half-headed curved arrow, 218, 218f, 220–221, 220t, 465, 571
Halides. *See also* Alkyl halides; Aryl halides; Hydrogen halides; Vinyl halides
  allylic, 248, 248f, 586
  benzylic, 710–711
  methyl
    S_N2 substitution reactions, 272–273, 279
  organic
    coupling reaction with organocuprate reagents, 1050–1052
    coupling with alkenes (Heck reaction), 1050, 1056–1058
    coupling with organoboron reagents (Suzuki reaction), 1050, 1053–1056
  unreactive, 686
α-Halo aldehyde or ketone, synthesis of, 937–941
Halo alkane, 249
α-Halo carbonyl compounds
  reactions of, 941
  synthesis of, 937–940
α-Halo carboxylic acids, amino acid synthesis from, 1156–1157, 1159f
Halocyclohexanes, E2 elimination in, 315–317, 315f
Haloethanes, 172
Haloform, 939–940, 940f
Haloform reaction, 939–940
Halogenation
  of activated benzenes, 701–702
  of alkanes, 574–583
    of achiral starting material, 582, 582t
    bromination, 578–580, 580f
    of chiral starting material, 582t, 583
    chlorination, 574–583, 581f
    energy changes during, 576–577, 577f
    mechanism of, 575–576
    in organic synthesis, 581
    stereochemistry of, 582–583, 582t

Halogenation (*Continued*)
of alkenes, 397f, 405–407, 408f
of alkyl benzenes, 709–711
of alkynes, 433f, 436–437
of benzene, 679f, 681–682, 705
of carbonyl compounds, at the α carbon, 937–941, 940f
in acid, 938, 940f
in base, 938–939, 940f
electrophilic aromatic substitution reaction, 679f, 681–682, 701–704
haloform reaction, 939–940
monohalogenation, 574
polyhalogenation, 701
stereochemistry of, 406–407, 408f
transition states in, 577, 577f, 579, 581, 581f
Halogens
in alkyl halides. *See* Alkyl halides
carbon–halogen bond, polar, 252–253, 253f
common nucleophiles, 261t
electronegativity, 707
as electron-withdrawing groups, 692–693, 697
halo group, 96t
polarizable, 405
X as representation for, 13
Halo group, 96t
Halohydrin
epoxide synthesis from, 343
synthesis from alkenes, 397f, 408–410, 410t
mechanism of, 408–409, 410t
regioselectivity of, 409–410, 410t
stereochemistry of, 409–410, 410t
use in organic synthesis, 410
Halomethanes, electrostatic potential map of, 253f
Halomon, 248
Halonium ion, bridged, 406–410, 410t, 436
Halothane, 251f
Hammond postulate, 275–278, 312, 400, 401f, 435, 580, 707
electrophilic aromatic substitution reaction, 696
features of, 275–277
S_N1 substitution reactions, 275, 277–278
Handedness, 175, 178, 180
Hansen's disease (leprosy), 184
Hardener, 1214–1215, 1215f
Hardening, 462, 1236
Haworth projections, 1118, 1119–1121
converting to chair form, 1122
HCFCs (hydrochlorofluorocarbons), 584
HDL particles (high density lipoproteins), 905
HDPE (high-density polyethylene), 593, 1202, 1209, 1221, 1221t
Heat of hydrogenation, 458–459, 615–616
benzene, 650–651, 650f
Heat of reaction, 221. *See also* Enthalpy change
Heat symbol, 214
Heck, Richard, 1056
Heck reaction, 1050, 1056–1058
Helicene, 648, 649f
Helium, electronic configuration of, 10
Heme, 1152, 1187–1188
Hemiacetal(s)
acetals compared to, 1124
in carbohydrates, 855–856
conversion to acetals, 848–849
conversion to carbonyl compounds, 850
cyclic
conversion to cyclic acetals, 854–855
structure, 852
synthesis of, 853–854
monosaccharides, 1116–1123
structure of, 852
synthesis of, 848–850
Hemibrevetoxin B, 96
Hemlock water dropwort, 30
Hemoglobin, 1153
sickle cell, 1188–1189
structure, 1185, 1187–1188, 1188f
Heptalene, 669
Heptanal, 969
Heptane
constitutional isomers of, 131t
molecular formula, 131t
Heptane-2,6-dione, 974
Heptan-2-one, synthesis of, 951
trans-Hept-2-ene, 1051
Herbicides, 682, 682f
Heroin, 113–114, 207, 523, 889
Heteroatom(s), 4
in condensed structures, 31
definition of, 23, 92
in functional groups, 92, 95, 116
in resonance structures, 23
in skeletal structures, 32, 35
stereogenic centers, 182
Heterocycle(s), 336
aromatic, 655–658, 657f
nitrogen, 1000, 1001f
Heterocyclic aromatic amines, basicity of, 1020, 1021t
Heterogeneous reaction mixture, 458
Heterolysis (heterolytic cleavage)
carbocation formation, 269, 275–277, 281, 286
description, 217–218
in E1 elimination reactions, 310
reactive intermediates of, 218f, 219
in two-step reaction mechanism, 231
Hexachloroethane, 203
Hexa-2,4-diene, 613, 653
(2*E*,4*Z*)-Hexa-2,4-diene, 1082–1084, 1086, 1090
Hexa-1,3-diyne, 429f
Hexamethylenediamine, 902
Hexamethylphosphoramide (HMPA), 260f
Hexane, 128, 405, 466f
constitutional isomers of, 131t
infrared (IR) spectrum, 513
mass spectrum of, 498–499, 498f, 501–502, 502f
molecular formula, 131t
skeletal structures, 32
as solvent, 106
solvent in extraction procedure, 749
Hexane-2,5-dione, aldol reactions of, 973–974
Hexanoic acid, 731, 732, 760
Hexan-2-ol, conversion to other compounds, 802, 803f
Hexanoyl chloride, 876
Hexaphenylethane, 602
Hexa-1,3,5-triene, 651, 653, 1077, 1081
molecular orbitals of, 1080, 1081f
Hex-1-ene, 405, 513
Hex-2-ene, 387
Hex-3-ene, 466f
cis-hex-3-ene, 388–389, 1062
trans-hex-3-ene, 389
Hex-1-yne, 514, 1056
Hex-3-yne, 466f
HFCs (hydrofluorocarbons), 584
High density lipoproteins (HDLs), 905
High-density polyethylene (HDPE), 593, 1202, 1209, 1221, 1221t
Highest occupied molecular orbital (HOMO), 663, 1080, 1083–1086, 1089–1091
High-performance liquid chromatography (HPLC), 1169
Histamine, 657–658, 1004–1005, 1042
Histidine
isoelectric point for, 1156t
pK_a values for, 1156t
structure of, 1154f
Histrionicotoxin, 121, 431, 431f
HMPA (hexamethylphosphoramide), 260f
^{1}H NMR. *See* Proton NMR
Hoffman, Albert, 689f
Hoffman, Felix, 737–738
Hoffmann, Roald, 1078
Hofmann elimination, 1024–1027
HOMO (highest occupied molecular orbital), 663, 1080, 1083–1086, 1089–1091
Homogeneous catalysts, 1209
Homologous series, 131, 131t
Homolysis (homolytic cleavage)
of allylic C–H bonds, 586
bond dissociation energy for, 221–222
chlorination of alkanes, 575, 576, 578
description, 217–218
endothermic nature of, 221–222
entropy increase with, 228
radical formation, 572
reactive intermediates of, 218–219, 218f
Homopolymers, 1207
Homotopic protons, 534–535
Honey, 1123
Honeycomb, 164
Hooke's law, 511, 511f
Hormones. *See also specific hormones*
local mediators distinct from, 1242
peptide, 1167–1168, 1168f
sex, 1253–1254, 1254t
Host–guest complex, 340
Housane, 141f
5-HPETE-(5-hydroperoxyeicosatetraenoic acid), 372, 600
Hückel, Erich, 652
Hückel's rule, 651–654
basis of, 660–663
Humulene, 325, 1055, 1055f
Hybridization, 36–45, 644
conjugated systems and, 611–612
definition of, 37
Hybridization effects
on acidity, 75–76, 76f, 77f
on basicity of amines, 1020–1021, 1021t
Hybrid orbitals, 36–45
percent *s*-character, 46, 76, 302
sp orbitals, 38–39, 39t, 42–43, 43f, 44t
in alkynes, 427, 428, 432, 433
percent *s*-character, 76
sp^2 orbitals, 38–39, 39t, 41–42, 42f, 44t
in alkenes, 299–300, 302
in aromatic heterocycles, 655–656, 657f
in benzene, 644
carbon radical, 571
carboxy group, 730
percent *s*-character, 76
sp^3 orbitals, 37–38, 38f, 39, 39t, 40, 44t
in acyclic alkanes, 129
in alkenes, 302
percent *s*-character, 76
superscripts, meaning of, 39
Hydrates
oxidation of, 478
synthesis from aldehydes and ketones, 844–847
kinetics, 846–847
thermodynamics, 845
Hydration
of aldehydes and ketones, 844–847
kinetics, 846–847
thermodynamics, 845
of alkenes, 397f, 404–405, 415
definition of, 834
hydroboration–oxidation compared, 415
reactions, 433f, 437–439
Hydrazine, in Wolff–Kishner reduction, 712
Hydrides
as base, 79, 79f
metal hydride reagents
reducing agents, 457, 466, 771–773, 777–781, 781t
reduction of aldehydes and ketones, 771–773
reduction of nitriles, 909
nucleophilic addition with, 767, 772
nucleophilic substitution with, 768
reduction of ketones and aldehydes, 771–773, 773f, 781t, 831–832
1,2-Hydride shift, 348–350, 687–688
Hydroboration
of alkenes, 411–413
of alkynes, 1054, 1056
definition of, 411
mechanism of, 411–412
regioselectivity of, 412, 414t
Hydroboration–oxidation reactions, 439–441
of alkenes, 397f, 411–415
mechanism of, 411–413, 414t
regioselectivity of, 412, 414t
stereochemistry of, 413–414, 414t
hydration compared, 415
synthesis of artemisinin, 414–415, 415f
Hydrocarbons
aliphatic, 93, 129
aromatic, 93–94, 94t, 641–676
definition of, 1, 93
functional groups in, 93–95, 94t
infrared (IR) absorptions in, 513–514
oxidation and reduction of, 456, 456f
polycyclic aromatic (PAH), 373
saturated, 129
synthesis in organocuprate couplings, 1051–1052
unsaturated, 385. *See also* Alkenes

Hydrochloric acid
  in amine extraction, 1015, 1015f
  elimination in nylon synthesis, 902
  in epoxide ring opening reactions, 369–370, 370f
  as Lewis acid, 83
  pK_a of, 70, 78
  for removal of Boc protecting group, 1174
Hydrochlorofluorocarbons (HCFCs), 584
Hydrofluorocarbons (HFCs), 584
Hydrogen
  addition to alkenes, 458–463
  addition to alkynes, 463–465, 466f
  alpha (α), 925, 929, 938–939, 968–969, 971, 977
  classification, 94–95
  covalent bond formation by, 12, 13f
  electronic configuration, 10
  isotopes of, 8, 504t
  mass of, 504t
  octet rule exceptions, 19
  primary (1°), 134
  as reducing agent, 457. *See also* Hydrogenation
  secondary (2°), 134
  sigma (σ) bond in molecular hydrogen, 36
  tertiary (3°), 134
Hydrogenation
  of aldehydes and ketones, 772
  of alkenes, 458–463
    alkene stability and, 458–459, 459f
    cis and trans isomers compared, 458, 459f
    degrees of unsaturation, determination of, 460–461
    mechanism of catalytic hydrogenation, 459–460
  alkene stability and, 458–459, 459f
  catalytic, 458–460
  of conjugated dienes, 615–616
  enantioselective synthesis of amino acids, 1163–1164
  as exothermic reactions, 458
  heat of, 458–459, 615–616, 650–651, 650f
  of oils, 461–463, 462f
  of unsaturated fatty acids, 1236
Hydrogen bonding
  in alcohols, 337
  in amides, 877, 878f
  in amines, 1001
  anion solvation by, 259
  in carboxylic acids, 734, 734f
  in DNA, 118, 118f, 1144, 1144f
  intramolecular in β-dicarbonyl compounds, 926
  in polar protic solvents, 259, 281
  in protein secondary structure, 1180–1182, 1181f, 1182f
  solubility and, 107, 108t
  strength of, 101, 102t
  in water, 101
Hydrogen bromide, for removal of Boc protecting group, 1174
Hydrogen chloride. *See* Hydrochloric acid
Hydrogen cyanide, 16, 834
Hydrogen fluoride, 13
Hydrogen halides
  addition to alkenes, 397–404, 397t, 398t, 399t, 403t
  addition to alkynes, 433f, 434–436
  alcohol conversion to alkyl halides in, 352–355
  reactivity of, 354
Hydrogen ions (H^+), Brønsted–Lowry acid definition and, 62
Hydrogenolysis
  of benzyl esters, 1175
  in peptide synthesis, 1175–1176
Hydrogen peroxide, as oxidizing agent, 467
Hydrohalogenation
  of alkenes, 397–404, 397f
    Markovnikov's rule, 400–402, 401f, 403t
    mechanism of, 397–400, 398f, 399f, 403t
    stereochemistry of, 402–404, 403t, 404f
  of alkynes, 433f, 434–436
  Markovnikov's rule, 434–436
Hydrolysis
  of acetals, 850–851
  in acetoacetic ester synthesis, 949, 950
  of amides, 899
  in amino acid synthesis, 1157
  base-promoted, 895–896
  definition of, 834
  of esters, 894–898
    acid-catalyzed, 894–895
    applications of, 896–898
    base-promoted, 895–896
  glucose formation by, 1114, 1125–1126, 1135, 1138, 1139
  of imines and enamines, 844
  of lipids, 896–898, 1233
  in malonic ester synthesis, 945–947
  of nitriles, 907–908
  partial of peptides, 1171–1172
  of starch, 339, 339f
  of triacylglycerols, 1236
Hydronium ion, formal charge on, 17
Hydroperoxides, 588, 589f, 600, 1236
5-Hydroperoxyeicosatetraenoic acid (5-HPETE), 372, 600
Hydrophilic, 108, 111, 113
Hydrophobic, 108, 111, 113, 164, 1233
Hydroxide ($^-$OH)
  as Brønsted–Lowry base, 79
  nucleophile in alcohol synthesis, 342
  oxygen bases, 798, 798f
α-Hydroxy acids, 74
Hydroxy aldehyde
  cyclization of, 853, 855–856
  β-hydroxy, synthesis in aldol reactions, 963–975
o-Hydroxybenzaldehyde, 992
Hydroxybenzene, 645. *See also* Phenol
o-Hydroxybenzoic acid, 763
p-Hydroxybenzoic acid, 763
3-Hydroxybutanal, 963
4-Hydroxybutanal, 853
2-Hydroxybutanedioic acid, 763
4-Hydroxybutanoic acid, 737, 917
1-Hydroxybutan-2-one, 564
4-Hydroxybutan-2-one, 564
3-Hydroxybutyric acid, 1222
γ-Hydroxybutyric acid (GHB), 737
β-Hydroxy carbonyl compounds
  synthesis of, in aldol reactions, 963–975
  useful transformations of, 970, 971f
3-Hydroxy carboxylic acids, 1222
Hydroxy group, 96t
  of alcohols, 332, 344
  of enols, 332
  of ethanol, 93
  hydrophilic nature of, 108
  IR absorption, 514, 735, 735f
  of monosaccharides, 1108, 1124, 1127–1128
  NMR absorption, 735, 736f
  OH proton, NMR spectra of, 551–552, 551f
  of phenols, 332
6-Hydroxyhexan-2-one, 851
Hydroxylamine, 1132
5-Hydroxypentanal, 853
5-Hydroxypentan-2-one, 792–794, 793f
3-Hydroxyvaleric acid, 1222
(–)-Hyoscyamine, 961
Hyperconjugation, 273, 274–275
Hypertension, 99
Hypophosphorus acid, 1031

I

Ibufenac, 726
Ibuprofen, 121, 957, 958
  anti-inflammatory action of, 164, 962
  enantiomers, 202
  mode of action, 1244
  synthesis of, 713, 773f, 962, 979
  systematic name of, 133
-ic acid (suffix), 732, 873, 874–875
Icosane, 131t
-ide (suffix), 249
D-Idose, 1115f
Iejimalide B, 421
Illudin-S, 951
Imatinib mesylate, 881
Imidazole, 793, 794
Imidazolide, 915
Imides, in Gabriel synthesis of primary amines, 1008
Imidic amide tautomer, 907–908
Imines, 450, 781, 1133
  hydrolysis of, 844
  reduction to amines, 1011
  synthesis of
    from aldehydes and ketones, 840–842, 842f, 843f, 1023, 1158
    from nucleophilic addition reactions, 830, 831f
    reduction of nitriles, 909–910
Iminium ion, 841, 843, 843f
Indene, 671
Indigo, 1036
Indinavir, 122
Indole, 728
Inductive effects
  on acidity, 72–74, 73f, 77f
  in aliphatic carboxylic acids, 744–746
  on basicity of amines, 1021t
  in carbocation stability, 273
  definition of, 73
  electron-donating, 691–693
  electron-withdrawing, 73, 691–693
  on substituted benzenes, 691–693
-ine (suffix), 249
Influenza, 99
Infrared (IR) absorptions/spectra
  of aldehydes, 823–824, 824f
  of amines, 1002, 1003f
  atom mass and, 510–511, 511f
  of benzene derivatives, 647t
  bond strength and, 510–512, 511f, 512t
  of carboxylic acid derivatives, 878–879, 879t
  of carboxylic acids, 735, 735f
  characteristic frequencies, A–9
  functional groups and, 508–517, 510f, 511f, 512t
  of hydrocarbons, 513–514
  important, 512, 512t
  of ketones, 823–824, 824f
  of nitriles, 878
  of nitrogen-containing compounds, 515–516
  of oxygen-containing compounds, 514–515
  percent *s*-character and, 512
  vibration of bonds and, 508, 510–513
  wavenumber scale of, 508, 509
Infrared (IR) spectroscopy, 496, 506, 508–519
  background on, 508–509
  functional group identification, 508–517, 510f, 511f, 512t
  IR inactive vibration, 513
  isomers distinguished by, 516
  structure determination in, 517–519
  vibration modes, 509
  wavenumber scale, 508, 509
Infrared (IR) spectrum
  characteristics of, 509–510, 510f
  fingerprint region, 509–511, 510f, 511f
  functional group region, 509–511, 510f
Ingenol, 1049, 1066
Initiator, 1200, 1201, 1205, 1206
Inorganic compounds, 1
Inscribed polygon method for predicting aromaticity, 663–666, 665f
Insecticide, 252, 1059
Inspra. *See* Eplerenone
Insulin, 855, 1153, 1184, 1184f
Integral, NMR spectrum, 541
Intermolecular forces, 99–102
  of aldehydes and ketones, 822, 823t
  boiling point and, 103–104, 104f
  in carboxylic acids, 734, 734f, 734t
  dipole–dipole interactions, 101, 102t
  electrostatic interactions, 99
  hydrogen bonding, 101, 102t, 1001
  ion–ion interactions, 99, 102t
  melting point and, 105–106
  solubility and, 106–109, 107f, 108t
  van der Waals forces, 100–101, 100f, 102t
International Union of Pure and Applied Chemistry, 132. *See also* IUPAC nomenclature
Intramolecular cyclization, of hydroxy aldehyde, 853
Intramolecular hydrogen bonding, in β-dicarbonyl compounds, 926
Intramolecular malonic ester synthesis, 948
Intramolecular reactions
  aldol reactions, 973–975, 975f
    Robinson annulation, 982–986
  electrocyclic ring closure, 1077
  Friedel–Crafts reactions, 689–690, 689f
  intramolecular Claisen (Diekmann reaction), 979–980
  sigmatropic rearrangements, 1091–1097
Inversion of configuration
  in alcohol reactions with HX, 354
  in alkyl tosylate reactions, 360, 361
  in S_N2 substitution reactions, 265–266, 266f, 267t, 280

Iodine, polarizability of, 100–101
Iodoform, 939–940
Iodoform test, 939
Iodomethane, polarizability and boiling point of, 103, 104f
1-Iodo-1-methylcyclohexane, 308
Ion(s)
  anions
    acetate, 74–75
    carbanion, 76, 76f, 79, 218f, 219
    cyclopentadienyl, 658–660, 664
    definition of, 8
    enolate, 22
    naked, 260
    radical, 465
    salt formation, 12
    solvation by hydrogen bonding, 259
  bond formation by, 219
  cations
    carbocation. *See* Carbocation(s)
    counterions, 62
    cyclopentadienyl, 659
    definition of, 8
    radical, 497
    salt formation, 12
    solvation by ion–dipole interactions, 259–260, 280
    spectator ions, 62
    tropylium, 660, 664
  counterions, 62
  polyatomic, 17
  spectator ions, 62
  transport across cell membranes, 115–116, 116f
Ion–dipole interactions
  cation solvation by, 259–260, 281
  description of, 106, 107f
Ionic addition, 592
Ionic bonding, 11–12
Ionic compounds
  boiling points of, 103
  formation of, 11–12
  intermolecular forces, 99, 102t
  melting points of, 105
  solubility of, 106, 107f, 108t
Ionic intermediates, 587
Ionic polymerization, 1204–1206, 1205f
Ion–ion interactions, 99, 102t
Ionophores, 115–116, 116f
IR spectroscopy. *See* Infrared (IR) spectroscopy
Iso- (prefix), 134
Isoamyl acetate, 880
Isobutane, 130
p-Isobutylacetophenone, 713
p-Isobutylbenzaldehyde, 957
Isobutyl chloride, 250f
Isobutylene, 1205f
Isobutyl group, 134
Isocomene, 424
Isocyanate, 1213
Isoelectric point (p*I*), for amino acids, 755, 1156, 1156t
Isoleucine
  isoelectric point for, 1156t
  pK_a values for, 1156t
  structure of, 1154f, 1155
Isomers
  constitutional. *See* Constitutional isomers
  definition of, 18, 176, 177
  infrared (IR) spectroscopy of, 516
  Lewis structures, 18–19
  meta, ortho, and para, 705–706
  resonance structures compared, 20
  stereoisomers. *See* Stereoisomers
  types of, 177, 177f
Isoniazid, 1042
Isooctane, 53
  combustion of, 162
  oxidation of, 224
Isopentane, 130, 141 *See also* 2-Methylbutane
Isopentenyl diphosphate, 608, 1247, 1248, 1250
Isophorone, 994
Isophthalic acid, 759
Isoprene, 614, 631, 631f, 638, 1210, 1226
Isoprene unit, 1245–1247, 1246t, 1258
Isopropyl alcohol, 335
Isopropylamine, 380
Isopropylbenzene
  oxidation to benzoic acid, 712
  synthesis from benzene, 713
Isopropylcyclopentane, 804–805
Isopropyl group, 134
3-Isopropyl-4-methylcyclohexanone, 821
Isopropyl 2-methylpentanoate, 876
Isoproterenol, 811
Isopulegone, 526
Isotactic polymer, 1208–1209
Isotopes, 8
  of bromine, 500
  of carbon, 8, 504t
  of chlorine, 500
  definition of, 8
  hydrogen, 504t
  masses of common, 504t
  of nitrogen, 10, 504t
  of oxygen, 504t
Isotretinoin, 737, 863
IUPAC nomenclature
  of alcohols, 334, 334f, 335
  of aldehydes, 819, 820f
  of alkanes, 135–140, 140f
  of alkenes, 387–388
  of alkyl halides, 249, 250f
  of alkynes, 428–429
  of benzene derivatives, 644–647
  of carboxylic acids, 731–734, 733f
  of cycloalkanes, 138–140, 140f
  description of, 132–133
  of epoxides, 336–337
  of ethers, 335–336
  of ketones, 820–821, 822f

J

Jack-o'-lantern mushroom, 951
Januvia. *See* Sitaglipitin
cis-Jasmone, 464
Johnson, W. S., 975
Jojoba, 1233
Joule (J), 45
Juvenile hormones, 789, 789f, 1051
  mimics of, 789

K

Kavain, 390
Keflex. *See* Cephalexin
Kekulé, August, 642
Kekulé structures, 642, 643
Kelsey, Frances Oldham, 184
Keratins, 1153
  α-keratins, 1180, 1186, 1186f, 1187f
Kerosene, composition of hydrocarbons in, 142
Ketal, 847
Ketene, 57
Keto–enol tautomers, 437–439
β-Keto esters
  as active methylene compounds, 970
  conversion to ketones, 949–950
  pK_a, 931t
  synthesis of
    Claisen reaction, 975–979, 975f, 976f
    Michael reaction, 982
D-Ketohexoses, 1115, 1116f, 1122–1123
Ketones. *See also* Carbonyl compounds
  aldol reactions, 963–975
  boiling point of, 823t
  carbohydrates, 855–856
  in crossed Claisen reactions, 978–979
  fragmentation pattern in the mass spectrum, 503
  functional group in, 98t
  hydration of, 844–847
    kinetics of hydrate formation, 846–847
    thermodynamics of hydrate formation, 845
  interesting examples, 825–826
  iodoform test of, 939
  IR spectra of, 514, 823–824, 824f
  melting point of, 823t
  NMR spectra of, 825
  nomenclature, 820–821, 822
    common names, 821, 822f
    IUPAC system, 820–821, 822f
  nucleophilic addition reactions, 767, 772, 817–867
    carbanion addition, 832
    cyanide addition, 833–834
    hydride addition, 831–832
    mechanism of, 829–830
    nucleophiles in, 830, 831f
    primary amine addition, 840–842, 843f
    secondary amine addition, 842–844, 843f
  physical properties, 822, 823t
  pK_a, 931t
  protecting groups for, 851–852
  reactions of
    aldol reactions, 963–975
    with amines, 1023
    at the α carbon, 828–829
    at the carbonyl carbon, 828, 829
    crossed Claisen reactions, 978–979
    halogenation at the α carbon, 937–941, 940f
    hydration, 844–847
    nucleophilic addition reactions, 767, 817–867
    with organometallic reagents, 785–789, 789f, 832
  reactivity of, 767
  reduction of, 831–832
    to alcohols, 770, 771, 772
    of aryl ketones to alkyl benzenes, 712–713
    reductive amination to amines, 1011–1013
    to secondary alcohols, 461
  solubility of, 823t
  structure of, 765, 818
  synthesis
    from carboxylic acid derivatives, 796–797
  synthesis of, 826–828
    acetotacetic ester synthesis, 945, 949–952
    alcohol oxidation, 477–478
    Friedel–Crafts acylation, 679f, 684, 712, 827
    hydration of alkynes, 437–439, 827
    hydroboration–oxidation of alkyne, 440
    hydrolysis of acetals, 850–851
    hydrolysis of imines and enamines, 844
    oxidation of secondary alcohols, 827
    oxidative cleavage of alkenes, 474–475, 827
    reactions of acid chlorides with organocuprates, 827
  unreactivity to nucleophilic substitution, 769
  unsymmetrical, alkylation of, 943–944
  Wittig reaction, 835–839
Ketoprofen, 78, 207
Ketoses
  D-ketose family, 1115, 1116f
  structure, 1108–1109
Keto tautomers, 908, 926–927
Kevlar, 1212, 1226
Kiliani–Fischer synthesis, 1131, 1132–1134
Kinetic enolates, 934, 935, 943, 971
Kinetic product, in electrophilic addition reactions of conjugated dienes, 619–621, 620f
Kinetic resolution, of amino acids using enzymes, 1162–1163
Kinetics
  definition of, 225, 233
  E1 elimination reactions, 310, 312t
  E2 elimination reactions, 303, 307t
  first-order, 235
  rate equations, 234–236
  second-order, 235
  S_N1 substitution reactions, 269, 273t
  S_N2 substitution reactions, 263, 267t
Kukui nuts, 383

L

Lactams, 870, 1212
β-Lactam(s), 870, 881, 900–901
β-Lactam family of antibiotics, 495, 517–518
Lactase, 238, 238f, 1136
Lactate dehydrogenase, 777
Lactic acid, 32, 736, 777
  copolymerization with glycolic acid, 1212
  optical activity of, 199
  specific rotation of, 201
Lactide, 1219
Lactols, 852
Lactone(s), 870
  synthesis by esterification of carboxylic acids, 892
Lactose, 855, 1130, 1136
  breakdown of, 238, 238f
Lactose intolerance, 1136
Laetrile, 834
Lanolin, 1233
Lanosterol, 1252f, 1253
Lard, 395

Latanoprost, 738
Latex, 1210, 1211
Laurencia cartilaginea, 252
Lauric acid, 1234t
Lavandulol, 791
Lavender oil, 331
l (levorotatory) compounds, 199, 1113
LDA. *See* Lithium diisopropylamide (LDA)
LDL particles (low density lipoproteins), 905
L-Dopa (levodopa), 7, 50–51, 1194
LDPE (low-density polyethylene), 593, 1202–1203, 1221t
Lead(II) acetate, 464
Leaving group
  of alcohols, 344, 346–347, 349–354, 356–361
  of carbonyl compounds, 765, 767, 768–769
  of carboxylic acid derivatives, 870, 872t, 882–883
  diphosphate, 608, 1247, 1249
  in E1 elimination reactions, 310, 312t
  in E2 elimination reactions, 304–306, 307t
  in nucleophilic acyl substitution reactions, 870, 882–883
  in nucleophilic substitution reactions, 253–257, 265–266, 280–281
    derived from phosphorus, 283–284
    good, 255–257, 256t
    periodic trends in ability, 255–256
    poor, 256t
    reaction rate and, 275
    weak bases, 255–257
  sulfonate anion, 752
  in terpene biosynthesis, 1247–1249
Le Châtelier's principle, 347, 891, 1124
Lecithin, 1238, 1239f
Length, units used to report wavelength, 507
Leprosy, 184
Letrozole, 906
Leucine
  isoelectric point for, 1156t
  pK_a values for, 1156t
  structure of, 1154f
Leukotrienes, 1242, 1243f, 1243t
  biological activity of, 372
  leukotriene A_4, 372
  leukotriene C_4, 372, 600
  synthesis of, 372
Levonorgestrel, 431
Levorotatory (*l*) compounds, 199, 1113
Lewis acid, 81–83
  borane as, 411
  carbocation as, 399
  catalyst, 354
Lewis acid–base reactions, 82–83, 254. *See also* Nucleophilic substitution reactions
  electrophilic addition, 399
  S_N1 substitution reactions, 269
Lewis base, 81–83
  alkene as, 399
  nucleophile as, 253, 257, 269
  tetrahydrofuran (THF) as, 411
Lewis structures, 13–18
  of acyclic alkanes, 129–130
  of benzene, 643
  condensed formula conversion to, 32
  drawing of, 13–15
  formal charge, 16–18, 18t
  isomers, 18–19
  multiple bonds in, 15–16
  octet rule exceptions, 19
  resonance structures, 19–25
Lexan, 1199f, 1214, 1220
Ligands, in palladium-catalyzed reactions, 1052–1053
Light
  polarized, 198–199
  speed of, 507
  symbol, 214
Lightning, 474
"Like dissolves like," 106–107
Limonene, 56, 475, 638, 1249
  (*R*)-limonene, 392f
Linalool, 331, 791
Linalyl diphosphate, 638, 1249
Linamarin, 834
Lindlar catalyst, 464, 466f
Linear molecule, 26, 27, 29t, 49
  alkynes, 427
Linen, 902, 902f
Linezolid, 923
Linoleic acid, 589, 1234, 1234t, 1235t
  as essential fatty acid, 393
  melting point, 394t
  sources of, 383, 393
  structure, 394f, 394t
Linolenic acid, 490, 1234, 1234t
  as essential fatty acid, 393
  melting point, 394t
  as omega-3-fatty acid, 394–395
  sources of, 383, 393, 395
  structure, 394f, 394t
Lipases, 897
Lipid bilayer, 113, 114f, 116f, 1239–1240
Lipids, 129, 164–165, 393–395, 1231–1260
  bonding in, 1232
  definition of, 164
  eicosanoids, 1242–1245, 1243f, 1244t
  energy in, 165, 1237
  examples of, 164–165, 164f, 1232f
  fat-soluble vitamins, 1241–1242, 1241f, 1242f
  hydrolysis of, 896–898, 1233
  hydrolyzable, 1233
  metabolism of, 1237
  nonhydrolyzable, 1233
  origin of word, 1232
  oxidation of, 588–589, 589f, 590
  phospholipids. *See* Phospholipids
  solubility of, 164, 1232
  steroids. *See* Steroids
  structure of, 164f
  terpenes, 1245–1250, 1246f, 1246t, 1247f
  triacylglycerols. *See* Triacylglycerols
  unsaturated, oxidation of, 588–589, 589f
  waxes, 1233
Lipitor. *See* Atorvastatin
Lipoprotein, 905, 905f
Lipoxygenase, 369, 1243, 1243f
Lisinopril, 190
Lithium
  electronegativity value of, 783
  valence electrons of, 12
Lithium acetylide, 784, 788
Lithium aluminum hydride ($LiAlH_4$), 771–773, 777–781, 781t
  as reducing agent, 457
  reduction of aldehydes and ketones, 831
  reduction of alkyl halides, 466–467, 467f
  reduction of a nitrile, 909
  reduction of epoxides, 466–467, 467f
  reduction of nitriles to primary amines, 1010
Lithium diisopropylamide (LDA), 942–944, 971
  enolate formation with, 932–935
  pK_a, 932t
  preparation of, 933
Lithium dimethylcuprate, 784, 1061
Lithium fluoride, 12
Lithium tri-*sec*-butylborohydride (L-selectride), 815
Lithium tri-*tert*-butoxyaluminum hydride, 777, 778, 781t
(–)-Littorine, 961
Living polymerization, 1206
Local anesthetic, 714, 726
Local mediators, 1242
Log, 66
London forces, 100. *See also* van der Waals forces
Lone pairs
  in amines, 1014, 1020, 1023
  in aromatic heterocycles, 655–657
  Brønsted–Lowry bases and, 62–63, 62f
  in carbanions, 219
  in heteroatoms, 35, 92, 116
  in hydrogen bonding, 101
  in Lewis structures, 14–16
  of negatively charged carbon atoms, 35
  number of, 35
  in resonance structures, 21, 23–24, 74, 609, 612
  second-row elements and, 13
Long, Dr. Crawford, 340
Loratadine, 649f
Losartan, 1069
Low density lipoproteins (LDLs), 905
Low-density polyethylene (LDPE), 1202–1203, 1221t
Lowest unoccupied molecular orbital (LUMO), 663, 1080, 1085, 1089–1091
LSD, 941
  intramolecular Friedel–Crafts acylation in the synthesis of, 689, 689f
  structure of, 1007
  synthesis of, 689, 689f
LUMO (lowest unoccupied molecular orbital), 663, 1080, 1085, 1089–1091
Lupin, 1022, 1048
Lupinine, 1048
Lycopene, 614, 631, 631f, 1258
Lycra. *See* Spandex
Lynen, Feodor, 1252
Lysergic acid, 640
Lysergic acid diethyl amide. *See* LSD
Lysine, 729, 763
  abbreviation for, 753t
  ionizable side chain of, 1155, 1156t
  isoelectric point for, 1156t
  pK_a values for, 1156t
  structure of, 1154f
Lysozyme, 1182, 1183f
D-Lyxose, 1114, 1115f, 1132

M

Magnesium, electronegativity value of, 783
Magnetic field, 528–531, 556
Magnetic resonance imaging (MRI), 560, 560f
Mahimahi, 657
Ma huang, 181, 208
Ma'ilione, 252
Malaria, 4–6, 5f, 307
Maleic acid, 127, 624
Maleic anhydride, 624f
Malonic acid, 90, 733
Malonic ester synthesis, 945–949
  intramolecular, 948
  retrosynthetic analysis of, 948–949
  steps in, 946–948
Maltose, 867, 1135
Mandelic acid, 210, 744
(*R*)-Mandelic acid, 1193
Manganese, in oxidizing agents, 468
Mannose
  in Fischer proof, 1135–1136
  D-mannose, 1114, 1115f, 1132
    Haworth projections of, 1119–1120
Margarine, 461–462, 1034
Marijuana, 78
Markovnikov addition of water, 437
Markovnikov's rule, 400–402, 401f, 403t, 434–436, 439, 617
  in cationic polymerization, 1204
Mass number, 8
Mass spectrometer
  features of, 496
  high-resolution, 504
  schematic of, 496f
Mass spectrometry (MS)
  of alkyl halides, 500–501, 501f
  of amines, 1002, 1002f
  analysis of unknowns using the molecular ion, 498–500
  base peak, 497
  of biomolecules, 506
  definition of, 496
  electrospray ionization (ESI), 506
  features of, 496–498
  fragmentation in, 497, 501–504, 502f
  gas chromatography and, 505–506, 505f, 506f
  high-resolution, 504
  M peak, 497, 500
  M + 1 peak, 498
  M + 2 peak, 500–501
Mass spectrum, 497
Mass-to-charge (*m/z*) ratio, 496f, 497, 504
Matrine, 1022
Mauveine, 1036
Maytansine, 1074
MDMA (3,4-methylenedioxymethamphetamine), 1046
Mefloquine, 211
Melamine, 1230
Melanin, 632
Melmac, 1230
Melting point
  of alcohols, 338t
  of aldehydes and ketones, 823t
  of alkanes, 143, 143t
  of alkenes, 391, 393, 394t

Melting point (*Continued*)
of alkyl halides, 250t
of alkynes, 429
of amines, 1001t
of carboxylic acids, 734t
definition of, 105
of enantiomers, 199, 199t
of epoxides, 338t
of ethers, 338t
of fats, 395, 1234–1235
of fatty acids, 393, 394t
intermolecular forces and, 105–106
of monosaccharides, 1116
of oils, 395, 1234–1235
of racemic mixtures, 199, 199t
of triacylglycerols, 462f
Membrane proteins, 1153
Menthol, 56, 123, 1245, 1246
Menthone, 123
Meperidine, 97, 957
Mercaptans, 364–365
Mercapto group, 96t, 364–365
Mercury, in Clemmensen reduction, 712
Merrifield, R. Bruce, 1177
Merrifield method, of peptide synthesis, 1177–1179
Mescaline, 1005
Mesityl oxide, 815
Meso compounds, 193–194
achiral compounds, 193, 195, 407, 408f, 624, 1060
carbene addition and, 1060
disubstituted cycloalkanes, 195
from epoxidation of alkenes, 470
plane of symmetry, 193
Mestranol, 124
Meta (prefix), 645
Metabolism, 1108
Meta director, 694–696, 698, 700, 701f, 702–703, 705, 715, 717
Metal–carbenes, 1064
Metal catalyst, 236, 458
Metal hydride reagents
reducing agents, 457, 466, 771–773, 777–781, 781t
reduction of aldehydes and ketones, 771–773
reduction of nitriles, 909
Metals
as catalysts, 236–237
in oxidizing agent, 468
Metathesis, 1062–1067, 1063f, 1065f
derivation of word, 1062
drawing products of, 1063, 1063f
mechanism, 1064
ring-closing (RCM), 1065–1067, 1065f
Met-enkephalin, 881
Meth- (prefix), 133
Methamphetamine, 1005, 1012, 1012f
Methanal. *See* Formaldehyde
Methanamine. *See* Methylamine
Methane
bonding in, 36–38, 38f
combustion of, 162
description, 2
halogenation of, 574, 574f
Lewis structure, 14
mass spectrum of, 497–499
molecular formula, 129, 130, 131t
molecular shape of, 27–28
in natural gas, 141
oxidation of, 456, 456f
pK_a of, 67, 70
structure of, 129
temporary dipole, 100
whole-number mass of, 497
Methanesulfonic acid, 752, 881
Methanethiol, 141
Methanol
ether formation by addition to alkenes, 405
hybrid orbitals in, 40
Lewis structure, 14–15
oxidation of, 482
as polar protic solvent, 259f
structure of, 339f
toxicity of, 339f
Methionine, 366
abbreviation for, 753t
isoelectric point for, 1156t
pK_a values for, 1156t
structure of, 1154f
synthesis of, 1159f
Methoprene, 789
Methotrexate, 670
p-Methoxybenzoic acid, 748
Methoxycyclohexane, conformations of, 227
Methoxy group, 336
Methoxy methyl ether, 862
p-Methoxyphenyl ester, 1196
Methyl acetate, 244
formation of, 262–263
NMR spectrum of, 557, 558, 559f
structure of, 869
Methyl acrylate, 624f
Methylamine, 999
electrostatic potential plot of, 998, 998f
m-Methylaniline, 708
N-Methylaniline, NMR spectrum of, 1003, 1003f
p-Methylaniline, 708–709
basicity of, 1018
electrostatic potential plot of, 1019f
Methylation, 285
N-Methylbenzamide, 876t
Methylbenzene, 645. *See also* Toluene
Methyl benzoate, synthesis of, 891
(*R*)-α-Methylbenzylamine, 1160–1162
2-Methylbuta-1,3-diene, 614, 1210
2-Methylbutane, 106, 130, 141, 597
2-Methylbutanenitrile, structure of, 875f
3-Methylbutane-1-thiol, 364
2-Methylbutan-2-ol, 345
2-Methylbutanoyl chloride, structure of, 873
3-Methylbut-1-ene, 587
3-Methylbut-2-en-1-ol, 638
(*R*)-α-Methylbutyrophenone, 454
2-Methyl-CBS-oxazaborolidine, 774
Methyl chloride, electrostatic potential plot for, 49, 49f
7-Methylcyclohepta-1,3,5-triene, 671
3-Methylcycloheptene, 388f
Methylcyclohexane, 139, 157, 157f
2-Methylcyclohexane-1,3-dione, 983
2-Methylcyclohexanethiol, 364f
3-Methylcyclohexanol, 334f
cis-3-methylcyclohexanol, 361
trans-4-methylcyclohexanol, 280
2-Methylcyclohexanone, 439, 971–972
alkylation of, 943–944
enolates formed from, 934
1-Methylcyclohexene, 803, 1052
3-Methylcyclohexene, 183–184
5-Methylcyclopenta-1,3-diene, 671
2-Methylcyclopentanamine, in Hofmann reaction, 1026
Methylcyclopentane, 183
2-Methylcyclopentanecarboxamide, structure of, 874
(*R*)-3-Methylcyclopentanone, 832
1-Methylcyclopentene, 388f
Methylene, 1061
α-Methylene-γ-butyrolactone, 945
Methylene chloride, 251f
Methylenecyclohexane, 390f, 839
Methylene group, 131, 390
Methyl ester, 1175
3-Methylfentanyl, 500
Methyl α-D-glucopyranoside, 1125, 1130f
Methyl group, 134
angular, 1250–1251
as electron-donating group, 694, 699
inductive effect of, 691
as ortho, para director, 694, 698–699, 704, 706, 716
Methyl halides
coupling reaction with organocuprates, 1050
S_N2 substitution reactions, 267, 272–273, 279
6-Methylhept-6-en-2-ol, 388
4-Methylhexane-2-thiol, 364f
2-Methylhexanoic acid, synthesis of, 949
5-Methylhexan-3-ol, 334
5-Methylhexan-2-one, 510, 510f
5-Methylhex-4-en-1-yne, 429f
Methyl 5-hydroxyhexanoate, 851
Methyl ketones, 437, 440
halogenation of, 939, 940f
product of acetoacetic ester synthesis, 945, 949–952
Methyllithium, 783
Methylmagnesium bromide, 783
Methylmagnesium chloride, 792
Methyl 2-methylbutanoate, 880
N-Methylmorpholine N-oxide (NMO), 473
Methyl orange, 1035
3-Methylpentanal, 820f
4-Methylpentan-1-amine, 1000
2-Methylpentane, 177f
3-Methylpentane, 177f
4-Methyl-1,4-pentanediol, 792–794, 793f
2-Methylpentan-3-one, 822f
3-Methylpentan-2-one, 821
3-Methylpent-2-ene, 389
(*E*)-3-methylpent-2-ene, 389
(*Z*)-3-methylpent-2-ene, 389
4-Methylpent-3-en-2-one, 822
1-Methyl-4-phenyl-4-propionoxypiperidine, 500
2-Methylpropane, 130, 134
2-Methylpropan-1-ol, 288
2-Methylpropan-2-ol, 688
2-Methylpropene, 727, 1061, 1205f
2-Methylpropene oxide, 1227
2-[4-(2-Methylpropyl)phenyl]propanoic acid, 133
4-Methylpyridine, 994
1,2-Methyl shift, 402
(*S*)-3-Methyl-3-sulfanylhexan-1-ol, 364
Methyl *trans*-chrysanthemate, 1072
Methyltriphenylphosphonium bromide, 836
β-Methylvaleraldehyde, 820f
Methyl vinyl ketone, 624f, 983
Metoprolol, 88
Micelles, 111–112, 112f
Michael acceptor, 981
Michael reaction, 980–982, 982f
Mifepristone, 431
Mineral oil, 172
Minus (–) compounds, 199
Mirror images, 175, 178–180, 192
Miscibility, 107
Misoprostol, 1244
Mixed aldol reaction. *See* Crossed aldol reactions
Molecular formula
of acyclic alkanes, 129, 131, 131t
of alkenes, 385
of alkyl halides, 248
of alkynes, 427
of amines, 1002
of butane, 131t
of carbohydrates, 1107f
of cycloalkanes, 129, 132
of decane, 131t
of ethane, 130, 131t
of heptane, 131t
of hexane, 131t
of icosane, 131t
of methane, 129, 130, 131t
of nonane, 131t
of octane, 131t
of pentane, 131t
of propane, 130, 131t
Molecular ion, 497–501
Molecular orbitals
of buta-1,3-diene, 1079–1080, 1079f
of ethylene, 1078, 1079f
of hexa-1,3,5-triene, 1080, 1081f
Molecular orbitals (MOs), 661–666, 662f, 663f
antibonding, 661–662, 662f, 663f, 664, 666
bonding, 662–666, 662f, 663f, 665f
degenerate, 662
highest occupied MO (HOMO), 663
lowest unoccupied MO (LUMO), 663
patterns for cyclic, completely conjugated systems, 665f
prediction/determination of relative energies of, 663–666
Molecular orbital (MO) theory, 661, 1081
Molecular recognition, 340
Molecular shape
bent, 29, 49
determining
bond angle, 26–29
bond length, 26, 26t
four groups around an atom, 26–29, 29t, 44t
linear, 26, 27, 29t, 49
tetrahedral, 26, 27–28, 29t
three groups around an atom, 26, 27, 29t, 44t
trigonal planar, 26, 27, 29t
trigonal pyramid, 28
two groups around an atom, 26, 27, 29t, 44t
Molecules
definition of, 12
polarity of, 49, 50f
Molina, Mario, 584
Molozonide, 474
Molybdenum, 1062
Monensin, 423, 855
Monohalogenation, 574
Monomers, 1199–1200

Monosaccharides, 1108–1134
  D-aldoses, family of, 1113–1115, 1115f
  common names of, 1108, 1114
  constitutional isomers of, 1108
  conversion to esters, 1127–1128
  conversion to ethers, 1127–1128
  cyclic forms, 1116–1123
    anomers, 1118–1123, 1122f
    chair form, 1122
    drawing glucose as a cyclic hemiacetal, 1117–1119, 1119f
    furanoses, 1122–1123
    Haworth projections, 1118, 1119–1121
    three-dimensional representations for D-glucose, 1122, 1122f
  determining structure of unknown, 1133–1134
  examples of, 1108
  Fischer projection formulas of, 1109–1113, 1112f
  glycosides, 1124–1127
    formation, 1124–1125
    hydrolysis, 1125–1126
    naturally occurring, 1126–1127
  isomers (D and L) of, 1112–1113
  D-ketoses, family of, 1115, 1116f
  melting point of, 1116
  physical properties of, 1116
  reactions at the carbonyl group, 1128–1134
    addition or removal of one carbon, 1131–1134
    Kiliani–Fischer synthesis, 1131, 1132–1134
    oxidation, 1129–1131, 1129f
    reduction, 1129
    Wohl degradation, 1131–1132, 1133
  reactions at hydroxy groups, 1127–1128
  solubility of, 1116, 1128
  stereogenic centers in, 1109–1118, 1120–1123
    D and L designations, 1112–1113
    multiple, 1111–1112
  structure of, 1108
  sweetness of, 1116
Monosodium glutamate (MSG), specific rotation of, 201
Monoterpene, 1246, 1246t, 1248, 1249
Morphine, 62–63, 114, 523, 630, 759, 889, 1004
Morton, Dr. William, 340
Motrin. *See* Ibuprofen
M peak, in mass spectrum, 497, 500
M + 1 peak, in mass spectrum, 498
M + 2 peak, of mass spectrum, 500–501
MPPP, 500
MRI (magnetic resonance imaging), 560, 560f
MS. *See* Mass spectrometry (MS)
MSG (monosodium glutamate), specific rotation of, 201
MTBE. *See tert*-Butyl methyl ether (MTBE)
Multiple bond
  in condensed structures, 31
  Lewis structures, 15–16
  position of, 21, 25
Multiplet, NMR spectrum, 544, 544t, 548, 548f
α-Multistriatin, 484, 484f
Muscalure, 295, 492
Muscone, 773f
Mushroom
  jack-o'-lantern, 951
  Psilocybe, 1006
Musk deer, 773
Mutarotation, 1118, 1124, 1135, 1136
Mycomycin, 211
Myelin sheath, 1240
Myoglobin, 1152, 1188, 1188f
Myosin, 1180
Myrcene, 56, 491, 1245, 1246
Myristic acid, 1234t

N

N- (prefix), 875
Nabumetone, 952
NAD^+ (nicotinamide adenine dinucleotide), 481
NADH (nicotinamide adenine dinucleotide, reduced), 481, 776–777
NAG (*N*-acetyl-D-glucosamine), 906, 1140
Nandrolone, 1254
Naphthalene, 648, 655, 724
1-Naphthol, 380
Naprosyn. *See* Naproxen
Naproxen, 86, 203, 943
Natta, Giulio, 1209
Natural gas, 40, 141
Natural product, 287
NBS (*N*-bromosuccinimide), 409, 586–588, 710
Nelfinavir, 649f
Neopentane, 130 *See also* 2,2-Dimethylpropane
Neoprene, 1211
Nepheliosyne B, 428
Neral, 121, 822
Nerol, 425
Nerolidol, 325
Neryl diphosphate, 1249, 1259
Neuroprotectin D1, 613
Neurotoxin (hemibrevetoxin B), 96
Neurotransmitter
  acetylcholine, 904
  dopamine, 1005–1006, 1006f
  serotonin, 1006
Neutrons, 8
Newman projection, 145–146, 146f, 148f
Niacin, 111, 777
Nickel, as catalyst, 237
Nicotinamide adenine dinucleotide (NAD^+), 481
Nicotinamide adenine dinucleotide, reduced (NADH), 481, 776–777
Nicotine, 59, 113–114, 286, 294, 1004, 1004f, 1021
Niphatoxin B, 443, 443f
Nitration
  of benzene, 679f, 682–683, 705, 715
  electrophilic aromatic substitution reactions, 679f, 682–683, 705–706
Nitric oxide, 571, 584
Nitriles, 869, 906–911
  acidity of, 930
  addition of organometallic reagents, 910–911
  dipole–dipole interactions of, 877
  hydrolysis of, 907–908, 1158
  infrared (IR) spectra of, 515
  IR spectra of, 878
  NMR spectra of, 879
  nomenclature, 875, 875f, 876t
  physical properties, 877
  pK_a of, 930, 931t
  products of S_N2 substitution reactions, 286, 287t
  reduction of, 907, 909–910, 1010
  structure and bonding of, 870, 873, 906
  synthesis of, 906
p-Nitroaniline
  basicity of, 1018
  electrostatic potential plot of, 1019f
Nitrobenzene, 679f, 682, 683, 694, 700
  reduction to aniline, 714
p-Nitrobenzoic acid, 748
Nitrogen
  in aromatic heterocycles, 655–658, 657f
  bonding, 13f, 18t
  common nucleophiles, 261t
  infrared (IR) absorptions in nitrogen-containing compounds, 515–516
  isotopes, 10, 504t
  mass of, 504t
Nitrogen heterocycles, 1000, 1001f
Nitrogen–nitrogen double bond, 1029, 1034
Nitrogen nucleophiles, 882
Nitrogen rule, 500
Nitro group
  as electron-withdrawing group, 694, 706–707
  in Friedel–Crafts reactions, 702
  as meta director, 694, 701, 703, 705, 715
  reduction of, 714–715, 1009
Nitronium ion, 682
o-Nitrophenol, 645
p-Nitrophenol, 706
p-Nitrophenyl ester, 1196
Nitrosamines, 1027–1029
  formation of
    N-nitrosamine from primary amines, 1028
    N-nitrosamine from secondary amines, 1029
Nitrosonium ion, 1027–1029
o-Nitrotoluene, synthesis from benzene, 705–706
p-Nitrotoluene, 703
Nitrous acid, reaction of amines with, 1028–1029
N-Methylpropan-2-amine, 999
NMO (*N*-methylmorpholine *N*-oxide), 473
NMR. *See* Nuclear magnetic resonance (NMR) spectroscopy
NMR spectrometers, 529–530, 529f
N,N-Dimethylacetamide, 69
N,N-Dimethylformamide, 569
N-Nitrosodimethylamine, 1029
Noble gases, stability of, 11
Node of electron density, 9
Nolvadex. *See* Tamoxifen
Nomenclature, A–3–6. *See also* Common names; IUPAC nomenclature
  of acid chlorides, 873, 876t
  of alcohols, 334–335, 334f
  of aldehydes, 819–820, 820f, 822
  of alkanes, 133–141, 140f
    acyclic alkanes, 135–138, 137f
    common names, 141, 141f
    cycloalkanes, 138–140, 140f
    IUPAC system, 135–140, 140f
    substituents, 134
  of alkenes, 387–390
  of alkyl halides, 249–250, 250f
  of alkyl substituents that contain branching, A–3
  of alkynes, 428–429
  of amides, 874–875, 876t
  of amines, 999–1001
  of anhydrides, 876t, 8736
  benzene derivatives, 644–647
  of bicyclic compounds, A–5–6
  of carboxylic acids, 731–734, 732t, 733f
  E, Z system, 389–390
  of epoxides, 336–337
  of esters, 873–874, 876t
  of ethers, 335–336
  generic name, 133
  of ketones, 820–821, 822, 822f
  of monosaccharides, 1108–1109, 1114
  of nitriles, 875, 875f, 876t
  of polyfunctional compounds, A–4–5
  of sulfides, 365–366
  systematic name, 133
  of thiols, 364–365, 364f
  trade name, 133
Nomex, 1226
Nonactin, 115, 116
Nonane
  constitutional isomers of, 131t
  molecular formula, 131t
Nonbonded interactions. *See* Intermolecular forces
Nonbonding (unshared) electrons, 13, 13f
Noncovalent interactions. *See* Intermolecular forces
Nonnucleophilic bases, 258
Nonpolar bond, 48
Nonpolar molecule
  description of, 49, 50f
  solubility of, 106
Nonreducing sugars, 1130, 1130f, 1137
Nonrenewable energy, 142
Nonsteroidal anti-inflammatory drugs (NSAIDs), 1244–1245
Nootkatone, 500
Noradrenaline (norepinephrine)
  structure of, 1005
  synthesis of, 285, 285f
Norethindrone, 109, 207, 430, 430f, 1254
Novocain. *See* Procaine
Noyori, Ryoji, 1164
NSAIDs (nonsteroidal anti-inflammatory drugs), 1244–1245
Nuclear magnetic resonance (NMR) spectroscopy, 496, 506, 527–569
  basis of, 528–530
  ^{13}C (carbon), 528, 556–559. *See also* Carbon NMR
  ^{1}H (proton), 528, 530–556. *See also* Proton NMR
  medical uses of, 560, 560f
  nuclear spin in, 528–529
  reference signal, 530
  signal versus peak, 543
  spectrometers, 529–530, 529f
  variables in, 529
Nucleophiles, 97, 257–261
  acetylide anions as, 434, 441
  addition to carbonyl groups, 830, 831f

alkyl halide substitution reactions with, 253
ambident, 933
amines as, 997, 1014, 1022–1024
attack of electron-deficient atoms by, 257
biological, 283–286
carbanions as, 219
common, table of, 261t
enolates, 933, 937
enols, 928
hydroxide, in alcohol synthesis, 342
Lewis base, 82
meaning of term, 82
negatively charged, 254, 261t
neutral, 254–255, 261t
nitrogen, 882
opening of epoxide rings with strong, 367–369
organometallic reagents as, 782, 783, 785–786
oxygen, 882
pi (π) bonds and, 92, 116
reactivity of, 116–117
strength, 257–261, 279–281, 282t
in substitution reactions, 253–254, 279–283. *See also* Nucleophilic substitution reactions
Nucleophile strength. *See* Nucleophilicity
Nucleophilic acyl substitution reactions, 768, 882–901, A–12
biological, 904–906, 905f
examples of, 885
general reaction of, 925
leaving groups in, 870
mechanism of, 882–883
polyester synthesis, 1212–1213
reactions of acid chlorides, 885–887, 901t
reactions of amides, 899–901, 901t
reactions of anhydrides, 887–889, 901t
reactions of carboxylic acids, 889–894, 889f, 901t
conversion to acid chlorides, 889f, 890
conversion to amides, 889f, 892–894
conversion to cyclic anhydrides, 889f, 890
conversion to esters, 889f, 891–892
reactions of esters, 894–898, 901t
reactivity of carboxylic acids and derivatives in, 883–884
summary of reactions, 901, 901t
Nucleophilic addition reactions
aldehydes and ketones, 767, 772, 778, 786–789, 795, 817–867, A–13
aldol reactions. *See* Aldol reactions
of amines, 1022–1023
of carbonyl compounds, 925, 946
Michael reaction, 981
with organometallic reagents, 786–788, 789f, 795, 801–802
polyurethane synthesis, 1213
reduction of acid chlorides and esters, 778, 795
Strecker synthesis, 1158–1159
α,β-unsaturated carbonyl compounds, 799–801
Nucleophilic aromatic substitution, 706–709, A–12
by addition–elimination, 706–708
by elimination–addition, 708–709
Nucleophilicity
basicity compared to, 257–258
periodic trends in, 258, 259, 260
S_N1 versus S_N2 substitution reactions, 279–280
solvent effects, 258–261
polar aprotic solvents, 260, 260f
polar protic solvents, 259, 259f
steric effects and, 258
Nucleophilic substitution reactions, A–12
acetylide anion reactions with alkyl halides, 441–443
adrenaline synthesis, 285–286, 285f
of alcohols, 343–344, 352–361, 362f
alcohol synthesis by, 341–342
aldehyde formation by, 778
of alkyl halides, 253–288
amine formation by, 1007–1009
bimolecular, 263
biological, 283–286
of carbonyl compounds, 767, 768–769, 925, 940–942
Claisen reaction, 975–979, 976f
components of, 253
crown ether use in, 340, 340f
epoxide ring opening reactions, 367–371
equilibrium and, 257
ethers with strong acids, 362–363
ether synthesis by, 341–342
features of, 253–255
first-order rate equation, 262–263
leaving groups in, 253–257, 256t, 265–266, 280–281, 283–284
mechanisms, 261–288
distinguishing between S_N1 and S_N2, 278–283, 279f, 282t
one-step, 262, 264, 267t
order of bond breaking and bond making, 261–264, 269–270
S_N1, 269–278, 269f, 270f, 271f, 273t, 279f, 282t
S_N2, 263–268, 264f, 266f, 267f, 268f, 268t, 279f, 282t, 287t
two-step, 262, 263, 269–270, 273t
nucleophile strength (nucleophilicity), 257–261, 279–281, 282t
organic synthesis by, 286–288, 287t
in polycarbonate synthesis, 1214
rate-determining step in, 262, 263
second-order rate equation, 262–263
solvent effects on, 258–261, 259f, 260f, 281, 282t
unimolecular, 263, 269
vinyl halides and, 286
Nucleotides, 118
Nucleus, atomic, 8
Nuts, as vitamin E source, 590
Nylon
history of, 902–903
nylon 6, 1212, 1222
nylon 6,6, 902, 1199f, 1200, 1211, 1218, 1218f
synthesis of, 902, 1211–1212

O

-o (suffix), 249
Occidentalol, 640
(*E*)-Ocimene, 301
Octane
constitutional isomers of, 131t
gasoline and, 93, 108
molecular formula, 131t
naming, 135–136
Octanenitrile, infrared (IR) spectrum of, 516
(2*E*,4*Z*,6*E*)-Octa-2,4,6-triene, 1082–1085
Oct-4-ene, 422
Oct-1-en-3-ol, 812
Octet rule
described, 11
exceptions to, 19
Octinoxate, 33, 993
Oct-1-yne, 784
Octylamine, infrared (IR) spectrum of, 515
Odor(s)
of acetone, 825
of acid chlorides, 885
of aldehydes, 826, 826f
of amines, 1004
of caraway, 204
of esters, 880
of spearmint, 204
Ofloxacin, 1044
OH proton, NMR spectra, 551–552, 551f
-oic acid (suffix), 731, 874–875
Oil(s), 395
crude, 128, 141–142, 142f
degree of unsaturation, 461–462
essential, 1245
fatty acid composition, 1235, 1235t
fish, 1235
hardening, 1236
hydrogenation of, 461–463, 462f, 1236
insolubility of, 144
melting points of, 395, 1234–1235
mineral, 172
oxidation of, 588
partial hydrogenation of, 461–462, 462f
rancid, 1236
unsaturated, 395
vegetable, 896, 898, 1235, 1236
Oil of ginger, 392f
Oil refinery, 142f
Oil spill, 128, 144
-ol (suffix), 334
Oleandrin, 851
Olefin metathesis, 1062–1067, 1063f, 1065f
Olefins, 384. *See also* Alkenes
Oleic acid, 517, 1234t, 1235t
melting point, 394t
structure, 118, 394f, 394t
triacylglycerol formation, 118
Olestra, 897–898
Olfactory receptors, 203, 203f
Olive oil, 395
-ol suffix, 364
Omega carbon, 732
Omega-3 fatty acids, 394, 395
-one (suffix), 820
-onitrile (suffix), 875, 876t
Open arrow, 444
Ophrys sphegodes, 130
Opium, 889
Opsin, 841, 842f
Optical activity, 197–199
of achiral compounds, 198
of chiral compounds, 198
dextrorotatory (*d*) and levorotatory (*l*) compounds, 199
enantiomeric excess, 200–201
of racemic mixtures, 199
specific rotation, 199
Optical purity, 200–201
Orajel. *See* Benzocaine
Oral contraceptives, 426, 430–431, 430f
Orbitals. *See also* Hybrid orbitals; Molecular orbitals (MOs); *specific orbitals*
description, 9–10
number of electrons in, 10
Orbital symmetry
cycloadditions and, 1088–1089
of electrocyclic reactions, 1082–1083
sigmatropic rearrangements and, 1093–1094
Order, of a rate equation, 235
Organic chemistry
definition of, 1
importance of, 1–2, 2f
malaria and, 4–6, 5f
Organic diphosphate, 283, 608
Organic halides. *See also* Alkyl halides
coupling reaction with organocuprate reagents, 1050–1052
coupling with alkenes (Heck reaction), 1050, 1056–1058
coupling with organoboron reagents (Suzuki reaction), 1050, 1053–1056
Organic molecules
shape. *See* Molecular shape
Organic molecules/compounds
aromatic compounds, 641–676
reactions of, 677–728
common features of, 4
drawing structures of, 30–36
condensed structures, 30–32, 31f
skeletal structures, 32–36, 33f
examples of, 1–6, 2f
grouping by functional group, 93
naming, 132
solubility of, 107–108
Organic monophosphate, 283
Organic reactions, 213–246. *See also specific types of reactions*
acid–base, 63–65, 68–69, 82–83
arrows used in describing, 218, 220–221, 220t
asymmetric, 482
bimolecular, 235
bond breaking and bond making, 217–221, 218f
bond dissociation energy, 221–225, 222t
catalysts, 237, 237f
endergonic, 228
endothermic, 221–222
energy diagrams for, 228–233, 230f, 233f
energy of activation, 229–233, 230f, 233f, 237, 237f
enthalpy change in, 221–225, 227–228
entropy change in, 227–228
enzymes in, 238, 238f
equations for, 214, 214f
equilibrium constants for, 225–227, 226f, 226t
exergonic, 228
exothermic, 221–222, 224
free energy changes in, 225–228, 226f, 226t
general types, A–12–13

kinetics of, 233
mechanisms of. *See* Reaction mechanism
multistep, 231–233, 233f
oxidation and reduction reactions, 161–162, 162f
polar, 218f, 219
radical, 218, 218f
rate of. *See* Reaction rate
sequential, 214
thermodynamics of, 225–227, 226f, 226t
types of, 215–217. *See also specific types of reactions*
unimolecular, 235
writing, 214, 214f
Organic solvents, 106–107, 108t
Organic synthesis
alkenes in, 417–418
of aspirin, 287, 287f
carbon–carbon bond-forming reactions in, 1049–1075
definition of, 287
halogenation reactions, 581
halohydrin use in, 410
by nucleophilic substitution reaction, 286–288, 287t
reactions with organometallic reagents, 802–805
retrosynthetic analysis, 444–447
sodium borohydride reductions used in, 773f
Organic triphosphate, 283–284
Organoaluminum compounds, 1209
Organoborane, 439, 440
definition of, 412
in Suzuki reaction, 1050, 1053–1056
synthesis of, 1054
Organocopper reagents. *See* Organocuprate reagents
Organocuprate reagents, 783
coupling reactions of, 1050–1052
general features of, 1050–1051
hydrocarbon synthesis in, 1051–1052
stereospecificity of, 1051
preparation of, 784
reactions with α,β-unsaturated carbonyl compounds, 801
reactions with acid chlorides, 796–797
Organolithium reagents, 783
addition to nitriles, 910
in anionic polymerization, 1206
arylborane synthesis, 1054
in ethynylestradiol synthesis, 788, 789f
in lithium diisopropylamide preparation, 933
preparation of, 783
reactions
with aldehydes and ketones, 786, 832
as base, 784–785
with esters and acid chlorides, 795
with α,β-unsaturated carbonyl compounds, 801
Organomagnesium reagents, 783
Organometallic reagents, 782–802. *See also specific reagents*
acetylide anions, 784
definition of, 782
organic synthesis with, 802–805
preparation of, 783–784
protecting groups and, 792–794, 793f
reactions
with acid chlorides, 795–797
with aldehydes, 785–787, 832
as base, 783, 784–785
with carbon dioxide, 798
with carbonyl compounds, 785–789, 792–794, 793f
with carboxylic acid derivatives, 794–797
with epoxides, 798–799
with esters, 794–796
with ketones, 785–789, 789f, 832
as nucleophile, 782, 783, 785–786
rapid acid–base reactions, 792
summary of, 802
with α,β-unsaturated carbonyl compounds, 799–801
Organopalladium compounds, 1053, 1055, 1058
Organophosphorus reagents, 835–836
Organosodium reagents, 784
Orlistat, 1197
Ortho (prefix), 645
Ortho, para activator, 695–696
Ortho, para deactivator, 695
Ortho, para director, 694–696, 698–699, 701f, 703–706, 716–718, 1032
-ose (suffix), 1108
Oseltamivir, 99, 382, 916–917
Osmium tetroxide
as oxidizing agent, 468
in syn dihydroxylation, 473–474
Osprey, 297
Osteoporosis, 19
Oxalic acid, 733, 736, 737
Oxaphosphetane, 837
Oxazaborolidine, 774
Oxidation, 161–162, 162f, 455–494
of alcohols, 468f, 476–482
of aldoses, 1129–1131
of alkanes, 161–163
of alkenes, 456f, 468f, 474–476
of alkyl benzenes, 712
of alkylboranes, 413–414
of alkynes, 456f, 468f, 476
biological, 481–482
of carbonyl compounds, 770, 782
carboxylic acid synthesis
from alkyl benzenes, 739
from alkynes, 739
from primary alcohols, 739
definition of, 161, 456, 769
dihydroxylation, 472–474
anti, 472, 473
syn, 472, 473–474
energy released by, 224
epoxidation, 468f, 469–472
of ethanol, 481–482
Sharpless, 482–484
stereochemistry of, 469–470
in the synthesis of disparlure, 470–472, 471f
of fatty acids, 897
general scheme for, 456f
green chemistry, 479–480
hydroboration–oxidation reactions, 397f, 411–415, 412f, 413f, 414t, 439–441
of hydrocarbons, 456, 456f
of lipids, 165
of oils, 462
oxidative cleavage
of alkenes, 468f, 474–476
of alkynes, 468f, 476
of polycyclic aromatic hydrocarbons (PAHs), 373
reduction, relationship to, 456
of substituted benzenes, 712
of unsaturated fatty acids, 1236
of unsaturated lipids, 588–589, 589f
Oxidative addition
in Heck reaction, 1058
in palladium-catalyzed reactions, 1053, 1055, 1058
in Suzuki reactions, 1055
Oxidative cleavage
of alkenes, 468f, 474–476
of alkynes, 468f, 476, 739
Oxide, alkene, 336, 337
Oxidized state, 161
Oxidizing agents, 467–468. *See also specific agents*
NAD^+, 777
peroxyacids, 467–468
Oxime, 1132
Oximene, 491
Oxiranes, 333, 336, 337. *See also* Epoxides
6-Oxoheptanal, 991
Oxone, 480
-oxy (suffix), 336
Oxycodone, 126, 523
Oxy-Cope rearrangements, 1095–1096
Oxygen
bonding, 13f, 18t
common nucleophiles, 261t
in condensed structures, 31
formal charge, 18t
infrared (IR) absorptions in oxygen-containing compounds, 514–515
isotopes of, 504t
mass of, 504t
as radical scavenger, 573
tetrahedreal, 333
Oxygen nucleophiles, 882
Oxytocin, 871, 1167–1168, 1168f, 1184
-oyl (suffix), 822
Ozone
destruction of atmospheric, 252, 584, 585f
as oxidizing agent, 467, 474–476
production of, 474
stratospheric, 3
synthesis of, 584
Ozonide, 474, 476
Ozonolysis, 474–475

P

PABA (*para*-aminobenzoic acid), 632, 725, 1037–1038
Paclitaxel, 174, 184, 208. *See also* Taxol
Padimate O, 632
PAHs. *See* Polycyclic aromatic hydrocarbons (PAHs)
Pain reliever, 952. *See also* Analgesics
Palau'amine, 527, 553
Palladium-catalyzed reactions, 236, 237
alkene reduction, 458
alkyne reduction, 464, 466f
general features of, 1052–1053
Heck reaction, 1050, 1056–1058
hydrogenation of aldehydes and ketones, 772
oxidative addition in, 1053, 1055, 1058
in peptide synthesis, 1175–1176
reductive elimination in, 1053, 1055, 1058
Suzuki reactions, 1050, 1054–1056
Palmitic acid, 1234, 1234t
Palmitoleic acid, 1234t
Palm oil, 1235
Pantothenic acid, 111
Para (prefix), 645
Para red, 1037
Parent ion, 497. *See also* Molecular ion
Parent name, 133
Parkinson's disease, treatment of, 7, 50, 51, 1005, 1006f
Paroxetine, 88, 342
Parts per million (ppm), 530
Patchouli alcohol, 352f, 1258
PCB (polychlorinated biphenyl), 108
PCC (pyridinium chlorochromate), 468, 477–478
Peanut butter, 461, 462
PEG (polyethylene glycol), 1208
Penicillamine, 1192
Penicillin(s), 881, 900–901
discovery of, 495, 517
penicillin G, 121, 172, 495, 517–518
Pentacycloanammoxic acid methyl ester, 1102
Pentadecane, 131t
Penta-1,4-diene, 605–606, 606f
hydrogenation of, 616
(*E*)-Penta-1,3-diene, hydrogenation of, 616
Pentalene, 669
Pentan-2-amine, 1027, 1027f
Pentane, 498–499, 616
boiling point, 103
constitutional isomers of, 131t
melting point of, 105
molecular formula, 131t
structure, 130–131
van der Waals forces, 100, 100f, 102
Pentane-2,4-dione, 926, 930, 954, 955
Pentane-3-thiol, 364f
Pentan-3-ol, 790
Pentan-2-one, 87
mass spectrum of, 504
preparation from an acid chloride and an organocuprate reagent, 797
Pentan-3-one, boiling point of, 103, 104f
Pent-1-ene, 498–499
2-Pentylcinnamaldehyde, 969
Pent-1-yne, 498–499
Peptide bonds, 1165–1167
s-cis conformation, 1166–1167
planar geometry of, 1166–1167
s-trans conformation, 1166–1167
Peptides, 880–881, 1164–1179
constitutional isomers, 1165
interesting examples, 1167–1168, 1168f
N- and C-terminal amino acids, 1165–1166
peptide bond in, 1165–1167
sequencing, 1169–1172
amino acid analysis, 1169
Edman degradation, 1169–1170
partial hydrolysis of peptide, 1171–1172, 1171t

Peptides (*Continued*)
simple, 1165–1166
synthesis of, 1172–1179
automated, 1177–1179
protecting groups in, 1173–1177
solid phase technique, 1177–1179
Percent *s*-character, 46, 76
bond length and, 615
effect on basicity of amines, 1020
hybrid orbital in alkenes, 302
infrared (IR) absorption and, 512
Percent transmittance, 509
Perfume, 464
Pericyclic reactions, 1076–1105
defined, 1077
rules for, 1097, 1097t
types, 1077–1078
cycloaddition reactions, 1077, 1087–1091
electrocyclic reactions, 1077, 1081–1087
sigmatropic rearrangements, 1078, 1091–1097
Period, in periodic table, 9
Periodic table, 8–10
bond length trends, 26
element location and bonding type, 11
of elements common in organic chemistry, 9f
first-row elements, 9f, 10
organization of, 9
second-row elements, 9f, 10
bonding of, 12–13
exceptions to the octet rule, 19
shape arrangements, 26
third-row elements, 19
trends
acidity, 70–72
bond length, 26
electronegativity, 47, 47f
Periodic trends
in bond dissociation energies, 223
in leaving group ability, 255–256
nucleophilicity, 258, 259, 260
Periplanone B, 518, 972, 972f, 1076, 1096, 1258
Perkin, William Henry, 1036
Perlon. *See* Nylon, Nylon 6
Peroxide, as radical initiator, 572, 590–591
Peroxyacetic acid, 468–469
Peroxyacids
in epoxidation, 469–471
as oxidizing agents, 467–468
Peroxy radical, 588, 589f, 1201
Pesticide, 252
PET. *See* Polyethylene terephthalate (PET)
Petroleum, 128, 141–142, 142f, 144
ethylene preparation from, 391
as feedstock of polymer synthesis, 1218
Ph- (abbreviation), 646
PHAs (polyhydroxyalkanoates), 1222
PHB (polyhydroxybutyrate), 1222
PHBV (polyhydroxyvalerate), 1222
α-Phellandrene, 1246f
Phenanthrene, 655, 671, 676
Phenol(s), 645
acidity of, 742, 744f
as antioxidants, 590
in aspirin synthesis, 287, 287f
Bakelite preparation from, 1216–1217, 1217f
benzene conversion to, 171
formation from anisole, 363
homolysis reaction, 246
hydroxy group of, 332
naming, 646
pK_a of, 742, 744f
polyhalogenation of, 701
substituted, 749
synthesis by Claisen rearrangement, 1096
synthesis from aryl diazonium salts, 1030
synthesis from chlorobenzene, 708
Phenoxide, 742–743, 743f, 744f
Phentermine, 86
Phenylacetaldehyde, 820f, 928
Phenyl acetate, 915
Phenylacetic acid, 992
Phenylacetonitrile, 957
Phenylalanine
abbreviation for, 753t
isoelectric point for, 1156t
N-acetyl, 1164
pK_a values for, 1156t
structure of, 1154f
synthesis of, 1158, 1164
Phenylcyclohexane
conformations of, 227
phenyl group in, 93–94
Phenylethanal, 820f
2-Phenylethanamine, 1012–1013
derivatives of, 1005–1007
in Hofmann elimination reaction, 1025
Phenyl group, 93–94, 94t, 646
Phenyl isothiocyanate, 1169–1170
(1*Z*,3*E*)-1-Phenylocta-1,3-diene, 1056
2-Phenylpropanoic acid, 212
1-Phenylpropan-1-ol, 296
N-Phenylthiohydantoin (PTH), 1169–1170, 1197
N-Phenylthiourea, 1170
Pheromones, 129, 130, 470–472, 471f, 484, 484f, 887, 972, 972f, 1055
Phomallenic acid C, 448
Phosgene, 1214
Phosphate, 283
Phosphatidylcholine, 1238
Phosphatidylethanolamine, 1238
Phosphine, in palladium-catalyzed reactions, 1053
Phosphoacylglycerols, 1238–1239, 1239f, 1240f
Phosphodiester, 1238–1240
Phospholipids, 126
in cell membranes, 113–115, 114f, 116f, 1238–1240
phosphoacylglycerols, 1238–1239, 1239f, 1240f
sphingomyelins, 1240, 1240f
structure of, 113, 114f, 1238
Phosphoric acid, 1238
Phosphoric acid diester, 1238
Phosphorus
leaving groups derived from, 283–284
octet rule exceptions in, 19
Phosphorus oxychloride, dehydration of alcohols using, 345, 351, 352f
Phosphorus tribromide, conversion of alcohols to alkyl halides with, 356, 357, 357t
Photochemical cycloaddition, 1088
Photochemical electrocyclic reactions, 1085–1086
Photons, 506–508
Photosynthesis, 339, 339f, 1107–1108
Phthalic acid, 712, 759
Phthalimide, 1008
Phylloquinone, 111
Physical properties. *See* Boiling point; Melting point; Solubility
Phytyl diphosphate, 690, 690f
Pi* (π*) antibonding molecular orbital, 661–662, 662f
Pi (π) bond, 42–43, 42f, 43f, 44t
in alkene double bond, 298–299
in alkenes, 384, 385, 396–399, 416, 417
in alkynes, 318, 319f, 427–428, 432, 433f
in aromatic compounds, 93, 642, 643
of benzyne, 709
breakage in addition reactions, 216, 384, 396–399, 397f, 404, 405, 406, 408, 416, 417
bromination as test for, 405
Brønsted–Lowry bases and, 62–63, 62f
in carbonyl group, 97, 765–766, 767
in conjugated allyl carbocation, 607
in cycloaddition reactions, 1077, 1088–1091, 1091t
degrees of unsaturation and, 385, 460–461
delocalized, 607, 616
in dienes, 605–606
in electrocyclic reactions, 1077, 1082–1086, 1087t
formation in elimination reactions, 215, 298–299, 310, 313, 318, 319f
in functional groups, 92, 116
in Lewis base, 81, 83
radical reactions with, 572–573, 595
in resonance structure, 74
in sigmatropic rearrangements, 1078, 1093–1095
strength, 384
Pi (π) bonding molecular orbital, 661–662, 662f
Picometer (pm), 26
Pi (π) electrons
in aromatic compounds, 647, 651–653, 653t, 656–660, 663–666
in benzene, 643, 644, 678
Hückel's rule and, 651–653, 653t, 660, 665
in NMR spectroscopy, 539–540, 540t
Pinacol, 381
Pinacolone, 381
α-Pinene, 1259
Pioglitazone, 722
Piperidine
basicity of, 1020–1021, 1021t
structure, 1001f
Pitavastatin, 995
Pitocin. *See* Oxytocin
Pivalic acid, 68
pK_a values, 66–75, A–1–2
acid strength and, 66–68
of amines, 1014, 1016–1017, 1021t
for amino acids, 1155, 1156, 1156t
of carbonyl compounds, 929, 929t, 931t
definition, 66
of leaving groups in nucleophilic substitution reactions, 256, 256t
logarithmic scale of, 68
of selected organic compounds, 67t
of strong acids, 78–79
of strong bases, 79
PLA (polylactic acid), 903
Planar carbocation, 403, 1124–1125
Planar double bond, 406, 412, 414, 4013
Planar molecule
benzene, 643–644
Hückel's rule, 652–653
Planar radical, 582–583
Plan B, 431
Plane of symmetry, 180, 1133–1134
in disubstituted cycloalkanes, 195
of meso compound, 193
Plane-polarized light, 198–199
Plaque, atherosclerotic, 905, 905f
Plasticizers, 108, 1217
Plastics, 2f, 94
Platinum, as catalyst, 237
Plavix, 190. *See also* Clopidogrel
β-Pleated sheet, 1181–1182, 1181f, 1182f, 1183f, 1185f
Plocoralide B, 252
Plus (+) compounds, 199
Poison ivy, 749
Polar aprotic solvents, 260, 260f, 281, 282t, 305, 307t, 934
Polar bond
carbon–halogen, 252–253, 253f
description of, 48–49, 50f
Polarimeter, 198
Polarity, of carboxylic acids, 741
Polarizability
boiling point and, 103, 104f
inductive effects on substituted benzenes, 691
van der Waals forces and, 100
Polarized light, 198–199
Polar molecule
description, 49, 50f
dipole–dipole interactions between, 101
solubility of, 106
Polar protic solvents, 259, 259f, 281, 282t, 311t, 312t, 935
Polar reactions, 218f, 219
Poly(acrylic acid), 594, 1223
Polyacrylonitrile, 1205f
Polyalkylation, 702
Polyamides, 902, 1211–1212
Polyaspartate, 1223
PolyAspirin, 1229
Polyatomic ion, 17
(*Z*)-Poly(buta-1,3-diene), 1211
Polybutyrate adipate terephthalate (PBAT), 1226
Polycarbonates, 1199f, 1214
Polychlorinated biphenyl (PCB), 108
Polycyclic aromatic hydrocarbons (PAHs), 373, 648, 648f, 655
Polyenes, 475, 613
Poly(ester amide), 1229
Polyesters, 903, 1212–1213
Polyethers, 340, 1207
Poly(ethyl acrylate), 601
Poly(ethyl α-cyanoacrylate), 1205f
Polyethylene, 94, 391, 392f, 593, 601, 1199f, 1202–1203, 1209
Polyethylenefuranoate (PEF), 1213
Poly(ethylene glycol) (PEG), 124, 1208
Polyethylene terephthalate (PET), 903, 1198, 1200f, 1212–1213, 1216, 1221, 1221t
Polyfunctional compounds, nomenclature of, A–4–5
Polyhalogenation, 701

Polyhydroxyalkanoates (PHAs), 1222
Polyhydroxybutyrate (PHB), 1222
Polyhydroxyvalerate (PHBV), 1222
Polyisobutylene, 601, 1205f
Poly(lactic acid) (PLA), 903, 1219
Polymerization, 593–595
  anionic, 1205–1206, 1205f
    of epoxides, 1207–1208
  cationic, 1204–1205, 1205f
  chain branching in, 1202–1203
  coordination, 1209
  definition of, 593, 1199
  feedstock for, 1218
  head-to-tail, 595
  ionic, 1204–1206, 1205f
  living, 1206
  radical, 595, 1201–1202, 1202f
  Ziegler–Natta, 1208–1209
Polymers, 593–595, 1198–1230
  alternating, 1207
  amorphous regions of, 1216–1217
  atatic, 1208–1209
  "Big Six," 1220–1221, 1221t
  biodegradable, 1222–1223
  chain-growth, 1199, 1200–1207
  in consumer products, 1199, 1199f
  copolymers, 1207, 1212
  crystalline regions of, 1216–1217
  definition of, 175, 1199
  derivation of word, 1199
  green synthesis of, 1217–1220, 1218f, 1219f
  insoluble, 1177
  isotactic, 1208–1209
  in medicine and dentistry, 594t
  molecular weights of, 1200
  natural and synthetic fibers, 901f, 902–903
  nylon, 902–903, 1199f, 1200, 1211–1212, 1218, 1218f, 1222
  physical properties of, 1216–1217
  polyesters, 903
  random, 1207
  recycling of, 1198, 1220–1222, 1221t
  shorthand representation of, 1200, 1200f
  step-growth, 1200, 1211–1215
  stereochemistry of, 1208–1209
  structure of, 1199, 1216–1217
  syndiotactic, 1208–1209
  synthetic, 1198–1230. *See also* Synthetic polymers
Poly(methyl methacrylate) (PMMA), 601
Poly (prefix), 1205f
Polypropylene, 594t, 1221t
Polysaccharides, 1138–1140
  cellulose, 1138
  glycogen, 1139
  starch, 1139
Polystyrene, 108, 570, 593, 601, 1200f, 1216, 1221t
  preparation from ethylene, 392f
Polystyrene derivative, 1178
Polytetrafluoroethylene (Teflon), 594t
Poly(trimethylene terephthalate) (PTT), 1219, 1219f
Polyurethanes, 1213
Poly(vinyl acetate), 594, 1205f
  preparation from ethylene, 392f
Poly(vinyl alcohol), 1228
Poly(vinyl butyral), 1228
Poly(vinyl chloride) (PVC), 124, 251, 593, 594t, 1199, 1217, 1221t
  preparation from ethylene, 392f
p orbitals
  description, 9
  hybridization of, 36–45, 38f, 43f, 44t
  hyperconjugation, 274–275
  overlap of, 605–606, 612, 615, 616, 644, 644f, 651, 661–662, 663f
    molecular orbitals and, 1078–1080, 1079f, 1081f
  2*p* orbitals, 10, 36–38, 38f
    in alkenes, 299–300
    in alkynes, 427
Porphyrin, 1188
Potassium *tert*-butoxide, 258
  in dehydrohalogenation reactions, 298, 298t
Potassium dichromate, 468, 477–479, 479f
Potassium hydroxide, in dehydrohalogenation reactions, 298t
Potassium iodide, 11, 1030
Potassium permanganate ($KMnO_4$)
  as oxidizing agent, 468, 712, 716, 782
  in syn dihydroxylation, 473
Potassium peroxymonosulfate, 480
Potassium propanoate, 733f
Potassium sorbate, 737
Potassium sulfate, 480
Precor. *See* Methoprene
Prednisone, 826
Prefix, 133
  cis, 388–389
  cyclo-, 132, 138
  D-, 1112–1113
  deoxy, 1141
  di-, 136, 335, 875, 8999
  E, 389–390
  epoxy, 336
  eth-, 133
  iso-, 134
  L-, 1112–1113
  meta-, 645
  meth-, 133
  N-, 875
  ortho-, 645
  para-, 645
  poly, 1205f
  R, 185–190, 194–195, 1110, 1112–1113
  S, 185–190, 194–195, 1110, 1112–1113
  sec-, 134
  tert-, 134
  tetra-, 136
  trans, 388–389
  tri-, 136, 8999
  Z, 389–390
Pregabalin, 121
Prepolymer, 1214–1215, 1215f
Primary (1°) alcohols, 96, 415
  classification as, 332
  conversion to alkyl halides with HX, 352–354, 357t
  conversion to alkyl halides with phosphorus tribromide, 357, 357t
  conversion to alkyl halides with thionyl chloride, 356, 357t
  dehydration by E2 mechanism, 347
  hydrogen bonding extent, 337
  from nucleophilic addition to aldehydes, 778
  oxidation to carboxylic acids, 477, 478–479, 739
  from reactions of aldehydes with organometallic reagents, 785–787
  from reduction of acid chlorides, 777–778
  from reduction of aldehydes, 461, 770, 771, 772
  from reduction of carboxylic acid derivatives, 770
  from reduction of carboxylic acids, 780
  from reduction of esters, 777–778
  from reduction of monosaccharides, 1129
Primary (1°) alkyl halides, 96
  acetylide anion reactions with, 441–442
  classification as, 248
  E2 elimination reactions, 306, 320t, 321
  S_N2 substitution reactions, 267, 279, 279f, 320t, 321
  synthesis of, 352
Primary (1°) amides, 97, 870, 874, 878t, 879–878, 883, 886, 888, 894
  tautomers, 907
Primary (1°) amines, 97
  addition to aldehydes and ketones, 840–842, 842f, 843f
  amino acids, 1155
  from direct nucleophilic substitution, 1007–1008
  Gabriel synthesis, 1008–1009
  IR spectra of, 1002, 1003f
  nomenclature, 999
  reactions with acid chlorides and anhydrides, 1023
  reactions with aldehydes and ketones, 1023
  reactions with nitrous acid, 1028–1029
  from reduction of amides, 1010
  from reduction of nitriles, 1010
  from reduction of nitro compounds, 1009
  structure of, 997
  synthesis from reduction of nitriles, 909
Primary carbon, 94
Primary carbon radical, 571–572, 572f
Primary hydrogen (1°), 94, 134
Primary structure, of proteins, 1180, 1185f
Pristane, 165
Procaine, 649f
Procyclidine, 812
Progesterone
  analogs of, 430
  function in body, 1254t
  structure of, 430, 1232f, 1254t
  synthesis of, 974, 975, 975f
Progestins, 1254
Proline, 762
  isoelectric point for, 1156t
  pK_a values for, 1156t
  structure of, 1154f, 1155, 1180
Prontosil, 1037
Propanal
  aldol reaction of, 965, 968
  enolate formation from, 929
  IR spectrum of, 823, 823f
  NMR spectra of, 825, 825f
  Wittig reaction, 837
Propanamide
  boiling point of, 104
  infrared (IR) spectrum of, 515
Propan-1-amine, 1025
Propan-2-amine, 999
Propane
  bromination of, 579–580, 580f
  chlorination of, 578–581, 581f
  molecular formula of, 130, 131t
  naming, 134
  Newman projection, 146f
  as propellant, 584
  structure of, 130
  torsional energy of, 147
Propanedioic acid, 733
Propane-1,3-diol, 1219, 1219f
Propanenitrile, 875
Propane-1-thiol, 364, 366
Propane-1,2,3-triol, 335
Propanoic acid, NMR spectra of, 736f
Propan-1-ol, 80
  conversion to butanenitrile, 358
  IR spectrum of, 509
  NMR spectrum of, 558, 559f
Propan-2-ol, 339f
Proparacaine, 726, 1048
Propene
  borane addition to, 412
  hydrohalogenation, 400
  monomers in polymerization, 594t
  radical halogenation, 585–586, 591–592
  synthesis from propan-1-amine, 1025
Prop-2-en-1-ol, 388
Propionic acid, structure of, 732t
Propiophenone, 775
  halogenation of, 938–939
Propoxyphene, 181
Propranolol, 85, 380
Propylbenzene, 245, 713
Propyl group, 134
Prostacyclins, 1242, 1243, 1243f, 1243t
Prostaglandins, 165, 233, 1242–1245
  analogues of, 738, 1244
  inhibitors of synthesis, 1244–1245
  PGA_2, 305
  PGE_1, 1244
  PGE_2, 1244
  $PGF_{2\alpha}$, 164, 164f, 242, 424, 603, 738, 1232f, 1242, 1243f, 1257
  PGG_2, 603, 738, 1243, 1243f
  PGI_2, 1243f
  synthesis of, 738, 1243, 1243f
Prosthetic group, 1187
Protecting groups, 792–794
  acetals as, 851–852
  amides for amines, 1024, 1024f
  in amino acid synthesis, 1173–1177
  general strategy for using, 793f
Proteins, 1152–1197, 1164. *See also* Peptides
  amide linkages in, 880, 899
  conjugated, 1187
  fibrous, 1186
  folding, 122–123
  globular, 1186
  important examples, 1186–1189, 1186f, 1187f, 1188f
  membrane, 1153
  natural fibers, 902, 902f
  stability of, 899
  structure, 1179–1186
    primary, 1180, 1185f
    quaternary, 1185, 1185f

Proteins (*Continued*)
secondary, 1180–1183, 1181f, 1182f, 1183f, 1185f
tertiary, 1183–1185, 1185f
subunits, 1185
Protic solvents, polar, 259, 259f, 281, 282t, 312t, 935
Proton acceptor, 62
Protonation
in acid-catalyzed ester hydrolysis, 894, 895
of alcohols, 344, 346, 347, 349, 350, 353, 354, 355
in aldehyde/ketone nucleophilic addition reactions, 829–830, 846
of carboxylic acids, 740
of epoxides, 367, 369
of ethers, 363
in glycoside formation, 1125
in glycoside hydrolysis, 1126
in nucleophilic addition to aldehydes and ketones, 767, 772
in tautomerization, 438–439, 927
Proton donor, 62, 69
Proton NMR, 528, 530–556
of aldehydes, 825, 825f
of amines, 1003, 1003f
of aromatic compounds, 654
of benzene derivatives, 552–553, 552f, 647, 647t
of carboxylic acid derivatives, 879
of carboxylic acids, 735, 736f
characteristic absorptions, A–10–11
chemical shift
equation for, 531
predicting values of, 537–538
protons on benzene rings, 539, 540t
protons on carbon–carbon double bonds, 539, 540t
protons on carbon–carbon triple bonds, 540, 540t
scale, 530
values for common proton types, 538t
coupled protons, 544
of cyclohexane conformations, 552
general principles
chemical shifts, 530–531
downfield signals, 530–531
reference signal, 530
scale, 530
signal intensity, 530
upfield signals, 530–531
identification of an unknown, 553–556
integrals, 541
intensity of signals, 541–542
of ketones, 825
n + 1 rule, 547
number of signals, 531–535, 532f
equivalent protons in alkenes and cycloalkanes, 533–534
general principles, 531–532
homotopic, enantiotopic, and diastereotopic protons, 534–535
OH proton, 551–552, 551f
position of signals, 535–538
chemical shift values, 537–538, 538t
shielding and deshielding effects, 536–537, 536f, 537f, 540f, 540t
regions of spectra, 540, 540f
spin–spin splitting, 542–551
in alkenes, 549–551, 549f, 550f
common patterns of, 545t
complex examples of, 546–549
how a doublet arises, 542–543
how a triplet arises, 543–544
overview of, 542
rules and examples, 544–546, 544t
Protons
description of, 8
diastereotopic, 534–535
enantiotopic, 534–535
geminal, 549
homotopic, 534–535
number of, 8
relative acidity of, determining, 77–78
Proton transfer reaction
aspirin and, 80–81
description of, 63–65
Protosterol cation, 1252f, 1253
Proventil. *See* Albuterol
Proximity effect, 620
Prozac, 3, 202, 268, 268f
D-Pscicose, 1116f
Pseudoephedrine, 72, 208, 1013, 1046
Psilocin, 1006
Psilocybe mushroom, 1006
PTH (*N*-phenylthiohydantoin), 1169–1170
PTT. *See* Poly(trimethylene terephthalate)
Puffer fish, 622, 622f
Purification procedures
extraction, 749–751, 749f, 751f
Purine, 670, 1001f, 1142
Putrescine, 1004
PVC. *See* Poly(vinyl chloride)
Pyranose ring, 1117, 1122, 1135
Pyrethrin I, 121, 1059, 1072
Pyridine, 80, 395, 655–657, 727, 885, 887, 917
alcohol conversion to alkyl tosylates in, 359, 361, 362f
basicity of, 1020–1021, 1021t
conversion of alcohols to alkyl halides, 356
conversion of monosaccharides to esters using, 1128
dehydration of alcohols using, 351, 352f
electrostatic potential plot of, 656, 657f
structure of, 1001f
Pyridinium chlorochromate (PCC), 468, 477–478
Pyrimidine, 1001f, 1142
α-Pyrone, 670
Pyroxidine (vitamin B_6), 124
Pyrrole, 655–657, 671, 727
basicity of, 1020, 1021t
electrostatic potential plot of, 657, 657f
structure of, 1001f
Pyrrolidine, structure of, 1001f
Pyruvic acid, 777

Q

Qinghao, 414
Quantum, 506
Quartet, NMR spectrum, 544t, 545t, 546, 551
Quaternary ammonium salt. *See also* Ammonium salt
in Hofmann elimination reaction, 1024–1026
synthesis of, 1008
Quaternary carbon, 94
Quaternary structure, of proteins, 1185, 1185f
Quiana, 1226
Quinapril, 126, 294
Quinine, 210
as natural alkaloid, 1004
source of, 307
structure of, 658, 1022
synthesis of, 307, 307f, 941, 941f
Quinoline, 464, 728
o-Quinone, 1105
Quintet, NMR spectrum, 544t
Quinuclidine, 295

R

R (prefix), 185–190, 194–195
Racemic mixture (racemate), 199, 199t
of alcohols from hydroboration–oxidation of alkenes, 414
alkene halogenation, 407
of alkyl chlorides, 354
of amino acids, separation of, 1159–1162, 1160f
in carbene addition, 1060
in enolate alkylation, 936, 936f, 943
from epoxidation of alkenes, 470, 472
in epoxide ring opening reactions, 368
formation by hydrohalogenation, 402–403
halogenation reactions and, 407, 582–583
in hydride reduction of ketones, 773–774
in organometallic reagent reactions, 788
S_N1 substitution reactions and, 271–272, 271f
Racemization
at the α carbon of carbonyl compounds, 936
definition of, 271
in S_N1 substitution reactions, 271–272, 271f
Radiation
electromagnetic, 506–508, 507f
infrared, 508
RF (radiofrequency), 528–529, 560
ultraviolet, 630–632
Radical(s)
antioxidant reaction with, 590
bond formation from, 219
classification of, 571
definition of, 571
as electrophile, 219
as reactive intermediate, 218–219, 218f
resonance-stabilized, 587, 588
stability of, 571–572, 572f
Radical anion, 465
Radical cation, 497, 501–502
Radical inhibitor, 573
Radical initiator, 572
Radical intermediates, 571, 587, 591
Radical polymerization, 1201–1202, 1202f
Radical reactions, 218, 218f, 570–603. *See also specific reactions*
of alkanes, 573–583
of alkenes, 573, 590–593
applications of
antioxidants, 589–590
oxidation of unsaturated lipids, 588–589, 589f
ozone destruction by CFCs, 584, 585f
carbon–hydrogen bond, reaction of radical with, 573
double bonds, addition to, 573, 590–593
general features of, 572–573
halogenation at the allylic carbon, 585–588
bromination, 586–588
product mixtures in, 587–588
halogenation of alkanes, 574–583
of an achiral starting material, 582, 582t
bromination, 578–580, 580ff
of a chiral starting material, 582t, 583
chlorination, 574–583, 581f
energy changes during, 576–577, 577f
mechanism of, 575–576
in organic synthesis, 581
stereochemistry of, 582–583, 582t
halogenation of alkyl benzenes, 709–711
inhibitors of, 573
polymerization, 593–595, 594t
two radicals reacting with each other, 573
Radical scavenger, 573
Radical substitution reaction, 574, A–12
Randomness, entropy as measure of, 226
Rate constant (k), 235
Rate-determining step
in chlorination of ethane, 577, 577f
E1 elimination reactions, 310, 311f, 312
electrophilic additions, 399, 399f, 404
in halogenation reactions, 577, 579–581
bromination, 580
chlorination, 581
halogenation of alkenes, 407
in a multistep mechanism, 232, 233f, 235
in nucleophilic substitution reactions, 262, 263
rate equation for, 235
S_N1 substitution reactions, 273, 275, 277–278
Rate equation, 234–236
first-order, 235, 262–263
for nucleophilic substitution reactions, 262–263
second-order, 235, 262–263
Rate law, 234–236
RCM (ring-closing metathesis), 1065–1067, 1065f
Reaction arrow, 214, 220t
curved, 218, 220t
Reaction coordinate, 229, 230f, 231, 233f
Reaction mechanism
bond cleavage, 217–218
bond formation, 219
concerted, 217
definition of, 217
energy diagram for a two-step, 231–233, 233f

rate equations and, 234–236
stepwise, 217, 231–233, 235
Reaction rate
carbocation stability and, 275
catalysts and, 236–237, 237f
concentration effect on, 234–236
of endothermic reactions, 230f, 276
energy of activation (E_a) and, 234
of exothermic reactions, 230f, 277
nucleophilic substitution reactions, 263, 267, 269, 275–277, 276f
rate-determining step, 232, 235
temperature effect on, 234–235
Reactions. *See* Organic reactions
Reactive intermediate
carbanion, 218f, 219
carbocation, 218f, 219, 269, 271, 348, 350
description of, 217
radicals, 218–219, 218f
synthetic intermediate compared, 418
in a two-step reaction mechanism, 231
Reactivity, functional groups and, 116–117
Reagent, 214, 214f
Rebaudioside A, 1127
(*S*,*S*)-Reboxetine, 492
Receptors, olfactory, 203, 203f
Recycling of polymers, 1198, 1220–1222, 1221t
Red algae, 252
Red seaweed, 252
Reduced state, 161
Reducing agents, 457. *See also specific agents*
chiral, 774
metal hydride reagents, 457, 771–773, 777–781, 781t
NADH, 776–777
Reducing sugars, 1130, 1130f, 1135, 1136
Reduction, 161–162, 162f, 455–494
of acid chlorides and esters, 777–779
of alkenes, 456, 456f, 458–461
of alkyl halides, 466–467, 467f
of alkynes, 456, 456f, 463–466, 466f
to alkanes, 456f, 463–464, 466f
to cis alkenes, 463, 464, 466f
to trans alkenes, 463, 465–466, 466f
of amides to amines, 780, 1010
of aryl ketones to alkyl benzenes, 712–713
asymmetric, 774
of carbonyl compounds, 769–781
biological, 776–777
carboxylic acids and derivatives, 777–781
by catalytic hydrogenation, 772
enantioselective, 774–777, 775f
with metal hydride reagents, 771–773, 773f, 781t
stereochemistry of, 773–774
of the carbonyl group of monosaccharides, 1129
Clemmensen, 712
definition of, 161, 456, 769
dissolving metal, 457, 465–466
of epoxides, 466–467, 467f
general scheme for, 456f
of hydrocarbons, 456, 456f
hydrogenation, 458–463
of ketones to amines, 1011–1013
of nitriles, 907, 909–910, 1010
of nitro groups, 714–715
oxidation, relationship to, 456
of polar C–X sigma (σ) bonds, 466–467
reagents for, 781t
of substituted benzenes, 712–715
of triple bonds, 463–466, 466f
Wolff–Kishner, 712
Reductive amination, of aldehydes and ketones to amines, 1011–1013
Reductive elimination
in the Heck reaction, 1058
in palladium-catalyzed reactions, 1053, 1055, 1058
in Suzuki reactions, 1055
Refining, 141–142, 142f, 144
Refrigerant, 252
Regioselectivity
of alcohol dehydrations, 345
of E1 elimination reactions, 312
of E2 elimination reactions, 308, 315
of electrophilic addition of hydrogen halides, 400, 403t
of epoxide ring opening reactions, 369–370
of halohydrin formation, 409–410, 410t
of Hofmann elimination, 1026–1027
of hydroboration, 412, 414t
Reserpine, 923, 1090
Resolution, 202
Resolution of amino acids, 1160–1163, 1160f
Resolving agent, 1160–1161
Resonance, 19–25
of allylic carbocations, 607–609
allyl type, 609
benzene, 643
examples of, 608–610
principles, 20
usage of term, 607
Resonance delocalization, 75
Resonance effects
on acidity, 74–75, 75f, 77f
on basicity of amines, 1021t
electron-donating, 691–693
electron-withdrawing, 691–693
on substituted benzenes, 691–693
Resonance hybrid, 20, 24–25, 610–612
of benzene, 643
drawing, 25
electron delocalization, hybridization, and geometry, 612
major and minor contributors, 24–25, 610–611
Resonance stabilization, of carboxylic acids, 742–743, 743f, 744f
Resonance-stabilized allylic carbocations, 607, 607f, 617–618
Resonance-stabilized carbocations, 435–436, 438, 607, 607f, 617–618, 620
in electrophilic aromatic substitution reaction, 679, 681, 696–697
in glycoside formation, 1124–1125
in glycoside hydrolysis, 1126
Resonance-stabilized radicals, 587, 588
Resonance structures, 19–25, 607–612
of acetate, 742, 744f
acidity and, 74–75
of allyl radical, 586
of benzylic radical, 710
of carboxylic acid derivatives, 871
cations with a positive charge adjacent to a lone pair, 609
conjugated allyl carbocation, 607
of conjugated dienes, 615
double bond
conjugated, 609
major contributor, 610–611, 615
minor contributors, 610–611, 615
with one atom more electronegative than the other, 610
drawing, 20–24
of enolates, 928
of enols, 928
IR spectrum and, 824
isomers compared to, 20
of naphthalene, 655
peptide bond, 1166
of phenoxide, 742–743, 743f, 744f
stability of, 607, 611
of substituted benzenes, 691–692
three atom allyl system, 609, 610
of an α,β-unsaturated carbonyl compound, 980
of a Wittig reagent, 835
Resveratrol, 600
Retention of configuration
in chlorination, 583
in conversion of alcohols to alkyl tosylates, 359, 361
in oxidation of alkylboranes, 413
in S_N2 substitution reactions, 265
Retention time, in gas chromatography, 505
Retinal, 841–842, 842f
11-*cis*-Retinal, 390
Retinol, 110
Retro Diels–Alder reaction, 628
Retrosynthetic analysis, 417, 444–447
in acetoacetic ester synthesis, 950–951
in benzene derivative synthesis, 1032–1033
of Diels–Alder product, 627–628
of disparlure, 470–471
of Grignard products, 790–792
of Heck reaction, 1057–1058
in malonic acid synthesis, 948–949
organometallic reagent reactions, 803–805
in phenylethanamine preparation, 1012
in substituted benzene syntheses, 714–717
synthesis of *m*-bromoaniline from benzene, 714
using an aldol reaction, 966–967
Wittig reaction, 837–839
Reversible reaction, unequal equilibrium arrows and, 69
RF radiation, 528–529, 560
R group, 92, 134
identity in S_N1 substitution reactions, 272–273, 273t
identity in S_N2 substitution reactions, 267, 267t
Rhodium, 1163
Rhodopsin, 841–842, 842f
Ribbon diagrams, 1182
α-D-Ribofuranose, 1123, 1141
β-D-Ribofuranose, 1123
Ribonucleosides, 1141–1142
Ribonucleotides, 1142
D-Ribose, 1114, 1115f, 1123, 1141
D-Ribulose, 1116f
Rimantadine, 1013
Ring-closing metathesis (RCM), 1065–1067, 1065f
Ring expansion reactions, 974
Ring-flipping, 155–156, 155f, 160
Ring formation
in aldol reaction, 975, 975f
in Robinson annulation, 982–986
Ring-opening metathesis polymerization (ROMP), 1071
Ring strain, 824
Risperidone, 722
Rizatriptan, 674
RNA, 1141, 1142–1143, 1143f
Robinson, Sir Robert, 982
Robinson annulation, 982–986
Rocaltrol. *See* Calcitriol
Rofecoxib, 992, 1244–1245
Rosiglitazone, 722
Rosmarinic acid, 590
Rosuvastatin, 988
Rotation
around carbon–carbon bonds, 144–145
barrier to, 150
conrotatory, 1083–1087, 1087t
disrotatory, 1083–1087, 1087t
restricted, 42, 300–301, 301f, 385t
Row, in periodic table, 9
Rowland, F. Sherwood, 584
RU 486, 431
Rubber, 1199f, 1210–1211, 1210f
Rubbing alcohol, 339f
Ruthenium, 1062

S

S (prefix), 185–190, 194–195
Saccharin, 1137, 1137f
S-Adenosylmethionine (SAM), 366
Safflower oil, 395
Safrole, 851, 1046
Salicin, 80f, 737, 1126
Salicylamide, 919
Salicylates, 80
Salicylic acid, 737, 917, 1229
Salinosporamide A, 122
Salmeterol, 371, 371f, 775f, 815
Salsolinol, 863
Salts
Brønsted–Lowry bases and, 62
of carboxylic acids, 737
formation of, 12
negatively charged nucleophile, 254
SAM (*S*-adenosylmethionine), 212, 284–285, 366
Samuelsson, Bengt, 738
Sand bee, 130
Sandmeyer reaction, 1030
Saponification, 895, 898
Saponins, 111
Saquinavir, 211
Saran, 1207
Saturated fats, 395
Saturated fatty acid, 393
Saturated hydrocarbons, 129
Saytzeff rule, 308
SBR (styrene–butadiene rubber), polymerization of, 1207
Sch38516, 1065f
s-Character, of hybrid orbitals, 76
Schiff base, 840
Schrock, Richard, 1062
s-Cis conformation, 613, 623
Scombroid fish poisoning, 657
S configuration, 1153
Scopolamine, 996

Scurvy, 91
Seashells, chiral helical, 180
Seaweed, 251–252
Sebacic acid, 1229
Sebacoyl chloride, 1229
sec- (prefix), 134
(*R*)-*sec*-Butylamine, 212
Secondary (2°) alcohols, 96, 415
  classification as, 332
  conversion to alkyl halides with HX, 352–355, 357t
  conversion to alkyl halides with phosphorus tribromide, 357, 357t
  conversion to alkyl halides with thionyl chloride, 356, 357t
  dehydration by an E1 mechanism, 346–347
  hydrogen bonding extent, 337
  oxidation to ketones, 477, 478
  from reactions of aldehydes with organometallic reagents, 785–787
  from reduction of ketones, 461, 770, 771, 772
Secondary (2°) alkyl halides, 96
  acetylide anion reactions with, 441–442
  classification as, 248
  E1 elimination reactions, 311, 320t, 321
  E2 elimination reactions, 306, 320t, 321
  S_N1 substitution reactions, 273, 279, 279f, 320t, 321
  S_N2 substitution reactions, 267, 279, 279f, 320t, 321
  synthesis of, 352
Secondary (2°) amides, 97, 870, 874–875, 877–878, 878t, 879, 886, 888, 894
Secondary (2°) amines, 97
  addition to aldehydes and ketones, 842–844, 843f
  from direct nucleophilic substitution, 1007–1008
  IR spectra of, 1002, 1003f
  nomenclature, 999–1000
  proline, 1155
  reactions with acid chlorides and anhydrides, 1023
  reactions with aldehydes and ketones, 1023
  reactions with nitrous acid, 1029
  from reduction of amides, 1010
  structure of, 997
Secondary carbon, 94
Secondary carbon radical, 571–572, 572f
Secondary hydrogen (2°), 94, 134
Secondary structure, of proteins, 1180–1183, 1181f, 1182f, 1183f, 1185f
Second-order kinetics, 235
  of E2 elimination reactions, 303
  in nucleophilic substitution reactions, 263–264
Second-order rate equation, 235, 262–263
Selective serotonin reuptake inhibitors (SSRIs), 1006
Separation, of amino acids, 1159–1163
Separatory funnel, 749, 749f, 1015, 1015f
Septet, NMR spectrum, 544t, 547
Serevent. *See* Salmeterol
Serine
  abbreviation for, 753t
  isoelectric point for, 1156t
  pK_a values for, 1156t
  structure of, 1154f
Serotonin, 1006
Sertraline, 649f, 690
β-Sesquiphellandrene, 623
Sesquiterpene, 1246, 1246t, 1248
Sevoflurane, 114, 340
Sex hormones, 1253–1254, 1254t. *See also specific hormones*
Sex pheromone, 972, 972f, 1055
Sextet, NMR spectrum, 544t, 548f
Shark liver oil, 165
Sharpless, K. Barry, 482
Sharpless epoxidation, 482–484, 1163
Sharpless reagent, 482–483
Shell, of electrons, 9–10
Shielding effects, in NMR, 536–537, 536t, 537f, 540f, 540t
1,2-Shifts, 348–350
Shikimic acid, 99
Sickle cell anemia, 1188, 1189
Sickle cell hemoglobin, 1188–1189
Side chain, of an amino acid, 1153, 1155, 1156t
Sigma (σ) bond, 36, 38, 41–43, 42f, 43f, 44t
  in alkanes, 93
  in alkenes, 300, 384
  in alkynes, 427–428
  breakage in addition reactions, 216
  carbon–oxygen, 332
  in cycloaddition reactions, 1077, 1087
  in electrocyclic reactions, 1077, 1081, 1083
  formation in addition reactions, 396, 397f, 398
  formation in elimination reactions, 215
  functional groups that contain C–Z σ bonds, 95, 96t
  hyperconjugation, 274–275
  inductive effect and, 73, 273
  length in conjugated dienes, 614–615
  radical reactions with, 572–573
  reduction of polar C–X bonds, 466–467
  rotation around, 144–145
  in sigmatropic rearrangements, 1078, 1091–1095
  in substitution reactions, 215, 261–262
Sigmatropic rearrangement, 1091–1097
  Claisen rearrangement, 1095, 1096–1097
  Cope rearrangement, 1095–1096
  description of, 1078
  orbital symmetry and, 1093–1094
  Woodward–Hoffmann rules for, 1093t
Sildenafil, 649f
Silent Spring, 252
Silk, 902, 902f
  structure of, 1182, 1183f
Silver(I) oxide, 1127
  in aqueous ammonium hydroxide, 782
  as oxidizing agent, 468
Silyl ether, 793
Simmons, H. E., 1061
Simmons–Smith reaction, 1061–1062
Simmons–Smith reagent, 1062
Simvastatin, 185, 614, 614f, 1253, 1253f
Sinemat, 50
α-Sinensal, 766
Singlet, NMR spectrum, 542, 544t, 546, 551
Sirenin, 1072
Sitaglipitin, 658, 991
Six-membered ring, synthesis of
  in aldol reaction, 975, 975f
  in Dieckmann reaction, 979–980
  in Robinson annulation, 982–986
Skeletal structures, 32–36, 33f
  with charged carbon atoms, 35
  interpreting, 33, 33f
Smith, R. D., 1061
S_N1 substitution reactions, 269–278
  alcohol conversion to alkyl halide, 353–355
  alkyl halide identity in, 278–279, 279f
  carbocation stability, 273–275
    Hammond postulate and, 275–278
    hyperconjugation and, 274–275
    inductive effects and, 273
  E1 elimination reaction compared, 313, 319–323, 320t
  E2 elimination reaction compared, 319–323, 320t
  energy diagram for, 270, 270f, 277, 278f
  ethers with strong acids, 362–363
  features of, 269, 273t
  kinetics of, 269, 273t
  leaving group in, 274, 280–281
  nucleophile strength in, 279–280
  racemization in, 271–272, 271f
  rate-determining step, 273, 275, 277–278
  R group in, 272–273, 273t
  S_N2 reactions compared, 278–283, 279f, 282t, 319–323, 320t
  solvent effects on, 281
  stereochemistry, 270–272, 271f, 273t
  transition state in, 270, 270f, 277, 278f, 280
  two-step mechanism for, 269–270, 273t
  as unimolecular, 263, 269
S_N2 substitution reactions, 263–268
  in acetoacetic ester synthesis, 950
  acetylide anion reactions with alkyl halides, 441–443
  acetylide anion reaction with epoxides, 444
  alcohol conversion to alkyl halides with HX, 353–354
  with phosphorus tribromide, 357
  with thionyl chloride, 356
  alcohol synthesis from alkyl halides, 341–342
  alkyl halide identity in, 278–279, 279f
  alkyl halide with ammonia or amine, 1007–1008
  alkyl tosylate reactions, 360–361
  in automated peptide synthesis, 1178
  backside attack in, 265–266, 280
  direct enolate alkylation, 942
  E1 elimination reactions compared, 319–323, 320t
  E2 elimination reactions compared, 319–323, 320t
  energy diagrams for, 264, 264f
  epoxide ring opening reactions, 367–368
  epoxide synthesis from halohydrins, 343
  epoxy resin synthesis, 1214
  ethers with strong acids, 362–363
  ether synthesis from alkyl halides, 341–342
  features of, 263, 267t
  frontside attack in, 265
  of α-halo acids with ammonia, 1156–1157, 1159f
  of α-halo carbonyl compounds, 941, 941f
  kinetics of, 263, 267t
  leaving group in, 265–266, 280–281
  in malonic ester synthesis, 947
  nitrile synthesis, 906
  nucleophile strength in, 279–280
  one-step mechanism for, 264, 267t
  organic synthesis, 286, 287t, 288
  reduction of alkyl halides, 467, 467f
  reduction of epoxides, 467, 467f
  R group in, 267, 267t
  S_N1 reactions compared, 278–282, 279f, 282t, 319–323, 320t
  solvent effects on, 281
  stereochemistry of, 265–266, 266f, 267t, 280
  steric hindrance and, 267, 267f
  transition state, 264, 264f, 266, 266f, 280
Soap, 111–113, 112f, 895, 898
Sodium, in dissolving metal reductions, 457, 465
Sodium acetate, 106, 733f
Sodium amide, for alkyne synthesis by dehydrohalogenation, 319, 319f
Sodium benzoate, 737, 750
Sodium bisulfite, 473
Sodium borohydride, 457, 771–773, 773f, 781t, 831, 1129
Sodium chloride, 11
  boiling point, 103
  melting point, 105
Sodium cyanoborohydride, for reductive amination of aldehydes and ketones, 1011, 1012
Sodium dichromate, 468, 477
Sodium ethoxide, 342
  in dehydrohalogenation reactions, 298t
Sodium hydroxide
  in dehydrohalogenation reactions, 298t, 299
  use in extraction procedure, 749–750, 751f
Sodium hypochlorite, 480
Sodium iodide, 1030
Sodium methoxide, 342
  in dehydrohalogenation reactions, 298t
Sodium salicylate, 737
Sodium trichloroacetate, 1071
Soft coral (*Capnella imbricata*), 443
Solanine, 1106, 1126
Solid phase technique, of amino acid synthesis, 1177–1179
Solubility, 106–109, 108t
  of alcohols, 338t
  of aldehydes and ketones, 823t
  of alkanes, 143, 143t
  of alkenes, 391
  of alkyl halides, 250t

of alkynes, 429
of amines, 1001t
of carboxylic acids, 734t
definition of, 106
of epoxides, 338t
of ethers, 338t
intermolecular forces and, 106–109, 107f
"like dissolves like," 106–107
of lipids, 164, 393, 1232
of monosaccharides, 1116, 1128
soap and, 111–113, 112f
of vitamins, 109–111
Solute, 106
Solvent effects, on nucleophilic substitution reactions, 258–261, 259f, 260f, 281, 282t
Solvents, 106
alkyl halides, 251
in E1 elimination reactions, 312t
in E2 elimination reactions, 305, 307t
for extraction procedure, 749–751, 749f, 751f
liquid, 214
nonpolar, 106
nucleophilicity and, 258–261
organic, 106–107, 108t
polar, 106
polar aprotic, 260, 260f, 281, 282t, 305, 307t, 934
polar protic, 259, 259f, 281, 282t, 312t, 935
s orbitals
description, 9
hybridization of, 36–45, 38f, 42–43
1*s* orbital, 10, 36
2*s* orbital, 10, 36–38, 38f
Sorbitol, 195, 1129
D-Sorbose, 1116f
Sorona, 1219, 1219f
Spandex, 1213
Spearmint, odor of, 204
Specific rotation, 199–200
Spectator ions, 62
Spectrometer, NMR, 529–530, 529f
Spectroscopy, 497
Spelunkers, 42
Spermaceti wax, 1233
Spermine, 97
SPF rating (sun protection factor), 632
Sphingomyelins, 1240, 1240f
Sphingosine, 1240
sp hybrid orbitals, 38–39, 39t, 42–43, 43f, 44t
in alkynes, 427, 428, 432, 433
percent *s*-character, 46, 76
sp^2 hybrid orbitals, 38–39, 39t, 41–42, 42f, 44t, 332, 391
in alkenes, 299–300, 302, 384
in alkyl halides, 248, 248f
in aromatic heterocycles, 656–657, 657f
in benzene, 644, 654
in benzyme, 709
carbocation, 270, 275
carbon radical, 571, 582
carboxy group, 730
percent *s*-character, 46, 76
in vinyl halides, 286
sp^3 hybrid orbitals, 37–38, 38f, 39f, 39t, 40, 44t, 332, 333, 391
in acyclic alkanes, 129
in alkenes, 302
in alkyl halides, 248, 248f, 253, 263, 286
in nucleophilic substitution reactions, 253–254, 262–263, 286
percent *s*-character, 46, 76
Spiriva. *See* Tiotropium bromide
Spiro ring system, A–6
Splenda. *See* Sucralose
Squalene, 490, 1246f, 1247f, 1248, 1252–1253, 1252f
Squalene epoxidase, 1252f, 1253
Squalene oxide, 1252f, 1253
Stachyose, 1150
Staggered conformation, 144–151, 146f, 147f, 148f, 150f, 191, 314, 314f
Stanozolol, 675, 1254
Starch
hydrolysis of, 175f, 176, 339, 339f, 1139
structure of, 175–177, 175f, 176f, 1139
Stearic acid, 1234, 1234t
melting point, 394t
structure, 393, 394f, 394t
Stearidonic acid, 394, 488
Step-growth polymers, 1200, 1211–1215
epoxy resins, 1214–1215, 1215f
polyamides, 1211–1212
polycarbonates, 1214
polyesters, 1212–1213
polyurethanes, 1213
Stepwise reaction, 217
Stereocenter. *See* Stereogenic centers
Stereochemistry, 144, 174–212
of addition reactions, 397, 402–404, 403t
of carbonyl reduction, 773–774
chemical properties of enantiomers, 202–204
chiral and achiral molecules, 178–180, 180f
diastereomers, 190–192, 192f, 201–202, 201f
of the Diels–Alder reaction, 624–625
of dienophiles, 624–625
of dihydroxylation, 472–474
of disubstituted cycloalkanes, 195–196
of E2 elimination reactions, 314–317, 314f
of electrocyclic reactions, 1082–1083
of electrophilic addition reactions, 402–404, 403t
of enolate alkylation, 943
of epoxidation, 469–470
of halogenation, 406–407, 408f, 582–583, 582t
of halohydrin formation, 409–410, 410t
of hydroboration–oxidation reactions, 413–414, 414t
meso compounds, 193–194, 194f
organometallic reagent reactions with carbonyl compounds, 788
physical properties of stereoisomers, 197–202, 199t, 201f
polymer, 1208–1209
of sigmatropic rearrangements, 1093
of S_N1 substitution reactions, 270–272, 271f, 273t
of S_N2 substitution reactions, 265–266, 266f, 267t, 280
of starch and cellulose, 175–177, 175f, 176f
stereogenic centers, 181–195, 183f, 187f
Stereogenic centers, 181–195
in amines, 998–999
in amino acids, 752, 1153, 1155
on carbon atoms not part of a ring, 181–182
carbonyl reduction, 773–774
in cyclic compounds, 183–185
cyclic hemiacetals, 853
definition of, 179
in diastereomers, 190–192
drawing enantiomers, 182–183, 183f
enantiotopic and diastereotopic protons, 534–535
formation from hydrohalogenation, 402, 403
halogenation reactions and, 582–583, 582t
inversion of configuration around, 265–267, 266f, 267t, 280
labeling with *R* or *S*, 185–190, 187f, 189f, 194–195
in meso compounds, 193–194
in monosaccharides, 1109–1118, 1120–1123
number in a molecule, 179
organometallic reagent addition, 788
priority assignment, 185–189, 187f
retention of configuration around, 265
tetrahedral, 179, 181, 182, 183, 184, 190, 193, 194
Stereoisomers, 19. *See also* Enantiomers
of alkenes, 300–302, 385t
chemical properties of, 202–204
cis, 158–161
configuration of, 177
of conjugated dienes, 613
constitutional isomers compared, 177f
of cycloalkanes, 158–161
definition of, 158
from dehydrohalogenation, 309
diastereomers, 190–192, 192f, 201–202, 201f
disubstituted cycloalkanes, 195–196
enantiomers, 179
of monosaccharides, 1111–1114, 1114f
nomenclature for alkenes, 388–390
number of, maximum, 190–191
physical properties of, 197–202, 199t, 201f
starch and cellulose, 175–177
trans, 158–161
Stereoselective reactions
alkyne reduction to a cis alkene, 464
dissolving metal reduction of an alkyne, 466
of E2 elimination reactions, 309
Stereospecific reactions
carbene addition, 1060
coupling reactions with organocuprates, 1051
definition of, 407
epoxidation, 469
halogenation of alkenes, 407
Heck reaction, 1057
Simmons–Smith reaction, 1062
Suzuki reaction, 1054
Steric effects, nucleophilicity and, 258
Steric hindrance, 258
in alcohols, 337
in hydroboration reactions, 412
S_N2 substitution reactions and, 267, 267f
Steric strain, 149, 150t
Steroids, 1250–1255
adrenal cortical, 1253, 1255
anabolic, 1254
cholesterol. *See* Cholesterol
sex hormones, 1253–1254, 1254t
structure of, 1250–1251, 1251f
synthesis in Diels–Alder reaction, 629
Stone, Reverend Edmund, 80f
Strain
angle, 151–152, 333, 344
Baeyer strain theory, 151
energy increase with, 147
steric, 149, 150t
torsional, 147, 150t, 152
s-Trans conformation, 613, 623
Strecker synthesis, 1158–1159, 1159f
Structural isomers. *See* Constitutional isomers
Strychnine, 1162
Styrene, 570, 711, 1063, 1063f, 1105
polymerization of, 1200f, 1201, 1203, 1207
structure of, 1200f, 1202f
Styrene–butadiene rubber (SBR), polymerization of, 1207
Substituents, naming, 134, 645–646
naming aromatic rings as, 646
Substitution reactions, A–12
alcohols, 343–344, 352–361
allylic, 587
of aromatic compounds
of benzene, 678–690
electrophilic, 678–706
nucleophilic, 706–709
of substituted benzenes, 691–706
of aryl diazonium salts, 1029–1033
of benzene, 642, 651
at the α-carbon of carbonyl compounds, 924–961
description of, 215
electrophilic aromatic. *See* Electrophilic aromatic substitution reactions
Heck reaction, 1050, 1056–1058
nucleophilic, 253–288
radical, 574
Substrate, 238, 238f
Succinaldehyde, 992
Succinic acid, 733
Succinimide, 587
Sucralose, 1137, 1137f
Sucrose, 124, 1136–1137, 1136f
olestra synthesis from, 897–898
stereogenic centers, 184
structure of, 1136–1137, 1136f
Sudafed, 72. *See also* Pseudoephedrine
Suffix, 133
-adiene, 387
-al, 819
-aldehyde, 820
-amide, 874–875, 876t
-amine, 999
-ane, 131, 133, 134, 135, 139, 141, 387
-ate, 733, 876, 876t
-carbaldehyde, 819
-carbonyl chloride, 873, 876t
-carboxylic acid, 873
-dioic acid, 733

Suffix (*Continued*)
-diol, 335
-ene, 387
-enol, A–5
-enone, A–5
-enyne, A–5
-ic acid, 732, 873, 874–875
-ide, –249
-ine, 249
-o, 249
-oic acid, 731, 874–875
-ol, 334, 364
-one, 820
-onitrile, 875, 876t
-ose, 1108
-oxy, 336
-oyl, 822
-thiol, 364
-yl, 134, 336, 822
-yl chloride, 873, 876, 876t
-ylic acid, 874
-yne, 428
Sugar, simple, 118. *See also* Monosaccharides
Sugars. *See also* Carbohydrates
amino, 1140
nonreducing, 1130, 1130f, 1137
reducing, 1130, 1130f, 1135, 1136
simple. *See* Monosaccharides
Sulfa drugs, 1037–1038, 1038f
Sulfamethoxazole, 1038f
Sulfanilamide, 1037–1038
Sulfide (functional group), 96t
Sulfides, 365–366
Sulfide suffix, 365
Sulfisoxazole, 1038f
Sulfonate anion, 752
Sulfonation
of benzene, 679f, 682–683
electrophilic aromatic substitution reaction, 679f, 682–683
Sulfonic acids, 751–752
Sulfonium ion, 366
Sulfur
common nucleophiles, 261t
in fossil fuels, 79
octet rule exceptions in, 19
Sulfuric acid, 19, 78–79
dehydration of alcohols in, 345–350, 348f
in nitration and sulfonation reactions of benzene, 682–683
Sulfur nucleophile, 366, 372
Sulfur oxides, 79
Sulfur trioxide, 683
Sulfur ylides, 1072
Sunscreen, 34, 632, 725
Supercoil, 1186, 1186f
Superhelix, 1186–1187
Suprafacial bond formation, 1089–1091, 1091t
Suprafacial rearrangement, 1093–1095, 1093t
Surface area
boiling point and, 103, 104f
van der Waals forces and, 100, 100f
Suzuki, Akira, 1056
Suzuki reaction, 1050, 1052–1056, 1055f
Sweeteners, artificial, 1137, 1137f, 1168
Sweetness, of monosaccharides, 1116
Sweet'n Low. *See* Saccharin
Symbol, element, 8
Symmetrical anhydrides, 870, 873
Symmetry
melting points and, 106
plane of, 180, 193, 195
Syn additions, 397, 403, 403t, 406, 411, 413, 458, 460, 463, 469, 471, 471f, 472, 473–474, 1060
Syndiotactic polymer, 1208–1209
Syn periplanar geometry, 314, 314f
Synthetic intermediate, 418
Synthetic polymers, 593–594, 1198–1230
amorphous regions of, 1216–1217
atatic, 1208–1209
"Big Six," 1220–1221, 1221t
biodegradable, 1222–1223
chain-growth, 1199, 1200–1207
in consumer products, 1199, 1199f
crystalline regions of, 1216–1217
disposal of, 1220–1222
green synthesis of, 1217–1220, 1218f, 1219f
isotactic, 1208–1209
molecular weights of, 1200
nylon, 902–903
physical properties of, 1216–1217
plasticizer addition to, 1217
polyesters, 903
recycling of, 1198, 1220–1222, 1221t
shorthand representation of, 1200, 1200f
step-growth, 1200, 1211–1215
stereochemistry of, 1208–1209
structure of, 1199, 1216–1217
syndiotactic, 1208–1209
thermoplastics, 1216
thermosetting, 1216
Systematic name, 133. *See also* IUPAC nomenclature
of amines, 999–1000

T

2,4,5-T (2,4,5-trichlorophenoxyacetic acid herbicide), 682f
Tagamet. *See* Cimetidine
D-Tagatose, 1116f
Talose
D-talose, 1115f, 1132
Wohl degradation, 1132
Tamiflu, 99 *See also* Oseltamivir
Tamoxifen, 420, 924, 944
Target compound, 445
Tartaric acid, stereoisomers of, 201, 201f
Tautomerization
in acid, 438, 927
of amides, 907–908
in base, 927
of carbonyl compounds, 907–908, 926–928, 938
mechanism of, 927–928
in sigmatropic rearrangements, 1095–1096
Tautomers
definition of, 437
enol, 437–439, 908, 926–927
keto, 437–439, 908, 926–927
Taxol, 174, 184, 769 *See also* Paclitaxel
TBDMS (*tert*-butyldimethylsilyl ether), 793, 794
Teflon, 251. *See also* Polytetrafluoroethylene (Teflon)
Telfairine, 248
Temperature, reaction rate and, 234–235
Terephthalic acid, 903, 1198, 1200f, 1212, 1228
Terpenes, 1245–1250
biosynthesis of, 1247–1250, 1247f
classes of, 1246t
examples of, 1246f
locating isoprene units in, 1245–1247, 1246f
α-Terpinene, 1250
α-Terpineol, 425, 1249
tert- (prefix), 134
Tertiary (3°) alcohols, 96
classification as, 332
conversion to alkyl halides with HX, 352–355, 357t
dehydration by E1 mechanism, 346–347
hydrogen bonding extent, 337
from reaction of esters and acid chlorides with organometallic reagents, 794–796
from reaction of ketones with organometallic reagents, 785–787
Tertiary (3°) alkyl halides, 96
acetylide anion reaction with, 441–442
classification as, 248
E1 elimination reactions, 311, 320, 320t
E2 elimination reactions, 306, 320, 320t
S_N1 substitution reactions, 271, 273, 279, 279f, 320, 320t
synthesis of, 352, 355
Tertiary (3°) amides, 97, 870, 874–875, 886, 888, 894
enolate formation from, 930
Tertiary (3°) amines, 97
from direct nucleophilic substitution, 1007–1008
IR spectra of, 1002, 1003f
nomenclature, 999–1000
from reduction of amides, 1010
structure of, 997
Tertiary carbon, 94
Tertiary carbon radical, 571–572
Tertiary hydrogen (3°), 94, 134
Tertiary structure, of proteins, 1183–1185, 1185f
Testosterone, 1254t
Tetra- (prefix), 136
Tetrabutylammonium fluoride, 793
Tetrachloromethane, 251 *See also* Carbon tetrachloride
Tetradecane, 131t
Tetrafluoroethylene, monomers in polymerization, 594t
Tetrahalide, 436
Tetrahedral arrangement, 26, 27–28, 29t
Tetrahedral geometry, of acyclic alkanes, 129–130
Tetrahedral oxygen, 333
Tetrahedral stereogenic center, 179, 181, 183, 184, 190, 193, 194
Tetrahedrane, 152
Tetrahydrocannabinol (THC), 125
mass spectrum of, 505, 506, 506f
structure of, 78
Tetrahydrofuran (THF), 126, 336, 674, 932–935, 942–943, 943
complexed with borane, 411
as solvent, 260f, 362
Tetrahydrogestrinone, 1254
α-Tetralone, 689
Tetramethyl α-D-glucopyranose, 1130f
Tetramethylsilane (TMS), 530, 531, 556
Tetrapeptide, 1167
Tetraterpene, 1246t
Tetravalent carbon, 4, 28, 30, 32–34
Tetrodotoxin, synthesis of, 622, 622f
Thalidomide, 184
THC. *See* Tetrahydrocannabinol (THC)
Thermodynamic enolates, 934, 935, 943, 944
Thermodynamic product, in electrophilic addition reactions of conjugated dienes, 619–621, 620f
Thermodynamics, 225–227, 226f, 226t
of hydrate formation, 845
Thermoplastics, 1216
Thermosetting polymers, 1216
THF. *See* Tetrahydrofuran (THF)
Thiazolinone, 1170, 1197
Thioester, 904–905
Thiol (functional group), 96t
Thiols, 364–365, 364f
-thiol suffix, 364
Thionyl chloride
conversion of alcohols to alkyl halides with, 356
conversion of carboxylic acids to acid chlorides, 889f, 890
Three-dimensional representation, of acyclic alkanes, 129–130
Threonine, 762
isoelectric point for, 1156t
pK_a values for, 1156t
structure of, 1154f, 1155
Threose
D-threose, 1114, 1114f, 1115f
L-threose, 1114, 1114f
Thrombin, 1153
Thromboxanes, 1242, 1243, 1243f, 1243t
Thymine, 1142, 1143f
Thymol, 562, 673
Ticlopidine, 863
Tiotropium bromide, 341
Titanium(IV) isopropoxide, 482–484
Toads (*Bufo*), 1006
Tobacco, 648f
Tobramycin, 1140
Tollens reagent, 782, 1129–1130
Toluene, 123, 506
in BTX mixture, 648
electrophilic aromatic substitution reaction, 694, 698
oxidation to benzoic acid, 712
structure of, 645
synthesis by Friedel–Crafts alkylation, 716
Toluene 2,6-diisocyanate, 1213
p-Toluenensulfonate. *See* Tosylate
p-Toluenesulfonic acid (TsOH), 79, 751, 847
dehydration of alcohols to alkenes in, 345, 351, 352f
p-Toluenesulfonyl chloride (TsCl), 359, 361, 362f
p-Toluidine
basicity of, 1018
electrostatic potential plot of, 1019f
Torsional energy, 147
Torsional strain, 147, 150t, 152
Tosic acid, 751
Tosylate, 359–361, 362f
Tosyl group, 359, 752

TPA (thermal polyaspartate), 1223
Trade name, 133
Trans (prefix), 389–390
Trans diaxial geometry, 315–316, 315f
Trans dihalide, 436
Trans fats, 462–463
Trans geometry
  alkenes, 385, 385t, 388–389, 403, 406
  cycloalkenes, 385
Trans isomers
  of but-2-ene, 301–302, 301f
  of cycloalkanes, 158–161
  definition of, 158, 301
  of disubstituted cycloalkanes, 195–196
  stability, 303
Transition state
  definition of, 229
  drawing structure of, 230
  of E1 elimination reactions, 310
  of E2 elimination reactions, 304–306, 304f, 308, 314
  in electrophilic addition, 399, 399f
  endothermic reaction, 276, 276f
  in energy diagrams, 229–233, 233f
  exothermic reaction, 276–277, 277f
  four-centered, 412
  in halogenation reactions, 577, 577f, 579, 580f, 581, 581f
  Hammond postulate and, 275–278, 276f–278f, 580
  S_N1 substitution reactions, 270, 270f, 277, 278f, 280
  S_N2 substitution reactions, 264, 264f, 266, 266f, 280
  in two-step reactions, 231–233, 233f
Trans protons, 549, 549f
Tri- (prefix), 136, 999
Triacylglycerols, 1234–1237
  combustion of, 1237
  common fatty acids in, 1234t
  degrees of unsaturation in, 461–462
  energy storage in, 1237
  fats and oils, 395
  formation of, 118
  hydrogenation of, 461–463, 462f, 1236
  hydrolysis of, 393, 897, 898, 1236
  melting point, 393, 394t, 1234–1235
  mixed, 1234
  olestra as a substitute for, 897–898
  oxidation of, 588, 589f, 1236
  saponification of, 898
  saturated, 896, 897f, 1234–1235, 1234t, 1235f
  simple, 1234
  structure of, 393, 896, 897f, 1232f, 1234–1235, 1234t, 1235f, 1240f
  unsaturated, 896, 1234–1236, 1234t, 1235f
Trialkylborane, 412, 412f
Triarylphosphine, 1056–1057
1,3,5-Tribromobenzene, synthesis of, 1031, 1031f
Tributyltin hydride, 602
Trichlorofluoromethane, 3, 584
Trichloromethane, 251 *See also* Chloroform
2,4,5-Trichlorophenoxyacetic acid herbicide (2,4,5-T), 682f
Tricyclohexylphosphine, 1053
Tridecane, 131t
Trienes, 387
Triethylamine, 80, 295, 999, 1056–1057
1,2,4-Triethylcyclopentane, 140f
Trifluoroacetic acid, for removal of Boc protecting group, 1174
2,2,2-Trifluoroethanol, 72–73, 73f
Trifluoromethanesulfonic acid, 752
Trigonal planar carbocation, 270–271, 273t
Trigonal planar double bond, 299
Trigonal planar molecule, 26, 27, 29t
  carbon radical, 571, 583
  carbonyl group, 765–766, 788
  carboxy group, 730
Trigonal pyramidal molecule, 28
  amines, 997
Trihalomethanes, 1059
Trimethylamine, 1004
  electrostatic potential plot of, 998, 998f
Trimethyl borate, 1054
2,5,5-Trimethylcyclohexanecarboxylic acid, 731
1,2,2-Trimethylcyclohexanol, 381
2,5,5-Trimethylcyclohexanol, 334f
1,3,3-Trimethylcyclohexene, 402
2,3,5-Trimethylhex-2-ene, 387
2,2,4-Trimethylpentane, combustion of, 162 *See also* Isooctane
2,3,5-Trimethyl-4-propylheptane, 137f
Triols, 335
Tripeptide, 1165–1166, 1177
Triphenylene, 676
Triphenylmethyl radical, 602
Triphenylphosphine, 836, 1053, 1055
Triphenylphosphine oxide, 835, 837
Triphosphate, 283–284
Triple bonds
  in alkynes, 93, 94t, 427–428
  of benzyne, 708–709
  bond length and bond strength, 45, 46t
  components of carbon–carbon, 42–43
  as functional group, 93, 94t
  Lewis structure, 16
  NMR spectra of, 540, 540t
  reduction of, 463–466, 466f
Triple helix, of collagen, 1187, 1187f
Triplet, NMR spectrum, 542, 543–544, 544t, 545t, 548f, 551
Tristearin, 1237
Triterpene, 1246t, 1248, 1252
Tri(*o*-tolyl)phosphine, 1053, 1058
Triynes, 428
Tropylium cation, 660, 664
Truvia. *See* Rebaudioside A
Trypsin, 1171, 1171t
Tryptophan, 763
  isoelectric point for, 1156t
  pK_a values for, 1156t
  structure of, 1154f
TsCl (*p*-toluenesulfonyl chloride), 359, 361, 362f
TsOH. *See p*-Toluenesulfonic acid (TsOH)
Tuberculosis, 268
Tungsten, 1062
ar-Turmerone, 972
Twistoflex, 648, 649f
Two-electron bond, 12, 13, 15
Two-step reaction mechanism, 231–233, 233f, 235
Tylenol. *See* Acetaminophen
Tyrian purple, 1036
Tyrosine
  isoelectric point for, 1156t
  pK_a values for, 1156t
  structure of, 1154f
Tyvek, 1221

U

Ultraviolet light
  absorption by conjugated dienes, 630–632, 631f
  skin damage from, 632
Undecane, 129, 131t
Unimolecular elimination. *See* E1 elimination reactions
Unimolecular reactions
  elimination reactions, 303, 310
  in nucleophilic substitution reactions, 263, 269
  organic reactions, 235
Unknown
  analysis using mass spectrometry, 498–500
  using ^{1}H NMR to identify an, 553–556
Unoprostone isopropyl, 1244
α,β-Unsaturated carbonyl compounds, 799–801
  1,2-addition to, 799–801
  1,4-addition to, 800–801
  in the Michael reaction, 980–982, 982f
  resonance structures, 980
  in Robinson annulation, 982–986
  synthesis of
    in dehydration of aldol product, 965–967, 968, 971f
    from α-halo carbonyl compounds, 941
Unsaturated fatty acid. *See* Fatty acids, unsaturated
Unsaturated lipids, oxidation of, 588–589, 589f
Unsaturation, degrees of
  in alkenes, 460–461
  benzene, 642, 643
  calculation of, 385–387, 460–461
  hydrogenation data and, 460–461
  in triacylglycerols, 461–462
Unsaturation, in alkynes, 427
Uracil, 1142, 1143f
Urethanes, 1213, 1228
Uridine monophosphate, 293
Urushiols, 749

V

Valdecoxib, 1244–1245
Valence bond theory, 661
Valence electrons, 10, 11–13, 18t, 36
Valence shell electron pair repulsion (VSEPR) theory, 26
Valeric acid, structure of, 732t
Valine
  isoelectric point for, 1156t
  pK_a values for, 1156t
  structure of, 1154f
Valinomycin, 115
Valium. *See* Diazepam
Valproic acid, 88
Vancomycin, 126
van der Waals forces, 100–101, 100f, 102t
  in alkanes, 143
  surface area and, 100, 100f
Vane, John, 738
Vanillin, 34, 562, 826f
Vanilloids, 4
Vasopressin, 1167–1168, 1168f, 1184
Vegetable oil, 896, 898
  partial hydrogenation of, 461–462, 462f
Venlafaxine, 792, 816
Ventolin. *See* Albuterol
β-Vetivone, 958, 961, 989
Viagra. *See* Sildenafil
Vibration of bonds, infrared spectroscopy and, 508, 510–513, 510f
Vicia faba, 7
Vicinal dibromide, 431, 432
Vicinal dihalides, 318, 405
  alkyne synthesis from, 431–432
Vinegar, 736
Vinyl acetate
  NMR spectrum of, 550, 550f
  polymerization of, 1202, 1205f
  structure of, 1202f
Vinylboranes, 1054, 1056
Vinyl bromide, 1054, 1056
Vinyl carbanion, 466
Vinyl carbocation, 434, 435, 438
Vinyl chloride, 57
  monomers in polymerization, 594t
  polymerization of, 1199, 1207
  structure of, 1199, 1202f
1-Vinylcyclopentene, 390f
Vinyl group, 390
Vinyl halides, 248, 248f, 286
  coupling reactions with
    alkenes, 1056–1058
    organoboranes, 1054
    organocuprates, 1050–1051
  reactivity of, 686
  synthesis in hydrohalogenation of alkynes, 434–435
Vinylidene chloride, polymerization of, 1207
Vinyl iodide, 1057
Vioxx. *See* Rofecoxib
Viracept. *See* Nelfinavir
Vitamin(s), 109–111
  definition of, 109, 1241
  fat soluble, 110, 164, 1241–1242, 1242f
  origin of term, 110
  water soluble, 110, 111
Vitamin A, 110, 352f
  β-carotene conversion to, 110
  deficiency, 110, 1241f
  electrostatic potential plot of, 1242, 1242f
  role in the body, 1241f
  solubility, 110
  sources, 1241, 1241f
  structure, 110, 300, 1241f, 1246
Vitamin B_3, 111, 777
Vitamin B_5, 111
Vitamin B_6, 51
Vitamin C (ascorbic acid), 90, 91, 600, 923, 955
  solubility of, 111
  sources of, 110
  structure of, 110, 880
Vitamin D_3, 614, 1086
  deficiency, 1241f
  role in the body, 1241f
  sources of, 1241
  structure of, 300, 1241f
Vitamin E, 124
  as antioxidant, 589–590
  deficiency, 1241f
  electrostatic potential plot of, 1242, 1242f
  as radical scavenger, 573
  role in the body, 1241f
  sources of, 1241
  structure of, 589, 1241f

Vitamin K (phylloquinone), 111, 677, 690, 690f
  deficiency, 1241f
  role in the body, 1241f
  sources of, 1241
  structure of, 1241f
VSEPR (valence shell electron pair repulsion) theory, 26
Vulcanization, 1210, 1210f

W

Wald, George, 842
Walden, Dr. Paul, 265
Walden inversion, 265
Warfarin, 763
Water
  as an achiral molecule, 178
  electrophilic addition of, 404–405, 415
  electrostatic potential plot for, 50f
  hybrid orbitals in, 39, 39f
  hydration of alkynes, 433f, 437–439
  hydrogen bonding, 101
  as Lewis base, 82
  Markovnikov addition of, 437
  molecular shape of, 29
  nucleophilicity of, 279, 280
  pK_a of, 67t, 70
  as a polar molecule, 49
  removing from reaction mixture using Dean–Stark trap, 848, 848f
  as solvent, 106–109, 111–112, 112f, 259, 259f, 281
    in extraction procedure, 749, 749f
Wavelength
  definition of, 507
  electromagnetic spectrum and, 506–508, 507f
  length units for, 507
Wavenumber, 508, 509
Waxes, 164, 164f, 1233
Wedge symbol, 28, 158, 1109–1112
Welding torch, 42
Wellbutrin. *See* Bupropion
Williamson ether synthesis, 341, 342
Willow tree, 80, 80f
Wittig, Georg, 835
Wittig reaction, 835–839
  mechanism of, 836–837
  retrosynthetic analysis using, 837–839
Wittig reagent, 835–839
Wohl degradation, 1131–1132, 1133
Wolff–Kishner reduction, 712
Wood alcohol, 339f. *See also* Methanol
Woodward, R. B., 923, 991, 1078
Woodward–Hoffmann rules
  for cycloaddition reactions, 1091, 1091t
  for electrocyclic reactions, 1086, 1087t
  for sigmatropic rearrangements, 1093t
Wool, structure of, 902, 902f

X

Xalatan. *See* Latanoprost
m-Xylene, 704
p-Xylene, 506, 598, 648
D-Xylose, 1114, 1115f
D-Xylulose, 1116f

Y

-yl (suffix), 134, 336, 822
-yl chloride (suffix), 873, 876, 876t
-ylic acid (suffix), 874
Ylide, 835–836
-yne (suffix), 428

Z

Z (prefix), 389–390
Zaitsev rule, 308–309, 312, 316, 345, 1027
Z configuration, double bond, 393, 1210–1211
Zidovudine, 4. *See also* AZT (azidodeoxythymidine)
Ziegler, Karl, 1209
Ziegler–Natta catalysts, 1209, 1210, 1211
Ziegler–Natta polymerization, 1208–1209
Zigzag skeletal structures, 151
Zileuton, 372
Zinc
  in Clemmensen reduction, 712
  conversion of an ozonide to carbonyl compounds, 474
Zinc chloride, 354
Zinc–copper couple, 1061–1062
Zingerone, 53, 973
Zingiberene, 392f, 490, 623, 1246f
Zocor. *See* Simvastatin
Zoloft. *See* Sertraline
Zolpidem, 674, 1042
Zuclomiphene, 422
Zwitterion, 90, 753, 1155
Zyban. *See* Bupropion
Zyflor. *See* Zileuton